

Origins of the method to determine the CKM angle γ using $B^\pm \rightarrow DK^\pm$, $D \rightarrow K_S^0 \pi^+ \pi^-$ decays

A. Ceccucci^c, T. Gershon^g, M. Kenzie^g, Z. Ligeti^l, Y. Sakai^s, K. Trabelsi^t

^c *EP Department, CERN, Switzerland*

^g *Department of Physics, University of Warwick, Coventry, United Kingdom*

^l *Lawrence Berkeley National Laboratory, University of California, Berkeley, CA 94720, USA*

^s *Institute of Particle and Nuclear Studies, High Energy Accelerator Research Organization (KEK),
Tsukuba, 305-0801, Japan*

^t *Université Paris-Saclay, CNRS/IN2P3, IJCLab, 91405 Orsay, France*

Abstract

The angle γ of the Cabibbo–Kobayashi–Maskawa unitarity triangle is a benchmark parameter of the Standard Model of particle physics. A method to determine γ from $B^+ \rightarrow DK^+$ with subsequent $D \rightarrow K_S^0 \pi^+ \pi^-$ or similar multibody decays has been proven to provide good sensitivity. We review the first discussions on the use of this technique, and its impact subsequently. We propose that this approach should be referred to as the BPGGSZ method.

The authors thank Alex Bondar, Anton Poluektov, Anjan Giri, Yuval Grossman, Abner Soffer and Jure Zupan for endorsing this note.

The angle $\gamma = \arg(-V_{ud}V_{ub}^*/V_{cd}V_{cb}^*)$ of the unitarity triangle formed from elements of the Cabibbo–Kobayashi–Maskawa quark mixing matrix [1, 2] is a benchmark parameter of the Standard Model of particle physics.¹ The value of γ is a measure of the extent to which the CP symmetry between particles and antiparticles is violated in the weak interactions of quarks. It can be determined with negligible theoretical uncertainty from measurements of decay rates and CP -violating asymmetries in processes where interference between $b \rightarrow c\bar{u}s$ and $b \rightarrow u\bar{c}s$ transitions can occur [3]. The archetypal example, which is also the most experimentally accessible, is that of $B^+ \rightarrow DK^+$ decays, where D indicates a neutral D meson that is an admixture of D^0 and \bar{D}^0 states, but the same concepts are valid also for related channels such as $B^+ \rightarrow D^*K^+$, $B^+ \rightarrow DK^{*+}$ and $B^0 \rightarrow DK^{*0}$. The observables (relative decay rates and CP -violating asymmetries) can be expressed in terms of γ , r_B and δ_B , where r_B is the relative magnitude of the $b \rightarrow u\bar{c}s$ and $b \rightarrow c\bar{u}s$ transition amplitudes and δ_B is their relative strong (*i.e.* CP -conserving) phase. Detailed reviews on the determination of γ from such processes can be found, for example, in Refs. [4–8].

Initial discussions of the use of this approach to obtain experimental sensitivity to γ focussed on the case where the neutral D meson decays to a CP eigenstate, such as (CP -even) K^+K^- , $\pi^+\pi^-$ or (CP -odd) $K_s^0\pi^0$ [9, 10]. In this approach, now widely known as the GLW method, the amplitudes for D^0 and \bar{D}^0 decays to the final states of interest are related trivially, under the assumption of negligible effects of CP violation in the D system. The B decay observables can then be expressed in terms of the unknown parameters (γ , r_B , δ_B) only. To determine γ from this method alone, however, requires sensitivity to observe the relatively small CP asymmetries, expected to be comparable in magnitude to r_B , which is of order 0.1 for $B^+ \rightarrow DK^+$ decays. Moreover, such a determination of γ suffers from trigonometric ambiguities since there are only three linearly independent observables in the GLW method, even in the case that both CP -even and CP -odd D decay final states are used.

To overcome these issues, it is essential to include D decays to non- CP eigenstates. The use of $B^+ \rightarrow DK^+$ with subsequent $D \rightarrow K^\mp\pi^\pm$ decays was noted as being particularly valuable [11], since both doubly Cabibbo-suppressed (*e.g.* $\bar{D}^0 \rightarrow K^-\pi^+$) and Cabibbo-favoured (*e.g.* $D^0 \rightarrow K^-\pi^+$) amplitudes contribute. The smallness of the relative magnitude of these amplitudes, denoted r_D , complements the size of r_B , so that larger CP asymmetries are possible in $B^+ \rightarrow DK^+$ decays. Both r_D and the relative strong phase between the D decay amplitudes, δ_D , can be determined independently of the B decay data, so that the observables depend on the same set of unknown parameters (γ , r_B , δ_B) as in the GLW case. This approach is now widely known as the ADS method. While the two-body $D \rightarrow K^\mp\pi^\pm$ decays are particularly attractive experimentally, the ADS method can also be applied for multibody decays such as $D \rightarrow K^\mp\pi^\pm\pi^0$ and $D \rightarrow K^\mp\pi^\pm\pi^+\pi^-$ [11, 12].

Before the start of the BaBar and Belle experiments, γ was expected to be constrained only weakly by the anticipated data [13]. As the first results with the GLW and ADS methods were published [14–17], their precision confirmed that much larger data samples would be necessary to constrain γ to the 10° level or better.

¹An alternative notation, $\phi_3 \equiv \gamma$, is also in widespread use in the literature.

This situation motivated the investigation of further decay modes that could be usefully employed to determine γ . Several authors noted that the available statistics could be increased by using multibody final states, including those from doubly Cabibbo-suppressed [12] and singly Cabibbo-suppressed [18] D decays. However, these focussed either on inclusive approaches, in which the whole phase-space was integrated over, or on contributions from particular resonances. The key point of how interference between different resonances in the Dalitz plot of a multibody D decay could be exploited to measure γ was not realised until it was proposed to use decays into self-conjugate multibody final states, such as $K_s^0\pi^+\pi^-$. This method has proven over time to have very good sensitivity. The original work was performed separately and independently by two groups. Bondar and Poluektov developed their ideas within the Belle collaboration, while Giri, Grossman, Soffer and Zupan developed theirs for a theoretical paper.

The first presentation on such a method was made by Alex Bondar at an internal Belle collaboration workshop in September 2002 [19]. The slides of this presentation, which have not previously been available publicly, are included in Appendix A. The idea was inspired by the Dalitz plot analysis of $D^0 \rightarrow K_s^0\pi^+\pi^-$ decays by the CLEO collaboration that had been published earlier in 2002 [20], in which it was demonstrated that the relative magnitude and phase of the doubly Cabibbo-suppressed $D^0 \rightarrow K^{*+}\pi^-$ and Cabibbo-favoured $D^0 \rightarrow K^{*-}\pi^+$ decay amplitudes could be determined from a sample of flavour-tagged D mesons. If the same relative phase could be measured separately in samples of neutral D mesons from $B^+ \rightarrow DK^+$ and charge-conjugate decays, the difference between these two quantities could be used to obtain 2γ . The solution thus obtained would be unique in the range $[0, \pi]$. The concept was developed further, subsequently to the initial presentation, and implemented with the Belle data by Anton Poluektov. Due to the strong competition between Belle and BaBar at that time, it was decided not to publish the method as a standalone paper, but rather to describe the approach together with the experimental results in a Belle collaboration publication. The first results were presented in preliminary form at the Lepton Photon conference in August 2003 [21], and published not long thereafter [22].

The first publication of the idea was the paper by Anjan Giri, Yuval Grossman, Abner Soffer, and Jure Zupan [23]. This was made available on the arXiv preprint server in March 2003, and presented at the CKM 2003 Workshop in early April, prior to publication. In addition to discussing that γ could be determined from a model-dependent fit to the Dalitz plot distribution of $D \rightarrow K_s^0\pi^+\pi^-$ decays produced in $B^+ \rightarrow DK^+$ and charge conjugate processes, Ref. [23] proposed a model-independent approach based on binning the Dalitz plot. This model-independent approach removes a potentially large source of uncertainty originating from the description of the *a priori* unknown strong phase variation across the Dalitz plot, at the cost of some statistical precision due to the finite size of the bins. Each bin has associated with it a set of hadronic parameters, corresponding to the r_D and δ_D parameters of the ADS method, that can in principle be determined independently of the B decay data, so that again the only unknowns to be found are (γ, r_B, δ_B) . In particular, Ref. [23] introduced the c_i and s_i parameters, which are the amplitude-weighted averages of the cosine and sine of the strong phase difference between D^0 and \bar{D}^0 decay amplitudes

within Dalitz plot bin i . These parameters can be determined from quantum-correlated $\psi(3770) \rightarrow D\bar{D}$ decays.

Both model-dependent and model-independent variants of the method to determine γ from $B^\pm \rightarrow DK^\pm$ with multibody D decays, which we propose to refer to as the BPGGSZ method (see Appendix B for a recommendation on appropriate citations), have been intensively pursued by experiments.² The most recent results from the BaBar [24], Belle [25] and LHCb [26] collaborations with the model-dependent approach, using $D \rightarrow K_s^0 \pi^+ \pi^-$ decay models obtained in Refs. [27,28],³ have statistical uncertainties on γ as low as 12–15° and model uncertainties that vary in the range 3–9° depending on how conservative a range of model variations is considered. The latest results from the Belle [30] and LHCb [31,32] collaborations with the model-independent approach have uncertainties on γ of 10–15°. A small contribution to this, around 4°, is due to the limited precision with which the c_i and s_i parameters have been measured using data from the CLEOc experiment [33,34] following the scheme for binning of the Dalitz plot proposed in Refs. [35,36].⁴ Further improvement in precision can be anticipated as the existing LHCb data sample is analysed, and as much larger data samples are collected in the future with upgrades of the LHCb detector [39–42] and with the Belle II experiment [43,44].

In addition to its use in the $B^+ \rightarrow DK^+$, $D \rightarrow K_s^0 \pi^+ \pi^-$ decay chain, the BPGGSZ method has also been used with $B^+ \rightarrow D^* K^+$ [24,25], $B^+ \rightarrow DK^{*+}$ [24,28] and $B^0 \rightarrow DK^{*0}$ [45–47] decays. In addition, D decays to the $K_s^0 K^+ K^-$ [24,31,32] and $K_s^0 \pi^+ \pi^- \pi^0$ [48,49] final states have been exploited and application of the method with other self-conjugate multibody decays is likely in the future. Moreover, the BPGGSZ method has inspired similar methods to determine additional important quantities in heavy flavour physics. Analysis of $B \rightarrow D\pi^0$ and similar decays with $D \rightarrow K_s^0 \pi^+ \pi^-$ and other self-conjugate multibody final states can be used to determine the angle β of the unitarity triangle [50]; this method has been implemented in both model-dependent [29,51] and model-independent [52] variants. A “double Dalitz plot” analysis for $B \rightarrow D\pi^+ \pi^-$ with $D \rightarrow K_s^0 \pi^+ \pi^-$ decays has also been proposed [53], building on a similar concept for γ determination [54,55]. Binning of the $D \rightarrow K_s^0 \pi^+ \pi^-$ Dalitz plot has also been noted to have highly advantageous feature for the experimental determination of D meson mixing parameters from these decays [56–58]. Results obtained with these methods [59,60] currently provide the best sensitivity out of all charm mixing measurements to the mass difference between the neutral D meson eigenstates.

In conclusion, the development of the BPGGSZ method enabled significant improvement in the measurement of γ , removing ambiguities in its determination. This permitted global fits to the parameters of the CKM matrix, allowing for, and deriving stringent bounds on, contributions from physics beyond the SM in loop-dominated processes [61,62]. Such analyses proved that the dominant contributions to flavour-changing processes and to CP

²This approach has until now been referred to in a variety of ways, sometimes using the GGSZ acronym.

³An updated study of the $D \rightarrow K_s^0 \pi^+ \pi^-$ decay amplitude has been published in Ref. [29], but not yet used in any γ determination.

⁴More precise measurements of the c_i and s_i parameters have recently been reported by the BESIII collaboration [37,38], but not yet used in any γ determination.

violation in meson decays are those of the Standard Model, as recognised by the award of the 2008 Nobel Prize in Physics to Kobayashi and Maskawa. Results obtained with the BPGGSZ method continue to provide some of the most precise constraints on γ that enter today's global fits [4, 5], and are expected to continue to do so in the future.

Acknowledgements

The authors gratefully acknowledge Alex Bondar, Anton Poluektov, Anjan Giri, Yuval Grossman, Abner Soffer and Jure Zupan for providing background information, and for comments on the manuscript. The authors would also like to thank Sébastien Descotes-Genon, Jim Libby and Sneha Malde for helpful discussions.

References

- [1] N. Cabibbo, *Unitary symmetry and leptonic decays*, *Phys. Rev. Lett.* **10** (1963) 531.
- [2] M. Kobayashi and T. Maskawa, *CP violation in the renormalizable theory of weak interaction*, *Prog. Theor. Phys.* **49** (1973) 652.
- [3] I. I. Y. Bigi and A. I. Sanda, *On direct CP violation in $B \rightarrow D^0/\bar{D}^0 K\pi$'s versus $\bar{B} \rightarrow D^0/\bar{D}^0 \bar{K}\pi$'s decays*, *Phys. Lett.* **B211** (1988) 213.
- [4] Particle Data Group, P. A. Zyla *et al.*, *Review of particle physics*, *Prog. Theor. Exp. Phys.* (2020) 083C01.
- [5] Heavy Flavor Averaging Group (HFLAV), Y. S. Amhis *et al.*, *Averages of b-hadron, c-hadron, and τ -lepton properties as of 2018*, [arXiv:1909.12524](https://arxiv.org/abs/1909.12524), updated results and plots available at <http://www.slac.stanford.edu/xorg/hflav/>.
- [6] BaBar and Belle collaborations, A. J. Bevan *et al.*, *The physics of the B factories*, *Eur. Phys. J.* **C74** (2014) 3026, [arXiv:1406.6311](https://arxiv.org/abs/1406.6311).
- [7] LHCb collaboration, R. Aaij *et al.*, *Measurement of the CKM angle γ from a combination of LHCb results*, *JHEP* **12** (2016) 087, [arXiv:1611.03076](https://arxiv.org/abs/1611.03076).
- [8] T. Gershon and V. V. Gligorov, *CP violation in the B system*, *Rept. Prog. Phys.* **80** (2017) 046201, [arXiv:1607.06746](https://arxiv.org/abs/1607.06746).
- [9] M. Gronau and D. London, *How to determine all the angles of the unitarity triangle from $B^0 \rightarrow DK_s^0$ and $B_s^0 \rightarrow D\phi$* , *Phys. Lett.* **B253** (1991) 483.
- [10] M. Gronau and D. Wyler, *On determining a weak phase from CP asymmetries in charged B decays*, *Phys. Lett.* **B265** (1991) 172.

- [11] D. Atwood, I. Dunietz, and A. Soni, *Enhanced CP violation with $B \rightarrow KD^0(\bar{D}^0)$ modes and extraction of the CKM angle γ* , *Phys. Rev. Lett.* **78** (1997) 3257, [arXiv:hep-ph/9612433](#).
- [12] D. Atwood, I. Dunietz, and A. Soni, *Improved methods for observing CP violation in $B^\pm \rightarrow KD$ and measuring the CKM phase γ* , *Phys. Rev.* **D63** (2001) 036005, [arXiv:hep-ph/0008090](#).
- [13] BaBar collaboration, D. Boutigny *et al.*, *The BABAR physics book: Physics at an asymmetric B factory*, in *Report of the BABAR Physics Workshop* (P. F. Harrison and H. R. Quinn, eds.), 1998. Available as [SLAC report SLAC-R-504](#).
- [14] Belle collaboration, K. Abe *et al.*, *Studies of the decay $B^+ \rightarrow D_{CP}K^\pm$* , *Phys. Rev. Lett.* **90** (2003) 131803, [arXiv:hep-ex/0207012](#).
- [15] Belle collaboration, S. K. Swain *et al.*, *Measurement of branching fraction ratios and CP asymmetries in $B^\pm \rightarrow D_{CP}K^\pm$* , *Phys. Rev.* **D68** (2003) 051101, [arXiv:hep-ex/0304032](#).
- [16] BaBar collaboration, B. Aubert *et al.*, *Measurement of the branching fractions and CP-asymmetry of $B^- \rightarrow D_{CP}^0K^-$ decays with the BaBar detector*, *Phys. Rev. Lett.* **92** (2004) 202002, [arXiv:hep-ex/0311032](#).
- [17] BaBar collaboration, B. Aubert *et al.*, *Search for $B^\pm \rightarrow (K^\mp\pi^\pm)_DK^\pm$ and upper limit on the $b \rightarrow u$ amplitude in $B^\pm \rightarrow DK^\pm$* , *Phys. Rev. Lett.* **93** (2004) 131804, [arXiv:hep-ex/0402024](#).
- [18] Y. Grossman, Z. Ligeti, and A. Soffer, *Measuring γ in $B^\pm \rightarrow K^\pm(KK^*)_D$ decays*, *Phys. Rev.* **D67** (2003) 071301, [arXiv:hep-ph/0210433](#).
- [19] A. Bondar. *Proceedings of BINP special analysis meeting on Dalitz analysis*, 24–26 Sep. 2002, unpublished.
- [20] CLEO collaboration, H. Muramatsu *et al.*, *Dalitz analysis of $D^0 \rightarrow K_S^0\pi^+\pi^-$* , *Phys. Rev. Lett.* **89** (2002) 251802, Erratum *ibid.* **90** (2003) 059901, [arXiv:hep-ex/0207067](#).
- [21] Belle collaboration, K. Abe *et al.*, *Measurement of the angle ϕ_3 with Dalitz analysis of three body D^0 decay from $B \rightarrow D^0K$ process*, in *Proceedings, 21st International Symposium on Lepton and Photon Interactions at High Energies (LP 03)*, 2003, [arXiv:hep-ex/0308043](#).
- [22] Belle collaboration, A. Poluektov *et al.*, *Measurement of ϕ_3 with Dalitz plot analysis of $B^\pm \rightarrow D^{(*)}K^\pm$ decay*, *Phys. Rev.* **D70** (2004) 072003, [arXiv:hep-ex/0406067](#).
- [23] A. Giri, Y. Grossman, A. Soffer, and J. Zupan, *Determining γ using $B^\pm \rightarrow DK^\pm$ with multibody D decays*, *Phys. Rev.* **D68** (2003) 054018, [arXiv:hep-ph/0303187](#).

- [24] BaBar collaboration, P. del Amo Sanchez *et al.*, *Evidence for direct CP violation in the measurement of the Cabibbo-Kobayashi-Maskawa angle γ with $B^\mp \rightarrow D^{(*)}K^{(*)\mp}$ decays*, *Phys. Rev. Lett.* **105** (2010) 121801, [arXiv:1005.1096](#).
- [25] Belle collaboration, A. Poluektov *et al.*, *Evidence for direct CP violation in the decay $B \rightarrow D^{(*)}K$, $D \rightarrow K_S^0\pi^+\pi^-$ and measurement of the CKM phase ϕ_3* , *Phys. Rev.* **D81** (2010) 112002, [arXiv:1003.3360](#).
- [26] LHCb collaboration, R. Aaij *et al.*, *Measurement of CP violation and constraints on the CKM angle γ in $B^\pm \rightarrow DK^\pm$ with $D \rightarrow K_S^0\pi^+\pi^-$ decays*, *Nucl. Phys.* **B888** (2014) 169, [arXiv:1407.6211](#).
- [27] BaBar collaboration, P. del Amo Sanchez *et al.*, *Measurement of D^0 - \bar{D}^0 mixing parameters using $D^0 \rightarrow K_S^0\pi^+\pi^-$ and $D^0 \rightarrow K_S^0K^+K^-$ decays*, *Phys. Rev. Lett.* **105** (2010) 081803, [arXiv:1004.5053](#).
- [28] Belle collaboration, A. Poluektov *et al.*, *Measurement of ϕ_3 with Dalitz plot analysis of $B^+ \rightarrow D^{(*)}K^{(*)+}$ decay*, *Phys. Rev.* **D73** (2006) 112009, [arXiv:hep-ex/0604054](#).
- [29] BaBar and Belle collaborations, I. Adachi *et al.*, *Measurement of $\cos 2\beta$ in $B^0 \rightarrow D^{(*)}h^0$ with $D \rightarrow K_S^0\pi^+\pi^-$ decays by a combined time-dependent Dalitz plot analysis of BaBar and Belle data*, *Phys. Rev.* **D98** (2018) 112012, [arXiv:1804.06153](#).
- [30] Belle collaboration, H. Aihara *et al.*, *First measurement of ϕ_3 with a model-independent Dalitz plot analysis of $B \rightarrow DK$, $D \rightarrow K_S^0\pi^+\pi^-$ decay*, *Phys. Rev.* **D85** (2012) 112014, [arXiv:1204.6561](#).
- [31] LHCb collaboration, R. Aaij *et al.*, *Measurement of the CKM angle γ using $B^\pm \rightarrow DK^\pm$ with $D \rightarrow K_S^0\pi^+\pi^-$, $K_S^0K^+K^-$ decays*, *JHEP* **10** (2014) 097, [arXiv:1408.2748](#).
- [32] LHCb collaboration, R. Aaij *et al.*, *Measurement of the CKM angle γ using $B^\pm \rightarrow DK^\pm$ with $D \rightarrow K_S^0\pi^+\pi^-$, $K_S^0K^+K^-$ decays*, *JHEP* **08** (2018) 176, Erratum *ibid.* **10** (2018) 107, [arXiv:1806.01202](#).
- [33] CLEO collaboration, R. A. Briere *et al.*, *First model-independent determination of the relative strong phase between D^0 and $\bar{D}^0 \rightarrow K_S^0\pi^+\pi^-$ and its impact on the CKM angle γ/ϕ_3 measurement*, *Phys. Rev.* **D80** (2009) 032002, [arXiv:0903.1681](#).
- [34] CLEO collaboration, J. Libby *et al.*, *Model-independent determination of the strong-phase difference between D^0 and $\bar{D}^0 \rightarrow K_{S,L}^0h^+h^-$ ($h = \pi, K$) and its impact on the measurement of the CKM angle γ/ϕ_3* , *Phys. Rev.* **D82** (2010) 112006, [arXiv:1010.2817](#).
- [35] A. Bondar and A. Poluektov, *Feasibility study of model-independent approach to ϕ_3 measurement using Dalitz plot analysis*, *Eur. Phys. J.* **C47** (2006) 347, [arXiv:hep-ph/0510246](#).

- [36] A. Bondar and A. Poluektov, *The use of quantum-correlated D^0 decays for ϕ_3 measurement*, *Eur. Phys. J.* **C55** (2008) 51, [arXiv:0801.0840](#).
- [37] BESIII collaboration, M. Ablikim *et al.*, *Determination of strong-phase parameters in $D \rightarrow K_{S,L}^0 \pi^+ \pi^-$* , *Phys. Rev. Lett.* **124** (2020) 241802, [arXiv:2002.12791](#).
- [38] BESIII collaboration, M. Ablikim *et al.*, *Model-independent determination of the relative strong-phase difference between D^0 and $\bar{D}^0 \rightarrow K_{S,L}^0 \pi^+ \pi^-$ and its impact on the measurement of the CKM angle γ/ϕ_3* , *Phys. Rev.* **D101** (2020) 112002, [arXiv:2003.00091](#).
- [39] LHCb collaboration, I. Bediaga *et al.*, *Framework TDR for the LHCb upgrade: Technical design report*, LHCb-TDR-012, [CERN-LHCC-2012-007](#), 2012.
- [40] LHCb collaboration, R. Aaij *et al.*, and A. Bharucha *et al.*, *Implications of LHCb measurements and future prospects*, *Eur. Phys. J.* **C73** (2013) 2373, [arXiv:1208.3355](#).
- [41] LHCb collaboration, R. Aaij *et al.*, *Expression of Interest for a Phase-II LHCb Upgrade: Opportunities in flavour physics, and beyond, in the HL-LHC era*, [CERN-LHCC-2017-003](#), 2017.
- [42] LHCb collaboration, *Physics case for an LHCb Upgrade II — Opportunities in flavour physics, and beyond, in the HL-LHC era*, [arXiv:1808.08865](#), CERN-LHCC-2018-027.
- [43] Belle II collaboration, T. Abe *et al.*, *Belle II technical design report*, [arXiv:1011.0352](#).
- [44] Belle II collaboration, W. Altmannshofer *et al.*, *The Belle II Physics Book*, [PTEP \(2019\) 123C01](#), Erratum *ibid.* (2020) 029201, [arXiv:1808.10567](#).
- [45] Belle collaboration, K. Negishi *et al.*, *First model-independent Dalitz analysis of $B^0 \rightarrow DK^{*0}$, $D \rightarrow K_S^0 \pi^+ \pi^-$ decay*, *PTEP* **2016** (2016) 043C01, [arXiv:1509.01098](#).
- [46] LHCb collaboration, R. Aaij *et al.*, *Model-independent measurement of the CKM angle γ using $B^0 \rightarrow DK^{*0}$ decays with $D \rightarrow K_S^0 \pi^+ \pi^-$ and $K_S^0 K^+ K^-$* , *JHEP* **06** (2016) 131, [arXiv:1604.01525](#).
- [47] LHCb collaboration, R. Aaij *et al.*, *Measurement of the CKM angle γ using $B^0 \rightarrow DK^{*0}$ with $D \rightarrow K_S^0 \pi^+ \pi^-$ decays*, *JHEP* **08** (2016) 137, [arXiv:1605.01082](#).
- [48] P. K. Resmi, J. Libby, S. Malde, and G. Wilkinson, *Quantum-correlated measurements of $D \rightarrow K_S^0 \pi^+ \pi^- \pi^0$ decays and consequences for the determination of the CKM angle γ* , *JHEP* **01** (2018) 082, [arXiv:1710.10086](#).
- [49] Belle collaboration, P. K. Resmi *et al.*, *First measurement of the CKM angle ϕ_3 with $B^\pm \rightarrow D(K_S^0 \pi^+ \pi^- \pi^0) K^\pm$ decays*, *JHEP* **10** (2019) 178, [arXiv:1908.09499](#).
- [50] A. Bondar, T. Gershon, and P. Krokovny, *A method to measure ϕ_1 using $\bar{B}^0 \rightarrow Dh^0$ with multibody D decay*, *Phys. Lett.* **B624** (2005) 1, [arXiv:hep-ph/0503174](#).

- [51] BaBar and Belle collaborations, I. Adachi *et al.*, *First evidence for $\cos 2\beta > 0$ and resolution of the Cabibbo-Kobayashi-Maskawa quark-mixing Unitarity Triangle ambiguity*, *Phys. Rev. Lett.* **121** (2018) 261801, [arXiv:1804.06152](#).
- [52] Belle collaboration, V. Vorobyev *et al.*, *Measurement of the CKM angle φ_1 in $B^0 \rightarrow \bar{D}^{(*)0}h^0$, $\bar{D}^0 \rightarrow K_S^0\pi^+\pi^-$ decays with time-dependent binned Dalitz plot analysis*, *Phys. Rev.* **D94** (2016) 052004, [arXiv:1607.05813](#).
- [53] A. Bondar, A. Kuzmin, and V. Vorobyev, *A method for model-independent measurement of the CKM angle β via time-dependent analysis of the $B^0 \rightarrow D\pi^+\pi^-$, $D \rightarrow K_S^0\pi^+\pi^-$ decays*, *JHEP* **03** (2018) 195, [arXiv:1802.00200](#).
- [54] T. Gershon and A. Poluektov, *Double Dalitz plot analysis of the decay $B^0 \rightarrow DK^+\pi^-$, $D \rightarrow K_S^0\pi^+\pi^-$* , *Phys. Rev.* **D81** (2010) 014025, [arXiv:0910.5437](#).
- [55] D. Craik, T. Gershon, and A. Poluektov, *Optimising sensitivity to γ with $B^0 \rightarrow DK^+\pi^-$, $D \rightarrow K_S^0\pi^+\pi^-$ double Dalitz plot analysis*, *Phys. Rev.* **D97** (2018) 056002, [arXiv:1712.07853](#).
- [56] A. Bondar, A. Poluektov, and V. Vorobiev, *Charm mixing in the model-independent analysis of correlated $D^0\bar{D}^0$ decays*, *Phys. Rev.* **D82** (2010) 034033, [arXiv:1004.2350](#).
- [57] C. Thomas and G. Wilkinson, *Model-independent $D^0 - \bar{D}^0$ mixing and CP violation studies with $D^0 \rightarrow K_S^0\pi^+\pi^-$ and $D^0 \rightarrow K_S^0K^+K^-$* , *JHEP* **10** (2012) 185, [arXiv:1209.0172](#).
- [58] A. Di Canto *et al.*, *Novel method for measuring charm-mixing parameters using multibody decays*, *Phys. Rev.* **D99** (2019) 012007, [arXiv:1811.01032](#).
- [59] LHCb collaboration, R. Aaij *et al.*, *Model-independent measurement of mixing parameters in $D^0 \rightarrow K_S^0\pi^+\pi^-$ decays*, *JHEP* **04** (2016) 033, [arXiv:1510.01664](#).
- [60] LHCb collaboration, R. Aaij *et al.*, *Measurement of the mass difference between neutral char m -meson eigenstates*, *Phys. Rev. Lett.* **122** (2019) 231802, [arXiv:1903.03074](#).
- [61] CKMfitter Group, J. Charles *et al.*, *CP violation and the CKM matrix: Assessing the impact of the asymmetric B factories*, *Eur. Phys. J.* **C41** (2005) 1, [arXiv:hep-ph/0406184](#), updated results and plots available at: <http://ckmfitter.in2p3.fr>.
- [62] UTfit collaboration, M. Bona *et al.*, *The UTfit collaboration report on the status of the unitarity triangle beyond the standard model. I. Model-independent analysis and minimal flavor violation*, *JHEP* **03** (2006) 080, [arXiv:hep-ph/0509219](#), updated results and plots available at: <http://www.utfit.org/UTfit>.

A First presentation of method to determine γ from $B^+ \rightarrow DK^+, D \rightarrow K_S^0 \pi^+ \pi^-$ decays


Figures 1–5 contain the slides of the presentation at the Belle collaboration workshop in September 2002 [19], in which the concept of measuring γ (denoted ϕ_s in the slides) from $B^+ \rightarrow DK^+, D \rightarrow K_S^0 \pi^+ \pi^-$ decays was first set out.

Improved Gronau-Wyler method for ϕ_3 extraction

Alex Bondar, BINP, Novosibirsk

*Special analysis meeting
Novosibirsk, 24-26 Sep. 2002*

Gronau-Wyler method for ϕ_3 extraction



$$|D_+^0\rangle = \frac{1}{\sqrt{2}} (|D^0\rangle + |\bar{D}^0\rangle),$$

$$\sqrt{2} A(B^+ \rightarrow K^+ D_+^0) = A(B^+ \rightarrow K^+ D^0) + A(B^+ \rightarrow K^+ \bar{D}^0)$$

$$\sqrt{2} A(B^- \rightarrow K^- D_+^0) = A(B^- \rightarrow K^- \bar{D}^0) + A(B^- \rightarrow K^- D^0).$$

$$A(B^+ \rightarrow K^+ \bar{D}^0) = A(B^- \rightarrow K^- D^0)$$

$$A(B^+ \rightarrow K^+ D^0) = A(B^- \rightarrow K^- \bar{D}^0) \times e^{2i\gamma},$$





Figure 1: Slides 1–2 of the presentation of Ref. [19].



decay mode	$D\pi$	DK	Feed-down	Stat. Sig
$B^\pm \rightarrow D^0 h^\pm$	6052 ± 87.7	347.5 ± 21	134.4 ± 14.7	26.1
$B^\pm \rightarrow D_1 h^\pm$	666.9 ± 32.9	51 ± 9.3	12.9 ± 6.3	7.6
$B^\pm \rightarrow D_2 h^\pm$	639.3 ± 29.7	57.3 ± 9.6	9.9 ± 5.6	8.2

decay mode	B^+	B^-
$D^0\pi$	3085.9 ± 68.7	2981.7 ± 62
D^0K	165.4 ± 14.5	179.6 ± 15.0
Feed-across	66.4 ± 10.0	71.8 ± 10.6
$D_1\pi$	302 ± 19.4	339.3 ± 21.6
D_1K	24.5 ± 6.5	26.5 ± 7.9
Feed-across	5.2 ± 4.2	7.9 ± 4.6
$D_2\pi$	339 ± 22.5	291.5 ± 22.7
D_2K	31.8 ± 6.8	23.9 ± 6.3
Feed-across	4.6 ± 3.6	0.7 ± 3.2

Figure 2: Slides 3–4 of the presentation of Ref. [19].

TABLE I. Standard fit results. The errors shown are statistical, experimental systematic, and modeling systematic respectively. See the text for further discussion.

Component	Amplitude	Phase	Fit Fraction (%)
$K^*(892)^+\pi^-\times B(K^*(892)^+\rightarrow K^0\pi^+)$	$(11\pm 2\pm 1\pm 1)^{+4}_{-1}\times 10^{-2}$	$321\pm 10\pm 3\pm 15$	$0.34\pm 0.13\pm 0.03\pm 0.02$
$\overline{K}^0\rho^0$	1.0 (fixed)	0 (fixed)	$26.4\pm 0.9\pm 0.9\pm 0.4$
$\overline{K}^0\omega\times B(\omega\rightarrow\pi^+\pi^-)$	$(37\pm 5\pm 1\pm 3)\times 10^{-3}$	$114\pm 7\pm 4\pm 5$	$0.72\pm 0.18\pm 0.04\pm 0.10$
$K^*(892)^-\pi^+\times B(K^*(892)^-\rightarrow\overline{K}^0\pi^-)$	$1.56\pm 0.03\pm 0.02\pm 0.15$	$150\pm 2\pm 2\pm 2$	$65.7\pm 1.3\pm 1.1\pm 1.4$
$\overline{K}^0f_0(980)\times B(f_0(980)\rightarrow\pi^+\pi^-)$	$0.34\pm 0.02\pm 0.04\pm 0.04$	$188\pm 4\pm 5\pm 8$	$4.3\pm 0.5\pm 0.4\pm 0.5$
$\overline{K}^0f_2(1270)\times B(f_2(1270)\rightarrow\pi^+\pi^-)$	$0.7\pm 0.2\pm 0.3\pm 0.4$	$308\pm 12\pm 15\pm 66$	$0.27\pm 0.15\pm 0.24\pm 0.28$
$\overline{K}^0f_0(1370)\times B(f_0(1370)\rightarrow\pi^+\pi^-)$	$1.8\pm 0.1\pm 0.2\pm 0.2$	$85\pm 4\pm 4\pm 13$	$9.9\pm 1.1\pm 2.4\pm 1.4$
$K_0^*(1430)^-\pi^+\times B(K_0^*(1430)^-\rightarrow\overline{K}^0\pi^-)$	$2.0\pm 0.1\pm 0.1\pm 0.5$	$3\pm 4\pm 4\pm 7$	$7.3\pm 0.7\pm 0.4\pm 3.1$
$K_2^*(1430)^-\pi^+\times B(K_2^*(1430)^-\rightarrow\overline{K}^0\pi^-)$	$1.0\pm 0.1\pm 0.1\pm 0.3$	$155\pm 7\pm 1\pm 7$	$1.1\pm 0.2\pm 0.3\pm 0.6$
$K^*(1680)^-\pi^+\times B(K^*(1680)^-\rightarrow\overline{K}^0\pi^-)$	$5.6\pm 0.6\pm 0.7\pm 4.0$	$174\pm 6\pm 10\pm 13$	$2.2\pm 0.4\pm 0.5\pm 1.7$
$\overline{K}^0\pi^+\pi^-$ non-resonant	$1.1\pm 0.3\pm 0.5\pm 0.9$	$160\pm 11\pm 30\pm 55$	$0.9\pm 0.4\pm 1.0\pm 1.7$

Base on 5299 events CLEO got the relative phase of the double Cabibbo suppressed amplitude with accuracy of 10^0 . In case $B^+\rightarrow D^0(\overline{D}^0)K^+$ decay the ratio of the color suppressed and color allowed amplitudes is in order of $\frac{a_2|V_{ub}|V_{cs}^*}{a_1|V_{cb}|V_{cs}^*}\approx\frac{|V_{ub}|}{3|V_{cb}|\lambda}\approx\frac{1}{8}$, which is greater than in case of double Cabibbo suppressed amplitude by $1/8\lambda^2$ times. This results in the lower number of events (~ 800) necessary for the similar accuracy in ϕ_3 measurement.

The similar method applicable to the $\overline{B}^0\rightarrow D^0(\overline{D}^0)K^{*0}$, $D^0(\overline{D}^0)\rightarrow K_S^0\pi^+\pi^-$ decay. In this case the ratio of the amplitudes even more suitable ($\frac{|V_{ub}|}{|V_{cb}|\lambda}\approx 0.4$).





Figure 4: Slides 7–8 of the presentation of Ref. [19].

$\bar{B}^0 \rightarrow D^0 \bar{K}^{*0}$ decay (sum of three modes)



Conclusion

Suggested method of ϕ_3 extraction has the following advantages:

- Only one final state ($K^{(*)}K_S^0\pi^+\pi^-$) is used in the analysis.
- Therefore, the experimental systematic errors should cancel.
- The extraction of ϕ_3 is possible with any strong phase difference.
- Double Cabibbo suppressed amplitudes do not effect on ϕ_3 extraction.
- This method helps resolve discrete ambiguities that are usually present in measurements of the weak phase ϕ_3 .

Need to do:

- MC study of necessary statistics for different ϕ_3 and strong phase difference.
- Study of the possible model error due to parameterization of the $D^0 \rightarrow K_S^0\pi^+\pi^-$ decay amplitude.

Figure 5: Slides 9–10 of the presentation of Ref. [19].

B Recommendation for citation of BPGGSZ method

We recommend to use the following citations for the BPGGSZ method:

- [1] A. Bondar. Proceedings of BINP special analysis meeting on Dalitz analysis, 24–26 Sep. 2002, unpublished.
- [2] A. Giri, Y. Grossman, A. Soffer, and J. Zupan, Determining γ using $B^+ \rightarrow DK^\pm$ with multibody D decays, [Phys. Rev. **D68** \(2003\) 054018](#), [arXiv:hep-ph/0303187](#).
- [3] Belle collaboration, A. Poluektov et al., Measurement of ϕ_3 with Dalitz plot analysis of $B^\pm \rightarrow D^{(*)}K^\pm$ decay, [Phys. Rev. **D70** \(2004\) 072003](#), [arXiv:hep-ex/0406067](#).