

Numerical considerations about the SIR epidemic model with infection age

Ralph Brinks, Annika Hoyer

Department of Statistics, Ludwig-Maximilians-University
Munich, Germany

We analyse the infection-age-dependent SIR model from a numerical point of view. First, we present an algorithm for calculating the solution the infection-age-structured SIR model without demography of the background host. Second, we examine how and under which conditions, the conventional SIR model (without infection-age) serves as a practical approximation to the infection-age SIR model. Special emphasis is given on the effective reproduction number.

Introduction

We analyse the infection-age SIR model without demography of the background host, whose foundations date back to Kermack and McKendrick [Ker27]. The focus is primarily on numerical aspects. In the SIR model, the population is partitioned into three states, the *susceptible* state, the *infected* and the *removed* state (the initial letters of the three states give the model's name 'SIR'). The *removed* state comprises people recovered and deceased from the *infected* state. The numbers of the people in the susceptible and the removed states at time t are denoted by $S(t)$ and $R(t)$, respectively.

Furthermore, let $i(t, \tau)$ denote the density of infected people at time t and duration τ since infection (i.e., the *infection age*), such that the number $I(t)$ of infected at t is

$$I(t) = \int_0^\infty i(t, \tau) d\tau. \quad (1)$$

The transmission rate of the infected with infection age τ is $\beta(t, \tau)$ and the removal rate from the infectious stage is $\gamma(\tau)$. The rate γ comprises mortality as well as remission. We can formulate the following model equations for the infection-age SIR model [Ina17]:

$$\frac{dS(t)}{dt} = -\lambda(t) S(t) \quad (2)$$

$$\left(\frac{\partial}{\partial t} + \frac{\partial}{\partial \tau} \right) i(t, \tau) = -\gamma(\tau) i(t, \tau) \quad (3)$$

$$\frac{dR(t)}{dt} = \int_0^\infty \gamma(\tau) i(t, \tau) d\tau. \quad (4)$$

The incidence rate λ in Eq. (2) is given by

$$\lambda(t) = \int_0^\infty \beta(t, \tau) i(t, \tau) d\tau$$

and is usually called *force of infection* [Ina17].

System (2) – (4) is accompanied with initial conditions

$$S(0) = S_0 \quad (5)$$

$$i(t, 0) = \lambda(t) S(t) \quad (6)$$

$$i(0, \tau) = i_0(\tau) \quad (7)$$

$$i(0, 0) = S_0 \int_0^\infty \beta(0, \tau) i_0(\tau) d\tau \quad (8)$$

with positive S_0 and integrable i_0 . For later use, we additionally assume that $i(t, \infty) := \lim_{\tau \rightarrow \infty} i(t, \tau) = 0$. Condition (8) is called *coupling equation* and guarantees that system (2) – (4) is well-defined [Che16]. Note that system (2) – (4) is a generalisation of the SEIR model [Ina17, Section 5.5].

Detailed discussion of Equations (2) – (4) with initial conditions (5) – (8) can be found in [Ina17, Chapter 5.3]. Using the definition

$$\Gamma(\tau) := \exp\left(-\int_0^\tau \gamma(\sigma) d\sigma\right), \quad (9)$$

the effective reproduction number $\mathcal{R}(t)$ is given by

$$\mathcal{R}(t) = S(t) \int_0^\infty \beta(t, \tau) \Gamma(\tau) d\tau, \quad (10)$$

[Nis09, Eq. (22),(23)].

A typical situation is that the transmission rate $\beta(t, \tau)$ and the initial conditions (5) – (8) are given. Then, system (2) – (4) is solved and the effective reproduction number \mathcal{R} is calculated by Eq. (10). In a way, \mathcal{R} can be seen as an indirect parameter for the infection-age-structured SIR model because it follows via Eq. (10) from the governing equations (2) – (4) and (5) – (8). Sometimes, $\mathcal{R}(t)$ can be estimated more easily from population surveys than, for instance, the transmission rate $\beta(\tau, t)$. Then, the question arises if and how the infection-age-structured SIR model can be solved if the effective reproduction number \mathcal{R} is given instead of β .

This article is organised as followed: First, we describe a numerical algorithm to solve the system given by Equations (2) – (4) with initial conditions (5) – (8) on a rectangular grid. Then, we consider an important special case where the transmission rate β does not need to be known to solve system (2) – (4). Finally, we present an example to demonstrate the theoretical considerations.

Numerical solution of the infection-age-structured SIR model

Assumed $i(t, \tau)$ has to be calculated on a rectangular grid $(t_m, \tau_n) = (m \times \delta_h, n \times \delta_h)$, $m = 0, \dots, M$, $n = 0, \dots, N$, as depicted in Figure 1. The grid points are assumed to be equidistant in t - and τ -direction with distance $\delta_h > 0$. A practical strategy for solving Equations (2) – (4) with initial conditions (5) – (8) is given by the following algorithm:

1. Calculate $i(t_m, \tau_n) = i_0(\tau_n - t_m) \Gamma(\tau_n)$ for all $n \geq m$. These are the incidence densities at the grid points located on and above the diagonal of the grid (on

and above the dashed line in Figure 1).

2. Given that $i(t_m, \tau_n)$ have been calculated on and above the diagonal, set $\ell := 0$ and calculate $\lambda(t_{\ell+1})$ and $S(t_{\ell+1})$ to determine $i(t_{\ell+1}, 0)$.
3. Calculate $i(t_{\ell+1+k}, \tau_k)$, $k = 1, 2, \dots$. The grid points $(t_{\ell+1+k}, \tau_k)$ are the points on a subdiagonal. We have $i(t_{\ell+1+k}, \tau_k) = i(t_{\ell+1}, 0) \Gamma(\tau_k)$.
4. Set $\ell := \ell + 1$ and repeat steps 2 to 4 until the incidence density i has been calculated on all points (t_m, τ_n) $m = 0, \dots, M, n = 0, \dots, N$, on the grid.




Figure 1: Rectangular grid representing calendar time t (abscissa) and infection-age τ (ordinate). The grid point (t_m, τ_n) above the main diagonal (dashed line) is highlighted.

In case the transmission rate $\beta(t, \tau)$ only depends on calendar time t , i.e., $\beta(t, \tau) = \beta(t)$, the force of infection can be written as

$$\lambda(t) = \beta(t) I(t) = \frac{\mathcal{R}(t) I(t)}{S(t) \int_0^\infty \Gamma(\tau) d\tau}.$$

Then, System (2) – (4) becomes explicitly dependent on \mathcal{R} . This means for given \mathcal{R} , the system can be solved, for instance by the algorithm above, such that the resulting effective reproduction number equals the prescribed \mathcal{R} . This is advantageous in situations, when the effective reproduction number \mathcal{R} is known while the transmission rate β is not.

Approximation of the age-structured SIR model by the conventional SIR model

The situation described at the end of the last section, when the effective reproduction $\mathcal{R}(t)$ is known instead of the transmission rate $\beta(t, \tau)$, happens quite frequently. Note that there are a variety of methods for estimating \mathcal{R} from a time series of numbers of incident cases, see e.g. [Cor13, Fra07]. The question arises, under which conditions system (2) – (4) can be approximated by a simpler model that explicitly depends on \mathcal{R} .

A simpler model related to System (2) – (4) is the conventional SIR model without demography of the background host:

$$\frac{dS(t)}{dt} = -\lambda(t) S(t) \quad (11)$$

$$\frac{dI(t)}{dt} = \lambda(t) S(t) - r(t) I(t) \quad (12)$$

$$\frac{dR(t)}{dt} = r(t) I(t) \quad (13)$$

Using Leibniz' integral rule and Eq. (5), the temporal derivative $\frac{dI}{dt}$ of the number of infected from Eq. (1) can be expressed as

$$\begin{aligned} \frac{dI(t)}{dt} &= \frac{d}{dt} \int_0^\infty i(t, \tau) d\tau = \int_0^\infty \frac{\partial}{\partial t} i(t, \tau) d\tau \\ &= - \int_0^\infty \gamma(\tau) i(t, \tau) d\tau - \int_0^\infty \frac{\partial}{\partial \tau} i(t, \tau) d\tau \\ &= - \int_0^\infty \gamma(\tau) i(t, \tau) d\tau - i(t, \infty) + i(t, 0). \end{aligned} \quad (14)$$

As $i(t, \infty) = 0$ (see above) and $i(t, 0) = \lambda(t) S(t)$, Eq. (14) reads as

$$\frac{dI(t)}{dt} = - \int_0^\infty \gamma(\tau) i(t, \tau) d\tau + \lambda(t) S(t). \quad (15)$$

It is reasonable to assume that the integral in Eq. (15) has a finite upper bound $\omega < \infty$, because there are no infected people with infinite infection-age. As $i(t, \tau) \geq 0$, the Mean Value Theorem for Definite Integrals [Has17] guarantees existence of a $\tau^* = \tau^*(t) \in [0, \omega]$ such that

$$\frac{dI(t)}{dt} = -\gamma(\tau^*(t)) I(t) + \lambda(t) S(t). \quad (16)$$

So far, we could show that Eq. (3) from the age-structured SIR model can be approximated by Eq. (12) with $r(t) = \gamma(\tau^*(t))$. If we can furthermore show that

$$\lambda S = \mathcal{R} r I, \quad (17)$$

we can reformulate Eq. (12) with an explicit dependency on \mathcal{R} . Assumed Eq. (17) holds true, we find

$$\begin{aligned} \frac{dI(t)}{dt} &= -r(t) I(t) + \mathcal{R}(t) r(t) I(t) \\ &= r(t) (\mathcal{R}(t) - 1) I(t). \end{aligned} \quad (18)$$

With the usual smoothness assumptions, Eq. (18) has the unique solution

$$I(t) = I(0) \exp \left(\int_0^t [r(\sigma) (\mathcal{R}(\sigma) - 1)] d\sigma \right), \quad (19)$$

where $I(0) = \int_0^\infty i_0(\tau) d\tau$ (note that i_0 was assumed to be integrable).

Apart from their simplicity, Eqs. (18) and (19) allow the common interpretation of the effective reproduction number \mathcal{R} : the number $I(t)$ of infected increases if and only if $\mathcal{R}(t) > 1$.

We have to examine the conditions such that Eq. (17) holds true. As $i(t, \cdot)$ is non-negative, the Mean Value Theorem for Definite Integrals applied to the left hand side of Eq. (17) reads as

$$\begin{aligned} S(t) \lambda(t) &= S(t) \int_0^\infty \beta(t, \tau) i(t, \tau) d\tau \\ &= S(t) \beta(t, \tau^*(t)) I(t) \end{aligned} \quad (20)$$

for $\tau^*(t) \in [0, \omega]$.

On the right hand side of Eq. (17), we have

$$\mathcal{R}(t) I(t) r(t) = S(t) \beta(t, \tau^*(t)) I(t) r(t) \int_0^{\omega'} \Gamma(\tau) d\tau, \quad (21)$$

where we assumed that $\beta(t, \cdot) \Gamma$ has a compact support $[0, \omega']$ and $\tau^*(t) \in [0, \omega']$.

By comparing Eqs. (20) and (21), we see that $r(t) \int_0^{\omega'} \Gamma(\tau) d\tau = 1$ and $\beta(t, \tau^*(t)) = \beta(t, \tau^*(t))$ implies $\lambda S = \mathcal{R} r I$. Hence, for practical purposes if $r(t) = \gamma(\tau^*(t))$ is close to $(\int_0^{\omega'} \Gamma(\tau) d\tau)^{-1}$, we can expect that Eq. (19) is a reasonable approximation for the number $I(t)$ of infected in the age-structured SIR model.

Example

We calculate the incidence-density i on the grid $(t_m, \tau_n) = (m \times \delta_h, n \times \delta_h)$, $m = 0, \dots, M$, $n = 0, \dots, N$, starting with $(t_0, \tau_0) = (0, 0)$, ending with $(t_M, \tau_N) = (40, 30)$ and equidistant stepsize $\delta_h = \frac{1}{24}$ (in units *days*). The transmission rate $\beta(t, \tau)$ is assumed to be the product of two Gaussian functions:

$$\beta(t, \tau) = \nu_0 \times \left(\nu_1 + \nu_2 \exp(-[(t-12)/4]^2/2) \right) \times \left(\exp(-[(\tau-5)/1.5]^2/2) \right),$$

where $\nu_0 = 5 \times 10^{-6}$, $\nu_1 = 1$, $\nu_2 = 3$. The removal rate γ is assumed to be constant $\gamma(\tau) = \frac{1}{8}$. The initial distribution i_0 is assumed to be $i_0(\tau) = 20 \times \Gamma(\tau)$, where Γ is defined in Eq. (9) with $\gamma(\tau) = \frac{1}{8}$. Figure 2 shows the resulting incidence-density $i(t, \tau)$ calculated with the Algorithm of the previous section.

The resulting reproduction number \mathcal{R} as calculated by Eq. (10) is depicted in Figure 3. If we try to approximate I using \mathcal{R} directly by Eq. (19), we obtain the graph as presented in Figure 4. The black curve corresponds to the approximated I according to Eq. (19) with constant $r(t) = 0.1125$. For comparison, the exact I calculated by (1) is shown as blue curve. Although little effort has been spend to optimize the fit between the exact and approximate I , the approximation is reasonably well.




Figure 2: Incidence-density $i(t, \tau)$ in the numerical example.




Figure 3: Reproduction number \mathcal{R} (ordinate) over calendar time t in the example.




Figure 4: Number I of infectives (ordinate) over calendar time t in the example. The blue curve corresponds to the exact solution (Eq. (1)) while the black curve is the approximation via Eq. (19).

Contact:

`<firstname>.<lastname>@stat.uni-muenchen.de`

Ludwigstr. 33

D-80539 München

Germany

References

- [Che16] Chen Y, Zou S, Yang J (2016) Global Analysis of an SIR Epidemic Model with Infection Age and Saturated Incidence, *Nonlin Ana: Real World App* 30: 16-31.
- [Cor13] Cori A, Ferguson NM, Fraser C, Cauchemez S (2013) A new framework and software to estimate time-varying reproduction numbers during epidemics. *Amer J Epidem* 178(9), 1505-1512.
- [Fra07] Fraser C (2007) Estimating Individual and Household Reproduction Numbers in an Emerging Epidemic. *PLoS ONE* 2(8): e758.
- [Has17] Hass JR, Heil C, Weir MD (2017) *Thomas' Calculus*, 14th ed, Pearson, Harlow.
- [Ina17] Inaba H (2017) *Age-structured Population Dynamics in Demography and Epidemiology*, Springer, Singapore.
- [Ker27] Kermack WO, McKendrick AG (1927) Contributions to the Mathematical Theory of Epidemics, *Proc Royal Soc* 115A, 700-21.
- [Nis09] Nishiura H, Chowell G (2009) The Effective Reproduction Number as a Prelude to Statistical Estimation of Time-Dependent Epidemic Trends, in: Chowell G, Hyman JM, Bettencourt LMA, Castillo-Chavez C: *Mathematical and Statistical Estimation Approaches in Epidemiology*, Springer, Dordrecht.