

The geometric classification of nilpotent \mathcal{CD} -algebras ¹Ivan Kaygorodov^a & Mykola Khrypchenko^b^a CMCC, Universidade Federal do ABC, Santo André, Brazil^b Departamento de Matemática, Universidade Federal de Santa Catarina, Florianópolis, Brazil

E-mail addresses:

Ivan Kaygorodov (kaygorodov.ivan@gmail.com)

Mykola Khrypchenko (nshrypchenko@gmail.com)

Abstract: We give a geometric classification of complex 4-dimensional nilpotent \mathcal{CD} -algebras. The corresponding geometric variety has dimension 18 and decomposes into 2 irreducible components determined by the Zariski closures of a two-parameter family of algebras and a four-parameter family of algebras (see Theorem 2). In particular, there are no rigid 4-dimensional complex nilpotent \mathcal{CD} -algebras.

Keywords: Nilpotent algebra, Lie algebra, Jordan algebra, \mathcal{CD} -algebra, geometric classification, degeneration.

MSC2010: 17A30, 17D99, 17B30.

INTRODUCTION

There are many results related to the algebraic and geometric classification of low-dimensional algebras in the varieties of Jordan, Lie, Leibniz and Zinbiel algebras; for algebraic classifications see, for example, [3, 6–8, 18, 21, 26, 31]; for geometric classifications and descriptions of degenerations see, for example, [3–7, 11–14, 17–21, 23, 25–29, 31–33, 35, 37, 39, 40]. In the present paper, we give algebraic classification of nilpotent \mathcal{CD} -algebras. This is a new class of non-associative algebras introduced in [3]. The idea of the definition of a \mathcal{CD} -algebra comes from the following property of Jordan and Lie algebras: *the commutator of any pair of multiplication operators is a derivation*. This gives three identities of degree four, which reduce to only one identity of degree four in the commutative or anticommutative case. Commutative and anticommutative \mathcal{CD} -algebras are related to many interesting varieties of algebras. Thus, anticommutative \mathcal{CD} -algebras is a generalization of Lie algebras, containing the intersection of Malcev and Sagle algebras as a proper subvariety. Moreover, the following intersections of varieties coincide: Malcev and Sagle algebras; Malcev and anticommutative \mathcal{CD} -algebras; Sagle and anticommutative \mathcal{CD} -algebras. On the other hand, the variety of anticommutative \mathcal{CD} -algebras is a proper subvariety of the varieties of binary Lie algebras and almost Lie algebras [34]. The variety of anticommutative \mathcal{CD} -algebras coincides with the intersection of the varieties of binary Lie algebras and almost Lie algebras. Commutative \mathcal{CD} -algebras is a generalization of Jordan algebras, which is a generalization of associative commutative algebras. On the other hand, the variety of commutative \mathcal{CD} -algebras is the variety of almost-Jordan algebras (sometimes, called Lie triple algebras) [16, 36, 38] and the bigger variety of generalized almost-Jordan algebras [1, 2, 15]. The n -ary version of commutative \mathcal{CD} -algebras was introduced in a recent paper by Kaygorodov, Pozhidaev and Saraiva [30]. Commutative \mathcal{CD} -algebras are also related to assosymmetric algebras [9].

Motivation and contextualization. Given algebras \mathbf{A} and \mathbf{B} in the same variety, we write $\mathbf{A} \rightarrow \mathbf{B}$ and say that \mathbf{A} *degenerates* to \mathbf{B} , or that \mathbf{A} is a *deformation* of \mathbf{B} , if \mathbf{B} is in the Zariski closure of the orbit of \mathbf{A} (under the base-change action of the general linear group). The study of degenerations of algebras is very rich and closely related to deformation theory, in the sense of Gerstenhaber [10]. It offers an insightful geometric perspective on the subject and has been the object of a lot of research. In particular, there are many results concerning degenerations of algebras of small dimensions in a variety defined by a set of identities. One of the main problems of the *geometric classification* of a variety of algebras is a description of its irreducible components. In the case of finitely-many orbits (i.e., isomorphism classes), the irreducible components are determined by the rigid algebras — algebras whose orbit closure is an irreducible component of the variety under consideration. The algebraic classification of complex 4-dimensional nilpotent \mathcal{CD} -algebras was obtained in [22], and in the present paper we continue the study of the variety by giving its geometric classification.

¹ The authors thank Prof. Dr. Pasha Zusmanovich for discussions about \mathcal{CD} -algebras.

Main result. The main result of the paper is Theorem 2, in which we prove that the variety of complex 4-dimensional nilpotent $\mathcal{C}\mathcal{D}$ -algebras has 2 irreducible components of dimensions 15 and 18.

1. DEGENERATIONS OF ALGEBRAS

Given an n -dimensional vector space \mathbf{V} , the set $\text{Hom}(\mathbf{V} \otimes \mathbf{V}, \mathbf{V}) \cong \mathbf{V}^* \otimes \mathbf{V}^* \otimes \mathbf{V}$ is a vector space of dimension n^3 . This space inherits the structure of the affine variety \mathbb{C}^{n^3} . Indeed, let us fix a basis e_1, \dots, e_n of \mathbf{V} . Then any $\mu \in \text{Hom}(\mathbf{V} \otimes \mathbf{V}, \mathbf{V})$ is determined by n^3 structure constants $c_{i,j}^k \in \mathbb{C}$ such that $\mu(e_i \otimes e_j) = \sum_{k=1}^n c_{i,j}^k e_k$. A subset of $\text{Hom}(\mathbf{V} \otimes \mathbf{V}, \mathbf{V})$ is *Zariski-closed* if it can be defined by a set of polynomial equations in the variables $c_{i,j}^k$ ($1 \leq i, j, k \leq n$).

The general linear group $\text{GL}(\mathbf{V})$ acts by conjugation on the variety $\text{Hom}(\mathbf{V} \otimes \mathbf{V}, \mathbf{V})$ of all algebra structures on \mathbf{V} :

$$(g * \mu)(x \otimes y) = g\mu(g^{-1}x \otimes g^{-1}y),$$

for $x, y \in \mathbf{V}$, $\mu \in \text{Hom}(\mathbf{V} \otimes \mathbf{V}, \mathbf{V})$ and $g \in \text{GL}(\mathbf{V})$. Clearly, the $\text{GL}(\mathbf{V})$ -orbits correspond to the isomorphism classes of algebras structures on \mathbf{V} . Let T be a set of polynomial identities which is invariant under isomorphism. Then the subset $\mathbb{L}(T) \subset \text{Hom}(\mathbf{V} \otimes \mathbf{V}, \mathbf{V})$ of the algebra structures on \mathbf{V} which satisfy the identities in T is $\text{GL}(\mathbf{V})$ -invariant and Zariski-closed. It follows that $\mathbb{L}(T)$ decomposes into $\text{GL}(\mathbf{V})$ -orbits. The $\text{GL}(\mathbf{V})$ -orbit of $\mu \in \mathbb{L}(T)$ is denoted by $O(\mu)$ and its Zariski closure by $\overline{O(\mu)}$.

Let \mathbf{A} and \mathbf{B} be two n -dimensional algebras satisfying the identities from T and $\mu, \lambda \in \mathbb{L}(T)$ represent \mathbf{A} and \mathbf{B} respectively. We say that \mathbf{A} *degenerates* to \mathbf{B} and write $\mathbf{A} \rightarrow \mathbf{B}$ if $\lambda \in \overline{O(\mu)}$. Note that in this case we have $\overline{O(\lambda)} \subset \overline{O(\mu)}$. Hence, the definition of a degeneration does not depend on the choice of μ and λ . If $\mathbf{A} \rightarrow \mathbf{B}$ and $\mathbf{A} \not\cong \mathbf{B}$, then $\mathbf{A} \rightarrow \mathbf{B}$ is called a *proper degeneration*. We write $\mathbf{A} \not\rightarrow \mathbf{B}$ if $\lambda \notin \overline{O(\mu)}$ and call this a *non-degeneration*. Observe that the dimension of the subvariety $\overline{O(\mu)}$ equals $n^2 - \dim \text{Der}(\mathbf{A})$. Thus if $\mathbf{A} \rightarrow \mathbf{B}$ is a proper degeneration, then we must have $\dim \text{Der}(\mathbf{A}) > \dim \text{Der}(\mathbf{B})$.

Let \mathbf{A} be represented by $\mu \in \mathbb{L}(T)$. Then \mathbf{A} is *rigid* in $\mathbb{L}(T)$ if $O(\mu)$ is an open subset of $\mathbb{L}(T)$. Recall that a subset of a variety is called *irreducible* if it cannot be represented as a union of two non-trivial closed subsets. A maximal irreducible closed subset of a variety is called an *irreducible component*. It is well known that any affine variety can be represented as a finite union of its irreducible components in a unique way. The algebra \mathbf{A} is rigid in $\mathbb{L}(T)$ if and only if $\overline{O(\mu)}$ is an irreducible component of $\mathbb{L}(T)$.

In the present work we use the methods applied to Lie algebras in [13, 14]. To prove degenerations, we will construct families of matrices parametrized by t . Namely, let \mathbf{A} and \mathbf{B} be two algebras represented by the structures μ and λ from $\mathbb{L}(T)$, respectively. Let e_1, \dots, e_n be a basis of \mathbf{V} and $c_{i,j}^k$ ($1 \leq i, j, k \leq n$) be the structure constants of λ in this basis. If there exist $a_i^j(t) \in \mathbb{C}$ ($1 \leq i, j \leq n, t \in \mathbb{C}^*$) such that the elements $E_i^t = \sum_{j=1}^n a_i^j(t) e_j$ ($1 \leq i \leq n$) form a basis of \mathbf{V} for any $t \in \mathbb{C}^*$, and the structure constants $c_{i,j}^k(t)$ of μ in the basis E_1^t, \dots, E_n^t satisfy $\lim_{t \rightarrow 0} c_{i,j}^k(t) = c_{i,j}^k$, then $\mathbf{A} \rightarrow \mathbf{B}$. In this case E_1^t, \dots, E_n^t is called a *parametric basis* for $\mathbf{A} \rightarrow \mathbf{B}$.

To prove a non-degeneration $\mathbf{A} \not\rightarrow \mathbf{B}$ we will use the following lemma (see [13]).

Lemma 1. *Let \mathcal{B} be a Borel subgroup of $\text{GL}(\mathbf{V})$ and $\mathcal{R} \subset \mathbb{L}(T)$ be a \mathcal{B} -stable closed subset. If $\mathbf{A} \rightarrow \mathbf{B}$ and \mathbf{A} can be represented by $\mu \in \mathcal{R}$ then there is $\lambda \in \mathcal{R}$ that represents \mathbf{B} .*

In particular, it follows from Lemma 1 that $\mathbf{A} \not\rightarrow \mathbf{B}$, whenever $\dim(\mathbf{A}^2) < \dim(\mathbf{B}^2)$.

When the number of orbits under the action of $\text{GL}(\mathbf{V})$ on $\mathbb{L}(T)$ is finite, the graph of primary degenerations gives the whole picture. In particular, the description of rigid algebras and irreducible components can be easily obtained. Since the variety of 4-dimensional nilpotent $\mathcal{C}\mathcal{D}$ -algebras contains infinitely many non-isomorphic algebras, we have to fulfill some additional work. Let $\mathbf{A}(\alpha) := \{\mathbf{A}(\alpha)\}_{\alpha \in I}$ be a family of algebras and \mathbf{B} be another algebra. Suppose that, for $\alpha \in I$, $\mathbf{A}(\alpha)$ is represented by a structure $\mu(\alpha) \in \mathbb{L}(T)$ and \mathbf{B} is represented by a structure $\lambda \in \mathbb{L}(T)$. Then by $\mathbf{A}(\alpha) \rightarrow \mathbf{B}$ we mean $\lambda \in \overline{\cup\{O(\mu(\alpha))\}_{\alpha \in I}}$, and by $\mathbf{A}(\alpha) \not\rightarrow \mathbf{B}$ we mean $\lambda \notin \overline{\cup\{O(\mu(\alpha))\}_{\alpha \in I}}$.

Let $\mathbf{A}(\alpha)$, \mathbf{B} , $\mu(\alpha)$ ($\alpha \in I$) and λ be as above. To prove $\mathbf{A}(\alpha) \rightarrow \mathbf{B}$, it is enough to construct a family of pairs $(f(t), g(t))$ parametrized by $t \in \mathbb{C}^*$, where $f(t) \in I$ and $g(t) = \left(a_i^j(t)\right)_{i,j} \in \text{GL}(\mathbf{V})$. Namely, let e_1, \dots, e_n be a basis of \mathbf{V} and $c_{i,j}^k$ ($1 \leq i, j, k \leq n$) be the structure constants of λ in this basis. If we construct $a_i^j : \mathbb{C}^* \rightarrow \mathbb{C}$ ($1 \leq i, j \leq n$) and $f : \mathbb{C}^* \rightarrow I$ such that $E_i^t = \sum_{j=1}^n a_i^j(t) e_j$ ($1 \leq i \leq n$) form a basis of \mathbf{V} for any $t \in \mathbb{C}^*$, and the structure constants $c_{i,j}^k(t)$ of $\mu(f(t))$ in the

basis E_1^t, \dots, E_n^t satisfy $\lim_{t \rightarrow 0} c_{i,j}^k(t) = c_{i,j}^k$, then $\mathbf{A}(\ast) \rightarrow \mathbf{B}$. In this case, E_1^t, \dots, E_n^t and $f(t)$ are called a *parametric basis* and a *parametric index* for $\mathbf{A}(\ast) \rightarrow \mathbf{B}$, respectively. In the construction of degenerations of this sort, we will write $\mu(f(t)) \rightarrow \lambda$, emphasizing that we are proving the assertion $\mu(\ast) \rightarrow \lambda$ using the parametric index $f(t)$.

2. THE ALGEBRAIC CLASSIFICATION OF COMPLEX 4-DIMENSIONAL NILPOTENT \mathcal{CD} -ALGEBRAS

The algebraic classification of complex 4-dimensional nilpotent \mathcal{CD} -algebras was obtained in the following three steps.

- (1) Trivial non-anticommutative \mathcal{CD} -algebras were classified in [8, Theorem 2.1, Theorem 2.3 and Theorem 2.5], and the only trivial anticommutative \mathcal{CD} -algebra is $\mathcal{CD}_{03}^{4\ast}$ from [22];
- (2) Non-trivial non-Leibniz terminal extensions of the algebra $\mathcal{CD}_{04}^{3\ast}(\lambda)$ were classified in [24, 1.4.5. 1-dimensional central extensions of $\mathbf{T}_{04}^{3\ast}$]. All of them are \mathcal{CD} -algebras.
- (3) Non-trivial Leibniz terminal extensions of $\mathcal{CD}_{04}^{3\ast}(\lambda)$ were found in [28]. All of them are \mathcal{CD} -algebras.
- (4) The rest of algebras were found in [22] and are listed below:

\mathcal{CD}_{11}^4	: $e_1 e_1 = e_2$	$e_2 e_2 = e_3$		
\mathcal{CD}_{12}^4	: $e_1 e_1 = e_2$	$e_2 e_1 = e_3$	$e_2 e_2 = e_3$	
\mathcal{CD}_{13}^4	: $e_1 e_1 = e_2$	$e_2 e_1 = e_3$		
$\mathcal{CD}_{14}^4(\lambda)$: $e_1 e_1 = e_2$	$e_1 e_2 = e_3$	$e_2 e_1 = \lambda e_3$	
\mathcal{CD}_{15}^4	: $e_1 e_1 = e_2$	$e_2 e_1 = e_4$	$e_2 e_2 = e_3$	
\mathcal{CD}_{16}^4	: $e_1 e_1 = e_2$	$e_1 e_2 = e_4$	$e_2 e_1 = e_3$	
$\mathcal{CD}_{17}^4(\lambda)$: $e_1 e_1 = e_2$	$e_1 e_2 = e_4$	$e_2 e_1 = \lambda e_4$	$e_2 e_2 = e_3$
$\mathcal{CD}_{18}^4(\alpha)$: $e_1 e_1 = e_2$	$e_1 e_3 = e_4$	$e_2 e_1 = e_3$	$e_2 e_2 = \alpha e_4$ $e_3 e_1 = -2e_4$
\mathcal{CD}_{19}^4	: $e_1 e_1 = e_2$	$e_1 e_2 = e_4$	$e_1 e_3 = e_4$	$e_2 e_1 = -2e_4$
$\mathcal{CD}_{20}^4(\alpha)$: $e_1 e_1 = e_2$	$e_1 e_2 = e_3$	$e_1 e_3 = -e_4$	
	$e_2 e_1 = e_3 + e_4$	$e_2 e_2 = \alpha e_4$	$e_3 e_1 = -e_4$	
$\mathcal{CD}_{21}^4(\lambda)$: $e_1 e_1 = e_2$	$e_1 e_2 = e_3$	$e_1 e_3 = (\lambda - 2)e_4$	
$\lambda \neq 1$	$e_2 e_1 = \lambda e_3 + e_4$	$e_2 e_2 = -(\lambda + 1)^2 e_4$	$e_3 e_1 = (1 - 2\lambda)e_4$	
$\mathcal{CD}_{22}^4(\alpha, \lambda)$: $e_1 e_1 = e_2$	$e_1 e_2 = e_3$	$e_1 e_3 = (\lambda - 2)e_4$	
	$e_2 e_1 = \lambda e_3$	$e_2 e_2 = \alpha e_4$	$e_3 e_1 = (1 - 2\lambda)e_4$	
$\mathcal{CD}_{23}^4(\alpha)$: $e_1 e_1 = e_2$	$e_1 e_2 = e_4$	$e_1 e_3 = e_4$	$e_2 e_1 = e_4$
$\alpha \neq \frac{1}{2}$	$e_2 e_3 = \alpha e_4$	$e_3 e_1 = e_4$	$e_3 e_2 = (\alpha + 1)e_4$	
$\mathcal{CD}_{24}^4(\alpha, \beta)$: $e_1 e_1 = e_2$	$e_1 e_2 = e_4$	$e_1 e_3 = \alpha e_4$	$e_2 e_1 = e_4$
	$e_2 e_2 = e_4$	$e_3 e_1 = \alpha e_4$	$e_3 e_2 = e_4$	$e_3 e_3 = \beta e_4$
$\mathcal{CD}_{25}^4(\alpha)$: $e_1 e_1 = e_2$	$e_1 e_2 = \alpha e_4$	$e_1 e_3 = e_4$	$e_2 e_1 = (\alpha + 1)e_4$ $e_3 e_1 = e_4$
\mathcal{CD}_{26}^4	: $e_1 e_1 = e_2$	$e_1 e_2 = e_4$	$e_1 e_3 = e_4$	$e_2 e_1 = e_4$
	$e_2 e_3 = -\frac{1}{2}e_4$	$e_3 e_2 = \frac{1}{2}e_4$	$e_3 e_3 = e_4$	
$\mathcal{CD}_{27}^4(\alpha)$: $e_1 e_1 = e_2$	$e_1 e_2 = e_4$	$e_2 e_1 = e_4$	$e_2 e_3 = \alpha e_4$ $e_3 e_2 = (\alpha + 1)e_4$
$\mathcal{CD}_{28}^4(\alpha)$: $e_1 e_1 = e_2$	$e_1 e_2 = \alpha e_4$	$e_2 e_1 = (\alpha + 1)e_4$	$e_3 e_3 = e_4$
\mathcal{CD}_{29}^4	: $e_1 e_1 = e_2$	$e_1 e_2 = e_4$	$e_2 e_1 = e_4$	$e_3 e_3 = e_4$
\mathcal{CD}_{30}^4	: $e_1 e_1 = e_2$	$e_1 e_2 = e_4$	$e_2 e_1 = e_4$	$e_3 e_3 = e_4$
$\mathcal{CD}_{31}^4(\alpha)$: $e_1 e_1 = e_2$	$e_1 e_3 = \alpha e_4$	$e_2 e_1 = e_4$	$e_2 e_2 = e_4$
	$e_2 e_3 = e_4$	$e_3 e_1 = \alpha e_4$	$e_3 e_2 = e_4$	$e_3 e_3 = e_4$
\mathcal{CD}_{32}^4	: $e_1 e_1 = e_2$	$e_1 e_3 = e_4$	$e_2 e_3 = -\frac{1}{2}e_4$	
	$e_3 e_1 = e_4$	$e_3 e_2 = \frac{1}{2}e_4$	$e_3 e_3 = e_4$	
$\mathcal{CD}_{33}^4(\alpha)$: $e_1 e_1 = e_2$	$e_1 e_3 = e_4$	$e_2 e_3 = \alpha e_4$	$e_3 e_1 = e_4$ $e_3 e_2 = (\alpha + 1)e_4$
$\mathcal{CD}_{34}^4(\alpha)$: $e_1 e_1 = e_2$	$e_1 e_3 = e_4$	$e_2 e_2 = e_4$	
	$e_3 e_1 = e_4$	$e_3 e_2 = e_4$	$e_3 e_3 = \alpha e_4$	
\mathcal{CD}_{35}^4	: $e_1 e_1 = e_2$	$e_1 e_3 = e_4$	$e_2 e_1 = e_4$	$e_3 e_1 = e_4$ $e_2 e_2 = e_4$
$\mathcal{CD}_{36}^4(\alpha)$: $e_1 e_1 = e_2$	$e_1 e_3 = \alpha e_4$	$e_2 e_2 = e_4$	$e_3 e_1 = (\alpha + 1)e_4$
\mathcal{CD}_{37}^4	: $e_1 e_1 = e_2$	$e_2 e_1 = e_4$	$e_2 e_3 = e_4$	$e_3 e_2 = e_4$ $e_3 e_3 = e_4$
$\mathcal{CD}_{38}^4(\alpha)$: $e_1 e_1 = e_2$	$e_2 e_1 = e_4$	$e_2 e_2 = e_4$	
$\alpha \neq 1$	$e_2 e_3 = e_4$	$e_3 e_2 = e_4$	$e_3 e_3 = \alpha e_4$	
\mathcal{CD}_{39}^4	: $e_1 e_1 = e_2$	$e_2 e_3 = -\frac{1}{2}e_4$	$e_3 e_2 = \frac{1}{2}e_4$	$e_3 e_3 = e_4$
\mathcal{CD}_{40}^4	: $e_1 e_1 = e_2$	$e_2 e_1 = e_4$	$e_2 e_3 = e_4$	$e_3 e_2 = e_4$
	$e_2 e_3 = e_4$	$e_3 e_1 = 2e_4$	$e_3 e_2 = e_4$	
$\mathcal{CD}_{41}^4(\alpha)$: $e_1 e_1 = e_2$	$e_2 e_3 = \alpha e_4$	$e_3 e_2 = (\alpha + 1)e_4$	
\mathcal{CD}_{42}^4	: $e_1 e_1 = e_2$	$e_2 e_1 = e_4$	$e_2 e_2 = e_4$	$e_3 e_3 = e_4$
\mathcal{CD}_{43}^4	: $e_1 e_1 = e_2$	$e_2 e_2 = e_4$	$e_3 e_3 = e_4$	
\mathcal{CD}_{44}^4	: $e_1 e_1 = e_2$	$e_2 e_2 = e_4$	$e_3 e_1 = e_4$	$e_3 e_3 = e_4$
$\mathcal{CD}_{45}^4(\alpha)$: $e_1 e_1 = e_2$	$e_2 e_2 = e_4$	$e_3 e_2 = e_4$	$e_3 e_3 = \alpha e_4$
\mathcal{CD}_{46}^4	: $e_1 e_1 = e_2$	$e_1 e_2 = e_4$	$e_2 e_1 = e_4$	$e_3 e_3 = e_4$
\mathcal{CD}_{47}^4	: $e_1 e_1 = e_2$	$e_2 e_3 = e_4$	$e_3 e_2 = e_4$	
\mathcal{CD}_{48}^4	: $e_1 e_1 = e_2$	$e_1 e_2 = e_3 + e_4$	$e_1 e_3 = \frac{1}{2}e_4$	$e_2 e_1 = \frac{1}{2}e_4$
	$e_2 e_2 = e_3$	$e_2 e_3 = -2ie_4$	$e_3 e_1 = 2e_4$	$e_3 e_2 = -ie_4$
\mathcal{CD}_{49}^4	: $e_1 e_1 = e_3 + e_4$	$e_1 e_2 = \frac{1}{2}e_4$	$e_1 e_3 = -\frac{1}{2}e_4$	$e_2 e_1 = \frac{1}{2}e_4$
	$e_2 e_2 = e_3$	$e_2 e_3 = -\frac{1}{2}e_4$	$e_3 e_1 = \frac{1}{2}e_4$	$e_3 e_2 = \frac{1}{2}e_4$

\mathcal{D}_{41}^4	:	$e_1 e_1 = e_3 + e_4$ $e_2 e_2 = e_3$	$e_1 e_2 = -\frac{1}{2}e_4$ $e_2 e_3 = -2ie_4$	$e_1 e_3 = e_4$ $e_3 e_1 = 2e_4$	$e_2 e_1 = -\frac{1}{2}e_4$ $e_3 e_2 = -ie_4$
\mathcal{D}_{42}^4	:	$e_1 e_1 = e_3 + e_4$ $e_2 e_2 = e_3$	$e_1 e_2 = -\frac{1}{2}e_4$ $e_2 e_3 = -\frac{1}{2}e_4$	$e_1 e_3 = -\frac{1}{2}e_4$ $e_3 e_1 = \frac{1}{2}e_4$	$e_2 e_1 = -\frac{1}{2}e_4$ $e_3 e_2 = \frac{1}{2}e_4$
$\mathcal{D}_{43}^4(\alpha)$:	$e_1 e_1 = e_3 + e_4$ $e_2 e_1 = \alpha e_4$	$e_1 e_2 = \alpha e_4$ $e_2 e_2 = e_3$	$e_1 e_3 = -\frac{1}{2}e_4$ $e_3 e_1 = \frac{1}{2}e_4$	
$\mathcal{D}_{44}^4(\alpha, \beta, \gamma)$:	$e_1 e_1 = e_3 + \alpha e_4$ $e_2 e_2 = e_3$	$e_1 e_2 = \beta e_4$ $e_3 e_1 = e_4$	$e_2 e_1 = (\beta + \gamma)e_4$ $e_3 e_3 = e_4$	
\mathcal{D}_{45}^4	:	$e_1 e_1 = e_3 + 2ie_4$ $e_3 e_1 = e_4$	$e_1 e_2 = e_4$ $e_3 e_2 = ie_4$	$e_2 e_1 = e_4$ $e_3 e_3 = e_4$	$e_2 e_2 = e_3$
$\mathcal{D}_{46}^4(\alpha)$ $\alpha \neq 0$:	$e_1 e_1 = e_3 - 2i\alpha e_4$ $e_3 e_1 = e_4$	$e_1 e_2 = \alpha e_4$ $e_3 e_2 = ie_4$	$e_2 e_1 = \alpha e_4$ $e_3 e_3 = e_4$	$e_2 e_2 = e_3$
$\mathcal{D}_{47}^4(\alpha, \beta)$ $\beta \neq 0$:	$e_1 e_1 = e_3 + e_4$ $e_2 e_3 = \beta e_4$	$e_1 e_3 = \alpha e_4$ $e_3 e_1 = (\alpha + 1)e_4$	$e_2 e_2 = e_3$ $e_3 e_2 = \beta e_4$	
$\mathcal{D}_{48}^4(\alpha)$ $\alpha \neq 0$:	$e_1 e_1 = e_3 + \alpha e_4$ $e_3 e_1 = e_4$	$e_2 e_1 = i\alpha e_4$ $e_3 e_2 = ie_4$	$e_2 e_2 = e_3$ $e_3 e_3 = e_4$	
$\mathcal{D}_{49}^4(\alpha)$ $\alpha \neq 0$:	$e_1 e_1 = e_3 + \alpha e_4$ $e_3 e_1 = e_4$	$e_2 e_1 = -i\alpha e_4$ $e_3 e_2 = ie_4$	$e_2 e_2 = e_3$ $e_3 e_3 = e_4$	
$\mathcal{D}_{50}^4(\alpha)$:	$e_1 e_1 = e_3 + \alpha e_4$	$e_2 e_1 = e_4$	$e_2 e_2 = e_3$	$e_3 e_3 = e_4$
$\mathcal{D}_{51}^4(\alpha)$:	$e_1 e_1 = e_3 + \alpha e_4$ $e_3 e_2 = ie_4$	$e_2 e_2 = e_3$ $e_3 e_3 = e_4$	$e_3 e_1 = e_4$	
\mathcal{D}_{52}^4	:	$e_1 e_1 = e_3 + e_4$	$e_2 e_2 = e_3$	$e_3 e_3 = e_4$	
\mathcal{D}_{53}^4	:	$e_1 e_1 = e_3$ $e_2 e_2 = e_3$	$e_1 e_2 = -\frac{1}{2}e_4$ $e_2 e_3 = ie_4$	$e_1 e_3 = e_4$ $e_3 e_1 = e_4$	$e_2 e_1 = \frac{1}{2}e_4$ $e_3 e_2 = ie_4$
$\mathcal{D}_{54}^4(\alpha)$:	$e_1 e_1 = e_3$ $e_2 e_2 = e_3$	$e_1 e_2 = e_4$ $e_2 e_3 = -i(\alpha + 1)e_4$	$e_1 e_3 = \alpha e_4$ $e_3 e_1 = (\alpha + 1)e_4$	$e_2 e_1 = e_4$ $e_3 e_2 = -i\alpha e_4$
$\mathcal{D}_{55}^4(\alpha)$:	$e_1 e_1 = e_3$ $e_2 e_1 = e_4$	$e_1 e_2 = e_4$ $e_2 e_2 = e_3$	$e_1 e_3 = \alpha e_4$ $e_3 e_1 = (\alpha + 1)e_4$	
\mathcal{D}_{56}^4	:	$e_1 e_1 = e_3$ $e_3 e_1 = e_4$	$e_1 e_2 = e_4$ $e_3 e_2 = ie_4$	$e_2 e_1 = -e_4$ $e_3 e_3 = e_4$	$e_2 e_2 = e_3$
$\mathcal{D}_{57}^4(\alpha, \beta)$ $\beta \notin \{0, -2\alpha\}$:	$e_1 e_1 = e_3$ $e_3 e_1 = e_4$	$e_1 e_2 = \alpha e_4$ $e_3 e_2 = ie_4$	$e_2 e_1 = (\alpha + \beta)e_4$ $e_3 e_3 = e_4$	$e_2 e_2 = e_3$
\mathcal{D}_{58}^4	:	$e_1 e_1 = e_3$ $e_2 e_3 = ie_4$	$e_1 e_3 = e_4$ $e_3 e_1 = e_4$	$e_2 e_1 = e_4$ $e_3 e_2 = ie_4$	$e_2 e_2 = e_3$
$\mathcal{D}_{59}^4(\alpha, \beta)$ $\beta \neq 0$:	$e_1 e_1 = e_3$ $e_2 e_3 = \beta e_4$	$e_1 e_3 = \alpha e_4$ $e_3 e_1 = (\alpha + 1)e_4$	$e_2 e_2 = e_3$ $e_3 e_2 = \beta e_4$	
\mathcal{D}_{60}^4	:	$e_1 e_1 = e_3$ $e_2 e_3 = e_4$	$e_1 e_3 = ie_4$ $e_3 e_1 = (i + 1)e_4$	$e_2 e_2 = e_3$ $e_3 e_2 = (i + 1)e_4$	
$\mathcal{D}_{61}^4(\alpha)$:	$e_1 e_1 = e_3$ $e_2 e_3 = \alpha e_4$	$e_1 e_3 = -i\alpha e_4$ $e_3 e_1 = (1 - i\alpha)e_4$	$e_2 e_2 = e_3$ $e_3 e_2 = (\alpha + i)e_4$	
\mathcal{D}_{62}^4	:	$e_1 e_1 = e_3$ $e_2 e_3 = -2ie_4$	$e_1 e_3 = e_4$ $e_3 e_1 = 2e_4$	$e_2 e_2 = e_3$ $e_3 e_2 = -ie_4$	
\mathcal{D}_{63}^4	:	$e_1 e_1 = e_3$ $e_2 e_3 = -\frac{i}{2}e_4$	$e_1 e_3 = -\frac{1}{2}e_4$ $e_3 e_1 = \frac{1}{2}e_4$	$e_2 e_2 = e_3$ $e_3 e_2 = \frac{1}{2}e_4$	
$\mathcal{D}_{64}^4(\alpha)$:	$e_1 e_1 = e_3$	$e_1 e_3 = \alpha e_4$	$e_2 e_2 = e_3$	$e_3 e_1 = (\alpha + 1)e_4$
$\mathcal{D}_{65}^4(\alpha)$ $\alpha \neq 0$:	$e_1 e_1 = e_3$ $e_3 e_1 = (\alpha + 1)e_4$	$e_1 e_3 = \alpha e_4$ $e_3 e_2 = ie_4$	$e_2 e_2 = e_3$	
\mathcal{D}_{66}^4	:	$e_1 e_1 = e_3$ $e_2 e_3 = e_4$	$e_2 e_1 = e_4$ $e_3 e_2 = e_4$	$e_2 e_2 = e_3$	
\mathcal{D}_{67}^4	:	$e_1 e_1 = e_3$	$e_2 e_2 = e_3$	$e_3 e_3 = e_4$	
\mathcal{D}_{68}^4	:	$e_1 e_1 = e_3 + e_4$ $e_2 e_3 = e_4$	$e_1 e_3 = ie_4$ $e_3 e_1 = ie_4$	$e_2 e_2 = e_3$ $e_3 e_2 = e_4$	
\mathcal{D}_{69}^4	:	$e_1 e_1 = e_3$ $e_2 e_3 = e_4$	$e_1 e_3 = ie_4$ $e_3 e_1 = ie_4$	$e_2 e_2 = e_3$ $e_3 e_2 = e_4$	
\mathcal{D}_{70}^4	:	$e_1 e_1 = e_3$	$e_1 e_3 = e_4$	$e_2 e_2 = e_3$	$e_3 e_1 = e_4$
\mathcal{D}_{71}^4	:	$e_1 e_1 = e_4$	$e_1 e_2 = e_3$	$e_1 e_3 = e_4$	$e_2 e_1 = -e_3$ $e_3 e_1 = -e_4$
\mathcal{D}_{72}^4	:	$e_1 e_1 = e_4$ $e_2 e_1 = -e_3$	$e_1 e_2 = e_3$ $e_2 e_2 = e_4$	$e_1 e_3 = e_4$ $e_3 e_1 = -e_4$	
\mathcal{D}_{73}^4	:	$e_1 e_2 = e_3 + e_4$	$e_1 e_3 = e_4$	$e_2 e_1 = -e_3$	$e_3 e_1 = -e_4$
$\mathcal{D}_{74}^4(\alpha)$:	$e_1 e_2 = e_3$	$e_1 e_3 = (\alpha + 1)e_4$	$e_2 e_1 = -e_3$	$e_3 e_1 = -\alpha e_4$
$\mathcal{D}_{75}^4(\alpha)$:	$e_1 e_2 = e_3$	$e_1 e_3 = (\alpha + 1)e_4$	$e_2 e_1 = -e_3$	$e_2 e_2 = e_4$ $e_3 e_1 = -\alpha e_4$
\mathcal{D}_{76}^4	:	$e_1 e_2 = e_3$	$e_1 e_3 = e_4$	$e_2 e_1 = -e_3$	$e_2 e_2 = e_4$ $e_3 e_1 = -e_4$
\mathcal{D}_{77}^4	:	$e_1 e_2 = e_3$	$e_1 e_3 = e_4$	$e_2 e_1 = -e_3$	$e_2 e_3 = e_4$ $e_3 e_2 = -e_4$
\mathcal{D}_{78}^4	:	$e_1 e_2 = e_3$ $e_2 e_2 = e_4$	$e_1 e_3 = e_4$ $e_2 e_3 = e_4$	$e_2 e_1 = -e_3$ $e_3 e_2 = -e_4$	
$\mathcal{D}_{79}^4(\alpha)$:	$e_1 e_2 = e_3 + \alpha e_4$	$e_1 e_3 = e_4$	$e_2 e_1 = -e_3$	$e_2 e_3 = e_4$ $e_3 e_3 = e_4$
\mathcal{D}_{80}^4	:	$e_1 e_2 = e_3 + e_4$	$e_1 e_3 = e_4$	$e_2 e_1 = -e_3$	$e_3 e_3 = e_4$
\mathcal{D}_{81}^4	:	$e_1 e_2 = e_3 + e_4$	$e_2 e_1 = -e_3$	$e_3 e_3 = e_4$	
\mathcal{D}_{82}^4	:	$e_1 e_2 = e_3$	$e_1 e_3 = e_4$	$e_2 e_1 = -e_3$	$e_2 e_2 = e_4$ $e_3 e_3 = e_4$
$\mathcal{D}_{83}^4(\alpha \neq 0)$:	$e_1 e_2 = e_3$	$e_2 e_1 = -e_3$	$e_2 e_2 = \alpha e_4$	$e_2 e_3 = e_4$ $e_3 e_3 = e_4$
\mathcal{D}_{84}^4	:	$e_1 e_2 = e_3$	$e_2 e_1 = -e_3$	$e_2 e_2 = e_4$	$e_3 e_3 = e_4$
\mathcal{D}_{85}^4	:	$e_1 e_2 = e_3$	$e_2 e_1 = -e_3$	$e_3 e_3 = e_4$	
\mathcal{D}_{86}^4	:	$e_1 e_2 = e_3$	$e_2 e_1 = -e_3$	$e_2 e_3 = e_4$	$e_3 e_2 = -e_4$
$\mathcal{D}_{87}^4(\lambda)$:	$e_1 e_1 = \lambda e_3 + (2\Theta - 1)e_4$	$e_1 e_2 = e_4$	$e_1 e_3 = e_4$	$e_2 e_1 = e_3 - (1 - \Theta)^2 \lambda^{-1} e_4$

$\lambda \neq 0, \frac{1}{4}$	$e_2e_2 = e_3$	$e_2e_3 = \Theta\lambda^{-1}e_4$	$e_3e_3 = e_4$	
$\mathfrak{CD}_{88}^4(\lambda)$	$e_1e_1 = \lambda e_3 + (1-2\Theta)e_4$	$e_1e_2 = e_4$	$e_1e_3 = e_4$	$e_2e_1 = e_3 - \Theta^2\lambda^{-1}e_4$
$\lambda \neq 0, \frac{1}{4}$	$e_2e_2 = e_3$	$e_2e_3 = \Theta\lambda^{-1}e_4$	$e_3e_3 = e_4$	
$\mathfrak{CD}_{89}^4(\lambda)$	$e_1e_1 = \lambda e_3 + (2\Theta-1)e_4$	$e_1e_2 = e_4$	$e_1e_3 = e_4$	$e_2e_1 = e_3 - (1-\Theta)^2\lambda^{-1}e_4$
$\lambda \neq 0, \frac{1}{4}$	$e_2e_2 = e_3$	$e_2e_3 = (1-\Theta)\lambda^{-1}e_4$	$e_3e_3 = e_4$	
$\mathfrak{CD}_{90}^4(\lambda)$	$e_1e_1 = \lambda e_3 + (1-2\Theta)e_4$	$e_1e_2 = e_4$	$e_1e_3 = e_4$	$e_2e_1 = e_3 - \Theta^2\lambda^{-1}e_4$
$\lambda \neq 0, \frac{1}{4}$	$e_2e_2 = e_3$	$e_2e_3 = (1-\Theta)\lambda^{-1}e_4$	$e_3e_3 = e_4$	
$\mathfrak{CD}_{91}^4(\lambda, \alpha)$	$e_1e_1 = \lambda e_3 + (2\Theta-1)e_4$	$e_1e_2 = e_4$	$e_1e_3 = \alpha e_4$	
$\lambda \neq 0, \frac{1}{4}$	$e_2e_1 = e_3 - (1-\Theta)^2\lambda^{-1}e_4$	$e_2e_2 = e_3$	$e_3e_3 = e_4$	
$\mathfrak{CD}_{92}^4(\lambda, \alpha)$	$e_1e_1 = \lambda e_3 + (1-2\Theta)e_4$	$e_1e_2 = e_4$	$e_1e_3 = \alpha e_4$	
$\lambda \neq 0, \frac{1}{4}$	$e_2e_1 = e_3 - \Theta^2\lambda^{-1}e_4$	$e_2e_2 = e_3$	$e_3e_3 = e_4$	
$\mathfrak{CD}_{93}^4(\alpha)$	$e_1e_1 = e_4$	$e_1e_2 = e_4$	$e_1e_3 = \alpha e_4$	$e_2e_1 = e_3 + e_4$
	$e_2e_2 = e_3$	$e_2e_3 = \alpha e_4$	$e_3e_3 = e_4$	
$\mathfrak{CD}_{94}^4(\alpha, \beta)$	$e_1e_1 = e_4$	$e_1e_2 = e_4$	$e_1e_3 = \alpha e_4$	
$\alpha \neq 0$	$e_2e_1 = e_3 + \beta e_4$	$e_2e_2 = e_3$	$e_3e_3 = e_4$	
$\mathfrak{CD}_{95}^4(\alpha)$	$e_1e_1 = e_4$	$e_1e_2 = e_4$	$e_1e_3 = \alpha e_4$	$e_2e_1 = e_3$
	$e_2e_2 = e_3$	$e_2e_3 = \alpha e_4$	$e_3e_3 = e_4$	
$\mathfrak{CD}_{96}^4(\alpha)$	$e_1e_1 = e_4$	$e_1e_2 = e_4$	$e_2e_1 = e_3 + \alpha e_4$	
	$e_2e_2 = e_3$	$e_2e_3 = e_4$	$e_3e_3 = e_4$	
$\mathfrak{CD}_{97}^4(\lambda)$	$e_1e_1 = \lambda e_3$	$e_1e_2 = e_4$	$e_1e_3 = \Theta e_4$	$e_2e_1 = e_3 - e_4$
	$e_2e_2 = e_3$	$e_2e_3 = e_4$	$e_3e_3 = e_4$	
$\mathfrak{CD}_{98}^4(\lambda)$	$e_1e_1 = \lambda e_3$	$e_1e_2 = e_4$	$e_1e_3 = (1-\Theta)e_4$	$e_2e_1 = e_3 - e_4$
$\lambda \neq \frac{1}{4}$	$e_2e_2 = e_3$	$e_2e_3 = e_4$	$e_3e_3 = e_4$	
$\mathfrak{CD}_{99}^4(\alpha)$	$e_1e_2 = e_4$	$e_1e_3 = e_4$	$e_2e_1 = e_3 - e_4$	
$\alpha \neq 1$	$e_2e_2 = e_3$	$e_2e_3 = \alpha e_4$	$e_3e_3 = e_4$	
$\mathfrak{CD}_{100}^4(\alpha)$	$e_1e_1 = \frac{1}{4}e_3$	$e_1e_2 = e_4$	$e_1e_3 = \alpha e_4$	$e_2e_1 = e_3 - e_4$
$\alpha \notin \{0, \frac{1}{2}\}$	$e_2e_2 = e_3$	$e_2e_3 = 2\alpha e_4$	$e_3e_3 = e_4$	
$\mathfrak{CD}_{101}^4(\alpha, \beta)$	$e_1e_2 = e_4$	$e_1e_3 = \alpha e_4$	$e_2e_1 = e_3$	
	$e_2e_2 = e_3$	$e_2e_3 = \beta e_4$	$e_3e_3 = e_4$	
$\mathfrak{CD}_{102}^4(\lambda, \alpha)$	$e_1e_1 = \lambda e_3$	$e_1e_2 = e_4$	$e_2e_1 = e_3 - e_4$	
$\lambda \neq 0$	$e_2e_2 = e_3$	$e_2e_3 = \alpha e_4$	$e_3e_3 = e_4$	
\mathfrak{CD}_{103}^4	$e_1e_2 = e_4$	$e_2e_1 = e_3 - e_4$	$e_2e_2 = e_3$	$e_3e_3 = e_4$
\mathfrak{CD}_{104}^4	$e_1e_3 = e_4$	$e_2e_1 = e_3 + e_4$	$e_2e_2 = e_3$	$e_2e_3 = e_4$ $e_3e_3 = e_4$
$\mathfrak{CD}_{105}^4(\lambda, \alpha, \beta)$	$e_1e_1 = \lambda e_3$	$e_1e_3 = e_4$	$e_2e_1 = e_3 + \alpha e_4$	
$\lambda \neq 0, \alpha \neq 0$	$e_2e_2 = e_3$	$e_2e_3 = \beta e_4$	$e_3e_3 = e_4$	
$\mathfrak{CD}_{106}^4(\alpha)$	$e_1e_3 = e_4$	$e_2e_1 = e_3 + \alpha e_4$	$e_2e_2 = e_3$	$e_3e_3 = e_4$
$\mathfrak{CD}_{107}^4(\lambda)$	$e_1e_1 = \lambda e_3$	$e_1e_3 = \Theta e_4$	$e_2e_1 = e_3$	
	$e_2e_2 = e_3$	$e_2e_3 = e_4$	$e_3e_3 = e_4$	
$\mathfrak{CD}_{108}^4(\lambda)$	$e_1e_1 = \lambda e_3$	$e_1e_3 = (1-\Theta)e_4$	$e_2e_1 = e_3$	
$\lambda \notin \{0, \frac{1}{4}\}$	$e_2e_2 = e_3$	$e_2e_3 = e_4$	$e_3e_3 = e_4$	
$\mathfrak{CD}_{109}^4(\lambda, \alpha)$	$e_1e_1 = \lambda e_3$	$e_2e_1 = e_3 + e_4$	$e_2e_2 = e_3$	$e_2e_3 = \alpha e_4$ $e_3e_3 = e_4$
$\mathfrak{CD}_{110}^4(\lambda)$	$e_1e_1 = \lambda e_3$	$e_2e_1 = e_3$	$e_2e_2 = e_3$	$e_2e_3 = e_4$ $e_3e_3 = e_4$
$\mathfrak{CD}_{111}^4(\lambda)$	$e_1e_1 = \lambda e_3$	$e_2e_1 = e_3$	$e_2e_2 = e_3$	$e_3e_3 = e_4$
$\mathfrak{CD}_{112}^4(\lambda, \alpha, \beta, \gamma)$	$e_1e_1 = \lambda e_3 + e_4$	$e_1e_3 = \alpha e_4$	$e_2e_1 = e_3 + \beta e_4$	
	$e_2e_2 = e_3$	$e_2e_3 = \gamma e_4$	$e_3e_3 = e_4$	

3. THE GEOMETRIC CLASSIFICATION OF COMPLEX 4-DIMENSIONAL NILPOTENT \mathfrak{CD} -ALGEBRAS

Theorem 2. *The variety of complex 4-dimensional nilpotent terminal algebras has 2 irreducible components: one of dimension 18 determined by the family of algebras $\mathfrak{CD}_{112}^4(\lambda, \alpha, \beta, \gamma)$ and one of dimension 15 determined by the family of algebras $\mathfrak{CD}_{12}^4(\lambda, \alpha)$.*

Proof. Thanks to [27, 28] the algebras $\mathfrak{N}_2(\alpha)$ and $\mathfrak{N}_3(\alpha)$ determine the irreducible components in the variety of complex 4-dimensional 2-step nilpotent algebras (which are trivial \mathfrak{CD} -algebras in our terminology), and the orbit closure of $\mathbf{D}_{01}^4(\lambda, \alpha, \beta)$ contains all terminal (Leibniz and non-Leibniz) extensions of $\mathfrak{CD}_{04}^{3*}(\lambda)$ (see [24, 28]), where

$$\begin{aligned}
\mathbf{D}_{01}^4(\lambda, \alpha, \beta) &: e_1e_1 = \lambda e_3 + e_4 & e_1e_3 = \alpha e_4 & e_2e_1 = e_3 & e_2e_2 = e_3 & e_2e_3 = \beta e_4 & e_3e_1 = e_4 \\
\mathfrak{N}_2(\alpha) &: e_1e_1 = e_3 & e_1e_2 = e_4 & e_2e_1 = -\alpha e_3 & e_2e_2 = -e_4 & & \\
\mathfrak{N}_3(\alpha) &: e_1e_1 = e_4 & e_1e_2 = \alpha e_4 & e_2e_1 = -\alpha e_4 & e_2e_2 = e_4 & e_3e_3 = e_4 &
\end{aligned}$$

Each algebra of the family $\mathfrak{CD}_{112}^4(\lambda, \alpha, \beta, \gamma)$ has 2-dimensional square, while the square of $\mathfrak{CD}_{12}^4(\lambda, \alpha)$ is of dimension 3. Hence, $\mathfrak{CD}_{112}^4(\lambda, \alpha, \beta, \gamma) \not\rightarrow \mathfrak{CD}_{12}^4(\lambda, \alpha)$ by Lemma 1. We also have $\mathfrak{CD}_{12}^4(\lambda, \alpha) \not\rightarrow \mathfrak{CD}_{112}^4(\lambda, \alpha, \beta, \gamma)$, because $\bigcup \text{Orb}(\mathfrak{CD}_{112}^4(\lambda, \alpha, \beta, \gamma))$ has dimension 18, while $\bigcup \text{Orb}(\mathfrak{CD}_{12}^4(\lambda, \alpha))$ is of dimension 15. The rest of the algebras degenerate either from $\mathfrak{CD}_{12}^4(\lambda, \alpha)$, or from $\mathfrak{CD}_{112}^4(\lambda, \alpha, \beta, \gamma)$, as shown below. In some of the degenerations we use the short notation $\Theta = \frac{1+\sqrt{1-4\Lambda}}{2}$ and $\Psi = \frac{1-\sqrt{1-4\Lambda}}{2} = 1 - \Theta$.

$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{D}_{01}^4(\Lambda, \mathbf{A}, \mathbf{B})$ $\Xi = (t - \mathbf{A})^{-\frac{1}{2}}$	$\lambda = \Lambda$ $E_1^t = t\Xi(e_1 + \Xi e_3)$	$\alpha = (\mathbf{A} - 1)\Xi$ $E_2^t = t\Xi e_2$	$\beta = -\mathbf{B}\Xi^2$ $E_3^t = t^2\Xi^2 e_3$	$\gamma = \mathbf{B}\Xi$ $E_4^t = t^3\Xi^4 e_4$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{R}_2(\mathbf{A})$ $\Xi = (\mathbf{A}t(t-1)(t^2+3t+1))^{-1}$	$\lambda = t^{-1}$ $E_1^t = te_1 - t\Xi(t^2 - \mathbf{A}t - 1)e_2$ $E_2^t = e_2 + ((\mathbf{A} + 1)t + \mathbf{A} - 1)e_3 + \frac{\Xi^2}{t^2}(\mathbf{A}t^4 + 2(\mathbf{A} - 1)(t^3 + \frac{1}{t}) - 3(2\mathbf{A} + 1)t^2 + (\mathbf{A}^2 - 5\mathbf{A} + 10)t + 3(2\mathbf{A} - 1))e_4$ $E_3^t = -\mathbf{A}^{-1}e_3 - \Xi\mathbf{A}^{-1}(\mathbf{A}t^2 - 3t - 2\mathbf{A} + 3)e_4$	$\alpha = t^{-1}$ $E_2^t = t^2 e_2 - 3t^3\Xi e_3 + \Xi^2 e_4$ $E_4^t = e_3 - \Xi t^{-3} e_4$	$\beta = \mathbf{A}^2 - t^2$ $E_3^t = \mathbf{A}^2 t(e_3 - \mathbf{A}^2 e_4)$	$\gamma = t$ $E_4^t = \mathbf{A}^4 t^2 e_4$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{R}_3(\mathbf{A})$	$\lambda = 1$ $E_1^t = -\mathbf{A}^2 t(e_1 - e_2 + te_3)$	$\alpha = -t$ $E_2^t = \mathbf{A}t e_2$ $E_3^t = 3e_4$	$\beta = \mathbf{A}^2 - t^2$ $E_3^t = \mathbf{A}^2 t(e_3 - \mathbf{A}^2 e_4)$	$\gamma = t$ $E_4^t = \mathbf{A}^4 t^2 e_4$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{01}^4$	$\lambda = 2$ $E_1^t = te_1$	$\alpha = 3t^{-4}$ $E_2^t = t^2 e_2$	$E_3^t = 3e_4$	$E_4^t = e_3$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{02}^4$ $\Xi = (9t^4 + 1)^{-1}$	$\lambda = 2$ $E_1^t = t(e_1 - t\Xi)e_2$ $E_3^t = e_4$	$\alpha = t^{-4}$ $E_2^t = t^2 e_2 - 3t^3\Xi e_3 + \Xi^2 e_4$ $E_4^t = e_3 - \Xi t^{-3} e_4$		
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{03}^4$	$\lambda = -2$ $E_1^t = te_1 - \frac{1}{37^2}e_2$	$\alpha = 0$ $E_2^t = t^2 e_2 - \frac{1}{37^2}e_3$	$E_3^t = e_4$	$E_4^t = t^2 e_3 + \frac{4}{37}e_4$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{04}^4(\Lambda)$	$\lambda = \Lambda$ $E_1^t = te_1$	$\alpha = 0$ $E_2^t = t^2 e_2$	$E_3^t = e_4$	$E_4^t = e_3 - t^{-3} e_4$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{05}^4$	$\lambda = t^{-1}$ $E_1^t = te_1$	$\alpha = t^{-3}$ $E_2^t = t^2 e_2$	$E_3^t = t^2 e_4$	$E_4^t = te_3$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{06}^4$ $\Xi = (t-1)^{-1}$	$\lambda = t^{-1}$ $E_1^t = te_1 - e_2$ $E_3^t = t^2 e_3 + \Xi(3t^2 - 4)e_4$	$\alpha = -\Xi t^{-2}(t^3 + t^2 - t - 2)$ $E_2^t = t^2 e_2 - (t+1)e_3 - \Xi t^{-2}(t^3 + t^2 - t - 2)e_3$ $E_4^t = e_4$		
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{07}^4(\Lambda)$	$\lambda = \Lambda$ $E_1^t = te_1$	$\alpha = (\Lambda - 2)t^{-1}$ $E_2^t = t^2 e_2$	$E_3^t = t^3(\Lambda - 2)e_4$	$E_4^t = t^3 e_3$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{08}^4(\mathbf{A})$	$\lambda = t^{-1}$ $E_1^t = te_1$	$\alpha = \mathbf{A}(1 - 2t)t^{-2}$ $E_2^t = t^2 e_2$	$E_3^t = t^2 e_3$	$E_4^t = t^2(1 - 2t)e_4$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{09}^4$ $\Xi = ((t-1)(t+4))^{-1}$	$\lambda = t^{-1}$ $E_1^t = te_1 + \Xi(2t-1)(te_2 + \Xi(2t-1)^2(t+1)^{-1}e_3)$ $E_3^t = t^2 e_3 + \Xi t(2t-1)(t^2 + t - 3)e_4$	$\alpha = (2t-1)t^{-2}$	$E_2^t = t^2 e_2 + \Xi t(t+1)(2t-1)e_3$ $E_4^t = t^2(1-2t)e_4$	
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{10}^4(\mathbf{A})$ $\Xi = (t((t+2)^2 + \mathbf{A}))^{-1}$	$\lambda = 1 + t$ $E_1^t = e_1 + \Xi(t^3 - 1)(e_2 + \mathbf{A}\Xi(t+2)^{-1}(t^3 - 1)e_3)$ $E_3^t = e_3 + \Xi(1-t)((t+2)(2t^2 + 1) - \mathbf{A}(t+1))e_4$	$\alpha = \mathbf{A}$	$E_2^t = e_2 + \Xi(t+2)(t^3 - 1)e_3$ $E_4^t = (1-t)e_4$	
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{11}^4(\Lambda)$ $\Xi = \Lambda - 1$	$\lambda = \Lambda$ $E_1^t = t\Xi(e_1 + (t+1)e_2)$ $E_3^t = t^3\Xi^3(e_3 - (t+1)(t-3(\Lambda+1))e_4)$	$\alpha = t - (\Lambda+1)^2$ $E_2^t = t^2\Xi^2(e_2 - (t+1)((\Lambda+1)e_3 - (t+1)(t - (\Lambda+1)^2))e_4)$	$E_4^t = t^4\Xi^4 e_4$	
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{13}^4(\mathbf{A})$	$\Xi = \sqrt{(\mathbf{A} + \mathbf{A}^2)^3(t^3 + (2\mathbf{A} + 1)(t^2 + \mathbf{A}^2) + \mathbf{A}^4)}$ $\lambda = \frac{\mathbf{A}^2(1+\mathbf{A})^2}{t(\mathbf{A}+t)^2}$ $\beta = \frac{t(\mathbf{A}+t)(1+\mathbf{A}+t)}{\mathbf{A}^2 + 2\mathbf{A}^3 + \mathbf{A}^4 + t^2 + 2\mathbf{A}t^2 + t^3}$ $E_1^t = -\frac{\sqrt{t^5}(\mathbf{A}+t)^2}{\Xi} (e_1 - e_2) - \frac{t^3(\mathbf{A}(t+\mathbf{A})(1+\mathbf{A}))^2}{\Xi^2} e_3$ $E_2^t = \frac{t^4\mathbf{A}^2(t+\mathbf{A})^2(1+\mathbf{A})^2}{\Xi^2} \left(e_3 + \frac{t\mathbf{A}^3(\mathbf{A}+t)^2(1+\mathbf{A})^3}{\Xi^2} e_4 \right)$ $E_3^t = \frac{\mathbf{A}(1+\mathbf{A})\sqrt{t^5}(\mathbf{A}+t)}{\Xi} \left(e_2 - \sqrt{t}(\mathbf{A}+t)\mathbf{A}^2(1+\mathbf{A})e_3 \right)$	$\alpha = -\frac{\mathbf{A}(1+\mathbf{A})\sqrt{t}(\mathbf{A}+t)}{\Xi}$ $\gamma = -\frac{\mathbf{A}(1+\mathbf{A})\sqrt{t}(\mathbf{A}+t)}{\Xi}$	$E_4^t = -\frac{t^7(\mathbf{A}+t)^4}{\Xi^4} e_4$	
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{14}^4(\mathbf{A}, \mathbf{B})$	$\Xi = \sqrt{(\mathbf{B}-t)(\mathbf{B}^2 - \mathbf{B}t + (\mathbf{A}-1)(\mathbf{A}-t)t)}$ $\lambda = \frac{(\mathbf{B}-t)^2}{(\mathbf{A}-t)^2 t}$ $E_1^t = (\mathbf{A}-t)^2 \sqrt{t^3} \left(\frac{i}{(\mathbf{B}-t)\Xi} (e_1 - e_2) - \frac{\sqrt{t}}{\Xi^2} e_3 \right)$ $E_3^t = \frac{\sqrt{t^3}(t-\mathbf{A})}{\Xi} \left(ie_2 - \frac{(t-\mathbf{A})\sqrt{t^3}}{\Xi} e_3 \right)$	$\alpha = \frac{i(1-t)\sqrt{t}(t-\mathbf{A})}{\Xi}$	$\beta = \frac{(\mathbf{A}-1)(\mathbf{A}-t)(\mathbf{B}-t)t}{\Xi}$ $E_2^t = -\frac{(\mathbf{A}-t)^2 t^2}{\Xi^2} \left(e_3 + \frac{(\mathbf{A}-t)^2 t(\mathbf{B}-t)}{\Xi^2} e_4 \right)$ $E_4^t = \frac{(\mathbf{A}-t)^4 t^4}{\Xi^4} e_4$	$\gamma = \frac{i(1-t)\sqrt{t}(t-\mathbf{A})}{\Xi}$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{15}^4(\mathbf{A})$ $\Xi = i\sqrt{\mathbf{A} + \mathbf{A}^2 + t}$	$\lambda = \frac{1}{t}$ $E_1^t = -\frac{i^2}{\Xi^2} (e_1 - e_2) + \frac{(1+\mathbf{A})t^2}{\Xi^2} e_3$ $E_3^t = \frac{4\sqrt{t}(1+4t)}{\Xi}$	$\alpha = \frac{\mathbf{A}-t}{\mathbf{A}\Xi}$ $E_2^t = \frac{t^3}{\Xi^2} e_3$	$\beta = -\frac{t}{\Xi}$ $E_3^t = \frac{i^2}{\Xi^2} e_2 + \frac{t^3}{\mathbf{A}\Xi^2} e_3$ $E_4^t = \frac{4\sqrt{t}(1+4t)}{\Xi}$	$\gamma = -\frac{t}{\Xi}$ $E_4^t = \frac{t^5}{\Xi^4} e_4$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{16}^4$ $\Xi = i\sqrt{1 + 20t + 176t^2}$	$\lambda = 2 + \frac{1}{4t} + 4t$ $E_1^t = -\frac{8\sqrt{t^5}}{\Xi} (e_1 - e_2) + \frac{16t^3(1+4t)^2}{\Xi^2} e_3$ $E_3^t = \frac{4\sqrt{t^5}(1+4t)}{\Xi} \left(e_2 - \frac{2\sqrt{t}(1+4t)}{\Xi} e_3 \right)$	$\alpha = \frac{4\sqrt{t}(1+4t)}{\Xi}$	$\beta = \frac{4t(1+4t)^3}{\Xi^2}$ $E_2^t = \frac{16t^4(1+4t)^2}{\Xi^2} \left(e_3 - \frac{4t(1+4t)^3}{\Xi^2} e_4 \right)$ $E_4^t = \frac{256t^7(1+4t)^4}{\Xi^4} e_4$	$\gamma = \frac{4\sqrt{t}(1+4t)}{\Xi}$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{17}^4(\mathbf{A})$	$\lambda = \frac{(1+\mathbf{A})^2}{t}$ $E_1^t = -\frac{\sqrt{t^5}}{\sqrt{\mathbf{A}(1+\mathbf{A})^5}} (e_1 - e_2) + \frac{t^3}{\mathbf{A}(1+\mathbf{A})^3} e_3$ $E_3^t = \frac{\sqrt{t^5}}{\sqrt{\mathbf{A}(1+\mathbf{A})^3}} e_2 + \frac{t^3}{(1+\mathbf{A})^3} e_3$	$\alpha = \frac{\sqrt{t}}{\sqrt{\mathbf{A}(1+\mathbf{A})^3}}$	$\beta = \frac{t}{\mathbf{A} + \mathbf{A}^2}$ $E_2^t = \frac{t^4}{\mathbf{A}(1+\mathbf{A})^3} \left(e_3 + \frac{t}{(1+\mathbf{A})^2} e_4 \right)$ $E_4^t = \frac{t^7}{\mathbf{A}^2(1+\mathbf{A})^6} e_4$	$\gamma = \frac{\sqrt{t}}{\sqrt{\mathbf{A}(1+\mathbf{A})^3}}$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{18}^4(\mathbf{A})$ $\Xi = \sqrt{1 + \mathbf{A} + \mathbf{A}^2}$	$\lambda = -\frac{1}{\Xi^2 t}$ $E_1^t = -i\Xi^2 \Xi(e_1 - e_2) + (1 + \mathbf{A})t^2 e_3$ $E_3^t = t^2 \left(e_2 - \frac{i\Xi}{\mathbf{A}} e_3 \right)$	$\alpha = \frac{i(t\Xi^2 - \mathbf{A})}{\mathbf{A}\Xi}$	$\beta = 0$ $E_2^t = t^3(e_3 - te_4)$ $E_4^t = t^5 e_4$	$\gamma = \frac{i\Xi}{\mathbf{A}}$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{19}^4$ $\Xi = \sqrt{2-t}$	$\lambda = \frac{1}{t}$ $E_1^t = -\frac{t^2}{\Xi} e_1 + \frac{t^2}{\Xi} e_2 + \frac{t^2}{\Xi^2} e_3$	$\alpha = \frac{t}{\Xi}$ $E_2^t = \frac{t^3}{\Xi^2} e_3 - \frac{t^4}{\Xi^4} e_4$	$\beta = \frac{t}{\Xi^2}$ $E_3^t = -\frac{t^2}{\Xi} e_2 + \frac{t^3}{\Xi^2} e_3$	$\gamma = \frac{t}{\Xi}$ $E_4^t = \frac{t^5}{\Xi^4} e_4$

$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{20}^4$	$\lambda = \frac{t-1+it\sqrt{1+t}}{t^2+t}$ $E_1^t = t^2(i\sqrt{1+t}e_1 + e_2 + e_3)$ $E_2^t = t^3e_2$	$\alpha = 0$ $E_3^t = t^3e_2$	$\beta = -\frac{it}{\sqrt{1+t}}$ $E_4^t = t^5e_4$	$\gamma = 0$ $E_4^t = t^5e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{21}^4(\mathbf{A})$	$\Xi = 1 - 2t + 2t^2 - 2t^3 + 3t^5 - 2t^6 + t^7 + \mathbf{A}^2(1-t+t^2) + \mathbf{A}(-2+3t-3t^2+2t^3-2t^4)$ $\lambda = -\frac{t(1-2t+3t^2-2t^3+t^5-t^6+t^7+\mathbf{A}^2(1-t^3+t^4)+\mathbf{A}(-2+2t-2t^2+t^3+t^4-2t^5))}{(t-1)(\mathbf{A}-1+t-t^2)^2}$ $\alpha = t^2$ $E_1^t = t(t-1)^2 \left(\frac{1-\mathbf{A}-t+t^2}{\Xi} e_1 + \frac{(t-\mathbf{A})t^2}{\Xi} e_2 - \frac{(t-1)t^2}{\Xi} e_3 \right)$ $E_2^t = \frac{(1-t)^3t^2}{\Xi} (te_3 + (1-t)e_4)$	$\beta = \frac{(1-t)(1-\mathbf{A}-t+2t^2)}{t(1-\mathbf{A}-t+t^2)}$ $\gamma = 1$	$E_3^t = \frac{(t-1)^6t^6}{\Xi^2} e_4$ $E_4^t = \frac{(1-t)^3t^3}{\Xi} (te_3 + e_4)$	
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{22}^4$	$\Xi = \sqrt{t^3(-16-64t-64t^4+17t^5+72t^6+16t^7)}$ $\alpha = \frac{4t^3(-4-t+4t^2)}{t^4+4t^5-\Xi}$ $E_1^t = \frac{\Xi(\Xi+t^2)}{8} e_1 + t^4e_2$	$\beta = \frac{32t^3-9t^4-40t^5-16t^6+(1+4t)\Xi}{4t^4+16t^5-4\Xi}$ $E_2^t = t^3e_3 - \frac{t^3(1+4t)}{4} e_4$	$\gamma = 1$ $E_3^t = t^2e_2 - \frac{t^2}{2}e_3$ $E_4^t = t^5e_4$	
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{23}^4(\mathbf{A})$ $\Xi = \mathbf{A} + \mathbf{A}^2$	$\lambda = \frac{\Xi(\Xi+t^2)}{t(\Xi+t)^2}$ $\beta = \frac{t(\Xi+t)}{\Xi+t^2}$ $E_1^t = \frac{\sqrt{t^5(\Xi+t)}}{\Xi^2\sqrt{\Xi+t^2}} \left((\Xi+t)e_1 - te_2 + \frac{\sqrt{t^3(\Xi+t)}}{(1+\mathbf{A})\sqrt{\Xi+t^2}} e_3 \right)$ $E_3^t = \frac{\sqrt{t^5(\Xi+t)}}{\Xi\sqrt{\Xi+t^2}} \left(e_2 + \frac{\sqrt{t(\Xi+t)}}{(1+\mathbf{A})\sqrt{\Xi+t^2}} e_3 \right)$	$\alpha = \frac{(1-\mathbf{A})\sqrt{t^3}}{\Xi\sqrt{\Xi+t^2}}$ $\gamma = \frac{\sqrt{t(\Xi+t)}}{\Xi\sqrt{\Xi+t^2}}$ $E_2^t = \frac{t^4(\Xi+t)^2}{\Xi^2(\Xi+t^2)} \left(e_3 + \frac{t(\Xi+t)^2}{\Xi(\Xi+t^2)} e_4 \right)$ $E_4^t = \frac{t^7(\Xi+t)^4}{\Xi^4(\Xi+t^2)^2} e_4$		
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{24}^4(\mathbf{A})$	$\lambda = \frac{\mathbf{A}^2-t-\mathbf{A}t^2}{\mathbf{A}^2t^3}$ $\beta = \frac{t^3(2+2\mathbf{A}^2+(3+\sqrt{1-4\mathbf{A}})\mathbf{A}t)}{2\mathbf{A}(\mathbf{A}+t)}$ $E_1^t = -\frac{i\sqrt{t^5}}{\sqrt{\mathbf{A}+t}} \left(te_1 + \frac{1}{\mathbf{A}}e_2 - \frac{i\sqrt{t^4}}{\sqrt{\mathbf{A}+t}} e_3 \right)$ $E_3^t = \frac{\sqrt{t^5}}{\sqrt{\mathbf{A}+t}} \left(-ie_2 + \frac{(1-\sqrt{1-4\mathbf{A}})\sqrt{t^3}}{2\sqrt{\mathbf{A}+t}} e_3 \right)$	$\alpha = \frac{\sqrt{t(-2\mathbf{A}+t+\sqrt{1-4\mathbf{A}}t-2\mathbf{A}^2t^2)}}{2i\mathbf{A}^2\sqrt{\mathbf{A}+t}}$ $\gamma = \frac{\sqrt{t^3}}{i\sqrt{\mathbf{A}+t}}$ $E_2^t = -\frac{t^4}{\mathbf{A}+t} (e_3 + t^3e_4)$ $E_4^t = \frac{t^8}{(\mathbf{A}+t)^2} e_4$		
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{25}^4$	$\lambda = \frac{t-t^4-1}{t^8}$ $E_1^t = i\sqrt{\frac{t^4}{2}} \left(t^4e_1 + e_2 + i\sqrt{\frac{t^9}{2}} e_3 \right)$ $E_3^t = \sqrt{\frac{t^{11}}{2}} \left(ie_2 - \sqrt{\frac{t^7}{2}} e_3 \right)$	$\alpha = \frac{i(t-2)}{\sqrt{2t}}$ $\beta = \frac{t^4(t^3-1)}{2}$ $\gamma = -i\sqrt{2}t^7$	$E_2^t = -\frac{t^8}{2} \left(e_3 + \frac{t^7}{2} e_4 \right)$ $E_4^t = \frac{t^{16}}{4} e_4$	
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{26}^4(\mathbf{A})$	$E_1^t = t^{-1}(e_1 - e_3)$ $E_2^t = t^{-2}e_3$	$\alpha = 0$ $E_3^t = t^{-2}e_3$	$\beta = \mathbf{A}t^{-3}$ $E_4^t = t^{-4}e_4$	$\gamma = t^{-3}$ $E_4^t = t^{-4}e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{27}^4$ $\Xi = \sqrt{1-t}$	$\lambda = -2 + t^{-4}$ $E_1^t = \frac{\sqrt{t^{11}}}{\Xi} (e_1 + e_2) + \frac{t^3}{\Xi^2} e_3$	$\alpha = -\frac{\sqrt{t}}{\Xi}$ $E_2^t = \frac{t^7}{\Xi^2} (e_3 + t^4e_4)$	$\beta = \frac{2t^4-2t^5-\sqrt{t^5}}{\Xi^2}$ $E_3^t = \frac{t^4}{\Xi} \left(e_2 + \frac{t^2}{\Xi} e_3 \right)$	$\gamma = 0$ $E_4^t = \frac{t^{13}}{\Xi^4} e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{28}^4(\mathbf{A})$ $\Xi = \sqrt{1-\mathbf{A}}$	$\lambda = -2 + t^{-2}$ $E_1^t = \frac{\sqrt{t^5}}{\Xi} (e_1 + e_2) + \frac{t^3}{\Xi^2} e_3$	$\alpha = \frac{-\sqrt{t^3}-\mathbf{A}t}{\Xi}$ $E_2^t = \frac{t^3}{\Xi^2} (e_3 + t^2e_4)$	$\beta = -\frac{\sqrt{t^3}}{\Xi^2} + 2t^2$ $E_3^t = \frac{t^2}{\Xi} \left(e_2 + \frac{t}{\Xi} e_3 \right)$	$\gamma = 0$ $E_4^t = \frac{t^6}{\Xi^4} e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{29}^4$ $\Xi = \frac{t^5}{\sqrt{1+4t^2}}$	$\lambda = -t^{-5}$ $E_1^t = 2t^8\Xi(-e_1 + e_2 + 4\Xi e_3)$	$\alpha = 2\Xi$ $E_2^t = -4t^6\Xi^2(e_3 - t^{10}e_4)$	$\beta = -4\Xi^2$ $E_3^t = 2t^4\Xi(e_2 + \Xi e_3)$	$\gamma = -2\Xi$ $E_4^t = 16t^{10}\Xi^4 e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{30}^4$ $\Xi = \sqrt{2t^5-t^3-1}$	$\lambda = -2 + t^{-5}$ $E_1^t = \frac{i\sqrt{t^{13}}}{\Xi} (e_1 + e_2) - \frac{t^7}{\Xi^2} e_3$	$\alpha = -\frac{i\sqrt{t}}{\Xi}$ $E_2^t = -\frac{t^8}{\Xi^2} \left(e_2 - \frac{t^5}{\Xi^2} e_3 \right)$	$\beta = \frac{t^3-2t^5}{\Xi^2}$ $E_3^t = \frac{\sqrt{t^9}}{\Xi} \left(ie_2 - \frac{t^5}{\Xi} e_3 \right)$	$\gamma = 0$ $E_4^t = \frac{t^{15}}{\Xi^4} e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{31}^4$ $\Xi = \sqrt{2t^2-1}$	$\lambda = -2 + t^{-2}$ $E_1^t = \frac{i\sqrt{t^7}}{\Xi} (e_1 + e_2)$	$\alpha = \frac{it(1-\sqrt{t^3})}{\Xi}$ $E_2^t = -\frac{t^5}{\Xi^2} \left(e_3 - \frac{t^2}{\Xi^2} e_4 \right)$	$\beta = -\frac{2t^2+\sqrt{t^5}-\sqrt{t^7}}{\Xi^2}$ $E_3^t = \frac{t^3}{\Xi} \left(ie_2 - \frac{t}{\Xi} e_3 \right)$	$\gamma = 0$ $E_4^t = \frac{t^9}{\Xi^4} e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{32}^4(\mathbf{A})$ $\Xi = \sqrt{(1+\mathbf{A})(2\mathbf{A}t^5-t^5-\mathbf{A})}$	$\lambda = -2 + t^{-5}$ $E_1^t = \frac{i\sqrt{t^{13}}}{\Xi} (e_1 + e_2) - \frac{t^9}{\Xi^2} e_3$ $E_3^t = \frac{i\sqrt{t^9}}{\Xi} \left(ie_2 - \frac{\mathbf{A}\sqrt{t^5}}{\Xi} e_3 \right)$	$\alpha = -\frac{2i\sqrt{t^5}}{\Xi}$ $E_2^t = -\frac{t^8}{\Xi^2} \left(e_3 - \frac{\mathbf{A}(1+\mathbf{A})t^5}{\Xi^2} e_4 \right)$ $E_4^t = \frac{t^{15}}{\Xi^4} e_4$	$\beta = -\frac{(1-2\mathbf{A})t^5}{\mathbf{A}+t^5-2\mathbf{A}t^5}$ $\gamma = \frac{i\sqrt{t^5}}{\Xi}$	
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{33}^4$	$\lambda = -2 + t^{-2}$ $E_1^t = -i\sqrt{t^5}(e_1 + e_2) - t^3e_3$	$\alpha = -i(t^3 - \sqrt{t})$ $E_2^t = -t^3(e_3 + t^2e_4)$	$\beta = 2t^2$ $E_3^t = -t^2(ie_2 + t^3e_3)$	$\gamma = it^3$ $E_4^t = t^6e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{34}^4$ $\Xi = \sqrt{2t^2-1}$	$\lambda = -2 + t^{-2}$ $E_1^t = \frac{\sqrt{t^5}}{\Xi} (e_1 + e_2) + \frac{t^4}{\Xi^2} e_3$	$\alpha = \frac{-\sqrt{t^3}+t^2}{\Xi}$ $E_2^t = \frac{t^3}{\Xi^2} \left(e_3 - \frac{t^2}{\Xi^2} e_4 \right)$	$\beta = \frac{2t^2}{\Xi}$ $E_3^t = \frac{t^2}{\Xi} \left(e_2 + \frac{t}{\Xi} e_3 \right)$	$\gamma = -\frac{t^2}{\Xi}$ $E_4^t = \frac{t^6}{(1+t)\Xi^4} e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{35}^4$	$\lambda = t^{-1}$ $E_1^t = i\sqrt{t^3}e_1$	$\alpha = 0$ $E_2^t = -t^2(e_3 + te_4)$	$\beta = 0$ $E_3^t = \sqrt{t^3}(ie_2 - \sqrt{t^3}e_3)$	$\gamma = -i\sqrt{t^3}$ $E_4^t = t^4e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{36}^4(\mathbf{A})$ $\Xi = \sqrt{\mathbf{A}(-4+(4+\mathbf{A})t^3)}$	$\lambda = -\frac{1}{\mathbf{A}}$ $E_1^t = \frac{\mathbf{A}t^2+\sqrt{t\Xi}}{2} e_1 + t^2e_2$	$\alpha = -\frac{2\sqrt{t^3}}{\mathbf{A}\sqrt{t^3+\Xi}}$ $E_2^t = t(e_3 - \mathbf{A}e_4)$	$\beta = -\frac{\mathbf{A}((2+\mathbf{A})\sqrt{t^3+\Xi})}{\mathbf{A}\sqrt{t^3+\Xi}}$ $E_3^t = te_2 + \frac{(\sqrt{1-4\mathbf{A}}-1)\mathbf{A}t}{2} e_4$	$\gamma = 1$ $E_4^t = t^2e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{37}^4$	$\lambda = t^{-1}$ $E_1^t = -\frac{it^2}{\sqrt{2}} (e_1 - e_2 - \frac{i}{\sqrt{2}}e_3)$	$\alpha = \frac{it}{\sqrt{2}}$ $E_2^t = -\frac{t^3}{2} (e_3 + \frac{t}{2}e_4)$	$\beta = 0$ $E_3^t = \frac{t^2}{\sqrt{2}} (ie_2 + \frac{t}{\sqrt{2}}e_3)$	$\gamma = \frac{it}{\sqrt{2}}$ $E_4^t = \frac{t^5}{4} e_4$

$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{38}^4$	$\lambda = 1$ $E_1^t = \sqrt{t^3}e_1$	$\alpha = 0$ $E_2^t = t^3(e_3 + e_4)$	$\beta = 1$ $E_3^t = t^2(e_2 + e_3)$	$\gamma = 0$ $E_4^t = t^5e_4$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{39}^4$ $\Xi = t - 1$	$E_1^t = \Xi(e_1 + \frac{1}{\sqrt{\Xi}}e_2 + e_3)$ $E_3^t = \Xi(te_3 + \frac{1}{\sqrt{\Xi}}(t+3)e_4)$	$\alpha = \frac{1}{2t^2}(2t^3 + 3t^2 + 24t - 36)$ $E_2^t = i\Xi(e_1 - \sqrt{\Xi}e_2 - (\frac{t}{2} + 2)e_3 - \frac{3}{2t}(t+3)e_4)$ $E_4^t = t\Xi^2e_4$	$\beta = \frac{9}{t^2}\sqrt{\Xi}$ $\beta = 0$	$\gamma = -\frac{\sqrt{\Xi}}{2t}(t+6)$ $\gamma = \sqrt{\Xi}$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{40}^4$ $\Xi = t - 1$	$E_1^t = \Xi(e_1 + \frac{1}{\sqrt{\Xi}}e_2 - \frac{1}{2}e_3)$ $E_3^t = t\Xi e_3$	$\alpha = \frac{1}{2t^2}(2t^3 + 13t^2 + 8t - 36)$ $E_2^t = i\Xi(e_1 - \sqrt{\Xi}e_2 - (\frac{3}{2}t + 2)e_3 - \frac{3}{2t}(7t+9)e_4)$ $E_4^t = t\Xi^2e_4$	$\beta = \frac{1}{t^2}(4t+9)\sqrt{\Xi}$ $\beta = 0$	$\gamma = -\frac{3\sqrt{\Xi}}{2t}(t+2)$ $\gamma = 0$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{41}^4$ $\Xi = t - 1$	$E_1^t = \Xi(e_1 + \frac{1}{\sqrt{\Xi}}e_2 + e_3)$ $E_3^t = \Xi(te_3 + \frac{1}{\sqrt{\Xi}}(7t+9)e_4)$	$\alpha = \frac{1}{4t}(4t^2 - 19t + 16)$ $E_2^t = i\Xi(e_1 - \sqrt{\Xi}e_2 + (3t - \frac{1}{2})e_3 - 6\Xi e_4)$ $E_4^t = t\Xi^2e_4$	$\beta = -\frac{2}{t}\sqrt{\Xi}$ $\beta = 0$	$\gamma = 3\sqrt{\Xi}$ $\gamma = 0$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{42}^4$ $\Xi = t - 1$	$E_1^t = \Xi(e_1 + \frac{1}{\sqrt{\Xi}}e_2 - \frac{1}{2}e_3)$ $E_3^t = \Xi(te_3 - 2\Xi e_4)$	$\alpha = \frac{1}{4t}(4t^2 - 19t + 16)$ $E_2^t = i\Xi(e_1 - \sqrt{\Xi}e_2 + (3t - \frac{1}{2})e_3 - 6\Xi e_4)$ $E_4^t = t\Xi^2e_4$	$\beta = -\frac{2}{t}\sqrt{\Xi}$ $\beta = 0$	$\gamma = 3\sqrt{\Xi}$ $\gamma = 0$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{43}^4(\mathbf{A})$ $\Xi = t\sqrt{1-t^2}$	$E_1^t = \frac{\Xi^2}{t}(e_1 + \frac{t^2}{\Xi}e_2 - \frac{1}{2}e_3)$ $E_2^t = \Xi e_2$	$\alpha = \frac{1}{4} + t - 2\mathbf{A}t^2$ $E_3^t = \Xi e_3$	$\beta = \mathbf{A}t\sqrt{1-t^2}$ $E_4^t = \frac{\Xi^4}{t}e_4$	$\gamma = 0$ $\gamma = 0$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{44}^4(\mathbf{A}, \mathbf{B}, \mathbf{C})$ $\Xi = \frac{1}{\sqrt{\rho^2\mathbf{B}(t+2)+\rho\mathbf{A}(t+1)-\mathbf{B}t}}$ $\rho = -\frac{\mathbf{A}+\sqrt{\mathbf{A}^2+4\mathbf{B}(\mathbf{B}+\mathbf{C})}}{2\mathbf{B}}$	$\lambda = \frac{\rho^2-t}{t^2}$ $E_1^t = \frac{i\Xi t}{\sqrt{\rho(\rho^2+1)}}(te_1 - (t+1)e_2) - \frac{\Xi^2 t^2}{\rho}e_3$ $E_3^t = -\frac{\Xi^2 t^2}{\rho}(e_3 + \frac{\Xi^2(\mathbf{A}+\rho\mathbf{B})t^2}{\rho}e_4)$	$\alpha = \frac{i\Xi(t-\rho^2)}{\sqrt{\rho(\rho^2+1)}}$ $E_2^t = -\frac{i\Xi t}{\rho\sqrt{\rho(\rho^2+1)}}(te_1 - (t-\rho^2)e_2)$ $E_4^t = \frac{\Xi^4}{\rho^2}e_4$	$\beta = \Xi^2(\rho\mathbf{A} + (\rho^2-1)\mathbf{B})t$ $E_3^t = \Xi^2 e_3$	$\gamma = \frac{i\Xi t}{\sqrt{\rho(\rho^2+1)}}$ $\gamma = 0$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{45}^4$ $\Xi = t^{-1}(t-1)^{-1}$	$E_1^t = -t^{-1}e_1 - \sqrt{\Xi t^{-1}}e_2 + t\Xi e_3$ $E_3^t = \Xi(e_3 - 4i\Xi t^{-1}e_4)$	$\alpha = -\Xi^2(t^3 - 2it^2 + 10it - 8i)$ $E_2^t = i\Xi(e_1 - \sqrt{\Xi}te_2)$	$\beta = i\Xi^{\frac{3}{2}}t^{-\frac{1}{2}}(3t-4)$ $E_3^t = \Xi e_3$	$\gamma = 0$ $E_4^t = \Xi^2 e_4$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{46}^4(\mathbf{A})$ $\Xi = t^{-1}(t-1)$	$E_1^t = \Xi e_1 + \sqrt{\Xi t^{-1}}e_2$	$\alpha = -2i\mathbf{A}\Xi$ $E_2^t = i\Xi(e_1 - \sqrt{\Xi}te_2)$	$\beta = -i\mathbf{A}\sqrt{\Xi t^{-1}}$ $E_3^t = \Xi e_3$	$\gamma = 0$ $E_4^t = \Xi^2 e_4$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{47}^4(\mathbf{A}, \mathbf{B})$	$E_1^t = t(e_1 + \mathbf{A}e_3)$	$\alpha = -\mathbf{A} - \mathbf{A}^2 + \mathbf{B}^2 + t$ $E_2^t = t(e_2 + \mathbf{B}e_3)$	$\beta = -\mathbf{A}\mathbf{B}$ $E_3^t = t^2(e_3 + \mathbf{B}^2e_4)$	$\gamma = -\mathbf{B}$ $E_4^t = t^3e_4$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{48}^4(\mathbf{A})$ $\Xi = t^{-1}(t-1)$	$E_1^t = \Xi e_1 + \sqrt{\Xi t^{-1}}e_2$	$\alpha = \mathbf{A}\Xi$ $E_2^t = i\Xi(e_1 - \sqrt{\Xi}te_2)$	$\beta = \mathbf{A}\sqrt{\Xi t^{-1}}$ $E_3^t = \Xi e_3$	$\gamma = -\mathbf{A}\sqrt{\Xi t^{-1}}$ $E_4^t = \Xi^2 e_4$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{49}^4(\mathbf{A})$ $\Xi = t^{-1}(t-1)^{-1}$	$E_1^t = -t^{-1}e_1 + \sqrt{\Xi t^{-1}}e_2 + t\Xi e_3$ $E_3^t = \Xi e_3 - 2\mathbf{A}\Xi^2 t^{-1}e_4$	$\alpha = -\Xi^2(t^3 - \mathbf{A}t^2 + 5\mathbf{A}t - 4\mathbf{A})$ $E_2^t = i\Xi(e_1 - \sqrt{\Xi}te_2)$	$\beta = -\mathbf{A}(t-2)\Xi^{\frac{3}{2}}t^{-\frac{1}{2}}$ $E_3^t = -i(t^{-1}e_1 + \Xi^{-\frac{1}{2}}t^{-\frac{3}{2}}e_2 - t\Xi e_3)$ $E_4^t = \Xi^2 e_4$	$\gamma = -\mathbf{A}\Xi^{\frac{3}{2}}t^{\frac{1}{2}}$ $\gamma = 0$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{50}^4(\mathbf{A})$	$E_1^t = t^{-1}e_1$	$\alpha = \mathbf{A}t^{-2}$ $E_2^t = t^{-1}e_2$	$\beta = 0$ $E_3^t = t^{-2}e_3$	$\gamma = t^{-2}$ $E_4^t = t^{-4}e_4$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{51}^4(\mathbf{A})$ $\Xi = t^{-1}(t-1)$	$E_1^t = \Xi e_1 + \sqrt{\Xi t^{-1}}e_2$	$\alpha = \mathbf{A}\Xi t^{-1}(t-2)$ $E_2^t = i\Xi(e_1 - \sqrt{\Xi}te_2)$	$\beta = \mathbf{A}\Xi^{\frac{3}{2}}t^{-\frac{1}{2}}$ $E_3^t = \Xi(e_3 - \mathbf{A}\Xi t^{-1}e_4)$	$\gamma = 0$ $E_4^t = \Xi^2 e_4$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{52}^4$	$E_1^t = t^{-1}e_1$	$\alpha = t^{-2}$ $E_2^t = t^{-1}e_2$	$\beta = 0$ $E_3^t = t^{-2}e_3$	$\gamma = 0$ $E_4^t = t^{-4}e_4$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{53}^4$	$E_1^t = te_1 + e_3$	$\alpha = -\frac{2+t}{t^2}$ $E_2^t = te_2 + ie_3$	$\beta = -\frac{1}{2} - \frac{i}{t^2}$ $E_3^t = t^2e_3 - e_4$	$\gamma = 1 - \frac{i}{t}$ $E_4^t = t^2e_4$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{54}^4(\mathbf{A})$ $\Xi = t\sqrt{t-1}$	$E_1^t = -\frac{1}{t}(\Xi e_1 - te_2 + \Xi\mathbf{A}e_3)$ $E_3^t = te_3 + \frac{\Xi}{t^3}((2\mathbf{A}+1)^2\Xi + 2it^2)e_4$	$\alpha = -\frac{\mathbf{A}(\mathbf{A}+1)t^2 + 2(2\mathbf{A}+1)^2(t-1) + 4i\Xi}{t^2}$ $E_2^t = t e_2$	$\beta = \frac{(2\mathbf{A}+1)(\mathbf{A}(t-2)-1)\Xi + it^2(t-2)}{t^2}$ $E_3^t = -\frac{i}{t}(\Xi e_1 + t(t-1)e_2 - \Xi(\mathbf{A}+1)e_3)$ $E_4^t = -\Xi e_4$	$\gamma = \frac{(2\mathbf{A}+1)\Xi}{t^2}$ $\gamma = 0$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{55}^4(\mathbf{A})$	$E_1^t = te_1 + \mathbf{A}te_3$	$\alpha = -\mathbf{A}(1+\mathbf{A})$ $E_2^t = te_2$	$\beta = t$ $E_3^t = t^2e_3$	$\gamma = 0$ $E_4^t = t^3e_4$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{56}^4$ $\Xi = t^{-1}(t-1)$	$E_1^t = \Xi e_1 + \sqrt{\Xi t^{-1}}e_2$	$\alpha = 0$ $E_2^t = i\Xi(e_1 - \sqrt{\Xi}te_2)$	$\beta = i\sqrt{\Xi t^{-1}}$ $E_3^t = \Xi e_3$	$\gamma = -2i\sqrt{\Xi t^{-1}}$ $E_4^t = \Xi^2 e_4$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{57}^4(\mathbf{A}, \mathbf{B})$ $\Xi = t^{-1}(t-1)$	$E_1^t = \Xi e_1 + \sqrt{\Xi t^{-1}}e_2$ $E_3^t = \Xi(e_3 - i(2\mathbf{A}+\mathbf{B})\Xi t^{-1}e_4)$	$\alpha = -2i(2\mathbf{A}+\mathbf{B})\Xi t^{-1}$ $E_2^t = i\Xi(e_1 - \sqrt{\Xi}te_2)$	$\beta = i(\mathbf{A}(t-2) - \mathbf{B})\sqrt{\Xi t^{-3}}$ $E_3^t = \Xi e_3$	$\gamma = i\mathbf{A}\sqrt{\Xi t^{-1}}$ $E_4^t = \Xi^2 e_4$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{58}^4$	$E_1^t = te_1 + e_3$	$\alpha = -\frac{2+t}{t^2}$ $E_2^t = te_2 + ie_3$	$\beta = -\frac{i}{t}$ $E_3^t = t^2e_3 - e_4$	$\gamma = \frac{t-i}{t}$ $E_4^t = t^2e_4$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{59}^4(\mathbf{A}, \mathbf{B})$	$E_1^t = t(e_1 + \mathbf{A}e_3)$	$\alpha = -\mathbf{A} - \mathbf{A}^2 + \mathbf{B}^2$ $E_2^t = t(e_2 + \mathbf{B}e_3)$	$\beta = -\mathbf{A}\mathbf{B}$ $E_3^t = t^2(e_3 + \mathbf{B}^2e_4)$	$\gamma = -\mathbf{B}$ $E_4^t = t^3e_4$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{60}^4$ $\Xi = \sqrt{t-1}$	$E_1^t = \Xi e_1 + e_2 + i\Xi e_3$	$\alpha = t^{-2}((1-i)t^2 + 2(i-4)t + 8)$ $E_2^t = \Xi(ie_1 - i\Xi e_2 + e_3)$	$\beta = 2\Xi t^{-2}(t-2)$ $E_3^t = te_3 - 4t^{-1}\Xi^2 e_4$	$\gamma = -2i\Xi t^{-1}$ $E_4^t = t\Xi e_4$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{61}^4(\mathbf{A})$ $\Xi = \sqrt{t-1}$	$E_1^t = \Xi e_1 + e_2 - i\mathbf{A}\Xi e_3$	$\alpha = \mathbf{A}(\mathbf{A}+i)$ $E_2^t = \Xi(ie_1 - i\Xi e_2 + \mathbf{A}e_3)$	$\beta = 0$ $E_3^t = te_3$	$\gamma = 0$ $E_4^t = t\Xi e_4$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{62}^4$ $\Xi = \sqrt{t-1}$	$E_1^t = \Xi e_1 + e_2 + \Xi e_3$	$\alpha = -2t^{-2}(t^2 - 9t + 9)$ $E_2^t = i\Xi(e_1 - \Xi e_2 - 2e_3)$	$\beta = -3\Xi t^{-2}(t-3)$ $E_3^t = te_3 + 9\Xi^2 t^{-1}e_4$	$\gamma = -3\Xi t^{-1}$ $E_4^t = t\Xi e_4$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{63}^4$ $\Xi = \sqrt{t-1}$	$E_1^t = \Xi e_1 + e_2 - \frac{1}{2}\Xi e_3$	$\alpha = \frac{1}{4}$ $E_2^t = i\Xi(e_1 - \Xi e_2 - \frac{1}{2}e_3)$	$\beta = 0$ $E_3^t = te_3$	$\gamma = 0$ $E_4^t = t\Xi e_4$
$\mathcal{C}\mathcal{D}_{44}^4 \rightarrow \mathcal{C}\mathcal{D}_{64}^4(\mathbf{A})$	$E_1^t = t(e_1 + \mathbf{A}e_3)$	$\alpha = -\mathbf{A}^2 - \mathbf{A}$ $E_2^t = te_2$	$\beta = 0$ $E_3^t = t^2e_3$	$\gamma = 0$ $E_4^t = t^3e_4$

$\mathcal{E}\mathcal{D}_{44}^4 \rightarrow \mathcal{E}\mathcal{D}_{65}^4(\mathbf{A})$ $\Xi = \sqrt{t-1}$	$E_1^t = \Xi e_1 + e_2 + \mathbf{A}\Xi e_3$	$\alpha = -\mathbf{A}((\mathbf{A}+1)t-2\mathbf{A})t^{-2}\Xi^2$ $E_2^t = i\Xi(e_1 - \Xi e_2)$	$\beta = -\mathbf{A}^2\Xi^3t^{-2}$ $E_3^t = te_3 + \mathbf{A}^2\Xi^2t^{-1}e_4$	$\gamma = -\mathbf{A}\Xi t^{-1}$ $E_4^t = t\Xi e_4$
$\mathcal{E}\mathcal{D}_{44}^4 \rightarrow \mathcal{E}\mathcal{D}_{66}^4$	$E_1^t = e_1 - e_3$	$\alpha = t^{-2}$ $E_2^t = e_2 + t^{-1}e_3$	$\beta = t^{-1}$ $E_3^t = e_3 + t^{-2}e_4$	$\gamma = 0$ $E_4^t = t^{-1}e_4$
$\mathcal{E}\mathcal{D}_{44}^4 \rightarrow \mathcal{E}\mathcal{D}_{67}^4$	$E_1^t = t^{-1}e_1$	$\alpha = 0$ $E_2^t = t^{-1}e_2$	$\beta = 0$ $E_3^t = t^{-2}e_3$	$\gamma = 0$ $E_4^t = t^{-4}e_4$
$\mathcal{E}\mathcal{D}_{44}^4 \rightarrow \mathcal{E}\mathcal{D}_{68}^4$	$E_1^t = e_1 + ie_3$	$\alpha = \frac{2-it+t^2}{t^2}$ $E_2^t = te_2 + e_3$	$\beta = -\frac{i}{2}$ $E_3^t = t^2e_3 + e_4$	$\gamma = -\frac{1}{t}$ $E_4^t = t^2e_4$
$\mathcal{E}\mathcal{D}_{44}^4 \rightarrow \mathcal{E}\mathcal{D}_{69}^4$	$E_1^t = te_1 + ie_3$	$\alpha = \frac{2-it}{t^2}$ $E_2^t = te_2 + e_3$	$\beta = -\frac{i}{2}$ $E_3^t = t^2e_3 + e_4$	$\gamma = -\frac{1}{t}$ $E_4^t = t^2e_4$
$\mathcal{E}\mathcal{D}_{44}^4 \rightarrow \mathcal{E}\mathcal{D}_{70}^4$	$E_1^t = te_1 + e_3$	$\alpha = -\frac{1+t}{t^2}$ $E_2^t = te_2$	$\beta = 0$ $E_3^t = t^2e_3$	$\gamma = 0$ $E_4^t = t^2e_4$
$\mathcal{E}\mathcal{D}_{12}^4 \rightarrow \mathcal{E}\mathcal{D}_{71}^4$	$\lambda = -1 + t$ $E_1^t = t(e_1 + ie_2)$	$\alpha = -t^2$ $E_2^t = t(e_2 + ie_3 + 3te_4)$	$E_3^t = t^2(e_3 - 3ite_4)$	$E_4^t = (t-3)t^3e_4$
$\mathcal{E}\mathcal{D}_{12}^4 \rightarrow \mathcal{E}\mathcal{D}_{72}^4$	$\lambda = -1$ $E_1^t = te_1$	$\alpha = t^2$ $E_2^t = -3(e_2 - 3e_3 - 36e_4)$	$E_3^t = -3t(e_3 + 9e_4)$	$E_4^t = 9t^2e_4$
$\mathcal{E}\mathcal{D}_{12}^4 \rightarrow \mathcal{E}\mathcal{D}_{73}^4$	$\lambda = -1$ $E_1^t = te_1 - 6e_2$	$\alpha = \frac{t^2}{4}$ $E_2^t = t(e_2 + 9e_4)$	$E_3^t = t^2(e_3 + \frac{3t}{2}e_4)$	$E_4^t = -3t^3e_4$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{74}^4(\mathbf{A})$ $\Xi = \sqrt{\mathbf{A}(1+\mathbf{A})}$	$\lambda = \frac{1}{4}$ $E_1^t = te_1 - 6e_2$	$\alpha = -\frac{1+2\mathbf{A}}{\Xi}$ $E_2^t = t(e_2 + 9e_4)$	$\beta = 0$ $E_3^t = t^2(e_3 + \frac{3t}{2}e_4)$	$\gamma = 0$ $E_4^t = -3t^3e_4$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{75}^4(\mathbf{A})$ $\Xi = \sqrt{\mathbf{A}(1+\mathbf{A})}$	$\lambda = \frac{1}{4}$ $E_1^t = 8i\Xi t(2e_1 - e_2 + \frac{2i\mathbf{A}}{\Xi}e_3)$	$\alpha = -\frac{2}{\Xi}$ $E_2^t = 16t^2(i\Xi e_2 + 4\mathbf{A}e_3)$	$\beta = 4$ $E_3^t = 128\Xi^2t^3(e_3 + 8e_4)$	$\gamma = \frac{4i\mathbf{A}}{\Xi}$ $E_4^t = 2048\Xi^2t^4e_4$
$\mathcal{E}\mathcal{D}_{12}^4 \rightarrow \mathcal{E}\mathcal{D}_{76}^4$	$\lambda = -1$ $E_1^t = \frac{2}{\Xi}e_1 - \frac{1}{\Xi}e_2 + \frac{2}{1+\mathbf{A}}e_3$	$\alpha = -\frac{3t^2}{2}$ $E_2^t = \frac{2t}{\Xi}e_2$	$E_3^t = -\frac{2t}{\Xi}e_3$	$E_4^t = \frac{4t}{\Xi}e_4$
$\mathcal{E}\mathcal{D}_{78}^4 \rightarrow \mathcal{E}\mathcal{D}_{77}^4$	$E_1^t = t^{-1}e_1$	$E_2^t = t^{-1}e_2$	$E_3^t = t^{-2}e_3$	$E_4^t = t^{-3}e_4$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{78}^4$	$\lambda = \frac{1}{4}$ $E_1^t = -2i\sqrt{2t}(2e_1 - e_2)$ $E_3^t = -16t^2(e_3 + 4e_4)$	$\alpha = \frac{i}{\sqrt{2t}}(1-2t)$ $E_2^t = -2i\sqrt{2t}(2e_1 - (2t+1)e_2) + 8te_3$ $E_4^t = 128t^2e_4$	$\beta = 2$	$\gamma = i\sqrt{\frac{2}{t}}$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{79}^4(\mathbf{A})$ $\Xi = \sqrt{2t(\mathbf{A}-t)}$	$\lambda = \frac{1+2t^2}{4}$ $E_1^t = \frac{2}{\Xi}(2e_1 - e_2)$	$\alpha = -\frac{1+t}{\Xi}$ $E_2^t = \frac{4t}{\Xi}(e_2 + \frac{2t}{\Xi}e_3)$	$\beta = 2$ $E_3^t = \frac{8t}{\Xi^2}(e_3 + 4e_4)$	$\gamma = \frac{2(t-1)}{\Xi}$ $E_4^t = \frac{64t^2}{\Xi^4}e_4$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{80}^4$	$\lambda = \frac{1}{4}$ $E_1^t = \sqrt{\frac{2}{t}}(2e_1 - e_2)$	$\alpha = -\sqrt{2t}$ $E_2^t = 2\sqrt{2t}e_2$	$\beta = 2$ $E_3^t = -4(e_3 + 4e_4)$	$\gamma = 0$ $E_4^t = 16e_4$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{81}^4$	$\lambda = \frac{1}{4}$ $E_1^t = \sqrt{\frac{2}{t}}(2e_1 - e_2)$	$\alpha = 0$ $E_2^t = 2\sqrt{2t}e_2$	$\beta = 2$ $E_3^t = -4(e_3 + 4e_4)$	$\gamma = 0$ $E_4^t = 16e_4$
$\mathcal{E}\mathcal{D}_{102}^4 \rightarrow \mathcal{E}\mathcal{D}_{82}^4$	$\lambda = \frac{1}{4}$ $E_1^t = -\sqrt{2}(2e_1 - e_2)$	$\alpha = \sqrt{2t}$ $E_2^t = \sqrt{2t}e_2$	$E_3^t = 2t(e_3 - 2e_4)$	$E_4^t = 4t^2e_4$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{83}^4(\mathbf{A})$	$\lambda = \frac{1}{4}$ $E_1^t = 2\mathbf{A}^{-\frac{1}{2}}e_2$	$\alpha = \mathbf{A}^{-\frac{1}{2}}$ $E_2^t = t^{-1}(e_1 - \frac{1}{2}e_2)$	$\beta = 0$ $E_3^t = \mathbf{A}^{-\frac{1}{2}}t^{-1}e_3$	$\gamma = 0$ $E_4^t = t^{-2}\mathbf{A}^{-1}e_4$
$\mathcal{E}\mathcal{D}_{83}^4 \rightarrow \mathcal{E}\mathcal{D}_{84}^4$	$E_1^t = t^{-1}e_1 + te_2$	$\alpha = t^{-2}$ $E_2^t = e_2$	$E_3^t = t^{-1}e_3$	$E_4^t = t^{-2}e_4$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{85}^4$	$\lambda = \frac{1}{4}$ $E_1^t = t^{-2}e_1 - \frac{1}{2t}e_2$	$\alpha = 0$ $E_2^t = t^{-1}e_2$	$\beta = 0$ $E_3^t = -\frac{1}{2t^3}e_3$	$\gamma = 0$ $E_4^t = \frac{1}{4t^6}e_4$
$\mathcal{E}\mathcal{D}_{12}^4 \rightarrow \mathcal{E}\mathcal{D}_{86}^4$	$\lambda = -1$ $E_1^t = te_1 + e_2$	$\alpha = \frac{3t^3}{2}$ $E_2^t = 2e_2 + 3te_4$	$E_3^t = 2te_3 - 3t^3e_4$	$E_4^t = -6t^2e_4$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{87}^4(\Lambda)$ $\Xi = \Theta^{-\frac{1}{2}}$	$\lambda = \Lambda - \Psi t$ $E_1^t = \sqrt{\Theta}t^{-1}(e_1 - \Psi e_2)$	$\alpha = \Xi((\Theta - \Psi)t^{-1} - 1)$ $E_2^t = \Xi t^{-1}(e_1 - (t + \Psi)e_2)$	$\beta = 0$ $E_3^t = -t^{-2}(te_3 - \Xi^2e_4)$	$\gamma = \frac{\Xi}{\Psi t}(\Theta - \Psi)$ $E_4^t = t^{-2}e_4$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{88}^4(\Lambda)$	$\lambda = \Lambda - \Theta t$ $E_1^t = \sqrt{\Psi}t^{-1}(e_1 - \Theta e_2)$ $E_3^t = -t^{-2}(te_3 - \Psi^{-1}e_4)$	$\alpha = -\Psi^{-\frac{1}{2}}$ $E_2^t = \Psi^{-\frac{1}{2}}t^{-1}(e_1 - (t + \Theta)e_2)$ $E_4^t = t^{-2}e_4$	$\beta = 0$	$\gamma = 0$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{89}^4(\Lambda)$	$\lambda = \Lambda - \Psi t$ $E_1^t = \sqrt{\Theta}t^{-1}(e_1 - \Psi e_2)$ $E_3^t = -t^{-2}(te_3 - \Theta^{-1}e_4)$	$\alpha = -\Theta^{-\frac{1}{2}}$ $E_2^t = \Theta^{-\frac{1}{2}}t^{-1}(e_1 - (t + \Psi)e_2)$ $E_4^t = t^{-2}e_4$	$\beta = 0$	$\gamma = 0$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{90}^4(\Lambda)$ $\Xi = \Psi^{-\frac{1}{2}}$	$\lambda = \Lambda - \Theta t$ $E_1^t = \sqrt{\Psi}t^{-1}(e_1 - \Theta e_2)$	$\alpha = \Xi((\Psi - \Theta)t^{-1} - 1)$ $E_2^t = \Xi t^{-1}(e_1 - (t + \Theta)e_2)$	$\beta = 0$ $E_3^t = -t^{-2}(te_3 - \Xi^2e_4)$	$\gamma = \frac{\Xi}{\Theta t}(\Psi - \Theta)$ $E_4^t = t^{-2}e_4$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{91}^4(\Lambda, \mathbf{A})$ $\Xi = \Theta^{-\frac{1}{2}}$	$\lambda = \Lambda - \Psi t$ $E_1^t = \sqrt{\Theta}t^{-1}(e_1 - \Psi e_2)$	$\alpha = -\mathbf{A}\Xi(1 + \Psi t^{-1})$ $E_2^t = \Xi t^{-1}(e_1 - (t + \Psi)e_2)$	$\beta = 0$ $E_3^t = -t^{-2}(te_3 - \Xi^2e_4)$	$\gamma = -\mathbf{A}\Xi t^{-1}$ $E_4^t = t^{-2}e_4$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{92}^4(\Lambda, \mathbf{A})$ $\Xi = \Psi^{-\frac{1}{2}}$	$\lambda = \Lambda - \Theta t$ $E_1^t = \sqrt{\Psi}t^{-1}(e_1 - \Theta e_2)$	$\alpha = -\mathbf{A}\Xi(1 + \Theta t^{-1})$ $E_2^t = \Xi t^{-1}(e_1 - (t + \Theta)e_2)$	$\beta = 0$ $E_3^t = -t^{-2}(te_3 - \Xi^2e_4)$	$\gamma = -\mathbf{A}\Xi t^{-1}$ $E_4^t = t^{-2}e_4$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{93}^4(\mathbf{A})$	$\lambda = 0$ $E_1^t = t^{-1}e_1$	$\alpha = -\mathbf{A}$ $E_2^t = t^{-1}e_1 - e_2$	$\beta = t^{-2}$ $E_3^t = -t^{-1}e_3 - t^{-3}e_4$	$\gamma = 0$ $E_4^t = t^{-2}e_4$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{94}^4(\mathbf{A}, \mathbf{B})$	$\lambda = 0$ $E_1^t = t^{-1}e_1$	$\alpha = -\mathbf{A}$ $E_2^t = t^{-1}e_1 - e_2$	$\beta = -t^{-2}((\mathbf{B}-1)t - \mathbf{B})$ $E_3^t = -t^{-1}e_3 - \mathbf{B}t^{-3}e_4$	$\gamma = -\mathbf{A}t^{-1}$ $E_4^t = t^{-2}e_4$
$\mathcal{E}\mathcal{D}_{112}^4 \rightarrow \mathcal{E}\mathcal{D}_{95}^4(\mathbf{A})$	$\lambda = t^2$	$\alpha = \mathbf{A}$	$\beta = t^{-1}$	$\gamma = 0$

$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{96}^4(\mathbf{A})$	$E_1^t = -t^{-1}e_1$ $\lambda = 0$ $E_1^t = t^{-1}e_1$	$E_2^t = -t^{-1}e_1 + e_2$ $\alpha = t$ $E_2^t = t^{-1}e_1 - e_2$	$E_3^t = -t^{-1}e_3$ $\beta = -t^{-2}((\mathbf{A} - 1)t - \mathbf{A})$ $E_3^t = -t^{-1}e_3 - \mathbf{A}t^{-3}e_4$	$E_4^t = t^{-2}e_4$ $\gamma = t^{-1}$ $E_4^t = t^{-2}e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{97}^4(\Lambda)$ $\Xi = (t + \Psi - \Theta)^{-\frac{1}{2}}$	$\lambda = \Lambda - \Psi t$ $E_1^t = \Xi\sqrt{\Theta t^{-1}}(e_1 - \Psi e_2)$ $E_3^t = \Xi^2(-e_3 + \Xi^2\Psi^{-1}e_4)$	$\alpha = -\Xi\sqrt{\Theta t}$ $E_2^t = \Xi(\Theta t)^{-\frac{1}{2}}(e_1 - (t + \Psi)e_2)$ $E_4^t = \Xi^2e_4$	$\beta = (1 - \Theta\Psi^{-1})\Xi^2$	$\gamma = 0$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{98}^4(\Lambda)$ $\Xi = (t + \Theta - \Psi)^{-\frac{1}{2}}$	$\lambda = \Lambda - \Theta t$ $E_1^t = \Xi\sqrt{\Psi t^{-1}}(e_1 - \Theta e_2)$ $E_3^t = \Xi^2(-e_3 + \Xi^2\Theta^{-1}e_4)$	$\alpha = -\Xi\sqrt{\Psi t}$ $E_2^t = \Xi(\Psi t)^{-\frac{1}{2}}(e_1 - (t + \Theta)e_2)$ $E_4^t = \Xi^2e_4$	$\beta = (1 - \Psi\Theta^{-1})\Xi^2$	$\gamma = 0$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{99}^4(\mathbf{A})$ $\Xi = \frac{t+1}{\sqrt{t-1}}$	$\lambda = \frac{t}{(t+1)^2}$ $E_1^t = \Xi(-1 + \frac{1}{t})e_1 + e_2$	$\alpha = \frac{t-\mathbf{A}+1}{\sqrt{t-1}}$ $E_2^t = \Xi(t+1)(-\frac{1}{t}e_1 + e_2)$	$\beta = 1 + \frac{1}{t}$ $E_3^t = \Xi^2(-e_3 + \frac{1}{t}e_4)$	$\gamma = (1 - \mathbf{A})\frac{\Xi}{t}$ $E_4^t = \Xi^4e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{100}^4(\mathbf{A})$	$\lambda = -\frac{1}{2} + \frac{1}{4}$ $E_1^t = \sqrt{2}t^{-\frac{1}{4}}(\frac{1}{2}e_1 - \frac{1}{4}e_2)$ $E_3^t = -t^{-1}e_3 + 2t^{-2}e_4$	$\alpha = -\sqrt{2}\mathbf{A}$ $E_2^t = \sqrt{2}t^{-1}(e_1 - (t + \frac{1}{2})e_2)$ $E_4^t = t^{-2}e_4$	$\beta = 0$	$\gamma = 0$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{101}^4(\mathbf{A}, \mathbf{B})$ $\Xi = t + 1$	$\lambda = \Xi^{-2}t$ $E_1^t = \Xi t^{-\frac{1}{2}}(-\Xi e_1 + t e_2)$	$\alpha = -\mathbf{B}\sqrt{t}$ $E_2^t = -\Xi^2 t^{-\frac{1}{2}}e_1$	$\beta = \Xi t^{-1}$ $E_3^t = \Xi^2(e_3 + \Xi^2 t^{-1}e_4)$	$\gamma = (\mathbf{A} - \mathbf{B})\frac{\Xi}{\sqrt{t}}$ $E_4^t = \Xi^4e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{102}^4(\Lambda, \mathbf{A})$ $\Xi = (\Theta - 2\Lambda + t)^{-\frac{1}{2}}$	$\lambda = \Lambda + \Psi\Theta^{-1}t$ $E_1^t = \Theta^{\frac{1}{2}}\Xi t^{-\frac{1}{2}}(\Theta e_1 - \Lambda e_2)$ $E_3^t = \Theta\Xi^2(e_3 + \Theta\Psi^{-1}\Xi^2e_4)$	$\alpha = \mathbf{A}\Lambda\Theta^{\frac{1}{2}}\Xi t^{-\frac{1}{2}}$ $E_2^t = \Theta^{-\frac{1}{2}}\Xi t^{-\frac{1}{2}}(\Theta e_1 + (t - \Lambda)e_2)$ $E_4^t = \Theta^2\Xi^2e_4$	$\beta = (\Theta - 2\Lambda)\Xi^2\Psi^{-1}$	$\gamma = \mathbf{A}\Theta^{\frac{3}{2}}\Xi t^{-\frac{1}{2}}$
$\mathcal{C}\mathcal{D}_{102}^4 \rightarrow \mathcal{C}\mathcal{D}_{103}^4$	$\lambda = 0$ $E_1^t = -e_1 + t e_3$	$\alpha = 0$ $E_2^t = -e_2$	$E_3^t = e_3$	$E_4^t = e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{104}^4$	$\lambda = 0$ $E_1^t = t^{-1}e_1$	$\alpha = t^{-1}$ $E_2^t = t^{-1}e_2$	$\beta = t^{-2}$ $E_3^t = t^{-2}e_3$	$\gamma = t^{-1}$ $E_4^t = t^{-4}e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{105}^4(\Lambda, \mathbf{A}, \mathbf{B})$	$\lambda = \Lambda$ $E_1^t = t^{-1}e_1 - e_3$	$\alpha = t + t^{-1}$ $E_2^t = t^{-1}e_2$	$\beta = \mathbf{B} + \mathbf{A}t^{-2}$ $E_3^t = t^{-2}e_3$	$\gamma = \mathbf{B}t^{-1}$ $E_4^t = t^{-4}e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{106}^4(\mathbf{A})$	$\lambda = 0$ $E_1^t = t^{-1}e_1$	$\alpha = t^{-1}$ $E_2^t = t^{-1}e_2$	$\beta = \mathbf{A}t^{-2}$ $E_3^t = t^{-2}e_3$	$\gamma = 0$ $E_4^t = t^{-4}e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{107}^4(\Lambda)$	$\lambda = \Lambda$ $E_1^t = t^{-1}e_1$	$\alpha = \frac{1+\sqrt{1-4\Lambda}}{2t}$ $E_2^t = t^{-1}e_2$	$\beta = 0$ $E_3^t = t^{-2}e_3$	$\gamma = t^{-1}$ $E_4^t = t^{-4}e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{108}^4(\Lambda)$	$\lambda = \Lambda$ $E_1^t = t^{-1}e_1$	$\alpha = \frac{1-\sqrt{1-4\Lambda}}{2t}$ $E_2^t = t^{-1}e_2$	$\beta = 0$ $E_3^t = t^{-2}e_3$	$\gamma = t^{-1}$ $E_4^t = t^{-4}e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{109}^4(\Lambda, \mathbf{A})$	$\lambda = \Lambda$ $E_1^t = t^{-1}e_1$	$\alpha = 0$ $E_2^t = t^{-1}e_2$	$\beta = t^{-2}$ $E_3^t = t^{-2}e_3$	$\gamma = \mathbf{A}t^{-1}$ $E_4^t = t^{-4}e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{110}^4(\Lambda)$	$\lambda = \Lambda$ $E_1^t = t^{-1}e_1$	$\alpha = 0$ $E_2^t = t^{-1}e_2$	$\beta = 0$ $E_3^t = t^{-2}e_3$	$\gamma = t^{-1}$ $E_4^t = t^{-4}e_4$
$\mathcal{C}\mathcal{D}_{112}^4 \rightarrow \mathcal{C}\mathcal{D}_{111}^4(\Lambda)$	$\lambda = \Lambda$ $E_1^t = t^{-1}e_1$	$\alpha = 0$ $E_2^t = t^{-1}e_2$	$\beta = \Lambda^{-1}$ $E_3^t = t^{-2}e_3 + t^{-2}e_4$	$\gamma = 0$ $E_4^t = t^{-4}e_4$

□

REFERENCES

- [1] Arenas M., The Wedderburn principal theorem for generalized almost-Jordan algebras, *Communications in Algebra*, 35 (2007), 2, 675–688.
- [2] Arenas M., Labra A., On nilpotency of generalized almost-Jordan right-nilalgebras, *Algebra Colloquium*, 15 (2008), 1, 69–82.
- [3] Abdelwahab H., Calderón A.J., Kaygorodov I., The algebraic and geometric classification of nilpotent binary Lie algebras, *International Journal of Algebra and Computation*, 29 (2019), 6, 1113–1129.
- [4] Beneš T., Burde D., Classification of orbit closures in the variety of three-dimensional Novikov algebras, *Journal of Algebra Applications*, 13 (2014), 2, 1350081, 33 pp.
- [5] Burde D., Steinhoff C., Classification of orbit closures of 4–dimensional complex Lie algebras, *Journal of Algebra*, 214 (1999), 2, 729–739.
- [6] Calderon A., Fernández Ouaridi A., Kaygorodov I., The classification of 2-dimensional rigid algebras, *Linear and Multilinear Algebra*, 68 (2020), 4, 828–844.
- [7] Camacho L., Kaygorodov I., Lopatkin V., Salim M., The variety of dual Mock-Lie algebras, *Communications in Mathematics*, to appear, arXiv:1910.01484
- [8] Demir I., Misra K., Stitzinger E., On classification of four-dimensional nilpotent Leibniz algebras, *Communications in Algebra*, 45 (2017), 3, 1012–1018.
- [9] Dzhumadil'daev A., Assosymmetric algebras under Jordan product, *Communications in Algebra*, 46 (2018), 1, 129–142
- [10] Gerstenhaber M., The cohomology structure of an associative ring, *Annals of Mathematics (2)*, 78 (1963), 267–288.
- [11] Gorbatsevich V., Anticommutative finite-dimensional algebras of the first three levels of complexity, *St. Petersburg Mathematical Journal*, 5 (1994), 3, 505–521.
- [12] Gorshkov I., Kaygorodov I., Khrypchenko M., The geometric classification of Tortkara algebras, *Communications in Algebra*, 48 (2020), 1, 204–209.
- [13] Grunewald F., O'Halloran J., Varieties of nilpotent Lie algebras of dimension less than six, *Journal of Algebra*, 112 (1988), 315–325.
- [14] Grunewald F., O'Halloran J., A Characterization of orbit closure and applications, *Journal of Algebra*, 116 (1988), 163–175.
- [15] Hentzel I., Labra A., On left nilalgebras of left nilindex four satisfying an identity of degree four, *International Journal of Algebra and Computation*, 17 (2007), 1, 27–35.
- [16] Hentzel I., Peresi L., Almost Jordan rings, *Proceedings of the American Mathematical Society*, 104 (1988), 2, 343–348.

- [17] Jumaniyozov D., Kaygorodov I., Khudoyberdiyev A., The algebraic and geometric classification of nilpotent noncommutative Jordan algebras, *Journal of Algebra and its Applications*, 2020, DOI: 10.1142/S0219498821502029
- [18] Ismailov N., Kaygorodov I., Mashurov F., The algebraic and geometric classification of nilpotent assosymmetric algebras, *Algebras and Representation Theory*, 2020, DOI: 10.1007/s10468-019-09935-y
- [19] Ismailov N., Kaygorodov I., Volkov Yu., The geometric classification of Leibniz algebras, *International Journal of Mathematics*, 29 (2018), 5, 1850035.
- [20] Ismailov N., Kaygorodov I., Volkov Yu., Degenerations of Leibniz and anticommutative algebras, *Canadian Mathematical Bulletin*, 62 (2019), 3, 539–549.
- [21] Karimjanov I., Kaygorodov I., Khudoyberdiyev A., The algebraic and geometric classification of nilpotent Novikov algebras, *Journal of Geometry and Physics*, 143 (2019), 11–21.
- [22] Kaygorodov I., Khrypchenko M., The algebraic classification of nilpotent \mathcal{CD} -algebras, arXiv:2006.00734
- [23] Kaygorodov I., Khrypchenko M., Lopes S., The algebraic and geometric classification of nilpotent anticommutative algebras, *Journal of Pure and Applied Algebra*, 224 (2020), 8, 106337.
- [24] Kaygorodov I., Khrypchenko M., Popov Yu., The algebraic and geometric classification of nilpotent terminal algebras, arXiv:1909.00358.
- [25] Kaygorodov I., Lopes S., Popov Yu., Degenerations of nilpotent associative commutative algebras, *Communications in Algebra*, 48 (2020), 4, 1632–1639.
- [26] Kaygorodov I., Páez-Guillán P., Voronin V., The algebraic and geometric classification of nilpotent bicommutative algebras, *Algebras and Representation Theory*, 2020, DOI: 10.1007/s10468-019-09944-x
- [27] Kaygorodov I., Popov Yu., Pozhidaev A., Volkov Yu., Degenerations of Zinbiel and nilpotent Leibniz algebras, *Linear and Multilinear Algebra*, 66 (2018), 4, 704–716.
- [28] Kaygorodov I., Popov Yu., Pozhidaev A., Volkov Yu., Corrigendum to ‘Degenerations of Zinbiel and nilpotent Leibniz algebras’, *Linear and Multilinear Algebra*, 2020, DOI: 10.1080/03081087.2020.1749543
- [29] Kaygorodov I., Popov Yu., Volkov Yu., Degenerations of binary-Lie and nilpotent Malcev algebras, *Communications in Algebra*, 46 (2018), 11, 4929–4941.
- [30] Kaygorodov I., Pozhidaev A., Saraiva P., On a ternary generalization of Jordan algebras, *Linear and Multilinear Algebra*, 67 (2019), 6, 1074–1102.
- [31] Kaygorodov I., Volkov Yu., The variety of 2-dimensional algebras over an algebraically closed field, *Canadian Journal of Mathematics*, 71 (2019), 4, 819–842.
- [32] Kaygorodov I., Volkov Yu., Complete classification of algebras of level two, *Moscow Mathematical Journal*, 19 (2019), 3, 485–521.
- [33] Kaygorodov I., Volkov Yu., Degenerations of Filippov algebras, *Journal of Mathematical Physics*, 61 (2020), 2, 021701.
- [34] Kaygorodov I., Zusmanovich P., On anticommutative algebras for which $[R_a, R_b]$ is a derivation, in preparation.
- [35] Khudoyberdiyev A., The classification of algebras of level two, *Journal of Geometry and Physics*, 98 (2015), 13–20.
- [36] Osborn J., Lie triple algebras with one generator, *Mathematische Zeitschrift*, 110 (1969), 52–74.
- [37] Seeley C., Degenerations of 6-dimensional nilpotent Lie algebras over \mathbb{C} , *Communications in Algebra*, 18 (1990), 3493–3505.
- [38] Sidorov A., Lie triple algebras, *Algebra and Logic*, 87 (1981), 72–78.
- [39] Volkov Yu., Anticommutative Engel algebras of the first five levels, *Linear and Multilinear Algebra*, 2020, DOI: 10.1080/03081087.2020.1715333
- [40] Volkov Yu., n -ary algebras of the first level, arXiv:1910.10200