

Structural Gaussian mixture vector autoregressive model

Savi Virolainen
University of Helsinki

Abstract

A structural version of the Gaussian mixture vector autoregressive model is introduced. The shocks are identified by combining simultaneous diagonalization of the error term covariance matrices with zero and sign constraints. It turns out that this often leads to less restrictive identification conditions than in conventional SVAR models, while some of the constraints are also testable. The accompanying R-package gmvarkit provides easy-to-use tools for estimating the models and applying the introduced methods.

Keywords: structural nonlinear autoregression, structural mixture model, regime switching, Gaussian mixture

This work was supported by the Academy of Finland under Grant 308628. The author thanks Markku Lanne and Pentti Saikkonen for helpful discussions and comments.

Contact address: Savi Virolainen, Faculty of Social Sciences, University of Helsinki, P. O. Box 17, FI-00014 University of Helsinki, Finland; e-mail: savi.virolainen@helsinki.fi.

The author has no conflict of interest to declare.

1 Introduction

Tracing out the effects of an economic shock is a major task in econometrics. A popular approach is to consider a set of key variables and utilize a structural vector autoregressive (SVAR) or error correction (SVEC) model for the purpose. They have well established theoretical grounds (see Kilian and Lütkepohl, 2017, and the references therein) and are accommodated by many of the popular statistical software. Linear SVAR and SVEC models are not, however, suitable for modelling series in which the underlying data generating dynamics are nonlinear or the shocks have asymmetric effects in different states of the world. Models capable of capturing such behaviour include mixture models such as the MVAR model (Fong et al., 2010), the MPVAR model (Bentarzi and Djeddou, 2014), the GMVAR model (Kalliovirta et al., 2016), and the logit-MVAR model (Burgard et al., 2019).

In this paper, we introduce a structural version of the GMVAR model. In the SGMVAR model of autoregressive order p , the regime switching dynamics are endogeneously determined by the full distribution of the previous p observations in a way that leads to attractive properties such as ergodicity and fully known stationary distribution of p consecutive observations. Identification of the shocks requires that they are orthogonalized in all regimes. A suitable strategy is therefore to build on the one proposed by Lanne and Lütkepohl (2010) and Lanne et al. (2010) where a well known matrix decomposition is utilized to simulatenously diagonalize the (positive definite) error term covariance matrices.

Lanne and Lütkepohl (2010) point out that their identification strategy is a statistical one, but it allows to test for overidentifying constraints that lead structural shocks with economic interpretations. It turns out, however, that the structural shocks of a SGMVAR model can be uniquely identified by combining the simultaneous diagonalization of the covariance matrices with zero and sign constraints in way that is often less restrictive than in conventional SVAR models. While some of the constraints must be justified economically, some can also be validated statistically.

The rest of this paper is organized as follows. Section 2 defines the reduced form GMVAR model. In Section 3, the structural GMVAR model is first introduced. Then, identification

of the structural shocks and estimation of the model parameters are discussed. Section 4 discusses impulse response analysis and describes the generalized impulse response function (GIRF) (Koop et al., 1996). Appendices provide proofs for the stated propositions and a Monte Carlo algorithm for estimating the GIRF. Finally, we have accompanied this paper with the CRAN distributed R package "gmvarkit" (Virolainen, 2020a) which provides easy-to-use tools for estimating the models and applying the introduced methods.¹

2 Reduced form GMVAR model

To build theory and notation, consider first the reduced form GMVAR model introduced by Kalliovirta et al. (2016). Let y_t ($t = 1, 2, \dots$) be the d -dimensional time series of interest and \mathcal{F}_{t-1} denote the σ -algebra generated by random vectors $\{y_{t-j}, j > 0\}$. For a GMVAR model with M mixture components and autoregressive order p , we have

$$y_t = \sum_{m=1}^M s_{t,m}(\mu_{m,t} + u_{m,t}), \quad u_{m,t} \sim NID(0, \Omega_m) \quad (2.1)$$

$$\mu_{m,t} = \phi_{m,0} + \sum_{i=1}^p A_{m,i}y_{t-i}, \quad m = 1, \dots, M, \quad (2.2)$$

where $\phi_{m,0} \in \mathbb{R}^d$ is an intercept parameter, for each m the coefficient matrices $A_{m,i}$, $i = 1, \dots, p$, are assumed to satisfy the usual stability condition (given, for instance, in Kalliovirta et al., 2016, eq. (5)), and Ω_m are positive definite covariance matrices. The unobservable regime variables $s_{1,t}, \dots, s_{M,t}$ are such at each t exactly one of them takes the value one and the others take the value zero according to the conditional probabilities $P(s_{m,t} = 1 | \mathcal{F}_{t-1}) = \alpha_{m,t}$ that satisfy $\sum_{m=1}^M \alpha_{m,t} = 1$. The normally and independently distributed (NID) errors $u_{m,t}$ are assumed independent of \mathcal{F}_{t-1} , and conditional on \mathcal{F}_{t-1} , $s_{t,m}$ and $u_{m,t}$ are independent.

The definition (2.1)-(2.2) implies that at each t , the process generates an observations from one of its mixture components, a linear VAR process, that is randomly selected according to the

¹ The SGMVAR model is accommodated in gmvarkit from the version 1.20 onwards.

probabilities pointed by the mixing weights $\alpha_{m,t}$. Denoting $\mathbf{y}_{t-1} = (y_{t-1}, \dots, y_{t-p})$, the mixing weights are defined as (Kalliovirta et al., 2016, eq. (7))

$$\alpha_{m,t} = \frac{\alpha_m n_{dp}(\mathbf{y}_{t-1}; \mathbf{1}_p \otimes \mu_m, \Sigma_m)}{\sum_{n=1}^M \alpha_n n_{dp}(\mathbf{y}_{t-1}; \mathbf{1}_p \otimes \mu_n, \Sigma_n)}, \quad m = 1, \dots, M, \quad (2.3)$$

where $\alpha_1, \dots, \alpha_M$ are mixing weight parameters that satisfy $\sum_{i=1}^M \alpha_i = 1$ and $n_{dp}(\cdot; \mathbf{1}_p \otimes \mu_m, \Sigma_m)$ is the density function of the dp -dimensional normal distribution with mean $\mathbf{1}_p \otimes \mu_m$ and covariance matrix Σ_m . The symbol $\mathbf{1}_p$ denotes a p -dimensional vector of ones, \otimes is Kronecker product, $\mu_m = (I_d - \sum_{i=1}^p A_{m,i})^{-1} \phi_{m,0}$, and the covariance matrix Σ_m is given in Lütkepohl (2005), equation (2.1.39), but using the parameters of the m th component process.

The mixing weights are thus weighted ratios of the component process stationary densities corresponding to the previous p observations. Consequently, the researcher can associate specific characteristics to different regimes, so in addition the (generalized) impulse responses of the variables, responses of the mixing weights may also be of interest. The definition of the mixing weights also leads to attractive theoretical properties such as fully known stationary distribution of p consecutive observations and ergodicity of the process (Kalliovirta et al., 2016, Theorem 1). Specifically, the stationary distribution of the process $\mathbf{y}_t = (y_t, \dots, y_{t-p+1})$ is a mixture of dp -dimensional normal distributions that is characterized by the density

$$f(\mathbf{y}) = \sum_{m=1}^M \alpha_m n_{dp}(\mathbf{y}; \mathbf{1}_p \otimes \mu_m, \Sigma_m). \quad (2.4)$$

The knowledge of the stationary distribution is taken advantage of in the impulse response analysis in Section 4.

3 Structural GMVAR model

3.1 The model setup

Consider the GMVAR model defined in (2.1)-(2.2). We focus on the "B-model" setup and write the structural GMVAR model as

$$y_t = \sum_{m=1}^M s_{t,m} \left(\phi_{m,0} + \sum_{i=1}^p A_{m,i} y_{t-i} \right) + B_t e_t, \quad (3.1)$$

and

$$u_t \equiv B_t e_t = \begin{cases} u_{1,t} \sim N(0, \Omega_1) & \text{if } s_{t,1} = 1 & \text{(with probability } \alpha_{1,t}) \\ u_{2,t} \sim N(0, \Omega_2) & \text{if } s_{t,2} = 1 & \text{(with probability } \alpha_{2,t}) \\ \vdots & & \\ u_{M,t} \sim N(0, \Omega_M) & \text{if } s_{t,M} = 1 & \text{(with probability } \alpha_{M,t}) \end{cases} \quad (3.2)$$

where the probabilities are expressed conditionally on \mathcal{F}_{t-1} and e_t is an orthogonal structural error. Unlike in conventional SVAR analysis, the $(d \times d)$ "B-matrix" B_t , which governs the contemporaneous relations of the shocks, is time-varying. This enables to amplify a constant sized structural shock according to the conditional variance of the reduced form error which varies according to the mixing weights.

We have $\Omega_{u,t} \equiv \text{Cov}(u_t | \mathcal{F}_{t-1}) = \sum_{m=1}^M \alpha_{m,t} \Omega_m$, while the (conditional) covariance matrix of the structural errors $e_t = B_t^{-1} u_t$ (which have a mixture normal distribution and are not IID) is obtained as

$$\text{Cov}(e_t | \mathcal{F}_{t-1}) = \sum_{m=1}^M \alpha_{m,t} B_t^{-1} \Omega_m B_t'^{-1}. \quad (3.3)$$

The B-matrix B_t should thus be chosen so that the structural shocks are orthogonal regardless of which regime they come from. A suitable approach is thereby to build on the identification strategies proposed by Lanne and Lütkepohl (2010) and Lanne et al. (2010) where the error term covariance matrices are simultaneously diagonalized with a well known matrix decomposition.

Following Lanne et al. (2010), we decompose the error term covariance matrices as

$$\Omega_1 = WW' \quad \text{and} \quad \Omega_m = W\Lambda_mW', \quad m = 2, \dots, M, \quad (3.4)$$

where the diagonal of $\Lambda_m = \text{diag}(\lambda_{m1}, \dots, \lambda_{md})$, $\lambda_{mi} > 0$ ($i = 1, \dots, d$), contains the eigenvalues of the matrix $\Omega_m\Omega_1^{-1}$ and columns of the nonsingular W are the related eigenvectors. When $M = 2$, the decomposition (3.4) always exists (Muirhead, 1982, Theorem A9.9), but for $M \geq 3$ its existence requires that the matrices $\Omega_m\Omega_1^{-1}$ share the common eigenvectors in W . Whether imposing the structure (3.4) on the covariance matrices is appropriate with $M \geq 3$, can be evaluated by estimating the restricted and unrestricted SGMVAR models and conducting a likelihood ratio test, comparing values of information criteria, or examining quantile residual diagnostics (see Kalliovirta and Saikkonen, 2010), for example.

Lanne et al. (2010, Proposition 1) show that for a given ordering of the eigenvalues, W is unique apart from changing all signs in a column, as long as for all $i \neq j \in \{1, \dots, d\}$ there exists an $m \in \{2, \dots, M\}$ such that $\lambda_{mi} \neq \lambda_{mj}$. A locally unique B-matrix that amplifies a constant sized structural shock according to the conditional variance of the reduced form error is therefore obtained as

$$B_t = W(\alpha_{1,t}I_d + \sum_{m=2}^M \alpha_{m,t}\Lambda_m)^{1/2} \quad (3.5)$$

which simultaneously diagonalizes $\Omega_1, \dots, \Omega_M$, and $\Omega_{u,t}$ for each t so that $\text{Cov}(e_t|\mathcal{F}_{t-1}) = I_d$. Our specification of the B-matrix differs from Lanne et al. (2010) who assume that the instantaneous effects of the shocks are time-invariant (and specify $B_t = W$) but it extends the one in Lanne and Lütkepohl (2010) to accommodate time-varying mixing weights.

Even if all pairs of λ_{mi} are distinct for some m , unique identification of B_t requires deciding upon the ordering of λ_{mi} in the diagonals of Λ_m which fixes also the ordering of the columns of W . With an arbitrary ordering of the eigenvalues and eigenvectors, it is not guaranteed that the structural shocks have economic interpretations. Lanne and Lütkepohl (2010) and Lanne et al. (2010), however, demonstrate that after the statistical identification, overidentifying restrictions

that lead economically interpretable structural shocks can be tested and thus validated statistically. This method is well suited for the SGMVAR model as well. Furthermore, it turns out that globally unique identification of the structural shocks can often be achieved with less restrictive constraints than in conventional SVAR models. The following proposition gives sufficient and necessary conditions for global identification of the shocks when all pairs of λ_{mi} are distinct for some m .

Proposition 1. *Suppose $\Omega_1 = WW'$ and $\Omega_m = W\Lambda_m W'$, $m = 2, \dots, M$, where $\Lambda_m = \text{diag}(\lambda_{m1}, \dots, \lambda_{md})$, $\lambda_{mi} > 0$ ($i = 1, \dots, d$), contains the eigenvalues of $\Omega_m \Omega_1^{-1}$ in the diagonal and the columns of the nonsingular W are the related eigenvectors. Then, if (1) for all $i \neq j$ there exists an $m \in \{2, \dots, M\}$ such that $\lambda_{mi} \neq \lambda_{mj}$, (2) there is one (strict) sign constraint in each column of W , and (3) the columns of W are constrained in a way that their ordering cannot be changed, the above decomposition of $\Omega_1, \dots, \Omega_M$ is globally unique, provided that it exists.*

Notice that normalizing the diagonal of the B-matrix to be positive or negative is enough to fulfil condition (2) of Proposition (1). This kind of normalization is merely a matter of deciding upon whether positive or negative shocks are considered and therefore not restricting.

Because imposing zero or sign constraints on W equals to placing them on B_t , they may be justified economically. For example, if $d = 2$ and $\lambda_{m1} \neq \lambda_{m2}$ for some m , structural shocks can be uniquely identified by placing one (strict) negative and one (strict) positive sign constraint in the same row of B_t because reordering of the columns would violate the sign constraints. If the diagonal of B_t is normalized to be positive, the negative sign constraint becomes testable. Given that condition (1) holds, for $d = 3$ having one positive sign, one negative sign, and one zero constraint the same row of B_t , in addition to a single sign constraint in the column that has the zero constraint is enough for global identification of all the shocks. Moreover, because the zero constraint in the $d = 3$ example is overidentifying, it can be tested.

If $\lambda_{mi} = \lambda_{mj}$ for some $i \neq j$ and all $m \in \{2, \dots, M\}$ but they are placed on the first (or last) positions of Λ_m , the rest of the shocks with distinct λ_{mi} for some m are still globally identified as long as the conditions (2) and (3) of Proposition 1 are satisfied. Identification of the shocks with identical eigenvalues in all m requires stronger conditions, however. The following proposition

provides sufficient criteria for global identification of all the shocks when at most two eigenvalues are identical to each other in all m but there are possibly multiple pairs of such eigenvalues.

Proposition 2. *Consider the matrix decomposition of Proposition 1 and further suppose that $\lambda_{mi} = \lambda_{mj}$ for some $i \neq j$ and all m , but for all $l \neq k$ such that $l \notin \{i, j\}$ or $k \notin \{i, j\}$, $\lambda_{ml} \neq \lambda_{mk}$ for some m . Then, the decomposition of $\Omega_1, \dots, \Omega_M$ is globally unique if in addition to conditions (2) and (3) of Proposition 1, the following condition is satisfied:*

- (4) *each of the columns i, j of W (or equally B_t) such that $\lambda_{mi} = \lambda_{mj}$ for all m have at least one zero constraint where the other one does not and also have at least one (strict) sign constraint where the other one has a zero constraint.*

The result holds provided that the decomposition exists.

To exemplify, if $d = 4$, $\lambda_{m1} = \lambda_{m2}$ for all m , $\lambda_{m2} \neq \lambda_{m3}$ for some m , $\lambda_{m2} \neq \lambda_{m4}$ for some m , and $\lambda_{m3} \neq \lambda_{m4}$ for some m , the following constraints lead to global identification of all the shocks:

$$B_t = \begin{bmatrix} * & * & * & * \\ * & * & * & * \\ + & 0 & * & - \\ 0 & + & - & + \end{bmatrix} \text{ or } \begin{bmatrix} + & * & * & * \\ * & + & * & * \\ 0 & - & + & * \\ - & 0 & 0 & + \end{bmatrix} \text{ or } \begin{bmatrix} + & 0 & * & + \\ * & * & * & * \\ 0 & - & + & - \\ * & * & * & * \end{bmatrix} \quad (3.6)$$

and so on, where ” * ” signifies that the element is not constrained, ” + ” denotes strict positive and ” - ” a strict negative sign constraint, and ” 0 ” means that the element is constrained to zero. As is demonstrated above, the structural shocks can often be identified with less restrictive constraints than in a conventional SVAR model, even when some of the eigenvalues are identical for all covariance matrices.

The conditions of Proposition 2 are not in general necessary. If $d = 2$ and $\lambda_{m1} = \lambda_{m2}$ for all m , for instance, identification (of all regimes) with Cholesky decomposition leads to unique solution with less restrictive constraints than the ones required in Proposition 2. More generally, if all the eigenvalues are the same for each covariance matrix, then $\Omega_m = \lambda_{m1}\Omega_1$ and the necessary identification condition is the same as for the conventional SVAR model (which is given, for exam-

ple, in Lütkepohl, 2005, Section 9.1.2 for the B-model). If more than two eigenvalues are identical for all $m = 2, \dots, M$ but they are not all identical, it may still be possible to find less restrictive (or more suitable) conditions for identification of all the shocks than in a conventional SVAR model. Specifically, the idea utilized in the proof of Proposition 2 (presented in an Appendix) generalizes to larger numbers of identical eigenvalues, but the sufficient conditions become increasingly more complicated and are not presented for brevity. As a general remark, observe that constraining an element of B_t to be any constant other than zero is infeasible because all elements in the RHS of (3.5) are either zero or time varying due to the time-varying mixing weights.²

3.2 Estimation

The reduced form GMVAR model can be estimated by maximizing the exact or conditional likelihood function presented in Kalliovirta et al. (2016), equations (9) and (10). If $M = 2$, the structural GMVAR model is then obtained by simultaneously diagonalizing the error term covariance matrices, as discussed Section 3.1. However, should overidentifying restrictions be imposed on B_t or if $M \geq 3$, it is more convenient to reparametrize the model with W and Λ_m , $m = 2, \dots, M$, instead of $\Omega_1, \dots, \Omega_M$ and maximize the log-likelihood function subject to the new set of parameters and constraints.

When the model is reparametrized with W and Λ_m , they can be constrained in a way that leads to the desired restrictions in B_t . Specifically, the structure (3.5) implies that an element of B_t is constrained to zero (for all t) if and only if the corresponding element of W is also constrained to zero. Likewise, an element of B_t is constrained to have a specific sign if and only if an element of W is constrained to have the same sign. Constraining Λ_m does not have a role in the zero or sign constraints of B_t , but it might be appropriate to restrict some of its diagonal elements to be

² We have focused on the B-model where the structure is imposed on the contemporaneous relations of the shocks. Alternatively, one may consider the "A-model" setup in which the structure is placed on the contemporaneous relations of the observable variables that are governed by the "A-matrix" (see, e.g., Lütkepohl, 2005, Section 9.1.1). The A-model is obtained implicitly from the B-model (3.1) and (3.2) by defining the A-matrix as $A_t \equiv B_t^{-1}$ where B_t is given by (3.5). In practice, however, one needs to carefully derive how any specific constraint on A_t can be imposed by restricting W .

the same if their estimates are close to each other.

Maximizing the complex and highly multimodal log-likelihood function can be challenging in practice. Following Dorsey and Mayer (1995), Meitz et al. (2018a,b), and Virolainen (2020b,c), we employ a two-phase procedure where, in the first phase, a genetic algorithm is used to find starting values for a gradient based method which then, in the second phase, accurately converges to a nearby local maximum or saddle point. The genetic algorithm in the accompanying R package `gmvar` has been modified to improve its performance significantly, and it functions similarly to the one described in Virolainen (2020b) for the univariate GMAR (Kalliovirta et al., 2015), StMAR (Meitz et al., 2018a), and G-StMAR (Virolainen, 2020b) models. In order to obtain reliable results, a (sometimes large) number of estimation rounds should be performed, for which parallel computing may be taken use of.

4 Impulse response analysis

Following Koop et al. (1996) and Kilian and Lütkepohl (2017, Section 18.2.2), we consider the generalized impulse response function (GIRF) defined as

$$\text{GIRF}(n, \delta_j, \mathcal{F}_{t-1}) = \text{E}[y_{t+n} | \delta_j, \mathcal{F}_{t-1}] - \text{E}[y_{t+n} | \mathcal{F}_{t-1}], \quad (4.1)$$

where n is the chosen horizon, $\mathcal{F}_{t-1} = \sigma\{y_{t-j}, j > 0\}$ as before, the first term in the RHS is the expected realization of the process at time $t + n$ conditionally on a structural shock of magnitude $\delta_j \in \mathbb{R}$ in the j th variable at time t and the previous observations, and the second term in the RHS is the expected realization of the process conditionally on the previous observations only. GIRF thus expresses the expected difference in the future outcomes when the specific structural shock hits the system at time t as opposed to all shocks being random.

Due to the p -step Markov property of the SGMVAR model, conditioning on (the σ -algebra generated by) the p previous observations $\mathbf{y}_{t-1} \equiv (y_{t-1}, \dots, y_{t-p})$ is effectively the same as conditioning on \mathcal{F}_{t-1} at the time t and later. The history \mathbf{y}_{t-1} can be either fixed or random, but with

random history the GIRF becomes a random vector, however. Using fixed \mathbf{y}_{t-1} makes sense when one is interested in the effects of the shock in a particular point of time, whereas more general results are obtained by assuming that \mathbf{y}_{t-1} follows the stationary distribution of the process. If one is, on the other hand, concerned about a specific regime, \mathbf{y}_{t-1} can be assumed to follow the stationary distribution of the corresponding component model.

In practice, the GIRF and its distributional properties can be approximated with a Monte Carlo algorithm that generates independent realizations of the process and then takes the sample mean for point estimate. If \mathbf{y}_{t-1} is random and follows the distribution G , the GIRF should be estimated for different values of \mathbf{y}_{t-1} generated from G , and then the sample mean and sample quantiles can be taken to obtain the point estimate and confidence intervals. Such algorithm, which is adapted from Koop et al. (1996, pp. 135-136) and Kilian and Lütkepohl (2017, pp. 601-602), is given an Appendix.

Because the SGMVAR model allows to associate specific features or economic interpretations for different regimes, it might be interesting to also examine the effects of a structural shock to the mixing weights $\alpha_{m,t}$, $m = 1, \dots, M$. We then consider the related GIRFs $E[\alpha_{m,t+n} | \delta_j, \mathbf{y}_{t-1}] - E[\alpha_{m,t+n} | \mathbf{y}_{t-1}]$ for which point estimates and confidence intervals can be constructed similarly to (4.1).

References

- Bentarzi, M. and Djeddou, L. (2014). On mixture periodic vector autoregressive models. *Communications in Statistics - Simulation and Computation*, 43(10):2325–2352.
- Burgard, J., Neuenkirch, M., and Nöckel, M. (2019). State-dependent transmission of monetary policy in the euro area. *Journal of Money, Credit and Banking*, 51(7):2053–2070.
- Dorsey, R. and Mayer, W. (1995). Genetic algorithms for estimation problems with multiple optima, nondifferentiability, and other irregular features. *Journal of Business and Economic Statistics*, 13(1):53–66.

- Fong, P., Li, W., Yau, C., and Wong, C. (2010). On a mixture vector autoregressive model. *The Canadian Journal of Statistics*, 35(1):135–150.
- Kalliovirta, L., Meitz, M., and Saikkonen, P. (2015). A gaussian mixture autoregressive model for univariate time series. *Journal of Time Series Analysis*, 36(2):247–266.
- Kalliovirta, L., Meitz, M., and Saikkonen, P. (2016). Gaussian mixture vector autoregression. *Journal of Econometrics*, 192(2):465–498.
- Kalliovirta, L. and Saikkonen, P. (2010). Reliable residuals for multivariate nonlinear time series models. *Unpublished revision of HECER discussion paper No. 247*.
- Kilian, L. and Lütkepohl, H. (2017). *Structural Vector Autoregressive Analysis*. Cambridge University Press, Cambridge, 1st edition.
- Koop, G., Pesaran, M., and Potter, S. (1996). Impulse response analysis in nonlinear multivariate models. *Journal of Econometrics*, 74(1):119–147.
- Lanne, M. and Lütkepohl, H. (2010). Structural vector autoregressions with nonnormal residuals. *Journal of Business & Economic Statistics*, 28(1):159–168.
- Lanne, M., Lütkepohl, H., and Maciejowska, K. (2010). Structural vector autoregressions with markov switching. *Journal of Economic Dynamics and Control*, 34(2):121–131.
- Lütkepohl, H. (2005). *New Introduction to Multiple Time Series Analysis*. Springer, Berlin, 1st edition.
- Meitz, M., Preve, D., and Saikkonen, P. (2018a). A mixture autoregressive model based on student's t -distribution. *Unpublished working paper, available as arXiv:1805.04010*.
- Meitz, M., Preve, D., and Saikkonen, P. (2018b). *StMAR Toolbox: A MATLAB Toolbox for Student's t Mixture Autoregressive Models*.

Muirhead, R. (1982). *Aspects of Multivariate Statistical Theory*. John Wiley & Sons, Hoboken, New Jersey, 1nd edition.

Virolainen, S. (2020a). *gmvarKit: Estimate Gaussian Mixture Vector Autoregressive Model*. R package version 1.1.3, available at CRAN: <https://CRAN.R-project.org/package=gmvarKit>.

Virolainen, S. (2020b). A mixture autoregressive model based on gaussian and student's t -distributions. *Unpublished working paper, available as arXiv:2003.05221*.

Virolainen, S. (2020c). *uGMAR: Estimate Univariate Gaussian or Student's t Mixture Autoregressive Model*. R package version 3.2.5, available at CRAN: <https://CRAN.R-project.org/package=uGMAR>.

Appendix A Proofs

Let $\Omega_1, \dots, \Omega_M$ be positive definite covariance matrices. We consider the decomposition $\Omega_1 = WW'$ and $\Omega_m = W\Lambda_m W'$, $m = 2, \dots, M$, where $\Lambda_m = \text{diag}(\lambda_{m1}, \dots, \lambda_{md})$, $\lambda_{mi} > 0$ ($i = 1, \dots, d$), contains the eigenvalues of $\Omega_m \Omega_1^{-1}$ in the diagonal and the columns of the nonsingular W are the related eigenvectors. The decomposition always exists when $M = 2$ (see, e.g., Muirhead, 1982, Theorem A9.9) but not necessarily when $M \geq 3$. In the following, we assume the covariance matrices satisfy the decomposition.

A.1 Proof of Proposition 1

Repeating some of the proof in Lanne et al. (2010, p. 130, see also the proof of Theorem A9.9 in Muirhead, 1982) for convenience, suppose that we also have $\Omega_1 = MM'$ and $\Omega_m = M\Lambda_m M'$, $m = 2, \dots, M$, for some nonsingular ($d \times d$) matrix M . Because $M^{-1}WW'M'^{-1} = M^{-1}\Omega_1 M'^{-1} = I_d$, the matrix $Q' \equiv M^{-1}W$ is orthogonal, and hence, $M = WQ$ and $\Lambda_m Q = Q\Lambda_m$. It follows that $\lambda_{mi}q_{ij} = \lambda_{mj}q_{ij}$ where q_{ij} is the ij th element of Q . Thus, $q_{ij} = 0$ if $\lambda_{mi} \neq \lambda_{mj}$ for some m ,

indicating that Q is a diagonal matrix under condition (1) of Proposition 1. As the eigenvalues of an orthogonal matrix are ± 1 and the diagonal values of a diagonal matrix are its eigenvalues, it must be that $Q = \text{diag}(\pm 1, \dots, \pm 1)$.

The above result holds for any given Λ_m , $m = 2, \dots, M$, which are in general unique only up to ordering of the diagonal elements. Uniqueness, however, follows from condition (3) because it fixes an ordering for the eigenvectors (columns of W) and hence for the eigenvalues λ_{mi} as well. Finally, by condition (2) all signs in a column cannot be changed so $Q = \text{diag}(1, \dots, 1)$, implying that the decomposition globally unique. ■

A.2 Proof of Proposition 2

It is shown in the proof of Proposition 1 that any $(d \times d)$ matrix M that also satisfies $\Omega_1 = MM'$ and $\Omega_m = M\Lambda_m M'$, $m = 2, \dots, M$ can be presented as $M = WQ$ where Q is orthogonal and $q_{ij} = 0$ when $\lambda_{mi} \neq \lambda_{mj}$ for some m . Then observe that the j th column of WQ is a linear combination of the columns of W , with the multiplier of the i th column given by q_{ij} . It follows that if $\lambda_{mi} = \lambda_{mj}$ for some $i \neq j$ and all m , but for all $l \neq k$ such that $l \notin \{i, j\}$ or $k \notin \{i, j\}$, $\lambda_{ml} \neq \lambda_{mk}$ for some m , the i th and j th columns of WQ are linear combinations of the i th and j th columns of W . But if the j th column (of W and WQ) obeys a zero constraint where the i th column obeys a strict sign constraint, the multiplier q_{ij} must be zero (and similarly q_{ji} is as well). That is, under the conditions of Proposition 2, $q_{ij} = 0$ for all $i, j = 1, \dots, d$, and one may conclude global uniqueness of the decomposition with arguments similar to the proof of Proposition 1. ■

Appendix B Monte Carlo algorithm

We present a Monte Carlo algorithm that produces point estimates and with random initial value $\mathbf{y}_{t-1} = (y_{t-1}, \dots, y_{t-p})$ confidence intervals for the generalized impulse response function defined in (4.1). Our algorithm is adapted from Koop et al. (1996, pp. 135-136) and Kilian and Lütkepohl (2017, pp. 601-602). We assume that the history \mathbf{y}_{t-1} follows a known distribution G which

may be such that it produces a single outcome with probability one (corresponding to a fixed \mathbf{y}_{t-1}), or it can be the stationary distribution of the process or of a specific regime. In the following, $y_{t+n}^{(i)}(\delta_j, \mathbf{y}_{t-1})$ denotes a realization of the process at time $t+n$ conditional on the structural shock of magnitude δ_j in the j th variable hitting the system at time t and on the p observations $\mathbf{y}_{t-1} = (y_{t-1}, \dots, y_{t-p})$ preceding the time t , whereas $y_{t+n}^{(i)}(\mathbf{y}_{t-1})$ denotes an alternative realization conditional on the history \mathbf{y}_{t-1} only.

0. Decide the horizon N , the numbers of repetitions R_1 and R_2 , and the magnitude δ_j of the structural shock in the variable j of interest.
1. Draw initial value \mathbf{y}_{t-1} from G .
2. Draw $N+1$ realizations of a shock ε_t from $N(0, I_d)$. Also, draw an initial regime $m \in \{1, \dots, M\}$ according to the probabilities pointed by the mixing weights $\alpha_{1,t}, \dots, \alpha_{M,t}$ and compute the reduced form shock $u_t = W\Lambda_m^{1/2}\varepsilon_t$ where $\Lambda_1 = I_d$. Then, compute the structural shock $e_t = B_t^{-1}u_t$ and impose the magnitude δ_j on its j th element to obtain e_t^* . Finally, calculate the modified reduced form shock $u_t^* = B_t e_t^*$.³
3. Use the modified reduced form shock u_t^* and the rest N standard normal shocks ε_t obtained from step 2 to compute realizations $y_{t+n}^{(i)}(\delta_j, \mathbf{y}_{t-1})$ for $n = 0, 1, \dots, N$, iterating forward so that in each iteration the regime m that generates the observation is first drawn according to the probabilities pointed by the mixing weights. At $n = 0$, the initial regime and the modified reduced form shock u_t^* calculated from the structural shock in step 2 is used, and from $n = 1$ onwards the $n+1$ th shock ε_t is used to calculate the reduced form shock $u_{t+n} = W\Lambda_m^{1/2}\varepsilon_{t+n}$ where $\Lambda_1 = I_d$ and m is the selected regime.
4. Use the reduced form shock u_t and the rest N standard normal shocks ε_t obtained from step 2 to compute realizations $y_{t+n}^{(i)}(\mathbf{y}_{t-1})$ for $n = 0, 1, \dots, N$, so that the reduced form shock u_t

³ The independent standard normal shocks ε_t are introduced here to control random variation across the two sample paths $y_{t+n}^{(i)}(\delta_j, \mathbf{y}_{t-1})$ and $y_{t+n}^{(i)}(\mathbf{y}_{t-1})$.

(calculated in step 2) is used to compute the time $n = 0$ realization. Otherwise proceed similarly to the previous step.

5. Calculate $y_{t+n}^{(i)}(\delta_j, \mathbf{y}_{t-1}) - y_{t+n}^{(i)}(\mathbf{y}_{t-1})$.
6. Repeat steps 2-5 R_1 times and calculate the sample mean of $y_{t+n}^{(i)}(\delta_j, \mathbf{y}_{t-1}) - y_{t+n}^{(i)}(\mathbf{y}_{t-1})$ for $n = 0, 1, \dots, N$ to obtain an estimate of the $\text{GIRF}(n, \delta_j, \mathbf{y}_{t-1})$.
7. Repeat steps 1-6 R_2 times to obtain estimates of $\text{GIRF}(n, \delta_j, \mathbf{y}_{t-1})$ with different values of \mathbf{y}_{t-1} generated from the distribution G . Then take the sample mean and sample quantiles over the estimates to obtain point estimate and confidence intervals for the GIRF with random initial value.

Notice that if a fixed initial value \mathbf{y}_{t-1} is of interest, step 7 is redundant.