

# Shortened linear codes from APN and PN functions

Can Xiang, Chunming Tang and Cunsheng Ding

## Abstract

Linear codes generated by component functions of perfect nonlinear (PN for short) and almost perfect nonlinear (APN for short) functions and first-order Reed-Muller codes have been an object of intensive study by many coding theorists. In this paper, we investigate some binary shortened code of two families of linear codes from APN functions and some  $p$ -ary shortened codes from PN functions. The weight distributions of these shortened codes and the parameters of their duals are determined. The parameters of these binary codes and  $p$ -ary codes are flexible. Many of the codes presented in this paper are optimal or almost optimal in the sense that they meet some bound on linear codes. These results show high potential for shortening to be used in designing good codes.

## Index Terms

Linear code, shortened code, PN function, APN function,  $t$ -design

## I. INTRODUCTION

Let  $\text{GF}(q)$  denote the finite field with  $q = p^m$  elements, where  $p$  is a prime and  $m$  is a positive integer. An  $[v, k, d]$  linear code  $C$  over  $\text{GF}(q)$  is a  $k$ -dimensional subspace of  $\text{GF}(q)^v$  with minimum (Hamming) distance  $d$ . Let  $A_i$  denote the number of codewords with Hamming weight  $i$  in a code  $C$  of length  $v$ . The weight enumerator of  $C$  is defined by  $1 + A_1z + A_2z^2 + \cdots + A_vz^v$ . The sequence  $(1, A_1, \dots, A_v)$  is called the weight distribution of  $C$  and it is an important research topic in coding theory, as it contains crucial information to estimate the error correcting capability. Thus the study of the weight distribution attracts much attention in coding theory and much work focuses on the determination of the weight distributions of linear codes (see, for examples, [32], [34], [14], [15], [16], [17], [36], [45], [46], [39]). Denote by  $C^\perp$  and  $(A_0^\perp, A_1^\perp, \dots, A_v^\perp)$  the dual code of a linear code  $C$  and its weight distribution, respectively. It is known that the Pless power moments [28]

$$\sum_{i=0}^v i^t A_i = \sum_{i=0}^t (-1)^i A_i^\perp \left[ \sum_{j=i}^t j! S(t, j) q^{k-j} (q-1)^{j-i} \binom{v-i}{v-j} \right], \quad (1)$$

play an important role in calculating the weight distributions of linear codes, where  $A_0 = 1$ ,  $0 \leq t \leq v$  and  $S(t, j) = \frac{1}{j!} \sum_{i=0}^j (-1)^{j-i} \binom{j}{i} i^t$ . A code  $C$  is said to be a  $t$ -weight code if the number of nonzero  $A_i$  in the sequence  $(A_1, A_2, \dots, A_v)$  is equal to  $t$ . We call a  $[v, k, d]$  code *distance-optimal* if no  $[v, k, d+1]$  code exists and *dimension-optimal* if no  $[v, k+1, d]$  code exists. An  $[v, k, d]$  code is said to be length-optimal if there is no  $[v', k, d]$  code exists with  $v' < v$ . A code is said to be optimal if it is distance-optimal, dimension-optimal and length-optimal.

Let  $C$  be a  $[v, k, d]$  linear code over  $\text{GF}(q)$  and  $T$  a set of  $t$  coordinate positions in  $C$ . We use  $C^T$  to denote the code obtained by puncturing  $C$  on  $T$ , which is called the *punctured code* of  $C$  on  $T$ . Let

The research of C. Xiang was supported by the National Natural Science Foundation of China under grant number 11701187. The research of C. Tang was supported by National Natural Science Foundation of China under grant number 11871058 and China West Normal University (14E013, CXTD2014-4 and the Meritocracy Research Funds). The research of C. Ding was supported by the Hong Kong Research Grants Council, Project No. 16301020.

C. Xiang is with the College of Mathematics and Informatics, South China Agricultural University, Guangzhou, Guangdong 510642, China (email: cxiangxiang@hotmail.com).

C. Tang is with School of Mathematics and Information, China West Normal University, Nanchong, Sichuan 637002, China, and also with the Department of Computer Science and Engineering, The Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong, China (email: tangchunmingmath@163.com).

C. Ding is with the Department of Computer Science and Engineering, The Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong, China (email: cding@ust.hk).

$\mathcal{C}(T)$  be the set of codewords which are  $\mathbf{0}$  on  $T$ . We now puncture  $\mathcal{C}(T)$  on  $T$ , and obtain a linear code  $\mathcal{C}_T$ , which is called the *shortened code* of  $\mathcal{C}$  on  $T$ . The following property plays an important role in determining the parameters of the punctured and shortened codes of  $\mathcal{C}$  in [28, Theorem 1.5.7].

**Lemma 1.** [28] *Let  $\mathcal{C}$  be a  $[\nu, k, d]$  linear code over  $\text{GF}(q)$  and  $d^\perp$  the minimum distance of  $\mathcal{C}^\perp$ . Let  $T$  be any set of  $t$  coordinate positions. Then*

- $(\mathcal{C}_T)^\perp = (\mathcal{C}^\perp)^T$  and  $(\mathcal{C}^T)^\perp = (\mathcal{C}^\perp)_T$ .
- If  $t < \min\{d, d^\perp\}$ , then the codes  $\mathcal{C}_T$  and  $\mathcal{C}^T$  have dimension  $k - t$  and  $k$ , respectively.

It is worth noting that the shortening and puncturing technologies are two important approaches to constructing new linear codes. Very recently, Tang et al. obtained some ternary linear codes with few weights by shortening and puncturing a class of ternary codes in [37]. Afterwards, they presented a general theory for punctured and shortened codes of linear codes supporting  $t$ -designs and generalized Assmus-Mattson theorem in [38]. Some linear codes and  $t$ -designs can be obtained and their parameters can be also derived. However, till now few results on constructing punctured and shortened codes have been done and it is in general hard to determine their weight distributions. Motivated by this fact, we obtain some shortened codes of linear codes from almost perfect nonlinear (APN for short) and perfect nonlinear (PN for short) functions, and determine their parameters. Some of these codes are optimal or almost optimal.

The rest of this paper is arranged as follows. Section II introduces some notation and results related to group characters, gauss sums,  $t$ -designs and linear codes from APN and PN functions. Section III gives some general results on shortened codes. Section IV investigates some shortened codes of binary linear codes from APN functions. Section V studies some shortened codes of two classes of special linear codes from PN functions. Section VI concludes this paper and makes concluding remarks.

## II. PRELIMINARIES

In this section, we briefly recall some results on group characters, Gauss sums,  $t$ -designs, and linear codes from APN and PN functions. These results will be used later in this paper. We begin this section by fixing some notations throughout this paper.

- $p^* = (-1)^{(p-1)/2}p$ .
- $\zeta_p = e^{\frac{2\pi\sqrt{-1}}{p}}$  is the primitive  $p$ -th root of unity.
- $\text{GF}(q)^* = \text{GF}(q) \setminus \{0\}$ .
- $\text{Tr}_{q/p}$  is the trace function from  $\text{GF}(q)$  to  $\text{GF}(p)$ .
- SQ and NSQ denote the set of all squares and nonsquares in  $\text{GF}(p)^*$ , respectively.
- $\eta$  and  $\bar{\eta}$  are the quadratic characters of  $\text{GF}(q)^*$  and  $\text{GF}(p)^*$ , respectively. We extend these quadratic characters by letting  $\eta(0) = 0$  and  $\bar{\eta}(0) = 0$ .

### A. Group characters and Gauss sums

An additive character of  $\text{GF}(q)$  is a nonzero function  $\chi$  from  $\text{GF}(q)$  to the set of nonzero complex numbers such that  $\chi(x+y) = \chi(x)\chi(y)$  for any pair  $(x, y) \in \text{GF}(q)^2$ . For each  $b \in \text{GF}(q)$ , the function

$$\chi_b(c) = \zeta_p^{\text{Tr}(bc)} \quad \text{for all } c \in \text{GF}(q) \quad (2)$$

defines an additive character of  $\text{GF}(q)$ . When  $b = 0$ ,  $\chi_0(c) = 1$  for all  $c \in \text{GF}(q)$ , and  $\chi_0$  is called the *trivial additive character* of  $\text{GF}(q)$ . The character  $\chi_1$  in (2) is called the *canonical additive character* of  $\text{GF}(q)$ . It is well known that every additive character of  $\text{GF}(q)$  can be written as  $\chi_b(x) = \chi_1(bx)$  [30, Theorem 5.7]. The orthogonality relation of additive characters is given by

$$\sum_{x \in \text{GF}(q)} \chi_1(ax) = \begin{cases} q & \text{for } a = 0, \\ 0 & \text{for } a \in \text{GF}(q)^*. \end{cases}$$

The Gauss sum  $G(\eta, \chi_1)$  over  $\text{GF}(q)$  is defined by

$$G(\eta, \chi_1) = \sum_{c \in \text{GF}(q)^*} \eta(c) \chi_1(c) = \sum_{c \in \text{GF}(q)} \eta(c) \chi_1(c) \quad (3)$$

and the Gauss sum  $G(\bar{\eta}, \bar{\chi}_1)$  over  $\text{GF}(p)$  is defined by

$$G(\bar{\eta}, \bar{\chi}_1) = \sum_{c \in \text{GF}(p)^*} \bar{\eta}(c) \bar{\chi}_1(c) = \sum_{c \in \text{GF}(p)} \bar{\eta}(c) \bar{\chi}_1(c), \quad (4)$$

where  $\bar{\chi}_1$  is the canonical additive character of  $\text{GF}(p)$ .

The following four lemmas are proved in [30, Theorem 5.15, Theorem 5.33, Corollary 5.35] and [16, Lemma 7], respectively.

**Lemma 2.** [30] *Let  $q = p^m$  and  $p$  be an odd prime. Then*

$$\begin{aligned} G(\eta, \chi_1) &= (-1)^{m-1} (\sqrt{-1})^{(\frac{p-1}{2})^2 m} \sqrt{q} \\ &= \begin{cases} (-1)^{m-1} \sqrt{q} & \text{for } p \equiv 1 \pmod{4}, \\ (-1)^{m-1} (\sqrt{-1})^m \sqrt{q} & \text{for } p \equiv 3 \pmod{4}. \end{cases} \end{aligned}$$

and

$$G(\bar{\eta}, \bar{\chi}_1) = \sqrt{-1}^{(\frac{p-1}{2})^2} \sqrt{p} = \sqrt{p^*}.$$

**Lemma 3.** [30] *Let  $\chi$  be a nontrivial additive character of  $\text{GF}(q)$  with  $q$  odd, and let  $f(x) = a_2 x^2 + a_1 x + a_0 \in \text{GF}(q)[x]$  with  $a_2 \neq 0$ . Then*

$$\sum_{c \in \text{GF}(q)} \chi(f(c)) = \chi(a_0 - a_1^2 (4a_2)^{-1}) \eta(a_2) G(\eta, \chi).$$

**Lemma 4.** [30] *Let  $\chi_b$  be a nontrivial additive character of  $\text{GF}(q)$  with  $q$  even and  $f(x) = a_2 x^2 + a_1 x + a_0 \in \text{GF}(q)[x]$ , where  $b \in \text{GF}(q)^*$ . Then*

$$\sum_{c \in \text{GF}(q)} \chi_b(f(c)) = \begin{cases} \chi_b(a_0) q & \text{if } a_2 = b a_1^2, \\ 0 & \text{otherwise.} \end{cases}$$

**Lemma 5.** [16] *If  $m \geq 2$  is even, then  $\eta(y) = 1$  for each  $y \in \text{GF}(p)^*$ . If  $m \geq 1$  is odd, then  $\eta(y) = \bar{\eta}(y)$  for each  $y \in \text{GF}(p)$ .*

Let  $e$  be a positive integer and  $(a, b) \in \text{GF}(q)^2$ , define the exponential sum

$$S_e(a, b) = \sum_{x \in \text{GF}(q)} \chi_1(ax^{p^e+1} + bx). \quad (5)$$

Then we have the following results in [12], [8], [9] and [43].

**Lemma 6.** [12] *Let  $e$  be a positive integer and  $m$  be even with  $\gcd(m, e) = 1$ . Let  $p = 2$ ,  $q = 2^m$  and  $a \in \text{GF}(q)^*$ . Then*

$$S_e(a, 0) = \begin{cases} (-1)^{\frac{m}{2}} 2^{\frac{m}{2}} & \text{if } a \neq \alpha^{3t} \text{ for any } t, \\ -(-1)^{\frac{m}{2}} 2^{\frac{m}{2}+1} & \text{if } a = \alpha^{3t} \text{ for some } t, \end{cases}$$

where  $\alpha$  is a generator of  $\text{GF}(q)^*$ .

**Lemma 7.** [8] *Let  $e, h$  be positive integers and  $m$  be even with  $\gcd(m, e) = 1$ . Let  $p = 2$ ,  $q = 2^m$  and  $a \in \text{GF}(q)^*$ . Then*

$$\sum_{b \in \text{GF}(q)^*} (S_e(a, b))^h = \begin{cases} (2^m - 1) 2^{\frac{m}{2} \cdot h} & \text{if } h \text{ is even and } a \neq \alpha^{3t} \text{ for any } t, \\ (2^{m-2} - 1) 2^{(\frac{m}{2}+1) \cdot h} & \text{if } h \text{ is even and } a = \alpha^{3t} \text{ for some } t, \end{cases}$$

where  $\alpha$  is a generator of  $\text{GF}(q)^*$ .

**Lemma 8.** [9] Let  $p$  be an odd prime,  $q = p^m$ , and  $e$  be any positive integer such that  $m/\text{gcd}(m, e)$  is odd. Suppose  $a \in \text{GF}(q)^*$  and  $b \in \text{GF}(q)^*$ . Let  $x_{a,b}$  be the unique solution of the equation

$$a^{p^e} x^{p^{2e}} + ax + b^{p^e} = 0.$$

Then

$$S_e(a, b) = \begin{cases} (-1)^{m-1} \sqrt{q} \eta(-a) \chi_1(-ax_{a,b}^{p^e+1}), & \text{if } p \equiv 1 \pmod{4}, \\ (-1)^{m-1} i^{3m} \sqrt{q} \eta(-a) \chi_1(-ax_{a,b}^{p^e+1}), & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

**Lemma 9.** [43] Let  $p$  be an odd prime,  $q = p^m$ , and  $e$  be any positive integer such that  $m/\text{gcd}(m, e)$  is odd. Suppose  $a \in \text{GF}(q)^*$  and  $b \in \text{GF}(q)^*$ . Let  $x_{a,b}$  be the unique solution of the equation  $a^{p^e} x^{p^{2e}} + ax + b^{p^e} = 0$  and  $\Delta = \sum_{c \in \text{GF}(p)^*} S_e(ac, bc)$ . Then we have the following results.

- If  $m$  is odd, then

$$\Delta = \begin{cases} 0, & \text{if } \text{Tr}_{q/p}(a(x_{a,b})^{p^e+1}) = 0, \\ \eta(a) \eta(\text{Tr}_{q/p}(a(x_{a,b})^{p^e+1})) \sqrt{q} \sqrt{p^*}, & \text{if } p \equiv 1 \pmod{4} \text{ and } \text{Tr}_{q/p}(a(x_{a,b})^{p^e+1}) \neq 0, \\ \eta(a) \eta(\text{Tr}_{q/p}(a(x_{a,b})^{p^e+1})) i^{3m} \sqrt{q} \sqrt{p^*}, & \text{if } p \equiv 3 \pmod{4} \text{ and } \text{Tr}_{q/p}(a(x_{a,b})^{p^e+1}) \neq 0. \end{cases}$$

- If  $m$  is even, then

$$\Delta = \begin{cases} -(p-1) \eta(a) \sqrt{q}, & \text{if } p \equiv 1 \pmod{4} \text{ and } \text{Tr}_{q/p}(a(x_{a,b})^{p^e+1}) = 0, \\ \eta(a) \sqrt{q}, & \text{if } p \equiv 1 \pmod{4} \text{ and } \text{Tr}_{q/p}(a(x_{a,b})^{p^e+1}) \neq 0, \\ -i^m (p-1) \eta(a) \sqrt{q}, & \text{if } p \equiv 3 \pmod{4} \text{ and } \text{Tr}_{q/p}(a(x_{a,b})^{p^e+1}) = 0, \\ i^m \eta(a) \sqrt{q}, & \text{if } p \equiv 3 \pmod{4} \text{ and } \text{Tr}_{q/p}(a(x_{a,b})^{p^e+1}) \neq 0, \end{cases}$$

**Lemma 10.** [43] Let  $p$  be an odd prime,  $m$  and  $e$  be positive integers such that  $m/\text{gcd}(m, e)$  is odd. Let  $q = p^m$ . Define

$$\hat{N}_0(a, b) = \#\{x \in \text{GF}(q) : \text{Tr}_{q/p}(ax^{p^e+1} + bx) = 0\}.$$

Then we have the following results.

- If  $a = 0$  and  $b = 0$ , then  $\hat{N}_0(a, b) = q$ .
- If  $a = 0$  and  $b \neq 0$ , then  $\hat{N}_0(a, b) = p^{m-1}$ .
- If  $a \neq 0$  and  $b = 0$ , then

$$\hat{N}_0(a, b) = \begin{cases} p^{m-1}, & \text{if } m \text{ is odd,} \\ \frac{1}{p} (q - (p-1) \eta(a) \sqrt{q}), & \text{if } m \text{ is even and } p \equiv 1 \pmod{4}, \\ \frac{1}{p} (q - i^m (p-1) \eta(a) \sqrt{q}), & \text{if } m \text{ is even and } p \equiv 3 \pmod{4}. \end{cases}$$

- If  $a \neq 0$  and  $b \neq 0$ , then  $\hat{N}_0(a, b) = \frac{1}{p} (q + \Delta)$ , where  $R$  was given in Lemma 9.

## B. $t$ -designs and related results

Let  $k$ ,  $t$  and  $v$  be positive integers with  $1 \leq t \leq k \leq v$ . Let  $\mathcal{P}$  be a set of  $v \geq 1$  elements, and let  $\mathcal{B}$  be a set of  $k$ -subsets of  $\mathcal{P}$ . The *incidence structure*  $\mathbb{D} = (\mathcal{P}, \mathcal{B})$  is said to be a  $t$ - $(v, k, \lambda)$  *design* if every  $t$ -subset of  $\mathcal{P}$  is contained in exactly  $\lambda$  elements of  $\mathcal{B}$ . The elements of  $\mathcal{P}$  are called points, and those of  $\mathcal{B}$  are referred to as blocks. We usually use  $b$  to denote the number of blocks in  $\mathcal{B}$ . A  $t$ -design is called *simple* if  $\mathcal{B}$  has no repeated blocks. A  $t$ -design is called *symmetric* if  $v = b$  and *trivial* if  $k = t$  or  $k = v$ . When  $t \geq 2$  and  $\lambda = 1$ , a  $t$ -design is called a *Steiner system* and traditionally denoted by  $S(t, k, v)$ .

Linear codes and  $t$ -designs are companions. A  $t$ -design  $\mathbb{D} = (\mathcal{P}, \mathcal{B})$  induces a linear code over  $\text{GF}(p)$  for any prime  $p$ . Let  $\mathcal{P} = \{p_1, \dots, p_v\}$ . For any block  $B \in \mathcal{B}$ , the *characteristic vector* of  $B$  is defined by the vector  $\mathbf{c}_B = (c_1, \dots, c_v) \in \{0, 1\}^v$ , where

$$c_i = \begin{cases} 1, & \text{if } p_i \in B, \\ 0, & \text{if } p_i \notin B. \end{cases}$$

For a prime  $p$ , a *linear code*  $C_p(\mathbb{D})$  over the prime field  $\text{GF}(p)$  from the design  $\mathbb{D}$  is spanned by the characteristic vectors of blocks of  $\mathbb{B}$ , which is the subspace  $\text{Span}\{\mathbf{v}_B : B \in \mathcal{B}\}$  of the vector space  $\text{GF}(p)^v$ . Linear codes  $C_p(\mathbb{D})$  from designs  $\mathbb{D}$  have been studied and documented in the literature (see, for examples, [1], [17], [40], [41]).

On the other hand, a linear code  $C$  may induce a  $t$ -design under certain conditions, which is formed by supports of codewords of a fixed Hamming weight in  $C$ . Let  $\mathcal{P}(C)$  be the set of the coordinate positions of  $C$ , where  $\#\mathcal{P}(C) = v$  is the length of  $C$ . For a codeword  $\mathbf{c} = (c_i)_{i \in \mathcal{P}(C)}$  in  $C$ , the *support* of  $\mathbf{c}$  is defined by

$$\text{Supp}(\mathbf{c}) = \{i : c_i \neq 0, i \in \mathcal{P}(C)\}.$$

Let  $\mathcal{B}_w(C) = \{\text{Supp}(\mathbf{c}) : wt(\mathbf{c}) = w \text{ and } \mathbf{c} \in C\}$ . For some special  $C$ ,  $(\mathcal{P}(C), \mathcal{B}_w(C))$  is a  $t$ -design. In this way, many  $t$ -designs are derived from linear codes ( see, for examples, [1], [18], [19], [20], [25], [26], [29], [35], [40], [41]). A major approach to constructing  $t$ -designs from linear codes is the use of linear codes with  $t$ -homogeneous or  $t$ -transitive automorphism groups(see [14, Theorem 4.18]). Another major approach to constructing  $t$ -designs from codes is the use of the Assmus-Mattson Theorem [3], [28]. The following Assmus-Mattson Theorem for constructing simple  $t$ -designs is given in [2].

**Theorem 11.** *Let  $C$  be a linear code over  $\text{GF}(q)$  with length  $v$  and minimum weight  $d$ . Let  $C^\perp$  with minimum weight  $d^\perp$  denote the dual code of  $C$ . Let  $t$  ( $1 \leq t < \min\{d, d^\perp\}$ ) be an integer such that there are at most  $d^\perp - t$  weights of  $C$  in the range  $\{1, 2, \dots, v - t\}$ . Then the following holds:*

- $(\mathcal{P}(C), \mathcal{B}_k(C))$  is a simple  $t$ -design provided that  $A_k \neq 0$  and  $d \leq k \leq w$ , where  $w$  is defined to be the largest integer satisfying  $w \leq v$  and

$$w - \left\lfloor \frac{w+q-2}{q-1} \right\rfloor < d.$$

- $(\mathcal{P}(C^\perp), \mathcal{B}_k(C^\perp))$  is a simple  $t$ -design provided that  $A_k^\perp \neq 0$  and  $d^\perp \leq k \leq w^\perp$ , where  $w^\perp$  is defined to be the largest integer satisfying  $w^\perp \leq v$  and

$$w^\perp - \left\lfloor \frac{w^\perp+q-2}{q-1} \right\rfloor < d^\perp.$$

We will need the following results on the punctured and shortened codes of  $C$  in [38, Lemma 3.1, Theorem 3.2] .

**Lemma 12.** [38] *Let  $C$  be a linear code of length  $v$  and minimum distance  $d$  over  $\text{GF}(q)$  and  $d^\perp$  the minimum distance of  $C^\perp$ . Let  $t$  and  $k$  be two positive integers with  $0 < t < \min\{d, d^\perp\}$  and  $1 \leq k \leq v - t$ . Let  $T$  be a set of  $t$  coordinate positions in  $C$ . Suppose that  $(\mathcal{P}(C), \mathcal{B}_i(C))$  is a  $t$ -design for all  $i$  with  $k \leq i \leq k + t$ . Then*

$$A_k(C^T) = \sum_{i=0}^t \frac{\binom{v-t}{k} \binom{k+i}{t} \binom{t}{i}}{\binom{v-t}{k-t+i} \binom{v}{t}} A_{k+i}(C).$$

**Theorem 13.** [38] *Let  $C$  be a  $[v, \bar{k}, d]$  linear code over  $\text{GF}(q)$  and  $d^\perp$  be the minimum distance of  $C^\perp$ . Let  $t$  be a positive integer with  $0 < t < \min\{d, d^\perp\}$ . Let  $T$  be a set of  $t$  coordinate positions in  $C$ . Suppose that  $(\mathcal{P}(C), \mathcal{B}_i(C))$  is a  $t$ -design for any  $i$  with  $d \leq i \leq v - t$ . Then the shortened code  $C_T$  is a linear*

code of length  $v-t$  and dimension  $\bar{k}-t$ . The weight distribution  $(A_k(C_T))_{k=0}^{v-t}$  of  $C_T$  is independent of the specific choice of the elements in  $T$ . Specifically,

$$A_k(C_T) = \frac{\binom{k}{t} \binom{v-t}{k}}{\binom{v}{t} \binom{v-t}{k-t}} A_k(C).$$

### C. Linear codes from APN and PN functions

Let  $m, \tilde{m}$  be two positive integers with  $m \geq \tilde{m}$  and  $F$  be a mapping from  $\text{GF}(p^m)$  to  $\text{GF}(p^{\tilde{m}})$ . Define

$$\delta_F = \max\{\delta_F(a, b) : a \in \text{GF}(p^m)^*, b \in \text{GF}(p^{\tilde{m}})\},$$

where  $\delta_F(a, b) = \#\{x \in \text{GF}(p^m) : F(x+a) - F(x) = b\}$ ,  $a \in \text{GF}(p^m)$  and  $b \in \text{GF}(p^{\tilde{m}})$ . The function  $F(x)$  is called PN function if  $\delta_F = p^{m-\tilde{m}}$ , and it is called APN function if  $m = \tilde{m}$  and  $\delta_F = 2$ . From the above definition one immediately has that  $F(x)$  is PN if and only if  $F(x+a) - F(x)$  is balanced for each  $a \in \text{GF}(p^m)^*$ . Currently, all known PN and APN functions over  $\text{GF}(p^m)$  can be summarized in [14], [4], [10], [11], [13], [22], [44], [7], [6]. It is known that PN and APN functions are very important functions to constructing linear codes with good parameters (see, for examples, [43], [5], [33], [42]).

Let  $q = p^m$  and  $C$  denote the linear code of length  $q$  as follows:

$$C = \left\{ \left( (\text{Tr}_{q/p}(af(x) + bx + c))_{x \in \text{GF}(q)} : a, b, c \in \text{GF}(q) \right) \right\}, \quad (6)$$

where  $f(x)$  is a polynomial over  $\text{GF}(q)$ . Then we can regard  $\text{GF}(q)$  as the set of the coordinate positions  $\mathcal{P}(C)$  of  $C$ . It is known that  $C$  has dimension  $2m+1$  and the weight distribution in Table I when  $p=2$ ,  $m \geq 5$  is odd and  $f(x) = x^s$  is an APN function, where  $s$  takes the following values [14].

- $s = 2^e + 1$ , where  $\gcd(e, m) = 1$  and  $e$  is a positive integer.
- $s = 2^{2e} - 2^e + 1$ , where  $e$  is a positive integer and  $\gcd(e, m) = 1$ .
- $s = 2^{(m-1)/2} + 3$ .
- $s = 2^{(m-1)/2} + 2^{(m-1)/4} - 1$ , where  $m \equiv 1 \pmod{4}$ .
- $s = 2^{(m-1)/2} + 2^{(3m-1)/4} - 1$ , where  $m \equiv 3 \pmod{4}$ .

When  $f(x) = x^{2^e+1}$ ,  $p=2$  and  $m \geq 4$  is even with  $\gcd(e, m) = 1$ , the code  $C$  defined in (6) has dimension  $2m+1$  and the weight distribution in Table II [14].

TABLE I  
THE WEIGHT DISTRIBUTION OF  $C$  FOR  $m$  ODD

Weight	Multiplicity
0	1
$2^{m-1} - 2^{(m-1)/2}$	$(2^m - 1)2^{m-1}$
$2^{m-1}$	$(2^m - 1)(2^{m+1} - 2^m + 2)$
$2^{m-1} + 2^{(m-1)/2}$	$(2^m - 1)2^{m-1}$
$2^m$	1

It is known that the code  $C$  defined in (6) has dimension  $2m+1$  and a few weights when  $p$  is an odd prime and  $f(x) = x^s$  is a PN function. If  $s$  takes the following values [14], [31]

- $s = 2$ ,
- $s = p^e + 1$ , where  $m/\gcd(m, e)$  is odd.
- $s = (3^e + 1)/2$ , where  $p = 3$ ,  $e$  is odd and  $\gcd(m, e) = 1$ ,

then  $f(x) = x^s$  is a PN and also planar function,  $\text{Tr}_{q/p}(\beta f(x))$  is a weakly regular bent function [23], [27] for any  $\beta \in \text{GF}(q)^*$ , and the code  $C$  defined in (6) has four or six weights [31].

Let  $f(x)$  be a function from  $\text{GF}(q)$  to  $\text{GF}(p)$ , the Walsh transform of  $f$  at a point  $\beta \in \text{GF}(q)$  is defined by

$$\mathcal{W}_f(\beta) = \sum_{x \in \text{GF}(q)} \zeta_p^{f(x) - \text{Tr}_{q/p}(\beta x)}.$$

TABLE II  
THE WEIGHT DISTRIBUTION OF  $\mathcal{C}$  FOR  $m$  EVEN

Weight	Multiplicity
0	1
$2^{m-1} - 2^{m/2}$	$(2^m - 1)2^{m-2}/3$
$2^{m-1} - 2^{(m-2)/2}$	$(2^m - 1)2^{m+1}/3$
$2^{m-1}$	$2(2^m - 1)(2^{m-2} + 1)$
$2^{m-1} + 2^{(m-2)/2}$	$(2^m - 1)2^{m+1}/3$
$2^{m-1} + 2^{m/2}$	$(2^m - 1)2^{m-2}/3$
$2^m$	1

The function  $f(x)$  is said to be a  $p$ -ary bent function, if  $|\mathcal{W}_f(\beta)| = p^{m/2}$  for any  $\beta \in \mathbb{F}_q$ . A bent function  $f(x)$  is weakly regular if there exists a complex  $u$  with unit magnitude satisfying  $\mathcal{W}_f(\beta) = u p^{m/2} \zeta_p^{f^*(\beta)}$  for some function  $f^*(x)$ . Such function  $f^*(x)$  is called the dual of  $f(x)$ . A weakly regular bent function  $f(x)$  satisfies

$$\mathcal{W}_f(\beta) = \varepsilon \sqrt{p^*}^m \zeta_p^{f^*(\beta)},$$

where  $\varepsilon = \pm 1$  is called the sign of the Walsh Transform of  $f(x)$ . Let  $\mathcal{RF}$  be the set of  $p$ -ary weakly regular bent functions with the following two properties:

- $f(0) = 0$ ; and
- $f(ax) = a^h f(x)$  for any  $a \in \text{GF}(p)^*$  and  $x \in \text{GF}(q)$ , where  $h$  is a positive even integer with  $\gcd(h-1, p-1) = 1$ .

We will need the following results on  $p$ -ary weakly regular bent functions in [39].

**Lemma 14.** [39] Let  $\beta \in \text{GF}(q)^*$  and  $f(x) \in \mathcal{RF}$  with  $\mathcal{W}_f(0) = \varepsilon \sqrt{p^*}^m$ . Define

$$N_{f,\beta} = \#\{x \in \text{GF}(q) : f(x) = 0 \text{ and } \text{Tr}_{q/p}(\beta x) = 0\}.$$

If  $f^*(\beta) = 0$ , then

$$N_{f,\beta} = \begin{cases} p^{m-2} + \varepsilon \bar{\eta}^{m/2} (-1)(p-1)p^{(m-2)/2}, & \text{if } m \text{ is even;} \\ p^{m-2}, & \text{if } m \text{ is odd.} \end{cases}$$

**Lemma 15.** [39] Let  $\beta \in \text{GF}(q)^*$  and  $f(x) \in \mathcal{RF}$  with  $\mathcal{W}_f(0) = \varepsilon \sqrt{p^*}^m$ . Let

$$N_{sq,\beta} = \#\{x \in \text{GF}(q) : f(x) \in SQ \text{ and } \text{Tr}_{q/p}(\beta x) = 0\},$$

and

$$N_{nsq,\beta} = \#\{x \in \text{GF}(q) : f(x) \in NSQ \text{ and } \text{Tr}_{q/p}(\beta x) = 0\}.$$

We have the following results.

- If  $m$  is even and  $f^*(\beta) = 0$ , then

$$N_{sq,\beta} = N_{nsq,\beta} = \frac{p-1}{2} \left( p^{m-2} - \varepsilon \bar{\eta}^{m/2} (-1)p^{(m-2)/2} \right).$$

- If  $m$  is odd and  $f^*(\beta) = 0$ , then

$$N_{sq,\beta} = \frac{p-1}{2} \left( p^{m-2} + \varepsilon \sqrt{p^*}^{m-1} \right)$$

and

$$N_{nsq,\beta} = \frac{p-1}{2} \left( p^{m-2} - \varepsilon \sqrt{p^*}^{m-1} \right).$$

### III. SHORTENED BINARY LINEAR CODES WITH SPECIAL WEIGHT DISTRIBUTIONS

In this section, we give some general results on the shortened codes of linear codes with the weight distributions in Tables I and II.

Let  $T$  be a set of  $t$  coordinate positions in  $\mathcal{C}$  (i.e.,  $T$  is a  $t$ -subset of  $\mathcal{P}(\mathcal{C})$ ). Define

$$\Lambda_{T,w}(\mathcal{C}) = \{\text{Supp}(\mathbf{c}) : wt(\mathbf{c}) = w, \mathbf{c} \in \mathcal{C} \text{ and } T \subseteq \text{Supp}(\mathbf{c})\}.$$

and  $\lambda_{T,w}(\mathcal{C}) = \#\Lambda_{T,w}(\mathcal{C})$ .

#### A. Shortened linear codes holding $t$ -designs

Let  $p = 2$  and  $q = 2^m$ . Notice that if a binary code  $\mathcal{C}$  has length  $2^m$  and the weight distribution in Table I (resp. Table II), then the code  $\mathcal{C}$  holds 3-design (resp. 2-design) in [14], [21]. The following two theorems are easily derived from Theorem 13, Tables I and II, and we omit their proofs.

**Theorem 16.** *Let  $m \geq 5$  be odd, and  $\mathcal{C}$  be a binary linear code with length  $2^m$  and the weight distribution in Table I. Let  $T$  be a  $t$ -subset of  $\mathcal{P}(\mathcal{C})$ . We have the following results.*

- *If  $t = 1$ , then the shortened code  $C_T$  is a  $[2^m - 1, 2m, 2^{m-1} - 2^{(m-1)/2}]$  binary linear code with the weight distribution in Table III.*
- *If  $t = 2$ , then the shortened code  $C_T$  is a  $[2^m - 2, 2m - 1, 2^{m-1} - 2^{(m-1)/2}]$  binary linear code with the weight distribution in Table IV.*
- *If  $t = 3$ , then the shortened code  $C_T$  is a  $[2^m - 3, 2m - 2, 2^{m-1} - 2^{(m-1)/2}]$  binary linear code with the weight distribution in Table V.*

TABLE III  
THE WEIGHT DISTRIBUTION OF  $C_T$  FOR  $m$  ODD AND  $t = 1$

Weight	Multiplicity
0	1
$2^{m-1} - 2^{(m-1)/2}$	$2^{(m-5)/2}(2^m - 1)(2 + 2^{(1+m)/2})$
$2^{m-1}$	$-1 + 2^{m-1} + 2^{2m-1}$
$2^{m-1} + 2^{(m-1)/2}$	$2^{(m-5)/2}(2^m - 1)(-2 + 2^{(1+m)/2})$

TABLE IV  
THE WEIGHT DISTRIBUTION OF  $C_T$  FOR  $m$  ODD AND  $t = 2$

Weight	Multiplicity
0	1
$2^{m-1} - 2^{(m-1)/2}$	$2^{(m-7)/2}(-4 + 2^{2+m} + 2^{(1+3m)/2})$
$2^{m-1}$	$-1 + 2^{2m-2}$
$2^{m-1} + 2^{(m-1)/2}$	$2^{(m-7)/2}(4 - 2^{2+m} + 2^{(1+3m)/2})$

**Example 17.** *Let  $m = 5$  and  $T$  be a 1-subset of  $\mathcal{P}(\mathcal{C})$ . Then the shortened code  $C_T$  in Theorem 16 is a  $[31, 10, 12]$  binary linear code with the weight enumerator  $1 + 310z^{12} + 527z^{16} + 186z^{20}$ . The code  $C_T$  is optimal. The dual code of  $C_T$  has parameters  $[31, 21, 5]$  and is optimal according to the tables of best known codes maintained at <http://www.codetables.de>.*

**Example 18.** *Let  $m = 5$  and  $T$  be a 2-subset of  $\mathcal{P}(\mathcal{C})$ . Then the shortened code  $C_T$  in Theorem 16 is a  $[30, 9, 12]$  linear code with the weight enumerator  $1 + 190z^{12} + 255z^{16} + 66z^{20}$ . The code  $C_T$  is optimal.*



TABLE V  
THE WEIGHT DISTRIBUTION OF  $C_T$  FOR  $m$  ODD AND  $t = 3$

Weight	Multiplicity
0	1
$2^{m-1} - 2^{(m-1)/2}$	$-2^{(m-3)/2} + 3 \cdot 2^{(3m-7)/2} + 2^{m-3} + 2^{2m-4}$
$2^{m-1}$	$(-1 + 2^{m-2})(1 + 2^{m-1})$
$2^{m-1} + 2^{(m-1)/2}$	$2^{(m-3)/2} - 3 \cdot 2^{(3m-7)/2} + 2^{m-3} + 2^{2m-4}$

The dual code of  $C_T$  has parameters  $[30, 21, 4]$  and is optimal according to the tables of best known codes maintained at <http://www.codetables.de>.

**Example 19.** Let  $m = 5$  and  $T$  be a 3-subset of  $\mathcal{P}(C)$ . Then the shortened code  $C_T$  in Theorem 16 is a  $[29, 8, 12]$  binary linear code with the weight enumerator  $1 + 114z^{12} + 119z^{16} + 22z^{20}$ . The code  $C_T$  is optimal. The dual code of  $C_T$  has parameters  $[29, 21, 3]$  and is almost optimal according to the tables of best known codes maintained at <http://www.codetables.de>.

**Theorem 20.** Let  $m \geq 4$  be even, and  $C$  be a binary linear code with length  $2^m$  and the weight distribution in Table II. Let  $T$  be a  $t$ -subset of  $\mathcal{P}(C)$ . We have the following results.

- If  $t = 1$ , then the shortened code  $C_T$  is a  $[2^m - 1, 2m, 2^{m-1} - 2^{m/2}]$  binary linear code with the weight distribution in Table VI.
- If  $t = 2$ , then the shortened code  $C_T$  is a  $[2^m - 2, 2m - 1, 2^{m-1} - 2^{m/2}]$  binary linear code with the weight distribution in Table VII.

TABLE VI  
THE WEIGHT DISTRIBUTION OF  $C_T$  FOR  $m$  EVEN AND  $t = 1$

Weight	Multiplicity
0	1
$2^{m-1} - 2^{m/2}$	$1/3 \cdot 2^{-3+m/2} (2 + 2^{m/2}) (-1 + 2^m)$
$2^{m-1} - 2^{(m-2)/2}$	$1/3 \cdot 2^{m/2} (-1 + 2^{m/2}) (1 + 2^{m/2})^2$
$2^{m-1}$	$(2^m - 1)(1 + 2^{m-2})$
$2^{m-1} + 2^{(m-2)/2}$	$1/3 \cdot 2^{m/2} (-1 + 2^{m/2})^2 (1 + 2^{m/2})$
$2^{m-1} + 2^{m/2}$	$1/3 \cdot 2^{-3+m/2} (-2 + 2^{m/2}) (-1 + 2^m)$

TABLE VII  
THE WEIGHT DISTRIBUTION OF  $C_T$  FOR  $t = 2$  AND  $m$  EVEN

Weight	Multiplicity
0	1
$2^{m-1} - 2^{m/2}$	$1/3 \cdot 2^{m/2-4} (2 + 2^{m/2}) (2^m + 2^{1+m/2} - 2)$
$2^{m-1} - 2^{(m-2)/2}$	$1/3 \cdot 2^{m/2-1} (1 + 2^{m/2}) (2^m + 2^{m/2} - 2)$
$2^{m-1}$	$(2^{m-1} - 1)(1 + 2^{m-2})$
$2^{m-1} + 2^{(m-2)/2}$	$1/3 \cdot 2^{m/2-1} (-1 + 2^{m/2}) (2^m - 2^{m/2} - 2)$
$2^{m-1} + 2^{m/2}$	$1/3 \cdot 2^{m/2-4} (4 + 2^{1+m/2} + 2^{3m/2} - 2^{2+m})$

**Example 21.** Let  $m = 4$  and  $T$  be a 1-subset of  $\mathcal{P}(C)$ . Then the shortened code  $C_T$  in Theorem 20 is a  $[15, 8, 4]$  linear code with the weight enumerator  $1 + 15z^4 + 100z^6 + 75z^8 + 60z^{10} + 5z^{12}$ . This code  $C_T$  is optimal. Its dual  $C_T^\perp$  has parameters  $[15, 7, 5]$  and is optimal according to the tables of best known codes maintained at <http://www.codetables.de>.

**Example 22.** Let  $m = 4$  and  $T$  be a 2-subset of  $\mathcal{P}(C)$ . Then the shortened code  $C_T$  in Theorem 20 is a  $[14, 7, 4]$  binary linear code with the weight enumerator  $1 + 11z^4 + 60z^6 + 35z^8 + 20z^{10} + z^{12}$ . This code  $C_T$  is optimal. Its dual  $C_T^\perp$  has parameters  $[14, 7, 4]$  and is optimal according to the tables of best known codes maintained at <http://www.codetables.de>.

### B. Several general results on shortened codes

**Lemma 23.** Let  $m \geq 5$  be odd (resp.  $m \geq 4$  be even), and  $C$  be a binary linear code with the length  $2^m$  and the weight distribution in Table I (resp. Table II). Then the dual code  $C^\perp$  of  $C$  has parameters  $[2^m, 2^m - 2m - 1, 6]$ .

*Proof.* The weight distribution in Table I (or II) means that the dimension of  $C$  is  $2m + 1$ . Thus, the dual code  $C^\perp$  of  $C$  has dimension  $2^m - 2m - 1$ . Since the code length of  $C$  is  $2^m$ , from the weight distribution in Table I (or II) and the first seven Pless power moments in (1), it is easily obtain that  $A_6(C^\perp) > 0$  and  $A_i(C^\perp) = 0$  for any  $i \in \{1, 2, 3, 4, 5\}$ . The desired conclusions then follow.  $\square$

**Theorem 24.** Let  $m \geq 5$  be odd (resp.  $m \geq 4$  be even), and  $C$  be a binary linear code with length  $2^m$  and the weight distribution in Table I (resp. Table II). Let  $T$  be a 4-subset of  $\mathcal{P}(C)$  and  $\lambda_{T,6}(C^\perp) = \lambda$ , then  $\lambda = 0$  or 1. Furthermore, we have the following results.

- (I) If  $m \geq 5$  is odd and  $\lambda = 0$ , then the shortened code  $C_T$  is a  $[2^m - 4, 2m - 3, 2^{m-1} - 2^{(m-1)/2}]$  binary linear code with the weight distribution in Table VIII.
- (II) If  $m \geq 5$  is odd and  $\lambda = 1$ , then the shortened code  $C_T$  is a  $[2^m - 4, 2m - 3, 2^{m-1} - 2^{(m-1)/2}]$  binary linear code with the weight distribution in Table IX.

TABLE VIII  
THE WEIGHT DISTRIBUTION OF  $C_T$  FOR  $\lambda = 0$

Weight	Multiplicity
0	1
$2^{m-1} - 2^{(m-1)/2}$	$-2^{(m-3)/2} + 2^{m-3} + 2^{2m-5} + 2^{(3m-5)/2}$
$2^{m-1}$	$-1 - 2^{m-2} + 4^{m-2}$
$2^{m-1} + 2^{(m-1)/2}$	$2^{(m-3)/2} + 2^{m-3} + 2^{2m-5} - 2^{(3m-5)/2}$

TABLE IX  
THE WEIGHT DISTRIBUTION OF  $C_T$  FOR  $\lambda = 1$

Weight	Multiplicity
0	1
$2^{m-1} - 2^{(m-1)/2}$	$3 \times 2^{m-4} - 2^{(-3+m)/2} + 2^{-5+2m} + 2^{(-5+3m)/2}$
$2^{m-1}$	$2^{-4}(-8 + 2^m)(2 + 2^m)$
$2^{m-1} + 2^{(m-1)/2}$	$3 \times 2^{m-4} + 2^{(-3+m)/2} + 2^{-5+2m} - 2^{(-5+3m)/2}$

*Proof.* By definition, we have

$$\lambda = \lambda_{T,6}(C^\perp) = \#\Lambda_{T,6}(C^\perp) = \#\{\text{Supp}(\mathbf{c}) : wt(\mathbf{c}) = 6, \mathbf{c} \in C^\perp \text{ and } T \subseteq \text{Supp}(\mathbf{c})\}.$$

Suppose that  $\lambda \geq 2$ . There exist  $\text{Supp}(\mathbf{c}_1), \text{Supp}(\mathbf{c}_2) \in \Lambda_{T,6}(C^\perp)$ . Then  $\mathbf{c}_1 + \mathbf{c}_2 \in C^\perp$  and the weight  $\text{wt}(\mathbf{c}_1 + \mathbf{c}_2) \leq 4$ . This is a contradiction to the minimal distance 6 of  $C^\perp$  in Lemma 23. Thus,  $\lambda = 0$  or 1.

We now prove the two cases as follows.

(I) The case that  $\lambda = 0$  and  $m$  is odd.

By Lemma 23, the minimal distance of  $C^\perp$  is 6. Thus,

$$A_1 \left( (C^\perp)^T \right) = A_2 \left( (C^\perp)^T \right) = 0, \quad A_1 \left( (C_T)^\perp \right) = A_2 \left( (C_T)^\perp \right) = 0 \quad (7)$$

and the shortened code  $C_T$  has length  $n = 2^m - 4$  and dimension  $k = 2m - 3$  from  $\lambda_{T,6}(C^\perp) = 0$  and Lemma 1. By the definition and Lemma 23, we have  $A_i(C_T) = 0$  for  $i \notin \{0, i_1, i_2, i_3\}$ , where  $i_1 = 2^{m-1} - 2^{(m-1)/2}$ ,  $i_2 = 2^{m-1}$  and  $i_3 = 2^{m-1} + 2^{(m-1)/2}$ . Therefore, from (7) and (1), the first three Pless power moments

$$\begin{cases} A_{i_1} + A_{i_2} + A_{i_3} = 2^{2m-3} - 1, \\ i_1 A_{i_1} + i_2 A_{i_2} + i_3 A_{i_3} = 2^{2m-3-1}(2^m - 4), \\ i_1^2 A_{i_1} + i_2^2 A_{i_2} + i_3^2 A_{i_3} = 2^{2m-3-2}(2^m - 4)(2^m - 4 + 1). \end{cases}$$

yield the weight distribution in Table VIII. This completes the proof of (I).

(II) The case that  $\lambda = 1$  and  $m$  is odd.

The proof is similar to that of (I). Since  $\lambda_{T,6}(C^\perp) = 1$  and the minimal distance of  $C^\perp$  is 6, from Lemma 1 we have

$$\begin{aligned} A_1 \left( (C^\perp)^T \right) &= 0, \quad A_2 \left( (C^\perp)^T \right) = 1, \\ A_1 \left( (C_T)^\perp \right) &= 0, \quad A_2 \left( (C_T)^\perp \right) = 1. \end{aligned} \quad (8)$$

Then the desired conclusions follow from (8), the definitions and the first three Pless power moments of (1). This completes the proof.  $\square$

**Lemma 25.** *Let  $m \geq 4$  be even, and  $C$  be a binary linear code with length  $2^m$  and the weight distribution in Table II. Let  $T$  be a 3-subset of  $\mathcal{P}(C)$ . Suppose  $\lambda_{T,6}(C^\perp) = \lambda$ , then  $A_1 \left( (C^\perp)^T \right) = A_2 \left( (C^\perp)^T \right) = 0$ ,  $A_3 \left( (C^\perp)^T \right) = \lambda$  and  $A_4 \left( (C^\perp)^T \right) = 2 \cdot (2^{m-2} - 1)^2 - 3\lambda$ .*

*Proof.* By Lemma 23, the minimal distance of  $C^\perp$  is 6. Thus, from  $\#T = 3$  and the definitions, we have  $A_1 \left( (C^\perp)^T \right) = A_2 \left( (C^\perp)^T \right) = 0$  and  $A_3 \left( (C^\perp)^T \right) = \lambda$ . Note that the code  $C$  has length  $2^m$  and dimension  $2m + 1$ . By Lemma 23, Table II and the first seven Pless power moments of (1), we have

$$A_6(C^\perp) = \frac{1}{45} \cdot 2^{m-4}(2^m - 4)^2(2^m - 1).$$

Further, from Theorem 11, Lemmas 23 and 12, we have  $(\mathcal{P}(C^\perp), \mathcal{B}_6(C^\perp))$  is a 2-design and

$$A_4 \left( (C^\perp)^{\{t_1, t_2\}} \right) = \frac{\binom{6}{2}}{\binom{q}{2}} \cdot A_6(C^\perp) = \frac{2}{3} \cdot (2^{m-2} - 1)^2$$

for any  $\{t_1, t_2\} \subseteq \mathcal{P}(C)$ . Since  $\#T = 3$  and the minimal distance of  $C^\perp$  is 6,

$$A_4 \left( (C^\perp)^T \right) = \binom{3}{2} \left( A_4 \left( (C^\perp)^{\{t_1, t_2\}} \right) - \lambda \right).$$

Then the desired conclusions follow.  $\square$

**Theorem 26.** *Let  $m \geq 4$  be even, and  $C$  be a binary linear code with length  $2^m$  and the weight distribution in Table II. Let  $T$  be a 3-subset of  $\mathcal{P}(C)$ . Suppose  $\lambda_{T,6}(C^\perp) = \lambda$ , then the shortened code  $C_T$  is a  $[2^m - 3, 2m - 2, 2^{m-1} - 2^{m/2}]$  binary linear code with the weight distribution in Table X.*

*Proof.* The proof is similar to that of Theorem 24. From Lemma 1 and  $\#T = 3$ , the shortened code  $C_T$  has length  $n = 2^m - 3$  and dimension  $k = 2m - 2$ . By the definitions and the weight distribution in Table II, then  $A_i(C_T) = 0$  for  $i \notin \{0, i_1, i_2, i_3, i_4, i_5\}$ , where  $i_1 = 2^{m-1} - 2^{m/2}$ ,  $i_2 = 2^{m-1} - 2^{(m-2)/2}$ ,  $i_3 = 2^{m-1}$ ,  $i_4 = 2^{m-1} + 2^{(m-2)/2}$  and  $i_5 = 2^{m-1} + 2^{m/2}$ . Moreover, from Lemmas 1 and 25 we have  $A_1((C_T)^\perp) = A_2((C_T)^\perp) = 0$ ,  $A_3((C_T)^\perp) = \lambda$  and  $A_4((C_T)^\perp) = 2 \cdot (2^{m-2} - 1)^2 - 3\lambda$ . Therefore, the first five Pless power moments of (1) yield the weight distribution in Table X. This completes the proof.  $\square$

TABLE X  
THE WEIGHT DISTRIBUTION OF  $C_T$  FOR  $\lambda_{T,6}(C^\perp) = \lambda$

Weight	Multiplicity
0	1
$2^{m-1} - 2^{m/2}$	$1/3 \cdot 2^{m/2-5} (8 + 2^{3+m/2} + 2^{3m/2} + 2^{2+m} + 12\lambda)$
$2^{m-1} - 2^{(m-2)/2}$	$1/3 \cdot 2^{m/2-3} ((2 + 2^{m/2})(-8 + 3 \cdot 2^{m/2} + 2^{1+m}) - 6\lambda)$
$2^{m-1}$	$-1 + 4^{m-2}$
$2^{m-1} + 2^{(m-2)/2}$	$1/3 \cdot 2^{m/2-3} ((-2 + 2^{m/2})(-8 - 3 \cdot 2^{m/2} + 2^{1+m}) + 6\lambda)$
$2^{m-1} + 2^{m/2}$	$1/3 \cdot 2^{m/2-5} (-8 + 2^{3+m/2} + 2^{3m/2} - 2^{2+m} - 12\lambda)$

#### IV. SHORTENED LINEAR CODES FROM APN FUNCTIONS

Let  $p = 2$  and  $q = 2^m$ . In this section, we study some shortened codes  $C_T$  of linear codes  $C$  defined by (6) and determine their parameters, where  $f(x) = x^{2^e+1}$  is a special APN function,  $e$  is a positive integer, and  $\gcd(m, e) = 1$ . It is known that  $C$  has the weight distribution in Tables I (resp. Tables II) when  $m$  is odd (resp.  $m$  is even).

Let  $T$  be a  $t$ -subset of  $P(C)$ . We will consider some shortened code  $C_T$  of  $C$  for the case  $m \geq 4$  and  $t = 3$  (or  $t = 4$ ).

##### A. Some shortened codes for the case $t = 4$ and $m$ odd

We notice that it is difficult to determine the conditions satisfied by the four coordinate positions  $x_1, x_2, x_3, x_4 \in T$  such that  $\lambda_{T,6}(C^\perp) = 0$  (or 1) in Theorem 24. We start with the special code satisfying Theorem 24 and give the necessary and sufficient conditions in Theorem 28. In order to prove Theorem 28, we need the result in the following lemma.

**Lemma 27.** *Let  $e$  and  $m \geq 4$  be positive integers with  $\gcd(m, e) = 1$ . Let  $q = 2^m$  and  $\{x_1, x_2, x_3, x_4\} \subseteq \text{GF}(q)$ . Denote  $S_i = x_1^i + x_2^i + x_3^i + x_4^i$ . Let  $N$  be the number of solutions  $(x, y) \in (\text{GF}(q))^2$  of the system of equations*

$$\begin{cases} x_1 + x_2 + x_3 + x_4 + x + y = 0, \\ x_1^{2^e+1} + x_2^{2^e+1} + x_3^{2^e+1} + x_4^{2^e+1} + x^{2^e+1} + y^{2^e+1} = 0 \end{cases} \quad (9)$$

where  $x_1, x_2, x_3, x_4, x, y$  are pairwise distinct. Then  $N = 4$  if  $S_1 \neq 0$  and  $\text{Tr}_{q/2}(\frac{S_2^e+1}{S_1^2+1} + 1) = 0$ , and  $N = 0$  otherwise.

*Proof.* By definition, the system (9) can be written as follows :

$$\begin{cases} x + y = S_1, \\ x^{2^e+1} + y^{2^e+1} + S_{2^e+1} = 0 \end{cases} \quad (10)$$

Substituting  $y = x + S_1$ , the second equation of (10) leads to

$$\begin{aligned} & x^{2^e+1} + (x + S_1)^{2^e+1} + S_{2^e+1} \\ &= x^{2^e+1} + (x^{2^e} + S_1^{2^e})(x + S_1) + S_{2^e+1} \\ &= S_1 x^{2^e} + S_1^{2^e} \cdot x + S_1^{2^e+1} + S_{2^e+1} = 0. \end{aligned} \quad (11)$$

Further, Equation (11) with  $S_1 \neq 0$  is equivalent to

$$\frac{S_1 x^{2^e} + S_1^{2^e} \cdot x + S_1^{2^e+1} + S_{2^e+1}}{S_1^{2^e+1}} = \left(\frac{x}{S_1}\right)^{2^e} + \frac{x}{S_1} + 1 + \frac{S_{2^e+1}}{S_1^{2^e+1}} = 0. \quad (12)$$

This means that Equation (11) with  $S_1 \neq 0$  has only two different solutions  $x = u, v \in \text{GF}(q)$  if and only if  $\text{Tr}_{q/2}\left(\frac{S_{2^e+1}}{S_1^{2^e+1}} + 1\right) = 0$ , and no solution otherwise. Note that if  $(x, y)$  is the solution of (9), then so is  $(y, x)$ . Thus,  $N = 4$  when Equation (11) with  $S_1 \neq 0$  has solutions. Note that if (9) has nontrivial solution (i.e., not all being zero or pairwise equal), all elements  $x_1, x_2, x_3, x_4, x, y$  have to be distinct since the cyclic code with zero set  $\{1, 2^e + 1\}$  has minimum distance 5, where  $\text{gcd}(m, e) = 1$ . Thus,  $x_1, x_2, x_3, x_4, u, v$  are pairwise distinct. Then the desired conclusions follow.  $\square$

**Theorem 28.** *Let  $e$  and  $m \geq 4$  be positive integers with  $\text{gcd}(m, e) = 1$ . Let  $q = 2^m$ ,  $f(x) = x^{2^e+1}$  and  $C$  be defined in (6). Let  $\alpha$  be a primitive element of  $\text{GF}(q)$  and  $T = \{x_1, x_2, x_3, x_4\}$  be a 4-subset of  $\mathcal{P}(C)$ . Then  $\lambda_{T,6}(C^\perp) = 1$  iff  $\sum_{i=1}^4 x_i \neq 0$  and  $\text{Tr}_{q/2}\left(\sum_{i=1}^4 x_i^{2^e+1} / (\sum_{i=1}^4 x_i^{2^e+1} + 1)\right) = 0$ , and  $\lambda_{T,6}(C^\perp) = 0$  otherwise.*

*Proof.* By definitions, the dual code  $C^\perp$  of  $C$  has minimum distance 6. For any  $\{x_1, x_2, x_3, x_4\} \subseteq \text{GF}(q)$ , there exists a unique set  $\{x_1, x_2, x_3, x_4, x, y\}$  satisfying Equation (9) and  $0 \in \{x_1, x_2, x_3, x_4, x, y\}$ , since the code with zero set  $\{1, 2^e + 1\}$  has minimum distance 5, where  $\text{gcd}(m, e) = 1$ . From the definitions we have  $\lambda_{T,6}(C^\perp) = \frac{N}{2!} - 1$ , where  $N$  was defined in Lemma 27. Then the desired conclusions follow from Lemma 27.  $\square$

By Theorems 28 and 24, we have the following theorem, which is one of the main results in this paper.

**Theorem 29.** *Let  $m \geq 5$  be odd and  $e$  be a positive integer with  $\text{gcd}(m, e) = 1$ . Let  $q = 2^m$ ,  $f(x) = x^{2^e+1}$  and  $C$  be defined in (6). Let  $\alpha$  be a primitive element of  $\text{GF}(q)$  and  $T = \{x_1, x_2, x_3, x_4\}$  be a 4-subset of  $\mathcal{P}(C)$ . Then  $\lambda_{T,6}(C^\perp) = 1$  if  $\sum_{i=1}^4 x_i \neq 0$  and  $\text{Tr}_{q/2}\left(\sum_{i=1}^4 x_i^{2^e+1} / (\sum_{i=1}^4 x_i^{2^e+1} + 1)\right) = 1$ , and  $\lambda_{T,6}(C^\perp) = 0$  otherwise. The parameters of the shortened code  $C_T$  were given in Theorem 24.*

**Example 30.** *Let  $m = 5, e = 1$  and  $T = \{\alpha^1, \alpha^2, \alpha^4, \alpha^5\}$ . Then  $\gamma = \alpha^{17}$ ,  $\text{Tr}_{q/2}\left(\frac{\gamma}{\gamma^3}\right) = 0$  and  $\lambda_{T,6}(C^\perp) = 0$ , where  $\gamma = \alpha^1 + \alpha^2 + \alpha^4 + \alpha^5$ . The shortened code  $C_T$  in Theorem 29 is a  $[28, 7, 12]$  binary linear code with the weight enumerator  $1 + 66z^{12} + 55z^{16} + 6z^{20}$ . The code  $C_T$  is optimal according to the tables of best known codes maintained at <http://www.codetables.de>.*

**Example 31.** *Let  $m = 5, e = 1$  and  $T = \{\alpha^1, \alpha^2, \alpha^3, \alpha^4\}$ . Then  $\gamma = \alpha^{24}$ ,  $\text{Tr}_{q/2}\left(\frac{\gamma}{\gamma^3}\right) = 1$ , and  $\lambda_{T,6}(C^\perp) = 1$ , where  $\gamma = \alpha^1 + \alpha^2 + \alpha^3 + \alpha^4$ . The shortened code  $C_T$  in Theorem 29 is a  $[28, 7, 12]$  binary linear code with the weight enumerator  $1 + 68z^{12} + 51z^{16} + 8z^{20}$ . The code  $C_T$  is optimal according to the tables of best known codes maintained at <http://www.codetables.de>.*

### B. Some shortened codes for the case $t = 3$ and $m$ even

For any  $T = \{x_1, x_2, x_3\} \subseteq \mathcal{P}(C)$ , we notice that it is difficult to determine the value of  $\lambda$  in Theorem 26. We will study a class of special linear codes  $C$  defined by (6), where  $f(x) = x^{2^e+1}$  is an APN function,  $m \geq 4$  is even and  $\text{gcd}(e, m) = 1$ . Note that the code  $C$  satisfies Theorem 26. In the following, we will determine the value of  $\lambda$  in Theorem 33 and the parameters of the shortened code  $C_T$ . We need the result in the following lemma.

**Lemma 32.** Let  $e$  be a positive integer,  $m$  be even with  $\gcd(m, e) = 1$ , and  $q = 2^m$ . Let  $\hat{N}$  be the number of solutions  $(x, y, z) \in (\text{GF}(q))^3$  of the system of equations

$$\begin{cases} x + y + z = a, \\ x^{2^e+1} + y^{2^e+1} + z^{2^e+1} = b \end{cases} \quad (13)$$

where  $a, b \in \text{GF}(q)$  and  $a^{2^e+1} \neq b$ . Then  $\hat{N} = 2^m + (-2)^{m/2} - 2$  if  $a^{2^e+1} + b$  is not a cubic residue, and  $\hat{N} = 2^m + (-2)^{m/2+1} - 2$  if  $a^{2^e+1} + b$  is a cubic residue.

*Proof.* Replacing  $x$  with  $x + a$ ,  $y$  with  $y + a$  and  $z$  with  $z + a$ , we have

$$\begin{cases} x + y + z = 0, \\ x^{2^e+1} + y^{2^e+1} + z^{2^e+1} = a^{2^e+1} + b \end{cases} \quad (14)$$

By  $x + y + z = 0$ , we have the following equation

$$x^{2^e}y + y^{2^e}x = a^{2^e+1} + b. \quad (15)$$

Thus,  $\hat{N}$  equals the number of solutions  $(x, y) \in (\text{GF}(q))^2$  to Equation (15).

Replacing  $y$  with  $xy$  in Equation (15), we get

$$x^{2^e}y + y^{2^e}x = x^{2^e+1}y + y^{2^e}x^{2^e+1} = x^{2^e+1}(y + y^{2^e}) = a^{2^e+1} + b. \quad (16)$$

Since  $a^{2^e+1} + b \neq 0$ , a rearrangement of Equation (16) yields

$$y + y^{2^e} = (a^{2^e+1} + b)x^{-(2^e+1)}.$$

Then

$$\text{Tr}((a^{2^e+1} + b)x^{-(2^e+1)}) = 0. \quad (17)$$

Further, from Lemma 6 we get

$$\begin{aligned} & \#\{x \in \text{GF}(q)^* : \text{Tr}((a^{2^e+1} + b)x^{-(2^e+1)}) = 0\} \\ &= \#\{x \in \text{GF}(q)^* : \text{Tr}((a^{2^e+1} + b)x^{(2^e+1)}) = 0\} \\ &= \frac{1}{2} \cdot \sum_{u \in \text{GF}(2)} \sum_{x \in \text{GF}(q)^*} (-1)^{\text{Tr}(u(a^{2^e+1} + b)x^{(2^e+1)})} \\ &= \frac{1}{2} \cdot (q - 2 + \sum_{x \in \text{GF}(q)} (-1)^{\text{Tr}((a^{2^e+1} + b)x^{(2^e+1)})}) \\ &= \begin{cases} \frac{1}{2} \cdot (q - 2 + (-1)^{\frac{m}{2}} 2^{\frac{m}{2}}) & \text{if } a^{2^e+1} + b \text{ is not a cubic residue,} \\ \frac{1}{2} \cdot (q - 2 - (-1)^{\frac{m}{2}} 2^{\frac{m}{2}+1}) & \text{if } a^{2^e+1} + b \text{ is a cubic residue.} \end{cases} \end{aligned} \quad (18)$$

Note that if  $(x, y)$  is the solution of Equation (15), then so is  $(y, x)$ . This means that  $\hat{N}$  equals twice the number of solutions  $x \in \text{GF}(q)^*$  to Equation (17). Then the desired conclusions follow from Equation (18).  $\square$

**Theorem 33.** Let  $e$  be a positive integer and  $m \geq 4$  be even with  $\gcd(m, e) = 1$ . Let  $q = 2^m$ ,  $f(x) = x^{2^e+1}$  and  $C$  be defined in (6). Let  $\alpha$  be a primitive element of  $\text{GF}(q)$  and  $T = \{x_1, x_2, x_3\}$  be a 3-subset of  $\mathcal{P}(C)$ . Then

$$\lambda = \lambda_{T,6}(C^\perp) = \begin{cases} \frac{1}{6} \cdot (q - 2 + (-1)^{\frac{m}{2}} 2^{\frac{m}{2}}) - 1 & \text{if } a^{2^e+1} + b \text{ is not a cubic residue} \\ \frac{1}{6} \cdot (q - 2 - (-1)^{\frac{m}{2}} 2^{\frac{m}{2}+1}) - 1 & \text{if } a^{2^e+1} + b \text{ is a cubic residue,} \end{cases} \quad (19)$$

where  $a = \sum_{i=1}^3 x_i$  and  $b = \sum_{i=1}^3 x_i^{2^e+1}$ . Moreover, the shortened code  $C_T$  is a  $[2^m - 3, 2m - 2, 2^{m-1} - 2^{m/2}]$  binary linear code with the weight distribution in Table X.

*Proof.* By definitions, the code  $C$  has parameters  $[2^m, 2m+1, 2^{m-1} - 2^{m/2}]$  and the weight distribution in Table II. By Lemma 23, the minimal distance of  $C^\perp$  is 6. Consider the system of equations given by

$$\begin{cases} x_1 + x_2 + x_3 + x + y + z = 0, \\ x_1^{2^e+1} + x_2^{2^e+1} + x_3^{2^e+1} + x^{2^e+1} + y^{2^e+1} + z^{2^e+1} = 0. \end{cases} \quad (20)$$

There exists only one set  $\{x_1, x_2, x_3, x, y, z\}$  satisfying  $0 \in \{x_1, x_2, x_3, x, y, z\}$  and Equation (20), since the code with zero set  $\{1, 2^e + 1\}$  has minimum distance 5. Therefore, from the definitions we have  $\lambda_{T,6}(C^\perp) = \frac{\hat{N}}{3!} - 1$ , where  $\hat{N}$  was defined in Lemma 32. Then the desired conclusions follow from Lemma 32 and Theorem 26.  $\square$

**Example 34.** Let  $m = 4$ ,  $e = 1$  and  $T = \{\alpha^1, \alpha^2, \alpha^4\}$ . Then  $\lambda = 0$  and the shortened code  $C_T$  in Theorem 26 is a  $[13, 6, 4]$  binary linear code with the weight enumerator  $1 + 7z^4 + 36z^6 + 15z^8 + 4z^{10} + z^{12}$ . This code  $C_T$  is optimal. Its dual  $C_T^\perp$  has parameters  $[13, 7, 4]$  and is optimal according to the tables of best known codes maintained at <http://www.codetables.de>.

**Example 35.** Let  $m = 4$ ,  $e = 1$  and  $T = \{\alpha^2, \alpha^5, \alpha^7\}$ . Then  $\lambda = 2$  and the shortened code  $C_T$  in Theorem 26 is a  $[13, 6, 4]$  binary linear code with the weight enumerator  $1 + 8z^4 + 34z^6 + 15z^8 + 6z^{10}$ . This code  $C_T$  is optimal. Its dual  $C_T^\perp$  has parameters  $[13, 7, 3]$  and is almost optimal according to the tables of best known codes maintained at <http://www.codetables.de>.

### C. Some shortened codes for the case $t = 4$ and $m$ even

For any  $T = \{x_1, x_2, x_3, x_4\} \subseteq \mathcal{P}(C)$ , Magma programs show that the weight distribution of  $C_T$  for some code  $C$  is very complex. Thus, it is difficult to determine their parameters in general. In this subsection, we will study a class of special linear codes  $C$  with the weight distribution in Table II and determine parameters of  $C_T$  with the special set  $T$  and  $t = \#T = 4$  in Theorem 38.

In order to determine the parameters of  $C_T$ , we need the next two lemmas.

**Lemma 36.** Let  $m \geq 4$  be even and  $q = 2^m$ . Define

$$\begin{aligned} R_{(3,i)} &= \#\{x \in \text{GF}(q)^* : \text{Tr}_{q/2}(x) = i \text{ and } x \text{ is a cubic residue}\} \\ \bar{R}_{(3,i)} &= \#\{x \in \text{GF}(q)^* : \text{Tr}_{q/2}(x) = i \text{ and } x \text{ is not a cubic residue}\}, \end{aligned} \quad (21)$$

where  $i = 0$  or  $i = 1$ . Then

$$\begin{aligned} R_{(3,0)} &= \frac{1}{6}(2^m - 2 + (-2)^{m/2+1}) \\ R_{(3,1)} &= \frac{2^m - 1}{3} - \frac{1}{6}(2^m - 2 + (-2)^{m/2+1}) \\ \bar{R}_{(3,0)} &= (2^{m-1} - 1) - \frac{1}{6}(2^m - 2 + (-2)^{m/2+1}) \\ \bar{R}_{(3,1)} &= \frac{2^m}{3} - \frac{(-2)^{m/2}}{3}. \end{aligned}$$

*Proof.* By definitions, we get

$$\begin{aligned} R_{(3,0)} &= \frac{1}{3} \cdot \#\{x \in \text{GF}(q)^* : \text{Tr}_{q/2}(x^3) = 0\} \\ &= \frac{1}{6} \sum_{z \in \text{GF}(2)} \sum_{x \in \text{GF}(q)^*} (-1)^{z \text{Tr}_{q/2}(x^3)} \\ &= \frac{1}{6} \left( q - 2 + \sum_{x \in \text{GF}(q)} (-1)^{\text{Tr}_{q/2}(x^3)} \right). \end{aligned}$$

Then the value of  $R_{(3,0)}$  follows from Lemma 6. Note that it has  $\frac{q-1}{3}$  cubic residues in  $\text{GF}(q)^*$  and

$$\#\{x \in \text{GF}(q)^* : \text{Tr}_{q/2}(x) = 0\} = \frac{q}{2} - 1.$$

Then the desired conclusions follow from

$$\begin{cases} R_{(3,0)} + \bar{R}_{(3,0)} = \frac{q}{2} - 1, \\ R_{(3,0)} + R_{(3,1)} = \frac{q-1}{3}, \\ \bar{R}_{(3,0)} + \bar{R}_{(3,1)} = \frac{2(q-1)}{3} \end{cases}$$

This completes the proof.  $\square$

**Lemma 37.** *Let  $m \geq 4$  be even,  $e$  be positive integer with  $\gcd(m, e) = 1$ , and  $q = 2^m$ . Let  $N_{(0,1)}$  be the number of solutions  $(x, y, z, u) \in (\text{GF}(q))^4$  of the system of equations*

$$\begin{cases} x + y + z + u = 0, \\ x^{2^e+1} + y^{2^e+1} + z^{2^e+1} + u^{2^e+1} = 1 \end{cases} \quad (22)$$

where  $x, y, z, u$  are pairwise distinct. Then  $N_{(0,1)} = q(q - 2 - (-1)^{m/2} 2^{m/2+1})$ .

*Proof.* By definitions, we have

$$\begin{aligned} N_{(0,1)} &= \frac{1}{q^2} \sum_{x,y,z,u \in \text{GF}(q)} \sum_{a,b \in \text{GF}(q)} \chi_1(b(x+y+z+u)) \chi_1(a(x^{2^e+1} + y^{2^e+1} + z^{2^e+1} + u^{2^e+1} - 1)) \\ &= \frac{1}{q^2} \sum_{a,b \in \text{GF}(q)} \chi_1(-a) \left( \sum_{x \in \text{GF}(q)} \chi_1(ax^{2^e+1} + bx) \right)^4 \\ &= \frac{1}{q^2} \left( \sum_{a \in \text{GF}(q)} \chi_1(-a) \left( \sum_{x \in \text{GF}(q)} \chi_1(ax^{2^e+1}) \right)^4 + \right. \\ &\quad \left. \sum_{a \in \text{GF}(q)^*} \sum_{b \in \text{GF}(q)^*} \chi_1(-a) \left( \sum_{x \in \text{GF}(q)} \chi_1(ax^{2^e+1} + bx) \right)^4 \right) \\ &= \frac{1}{q^2} \left( \sum_{a \in \text{GF}(q)} (-1)^{\text{Tr}(-a)} \left( \sum_{x \in \text{GF}(q)} \chi_1(ax^{2^e+1}) \right)^4 + \right. \\ &\quad \left. \sum_{a \in \text{GF}(q)^*} (-1)^{\text{Tr}(-a)} \sum_{b \in \text{GF}(q)^*} \left( \sum_{x \in \text{GF}(q)} \chi_1(ax^{2^e+1} + bx) \right)^4 \right) \\ &= \frac{1}{q^2} \cdot (q^4 + (R_{(3,0)} - R_{(3,1)}) \cdot 2^{m+4} + (\bar{R}_{(3,0)} - \bar{R}_{(3,1)}) \cdot 2^m + \\ &\quad (R_{(3,0)} - R_{(3,1)}) \cdot (2^{3m+2} - 2^{2m+4}) + (\bar{R}_{(3,0)} - \bar{R}_{(3,1)}) \cdot (2^m - 1) 2^{2m}), \end{aligned}$$

where  $R_{(3,0)}, R_{(3,1)}, \bar{R}_{(3,0)}$  and  $\bar{R}_{(3,1)}$  were defined in (21) and the last equality holds from Lemmas 6 and 7. Then the desired conclusions follow from Lemma 36.  $\square$

**Theorem 38.** *Let  $m \geq 4$  be even and  $e$  be a positive integer with  $\gcd(m, e) = 1$ . Let  $q = 2^m$ ,  $f(x) = x^{2^e+1}$  and  $C$  be defined in (6). Let  $T = \text{GF}(4) = \{0, 1, w, w^2\} \subseteq \text{GF}(q)$ , where  $w$  is a generator of  $\text{GF}(4)^*$ . Then the shortened code  $C_T$  is a  $[2^m - 4, 2m - 3, 2^{m-1} - 2^{m/2}]$  binary linear code with the weight distribution in Table XI.*

*Proof.* By definitions and Lemma 23,  $C$  is a  $[2^m, 2m + 1, 2^{m-1} - 2^{m/2}]$  with the weight distribution in Table II and the minimal distance of  $C^\perp$  is 6. Note that  $\lambda_{T,6}(C^\perp) = 0$  from Theorem 28 and the definition of  $T$ . Thus, the minimal distance of  $(C^\perp)^T$  is at least 3. This means that

$$A_1 \left( (C^\perp)^T \right) = A_2 \left( (C^\perp)^T \right) = 0. \quad (23)$$



TABLE XI  
THE WEIGHT DISTRIBUTION OF  $C_T$  IN THEOREM 38

Weight	Multiplicity
0	1
$2^{m-1} - 2^{m/2}$	$1/3 \cdot 2^{m/2-6}(-16 + 2^{3m/2} - 2^{m+1}(-4 + (-1)^{m/2}) - 2^{4+m/2}(-1 + (-1)^{m/2}))$
$2^{m-1} - 2^{(m-2)/2}$	$1/24 \cdot (2^{m/2+2} + 2^m)(2^m + (-1)^{m/2}2^{m/2} - 2)$
$2^{m-1}$	$-1 + 2^{2m-5} - (-1)^{m/2}2^{3m/2-4}$
$2^{m-1} + 2^{(m-2)/2}$	$1/24 \cdot (-2^{m/2+2} + 2^m)(2^m + (-1)^{m/2}2^{m/2} - 2)$
$2^{m-1} + 2^{m/2}$	$1/3 \cdot 2^{m/2-6}(16 + 2^{3m/2} - 2^{m+1}(4 + (-1)^{m/2}) + 2^{4+m/2}(1 + (-1)^{m/2}))$

Further, from Theorem 33 and the definition of  $T$ , we have

$$\lambda_{\hat{T},6}(C^\perp) = \frac{1}{6} \cdot (q - 2 - (-1)^{\frac{m}{2}} 2^{\frac{m}{2}+1}) - 1 \quad (24)$$

for any  $\hat{T} = \{\hat{x}_1, \hat{x}_2, \hat{x}_3\} \subseteq T$ . Therefore,

$$A_3 \left( \left( C^\perp \right)^T \right) = \binom{4}{3} \cdot \left( \frac{1}{6} \cdot (q - 2 - (-1)^{\frac{m}{2}} 2^{\frac{m}{2}+1}) - 1 \right). \quad (25)$$

Note that the solutions of the system (22) have symmetrical property and  $(x, y, z, u) = (0, 1, w, w^2)$  is a solution of the system (22). Thus, from the definitions and Lemma 37 we get

$$\lambda_{T,8}(C^\perp) = \frac{N_{(0,1)}}{4!} - 1 - \binom{4}{3} \cdot \lambda_{\hat{T},6}(C^\perp), \quad (26)$$

where  $N_{(0,1)}$  was defined in Lemma 37 and  $\lambda_{\hat{T},6}(C^\perp)$  was defined in (24). By the proof of Lemma 25, we obtain

$$A_4 \left( \left( C^\perp \right)^{\{\bar{x}_1, \bar{x}_2\}} \right) = \frac{2}{3} \cdot (2^{m-2} - 1)^2 \quad (27)$$

for any  $\{\bar{x}_1, \bar{x}_2\} \subseteq T$ . It is obvious that there exists only two 3-subsets  $\bar{T}$  of  $T$  with  $\{\bar{x}_1, \bar{x}_2\} \subseteq \bar{T} \subseteq T$  for any  $\{\bar{x}_1, \bar{x}_2\} \subseteq T$ . Thus, from the definitions we have

$$A_4 \left( \left( C^\perp \right)^T \right) = \binom{4}{2} \cdot \left( A_4 \left( \left( C^\perp \right)^{\{\bar{i}_1, \bar{i}_2\}} \right) - 2 \cdot \lambda_{\hat{T},6}(C^\perp) \right) + \lambda_{T,8}(C^\perp). \quad (28)$$

Combining Equations (24), (26) and (27), Equation (28) yields

$$A_4 \left( \left( C^\perp \right)^T \right) = 4 \cdot (2^{m-2} - 1)^2 - \frac{8}{3} \cdot (2^m - 2 - (-1)^{\frac{m}{2}} 2^{\frac{m}{2}+1}) + \frac{N_{(0,1)}}{24} + 15. \quad (29)$$

Note that the shortened code  $C_T$  has length  $2^m - 4$  and dimension  $2m - 3$ , since  $\#T = 4$  and Lemma 1. By the definitions and the weight distribution in Table II, then  $A_i(C_T) = 0$  for  $i \notin \{0, i_1, i_2, i_3, i_4, i_5\}$ , where  $i_1 = 2^{m-1} - 2^{m/2}$ ,  $i_2 = 2^{m-1} - 2^{(m-2)/2}$ ,  $i_3 = 2^{m-1}$ ,  $i_4 = 2^{m-1} + 2^{(m-2)/2}$  and  $i_5 = 2^{m-1} + 2^{m/2}$ . Hence, from Lemma 1 and Equations (23), (25) and (29), the first five Pless power moments of (1) yield the weight distribution in Table XI. This completes the proof.  $\square$

**Example 39.** Let  $m = 4$  and  $e = 1$  (or 3). Then the shortened code  $C_T$  in Theorem 38 is a  $[12, 5, 4]$  binary linear code with the weight enumerator  $1 + 3z^4 + 24z^6 + 3z^8 + z^{12}$ . This code  $C_T$  is optimal. Its dual  $C_T^\perp$  has parameters  $[12, 7, 4]$  and is optimal according to the tables of best known codes maintained at <http://www.codetables.de>.

## V. SHORTENED LINEAR CODES FROM PN FUNCTIONS

In this section, we study some shortened codes of linear codes  $C$  from the special PN functions and determine their parameters.

Let  $p$  be odd prime and  $q = p^m$ . Let  $f(x) = x^2$  and  $C$  defined by (6). It is known that the code  $C$  is a  $[q, 2m + 1]$  linear code with the weight distribution in [31]. Note that the code  $C$  is affine invariant, and thus holds 2-designs. Then the following theorem is easily derived from the parameters of the code  $C$  in [31] and Theorem 13, and we omit its proof.

**Theorem 40.** *Let  $p$  be an odd prime,  $m$  and  $t$  be positive integers. Let  $q = p^m$ ,  $f(x) = x^2$  and  $C$  be defined in (6). Suppose  $T$  is a  $t$ -subset of  $\mathcal{P}(C)$ . We have the following results.*

- *It  $t = 1$ , then the shortened code  $C_T$  is a  $[2^m - 1, 2m]$  linear code with the weight distribution in Table XII (resp. Table XIV) when  $m$  is odd (resp. even).*
- *It  $t = 2$ , then the shortened code  $C_T$  is a  $[2^m - 2, 2m - 1]$  linear code with the weight distribution in Table XIII (resp. Table XV) when  $m$  is odd (resp. even).*

TABLE XII  
THE WEIGHT DISTRIBUTION OF  $C_T$  FOR  $m$  ODD AND  $t = 1$

Weight	Multiplicity
0	1
$p^{m-1}(p-1)$	$(p^m - 1)(1 + p^{m-1})$
$p^{m-1}(p-1) - p^{\frac{m-1}{2}}$	$1/2 \cdot (p-1)p^{(m-3)/2}(p^m - 1)(p + p^{(1+m)/2})$
$p^{m-1}(p-1) + p^{\frac{m-1}{2}}$	$1/2 \cdot (p-1)p^{(m-3)/2}(p^m - 1)(-p + p^{(1+m)/2})$

TABLE XIII  
THE WEIGHT DISTRIBUTION OF  $C_T$  FOR  $m$  ODD AND  $t = 2$

Weight	Multiplicity
0	1
$p^{m-1}(p-1)$	$p^{2m-2} - 1$
$p^{m-1}(p-1) - p^{\frac{m-1}{2}}$	$(p-1)(-p^{(m-1)/2} + p^{2m-2} + 2p^{(3m-3)/2})/2$
$p^{m-1}(p-1) + p^{\frac{m-1}{2}}$	$(p-1)(p^{(m-1)/2} + p^{2m-2} - 2p^{(3m-3)/2})/2$

TABLE XIV  
THE WEIGHT DISTRIBUTION OF  $C_T$  FOR  $m$  EVEN AND  $t = 1$

Weight	Multiplicity
0	1
$(p-1)(p^{m-1} - p^{\frac{m-2}{2}})$	$p^{m/2-1}(p + p^{m/2} - 1)(p^m - 1)/2$
$(p-1)p^{m-1} - p^{\frac{m-2}{2}}$	$(p-1)p^{m/2-1}(p^{m/2} - 1)(1 + p^{m/2})^2/2$
$(p-1)p^{m-1}$	$p^m - 1$
$(p-1)p^{m-1} + p^{\frac{m-2}{2}}$	$(p-1)p^{m/2-1}(p^{m/2} - 1)^2(1 + p^{m/2})/2$
$(p-1)(p^{m-1} + p^{\frac{m-2}{2}})$	$p^{m/2-1}(-p + p^{m/2} + 1)(p^m - 1)/2$

**Example 41.** *Let  $m = 3$ ,  $p = 3$  and  $T$  be a 1-subset of  $\mathcal{P}(C)$ . Then the shortened code  $C_T$  in Theorem 40 is a  $[26, 6, 15]$  linear code with the weight enumerator  $1 + 312z^{15} + 260z^{18} + 156z^{21}$ . This code  $C_T$  is*

TABLE XV  
THE WEIGHT DISTRIBUTION OF  $C_T$  FOR  $m$  EVEN AND  $t = 2$

Weight	Multiplicity
0	1
$(p-1)(p^{m-1} - p^{\frac{m-2}{2}})$	$p^{m/2-2}(p^{m/2} - 1)(-1 + p + p^{m/2})(p + p^{m/2})/2$
$(p-1)p^{m-1} - p^{\frac{m-2}{2}}$	$(p-1)p^{m/2-2}(1 + p^{m/2})(-p + p^{m/2} + p^m)/2$
$(p-1)p^{m-1}$	$p^{m-1} - 1$
$(p-1)p^{m-1} + p^{\frac{m-2}{2}}$	$(p-1)p^{m/2-2}(1 + p^{m/2})(-p - p^{m/2} + p^m)/2$
$(p-1)(p^{m-1} + p^{\frac{m-2}{2}})$	$p^{m/2-2}(p^{m/2} + 1)(1 - p + p^{m/2})(p^{m/2} - p)/2$

optimal. Its dual  $C_T^\perp$  has parameters  $[26, 20, 4]$  and is optimal according to the tables of best known codes maintained at <http://www.codetables.de>.

**Example 42.** Let  $m = 4$ ,  $p = 3$  and  $T$  be a 1-subset of  $\mathcal{P}(C)$ . Then the shortened code  $C_T$  in Theorem 40 is a  $[80, 8, 48]$  linear code with the weight enumerator  $1 + 1320z^{48} + 2400z^{51} + 80z^{54} + 1920z^{57} + 840z^{60}$ . This code  $C_T$  is optimal. Its dual  $C_T^\perp$  has parameters  $[80, 72, 4]$  and is optimal according to the tables of best known codes maintained at <http://www.codetables.de>.

**Example 43.** Let  $m = 5$ ,  $p = 3$  and  $T$  be a 2-subset of  $\mathcal{P}(C)$ . Then the shortened code  $C_T$  in Theorem 40 is a  $[241, 9, 153]$  linear code with the weight enumerator  $1 + 8010z^{153} + 6560z^{162} + 5112z^{171}$ . This code  $C_T$  is optimal. Its dual  $C_T^\perp$  has parameters  $[241, 232, 3]$  and is optimal according to the tables of best known codes maintained at <http://www.codetables.de>.

**Example 44.** Let  $m = 4$ ,  $p = 3$  and  $T$  be a 2-subset of  $\mathcal{P}(C)$ . Then the shortened code  $C_T$  in Theorem 40 is a  $[79, 7, 48]$  linear code with the weight enumerator  $1 + 528z^{48} + 870z^{51} + 26z^{54} + 552z^{57} + 210z^{60}$ . This code  $C_T$  is almost optimal. Its dual  $C_T^\perp$  has parameters  $[79, 72, 3]$  and is almost optimal according to the tables of best known codes maintained at <http://www.codetables.de>.

In the following, we will consider the shortened code  $C_T$  of  $C$  for the case  $\#T = p$ . In order to determine the parameters of  $C_T$ , we need the next lemmas.

**Lemma 45.** [24] Let  $q = p^m$  with  $p$  an odd prime. Then

$$\begin{aligned} & \#\{a \in \text{GF}(q)^* : \eta(a) = 1 \text{ and } \text{Tr}_{q/p}(a) = 0\} \\ &= \begin{cases} \frac{p^{m-1} - 1 - (p-1)p^{\frac{m-2}{2}}(\sqrt{-1})^{\frac{(p-1)m}{2}}}{2} & \text{for even } m, \\ \frac{p^{m-1} - 1}{2} & \text{for odd } m. \end{cases} \end{aligned}$$

and

$$\begin{aligned} & \#\{a \in \text{GF}(q)^* : \eta(a) = -1 \text{ and } \text{Tr}_{q/p}(a) = 0\} \\ &= \begin{cases} \frac{p^{m-1} - 1 + (p-1)p^{\frac{m-2}{2}}(\sqrt{-1})^{\frac{(p-1)m}{2}}}{2} & \text{for even } m, \\ \frac{p^{m-1} - 1}{2} & \text{for odd } m. \end{cases} \end{aligned}$$

**Lemma 46.** [24] Let  $q = p^m$  with  $p$  an odd prime and  $(a, b) \in \text{GF}(q)^2$ . Denote

$$N_0(a, b) = \#\{x \in \text{GF}(q) : \text{Tr}_{q/p}(ax^2 + bx) = 0\}. \quad (30)$$

Then the following results follow.

- If  $m$  is odd, then

$$N_0(a, b) = \begin{cases} p^m & \text{if } (a, b) = (0, 0), \\ p^{m-1} & \text{if } a = 0, b \neq 0, \text{ or } a \neq 0, 0 = \text{Tr}_{q/p}\left(\frac{b^2}{4a}\right), \\ p^{m-1} + p^{\frac{m-1}{2}}(-1)^{\frac{(p-1)(m+1)}{4}} & \text{if } a \neq 0, 0 \neq \text{Tr}_{q/p}\left(\frac{b^2}{4a}\right), \eta(a)\eta'\left(-\text{Tr}_{q/p}\left(\frac{b^2}{4a}\right)\right) = 1, \\ p^{m-1} + p^{\frac{m-1}{2}}(-1)^{\frac{(p-1)(m+1)+4}{4}} & \text{if } a \neq 0, 0 \neq \text{Tr}_{q/p}\left(\frac{b^2}{4a}\right), \eta(a)\eta'\left(-\text{Tr}_{q/p}\left(\frac{b^2}{4a}\right)\right) = -1, \end{cases} \quad (31)$$

- If  $m$  is even, then

$$N_0(a, b) = \begin{cases} p^m & \text{if } (a, b) = (0, 0), \\ p^{m-1} & \text{if } a = 0, b \neq 0, \\ p^{m-1} + (p-1)p^{\frac{m-2}{2}}(-1)^{\frac{m(p-1)+4}{4}} & \text{if } a \neq 0, 0 = \text{Tr}_{q/p}\left(\frac{b^2}{4a}\right), \eta(a) = 1, \\ p^{m-1} + (p-1)p^{\frac{m-2}{2}}(-1)^{\frac{m(p-1)}{4}} & \text{if } a \neq 0, 0 = \text{Tr}_{q/p}\left(\frac{b^2}{4a}\right), \eta(a) = -1, \\ p^{m-1} + p^{\frac{m-2}{2}}(-1)^{\frac{m(p-1)}{4}} & \text{if } a \neq 0, 0 \neq \text{Tr}_{q/p}\left(\frac{b^2}{4a}\right), \eta(a) = 1, \\ p^{m-1} + p^{\frac{m-2}{2}}(-1)^{\frac{m(p-1)+4}{4}} & \text{if } a \neq 0, 0 \neq \text{Tr}_{q/p}\left(\frac{b^2}{4a}\right), \eta(a) = -1, \end{cases} \quad (32)$$

**Lemma 47.** Let  $q = p^m$  with  $p$  an odd prime and  $a \in \text{GF}(q)^*$ . Define  $f(x) = \text{Tr}_{q/p}\left(-\frac{1}{4a}x^2\right)$ . Then the dual of  $f(x)$  is  $f^*(x) = \text{Tr}_{q/p}(ax^2)$  and the sign of the Walsh transform of  $f(x)$  is

$$\varepsilon = \eta\left(-\frac{1}{4a}\right)(-1)^{\frac{m(p-1)(p-3)}{8}+m-1} = \eta(a)(-1)^{\frac{m(p-1)(p-3)}{8}+m-1+\frac{q-1}{2}}.$$

*Proof.* Note that  $f(x)$  is a weakly regular bent function. Then the desired conclusions follow from the definitions and Lemmas 2 and 3.  $\square$

**Lemma 48.** Let  $q = p^m$  with  $p$  an odd prime,  $a \in \text{GF}(q)^*$  and  $e$  be any positive integer such that  $m/\text{gcd}(m, e)$  is odd. Define  $f(x) = \text{Tr}_{q/p}(ax^{p^e+1})$ . Then the dual of  $f(x)$  is  $f^*(\beta) = \text{Tr}_{q/p}(-a(x_{a,-\beta})^{p^e+1})$  for any  $\beta \in \text{GF}(q)$  and the sign of the Walsh transform of  $f(x)$  is

$$\varepsilon = \eta(a)(-1)^{\frac{m(p-1)(p-3)}{8}+m-1+\frac{q-1}{2}},$$

where  $x_{a,-\beta}$  is the unique solution of the equation

$$a^{p^e}x^{p^{2e}} + ax + (-\beta)^{p^e} = 0.$$

*Proof.* Note that  $f(x)$  is a weakly regular bent function. By definitions, we have

$$f^*(\beta) = \text{Tr}_{q/p}(-a(x_{a,-\beta})^{p^e+1}).$$

From the definitions and Lemma 8, we have

$$\varepsilon = \begin{cases} (-1)^{m-1+\frac{q-1}{2}+\frac{m(p-1)}{4}}\eta(a), & \text{if } p \equiv 1 \pmod{4}, \\ (-1)^{m-1+\frac{q-1}{2}+\frac{m(p-3)}{4}}\eta(a), & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

Then the desired conclusions follow.  $\square$

**Lemma 49.** Let  $q = p^m$  with  $p$  an odd prime,  $a \in \text{GF}(q)^*$ ,  $\gamma \in \text{GF}(p)$  and  $e$  be any positive integer such that  $m/\text{gcd}(m, e)$  is odd. Define

$$\tilde{N}_b = \#\{b \in \text{GF}(q) : \text{Tr}_{q/p}(a(x_{a,b})^{p^e+1}) = \gamma \text{ and } \text{Tr}_{q/p}(b) = 0\}$$

where  $x_{a,b}$  is the unique solution of the equation  $a^{p^e} x^{p^{2e}} + ax + (b)^{p^e} = 0$ . If  $\text{Tr}_{q/p}(a) = 0$ , then

$$\tilde{N}_b = \begin{cases} p^{m-2} + \varepsilon \bar{\eta}^{m/2} (-1) (p-1) p^{(m-2)/2}, & \text{if } \gamma = 0 \text{ and } m \text{ is even;} \\ p^{m-2}, & \text{if } \gamma = 0 \text{ and } m \text{ is odd.} \\ \frac{p-1}{2} \left( p^{m-2} - \varepsilon \bar{\eta}^{m/2} (-1) p^{(m-2)/2} \right) & \text{if } \gamma \neq 0 \text{ and } m \text{ is even;} \\ \frac{p-1}{2} \left( p^{m-2} + \varepsilon \sqrt{p^*}^{m-1} \right) & \text{if } \gamma \in SQ \text{ and } m \text{ is odd;} \\ \frac{p-1}{2} \left( p^{m-2} - \varepsilon \sqrt{p^*}^{m-1} \right). & \text{if } \gamma \in NSQ \text{ and } m \text{ is odd;} \end{cases}$$

where  $\varepsilon$  is given in Lemma 48.

*Proof.* By definitions, we have

$$a^{p^e} (x_{a,b})^{p^{2e}} + ax_{a,b} + (b)^{p^e} = (ax_{a,b}^{p^e})^{p^e} + ax_{a,b} + (b)^{p^e} = 0,$$

which deduces

$$\text{Tr}_{q/p}(ax_{a,b}^{p^e} + ax_{a,b} + (b)^{p^e}) = \text{Tr}_{q/p}(ax_{a,b}^{p^e} + ax_{a,b}) = \text{Tr}_{q/p}\left((a^{p^{-e}} + a)x_{a,b}\right) = 0.$$

When  $\text{Tr}_{q/p}(b) = 0$ , we have

$$\tilde{N}_b = \#\{x \in \text{GF}(q) : \text{Tr}_{q/p}(ax^{p^e+1}) = \gamma \text{ and } \text{Tr}_{q/p}\left((a^{p^{-e}} + a)x\right) = 0\}. \quad (33)$$

Note that  $x = 1$  is the unique solution of the equation

$$a^{p^e} (x)^{p^{2e}} + ax + (-a^{p^{-e}} - a)^{p^e} = 0.$$

By Lemma 48, we get

$$f^*(a^{p^{-e}} + a) = \text{Tr}_{q/p}(-a) = 0 \quad (34)$$

when  $\text{Tr}_{q/p}(a) = 0$ , where  $f^*$  is the dual of  $\text{Tr}_{q/p}(ax^{p^e+1})$ . By Equations (33) and (34), the desired conclusions follow from Lemmas 14, 15 and 48.  $\square$

**Theorem 50.** Let  $p$  be an odd prime and  $m$  be a positive integer. Let  $q = p^m$ ,  $f(x) = x^2$  and  $\mathcal{C}$  be defined in (6). Let  $T = \text{GF}(p)$ . Then the shortened code  $\mathcal{C}_T$  is a  $[2^m - p, 2m - 2]$  linear code. If  $m$  is odd, the weight distribution of  $\mathcal{C}_T$  is given in Table XVI; if  $m \geq 2$  is even, the weight distribution of  $\mathcal{C}_T$  is given in Table XVII, where  $B = (-1)^{\frac{m(p-1)(p-3)}{8} + m - 1 + \frac{q-1}{2}} \cdot \sqrt{p^*}^{m-1}$ ,  $B_1 = \frac{p^{m-1} - 1 - (p-1)p^{\frac{m-2}{2}}(\sqrt{-1})^{\frac{(p-1)m}{2}}}{2}$  and  $B_2 = (-1)^{\frac{m(p-1)(p-3)}{8} + m - 1 + \frac{q-1}{2}} \bar{\eta}^{m/2} (-1) \cdot (p-1)p^{(m-2)/2}$ .

TABLE XVI  
THE WEIGHT DISTRIBUTION OF  $\mathcal{C}_T$  FOR  $m$  ODD

Weight	Multiplicity
0	1
$p^{m-1}(p-1)$	$(p^{m-1} - 1)(p^{m-2} + 1)$
$p^{m-1}(p-1) - p^{\frac{m-1}{2}} (-1)^{\frac{(p-1)(m+1)}{4}}$	$\frac{p^{m-1} - 1}{2} \cdot (p-1)(p^{m-2} + B)$
$p^{m-1}(p-1) + p^{\frac{m-1}{2}} (-1)^{\frac{(p-1)(m+1)}{4}}$	$\frac{p^{m-1} - 1}{2} \cdot (p-1)(p^{m-2} - B)$

*Proof.* By definitions and Lemma 1, the shortened code  $\mathcal{C}_T$  has length  $2^m - p$  and dimension  $2m - 2$ . Since  $T = \text{GF}(p)$ , the weight distribution of  $\mathcal{C}_T$  is the same as the code

$$\hat{\mathcal{C}} = \left\{ (\text{Tr}_{q/p}(ax^2 + bx))_{x \in \text{GF}(q)} : a, b \in \text{GF}(q) \text{ and } \text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0 \right\}. \quad (35)$$

TABLE XVII  
THE WEIGHT DISTRIBUTION OF  $C_T$  FOR  $m$  EVEN

Weight	Multiplicity
0	1
$p^{m-1}(p-1)$	$p^{m-1} - 1$
$p^{m-1}(p-1) - (p-1)p^{\frac{m-2}{2}}(-1)^{\frac{m(p-1)+4}{4}}$	$B_1 \cdot (p^{m-2} + B_2)$
$p^{m-1}(p-1) - (p-1)p^{\frac{m-2}{2}}(-1)^{\frac{m(p-1)}{4}}$	$(p^{m-1} - 1 - B_1) \cdot (p^{m-2} - B_2)$
$p^{m-1}(p-1) - p^{\frac{m-2}{2}}(-1)^{\frac{m(p-1)}{4}}$	$B_1 \cdot (p^{m-1} - p^{m-2} - B_2)$
$p^{m-1}(p-1) - p^{\frac{m-2}{2}}(-1)^{\frac{m(p-1)+4}{4}}$	$(p^{m-1} - 1 - B_1) \cdot (p^{m-1} - p^{m-2} + B_2)$

For each  $(a, b) \in \text{GF}(q) \times \text{GF}(q)$  with  $\text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0$ , define the corresponding codeword

$$\mathbf{c}(a, b) = ((\text{Tr}_{q/p}(ax^2 + bx))_{x \in \text{GF}(q)}) \in \hat{\mathcal{C}}.$$

Then the Hamming weight of  $\mathbf{c}(a, b)$  is

$$\text{wt}(\mathbf{c}(a, b)) = q - N_0(a, b)$$

with  $\text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0$ , where  $N_0(a, b)$  was defined in Equation (30). We discuss the value of  $\text{wt}(\mathbf{c}(a, b))$  in the following two cases.

(I) The case that  $m$  is odd.

From Equation (31), we get

$$\begin{aligned} \text{wt}(\mathbf{c}(a, b)) &= q - N_0(a, b) \\ &= \begin{cases} 0 & \text{if } (a, b) = (0, 0), \text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0, \\ p^{m-1}(p-1) & \text{if } a = 0, b \neq 0, \text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0, \\ & \text{or } a \neq 0, \text{Tr}_{q/p}(\frac{b^2}{4a}) = 0, \text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0 \\ p^{m-1}(p-1) - p^{\frac{m-1}{2}}(-1)^{\frac{(p-1)(m+1)}{4}} & \text{if } a \neq 0, \text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0, \\ & \text{Tr}_{q/p}(\frac{b^2}{4a}) \neq 0, \eta(a)\bar{\eta}\left(-\text{Tr}_{q/p}(\frac{b^2}{4a})\right) = 1, \\ p^{m-1}(p-1) + p^{\frac{m-1}{2}}(-1)^{\frac{(p-1)(m+1)}{4}} & \text{if } a \neq 0, \text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0, \\ & \text{Tr}_{q/p}(\frac{b^2}{4a}) \neq 0, \eta(a)\bar{\eta}\left(-\text{Tr}_{q/p}(\frac{b^2}{4a})\right) = -1. \end{cases} \\ &= \begin{cases} 0 & \text{with 1 time,} \\ p^{m-1}(p-1) & \text{with } (p^{m-1} - 1)(p^{m-2} + 1) \text{ time,} \\ p^{m-1}(p-1) - p^{\frac{m-1}{2}}(-1)^{\frac{(p-1)(m+1)}{4}} & \text{with } \frac{p^{m-1}-1}{2} \cdot (p-1)(p^{m-2} + B) \text{ time,} \\ p^{m-1}(p-1) + p^{\frac{m-1}{2}}(-1)^{\frac{(p-1)(m+1)}{4}} & \text{with } \frac{p^{m-1}-1}{2} \cdot (p-1)(p^{m-2} - B) \text{ time,} \end{cases} \end{aligned}$$

when  $(a, b)$  runs through  $\text{GF}(q) \times \text{GF}(q)$ , where  $B = (-1)^{\frac{m(p-1)(p-3)}{8} + m - 1 + \frac{q-1}{2}} \cdot \sqrt{p}^{m-1}$  and the frequency is obtained based on Lemmas 45, 47, 14 and 15. We first compute the frequency  $A_w$  of the nonzero weight  $w$ , where

$$w = p^{m-1}(p-1) - p^{\frac{m-1}{2}}(-1)^{\frac{(p-1)(m+1)}{4}},$$

and

$$A_w = \#\{(a, b) : a \neq 0, \text{Tr}_{q/p}(\frac{b^2}{4a}) \neq 0, \eta(a)\bar{\eta}\left(-\text{Tr}_{q/p}(\frac{b^2}{4a})\right) = 1, \text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0\}.$$

Clearly, the number of  $a \in \text{GF}(q)^*$  such that  $\text{Tr}_{q/p}(a) = 0$  and  $\eta(a) = 1$ , is  $\bar{n}_a = \frac{p^{m-1}-1}{2}$  by Lemma 45; if we fix  $a$  with  $\text{Tr}_{q/p}(a) = 0$  and  $\eta(a) = 1$ , the number of  $b$  such that  $\text{Tr}_{q/p}(b) = 0$  and  $\bar{\eta}\left(-\text{Tr}_{q/p}(\frac{b^2}{4a})\right) = 1$ ,

is  $\bar{n}_b = \frac{p-1}{2}(p^{m-2} + (-1)^{\frac{m(p-1)(p-3)}{8} + m-1 + \frac{q-1}{2}} \cdot \sqrt{p^*}^{m-1})$  by Lemmas 47 and 15. Meanwhile, the number of  $a \in \text{GF}(q)^*$  such that  $\text{Tr}_{q/p}(a) = 0$  and  $\eta(a) = -1$ , is  $\hat{n}_a = \frac{p^{m-1}-1}{2}$  by Lemma 45; if we fix  $a$  with  $\text{Tr}_{q/p}(a) = 0$  and  $\eta(a) = -1$ , the number of  $b$  such that  $\text{Tr}_{q/p}(b) = 0$  and  $\bar{\eta}(-\text{Tr}_{q/p}(\frac{b^2}{4a})) = -1$ , is  $\hat{n}_b = \frac{p-1}{2}(p^{m-2} + (-1)^{\frac{m(p-1)(p-3)}{8} + m-1 + \frac{q-1}{2}} \cdot \sqrt{p^*}^{m-1})$  by Lemmas 47 and 15. Hence,

$$A_w = \bar{n}_a \bar{n}_b + \hat{n}_a \hat{n}_b = \frac{p^{m-1}-1}{2} \cdot (p-1)(p^{m-2} + (-1)^{\frac{m(p-1)(p-3)}{8} + m-1 + \frac{q-1}{2}} \cdot \sqrt{p^*}^{m-1}).$$

The frequencies of other nonzero weights can be similarly derived.

(II) The case that  $m$  is even.

From Equation (32), we get

$$\begin{aligned} \text{wt}(\mathbf{c}(a, b)) &= q - N_0(a, b) \\ &= \begin{cases} 0 & \text{if } (a, b) = (0, 0), \text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0, \\ p^{m-1}(p-1) & \text{if } a = 0, b \neq 0, \text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0, \\ p^{m-1}(p-1) - (p-1)p^{\frac{m-2}{2}}(-1)^{\frac{m(p-1)+4}{4}} & \text{if } a \neq 0, 0 = \text{Tr}_{q/p}(\frac{b^2}{4a}), \eta(a) = 1, \\ & \text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0, \\ p^{m-1}(p-1) - (p-1)p^{\frac{m-2}{2}}(-1)^{\frac{m(p-1)}{4}} & \text{if } a \neq 0, 0 = \text{Tr}_{q/p}(\frac{b^2}{4a}), \eta(a) = -1, \\ & \text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0, \\ p^{m-1}(p-1) - p^{\frac{m-2}{2}}(-1)^{\frac{m(p-1)}{4}} & \text{if } a \neq 0, 0 \neq \text{Tr}_{q/p}(\frac{b^2}{4a}), \eta(a) = 1, \\ & \text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0, \\ p^{m-1}(p-1) - p^{\frac{m-2}{2}}(-1)^{\frac{m(p-1)+4}{4}} & \text{if } a \neq 0, 0 \neq \text{Tr}_{q/p}(\frac{b^2}{4a}), \eta(a) = -1, \\ & \text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0. \end{cases} \\ &= \begin{cases} 0 & \text{with 1 time,} \\ p^{m-1}(p-1) & \text{with } p^{m-1} - 1 \text{ time,} \\ p^{m-1}(p-1) + (p-1)p^{\frac{m-2}{2}}(-1)^{\frac{m(p-1)}{4}} & \text{with } B_1 \cdot (p^{m-2} + B_2) \text{ time,} \\ p^{m-1}(p-1) - (p-1)p^{\frac{m-2}{2}}(-1)^{\frac{m(p-1)}{4}} & \text{with } (p^{m-1} - 1 - B_1) \cdot (p^{m-2} - B_2) \text{ time,} \\ p^{m-1}(p-1) - p^{\frac{m-2}{2}}(-1)^{\frac{m(p-1)}{4}} & \text{with } B_1 \cdot (p^{m-1} - p^{m-2} - B_2) \text{ time,} \\ p^{m-1}(p-1) + p^{\frac{m-2}{2}}(-1)^{\frac{m(p-1)}{4}} & \text{with } (p^{m-1} - 1 - B_1) \cdot (p^{m-1} - p^{m-2} + B_2) \text{ time,} \end{cases} \end{aligned}$$

where  $B_1 = \frac{p^{m-1}-1-(p-1)p^{\frac{m-2}{2}}(\sqrt{-1})^{\frac{(p-1)m}{2}}}{2}$ ,  $B_2 = (-1)^{\frac{m(p-1)(p-3)}{8} + m-1 + \frac{q-1}{2}} \bar{\eta}^{m/2}(-1) \cdot (p-1)p^{(m-2)/2}$ , and the frequency is easy to obtain based on Lemmas 45, 47, 14 and 15.

By the above two cases, the weight distributions in Tables XVI and XVII follow. This completes the proof.  $\square$

**Example 51.** Let  $m = 3$  and  $p = 3$ . Then the shortened code  $C_T$  in Theorem 50 is a  $[24, 4, 15]$  linear code with the weight enumerator  $1 + 48z^{15} + 32z^{18}$ . This code  $C_T$  is optimal. Its dual  $C_T^\perp$  has parameters  $[24, 20, 3]$  and is optimal according to the tables of best known codes maintained at <http://www.codetables.de>.

**Example 52.** Let  $m = 4$  and  $p = 3$ . Then the shortened code  $C_T$  in Theorem 50 is a  $[78, 6, 48]$  linear code with the weight enumerator  $1 + 240z^{48} + 240z^{51} + 26z^{54} + 192z^{57} + 30z^{60}$ . This code  $C_T$  is almost optimal. Its dual  $C_T^\perp$  has parameters  $[78, 72, 2]$  and is almost optimal according to the tables of best known codes maintained at <http://www.codetables.de>.

**Theorem 53.** Let  $p$  be an odd prime,  $m$  and  $e$  be positive integers such that  $m/\text{gcd}(m, e)$  is odd. Let  $q = p^m$ ,  $f(x) = x^{p^e+1}$  and  $C$  be defined in (6). Let  $T = \text{GF}(p)$ . Then the parameters of the shortened code  $C_T$  are the same as that of  $C_T$  in Theorem 50.

*Proof.* The proof is similar to that of Theorem 50. Recall that the code  $\mathcal{C}$  has length  $q$  and dimension  $2m+1$ . By Lemma 1, the shortened code  $\mathcal{C}_T$  has length  $2^m - p$  and dimension  $2m-2$ . Since  $T = \text{GF}(p)$ , the weight distribution of  $\mathcal{C}_T$  is the same as the code

$$\hat{\mathcal{C}} = \left\{ \left( \text{Tr}_{q/p}(ax^{p^e+1} + bx) \right)_{x \in \text{GF}(q)} : a, b \in \text{GF}(q) \text{ and } \text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0 \right\}. \quad (36)$$

For each  $(a, b) \in \text{GF}(q) \times \text{GF}(q)$  with  $\text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0$ , define the corresponding codeword

$$\mathbf{c}(a, b) = \left( (\text{Tr}_{q/p}(ax^{p^e+1} + bx))_{x \in \text{GF}(q)} \right) \in \hat{\mathcal{C}}.$$

Then the Hamming weight of  $\mathbf{c}(a, b)$  is

$$\text{wt}(\mathbf{c}(a, b)) = q - \hat{N}_0(a, b), \quad (37)$$

where  $\text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0$  and  $\hat{N}_0(a, b)$  was defined in Lemma 10.

We need prove the value of  $\text{wt}(\mathbf{c}(a, b))$  and its frequencies in the following four cases.

- (I)  $m$  is odd and  $p \equiv 1 \pmod{4}$ .
- (II)  $m$  is odd and  $p \equiv 3 \pmod{4}$ .
- (III)  $m$  is even and  $p \equiv 1 \pmod{4}$ .
- (IV)  $m$  is even and  $p \equiv 3 \pmod{4}$ .

Next we only give the proof for the case (I) and omit the proofs for the other three cases whose proofs are similar.

Suppose that  $m$  is odd and  $p \equiv 1 \pmod{4}$ . From Equation (37) and Lemma 10, we get

$$\begin{aligned} \text{wt}(\mathbf{c}(a, b)) &= q - \hat{N}_0(a, b) \\ &= \begin{cases} 0 & \text{if } (a, b) = (0, 0), \text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0, \\ p^{m-1}(p-1) & \text{if } a = 0, b \neq 0, \text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0, \\ & \text{or } a \neq 0, b = 0, \text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0, \\ & \text{or } a \neq 0, b \neq 0, \text{Tr}_{q/p}(a(x_{a,b})^{p^e+1}) = 0, \text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0, \\ p^{m-1}(p-1) - p^{m/2-1}\sqrt{p^*} & \text{if } a \neq 0, b \neq 0, \text{Tr}_{q/p}(a(x_{a,b})^{p^e+1}) \neq 0, \text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0, \\ & \eta(a)\eta(\text{Tr}_{q/p}(a(x_{a,b})^{p^e+1})) = 1. \\ p^{m-1}(p-1) + p^{m/2-1}\sqrt{p^*} & \text{if } a \neq 0, b \neq 0, \text{Tr}_{q/p}(a(x_{a,b})^{p^e+1}) \neq 0, \text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0, \\ & \eta(a)\eta(\text{Tr}_{q/p}(a(x_{a,b})^{p^e+1})) = -1. \end{cases} \\ &= \begin{cases} 0 & \text{with 1 time,} \\ p^{m-1}(p-1) & \text{with } (p^{m-1} - 1)(p^{m-2} + 1) \text{ time,} \\ p^{m-1}(p-1) - p^{\frac{m-1}{2}} & \text{with } \frac{p^{m-1}-1}{2} \cdot (p-1)(p^{m-2} + B) \text{ time,} \\ p^{m-1}(p-1) + p^{\frac{m-1}{2}} & \text{with } \frac{p^{m-1}-1}{2} \cdot (p-1)(p^{m-2} - B) \text{ time,} \end{cases} \end{aligned}$$

when  $(a, b)$  runs through  $\text{GF}(q) \times \text{GF}(q)$ , where  $B = (-1)^{\frac{m(p-1)(p-3)}{8} + m - 1 + \frac{q-1}{2}} \cdot \sqrt{p^*}^{m-1}$  and the frequency is obtained by Lemmas 48 and 45. As an example, we just compute the frequency  $A_w$  of the nonzero weight  $w$ , where

$$w = p^{m-1}(p-1) - p^{\frac{m-1}{2}}$$

and

$$A_w = \#\{(a, b) : a \neq 0, b \neq 0, \text{Tr}_{q/p}(a(x_{a,b})^{p^e+1}) \neq 0, \text{Tr}_{q/p}(a) = \text{Tr}_{q/p}(b) = 0, \eta(a)\eta(\text{Tr}_{q/p}(a(x_{a,b})^{p^e+1})) = 1\}.$$

Clearly, the number of  $a \in \text{GF}(q)^*$  such that  $\text{Tr}_{q/p}(a) = 0$  and  $\eta(a) = 1$ , is  $\bar{n}_a = \frac{p^{m-1}-1}{2}$  by Lemma 45; if we fix  $a$  with  $\text{Tr}_{q/p}(a) = 0$  and  $\eta(a) = 1$ , the number of  $b$  such that  $\text{Tr}_{q/p}(b) = 0$  and  $\eta(\text{Tr}_{q/p}(a(x_{a,b})^{p^e+1})) = 1$



, is  $\bar{n}_b = \frac{p-1}{2} \left( p^{m-2} + (-1)^{\frac{m(p-1)(p-3)}{8} + m-1 + \frac{q-1}{2}} \cdot \sqrt{p^*}^{m-1} \right)$  by Lemma 48. Meanwhile, the number of  $a \in \text{GF}(q)^*$  such that  $\text{Tr}_{q/p}(a) = 0$  and  $\eta(a) = -1$ , is  $\hat{n}_a = \frac{p^{m-1}-1}{2}$  by Lemma 45; if we fix  $a$  with  $\text{Tr}_{q/p}(a) = 0$  and  $\eta(a) = -1$ , the number of  $b$  such that  $\text{Tr}_{q/p}(b) = 0$  with  $b \neq 0$  and  $\eta(\text{Tr}_{q/p}(a(x_{a,b})^{p^e+1})) = -1$ , is  $\frac{p-1}{2} \left( p^{m-2} + (-1)^{\frac{m(p-1)(p-3)}{8} + m-1 + \frac{q-1}{2}} \sqrt{p^*}^{m-1} \right)$  by Lemma 48. Hence,

$$A_w = \bar{n}_a \bar{n}_b + \hat{n}_a \hat{n}_b = \frac{p^{m-1}-1}{2} \cdot (p-1) \left( p^{m-2} + (-1)^{\frac{m(p-1)(p-3)}{8} + m-1 + \frac{q-1}{2}} \cdot \sqrt{p^*}^{m-1} \right).$$

The frequencies of other nonzero weights can be similarly derived. This completes the proof of the weight distribution of Table in XVI for the case  $m$  odd and  $p \equiv 1 \pmod{4}$ .

The proofs of the other three cases are similar. The desired conclusions follow from Equation (37), Lemmas 10, 45 and 48. This completes the proof.  $\square$

## VI. CONCLUDING REMARKS

In this paper, we mainly investigated some shortened codes of linear codes from PN and APN functions and determined their parameters. The obtained codes have a few weights and some of these codes are optimal or almost optimal. Specifically, the main results are summarized as follows:

- For any binary linear code  $C$  with length  $q = 2^m$  and the weight distribution in Table I, we gave a general result on their shortened code  $C_T$  with  $\#T = 4$  in Theorem 24. Meanwhile, when  $m$  is odd, the parameters of the shortened code  $C_T$  of a class of binary linear codes from APN functions was determined in Theorem 28.
- For any binary linear code  $C$  with length  $q = 2^m$  and the weight distribution in Table II, we gave a general result on their shortened code  $C_T$  with  $\#T = 3$  in Theorem 26. Further, the parameters of the shortened code  $C_T$  of a special class of linear codes from APN functions were determined for  $\#T = 3$  and  $\#T = 4$  in Theorems 33 and 38.
- Two classes of  $p$ -ary shortened code  $C_T$  from PN functions were presented and their parameters were also determined in Theorems 50 and 53, where  $p$  is an odd prime.

Some of these shortened codes in this paper are optimal or almost optimal. Many codes with good parameters can be produced by using shortening and puncturing technologies. It seems hard to determine their weight distributions. The reader is cordially invited to settle these problems.

## REFERENCES

- [1] E. F. Assmus Jr. and J. D. Key, Designs and their Codes, Cambridge University Press, Cambridge, 1992.
- [2] E. F. Assmus Jr., H. F. Mattson Jr., New 5-designs, J. Comb. Theory Ser. A, 6(2): 122-151, 1969.
- [3] E. F. Assmus Jr and H. F. Mattson Jr, Coding and combinatorics, SIAM Rev 16: 349- 388, 1974.
- [4] L. Budaghyan and T. Helleseht, New perfect nonlinear multinomials over  $\mathbb{F}_{p^{2k}}$  for any odd prime  $p$ , Springer-Verlag, LNCS 5203, 403-414, 2018.
- [5] C. Carlet, C. Ding and J. Yuan, Linear codes from perfect nonlinear mappings and their secret sharing schemes. IEEE Trans. Inf. Theory, 51(6): 2089-2102, 2005.
- [6] C. Carlet, S. Mesnager: Four decades of research on bent functions. Des. Codes Cryptogr. 78(1): 5-50, 2016.
- [7] C. Carlet, Boolean and vectorial plateaued functions and APN functions. IEEE Trans. Inf. Theory, 61(11): 6272-6289, 2015.
- [8] X. Cao, W. Chou and J. Gu, On the number of solutions of certain diagonal equations over finite fields. Finite Fields Their Appl. 42: 225-252, 2016.
- [9] R. S. Coulter, The number of rational points of a class of Artin-chreier curves, Finite Fields Their Appl., 8: 397-413, 2002.
- [10] R. S. Coulter, M. Henderson, L. Hu and et al., Planar polynomials and commutative semifields two dimensional over their middle nucleus and four dimensional over their nucleus, Online at <http://www.math.udel.edu/~coulter/papers/d24.pdf>.
- [11] R. S. Coulter and R. W. Matthews, Planar functions and planes of Lenz-Barlotti class II, Designs, Codes Cryptogr. 10(2): 167-184, 1997.
- [12] R. S. Coulter, On the evaluation of a class of Weil sums in characteristic 2, New Zerland J. of Math. 28: 171-184, 1999.
- [13] P. Dembowski and T. G. Ostrom, Planes of order  $n$  with collineation groups of order  $n^2$ , Math. Z. 103(3): 239-258, 1968.
- [14] C. Ding, Designs from Linear Codes, World Scientific, Singapore, 2018.
- [15] C. Ding, A construction of binary linear codes from Boolean functions. Discret. Math. 339(9): 2288-2303, 2016.
- [16] K. Ding and C. Ding, A class of two-weight and three-weight codes and their applications in secret sharing, IEEE Trans. Inf. Theory, 61(11): 5835-5842, 2015.

- [17] C. Ding, Codes from Difference Sets, World Scientific, Singapore, 2015.
- [18] C. Ding, Infinite families of 3-designs from a type of five-weight code, Des. Codes Cryptogr. 86(3): 703-719, 2018.
- [19] C. Ding, An infinite family of Steiner systems from cyclic codes, Journal of Combinatorial Designs, 26: 127-144, 2018.
- [20] C. Ding, C. Li and Y. Xia, Another generalization of the binary Reed-Muller codes and its applications, Finite Fields Appl. 53: 144-174, 2018.
- [21] C. Ding and C. Tang, Combinatorial  $t$ -designs from special functions, Cryptography and Communications, doi.org/10.1007/s12095-020-00442-2.
- [22] C. Ding and J. Yuan, A family of skew Hadamard difference sets, J. Comb. Theory, Ser. A, 113(7): 1526-1535, 2006.
- [23] G. Gong, T. Helleseeth, H. Hu, and et al., On the dual of certain ternary weakly regular bent functions, IEEE Trans. Inf. Theory, 58(4): 2237-2243, 2012.
- [24] Z. Heng and C. Ding, The subfield codes of hyperoval and conic codes. Finite Fields Their Appl. 56: 308-331, 2019.
- [25] M. Harada, M. Kitazume and A. Munemasa, On a 5-design related to an extremal doubly-even self-dual code of length 72, J. Combin. Theory, Ser. A, 107(1): 143-146, 2004.
- [26] M. Harada, A. Munemasa and V. Tonchev, A characterization of designs related to an extremal doubly-even self-dual code of length 48, Annals of Combinatorics, 9: 189-198, 2005.
- [27] T. Helleseeth, H. Hollmann, A. Kholosha and et al., Proofs of two conjectures on ternary weakly regular bent functions, IEEE Trans. Inf. Theory, 55(11):5272-5283, 2009.
- [28] W. C. Huffman and V. Pless, Fundamentals of Error-Correcting Codes, Cambridge University Press, Cambridge, 2003.
- [29] J. H. Koolen and A. Munemasa, Tight 2-designs and perfect 1-codes in Doob graphs, J. Stat. Planning and Inference, 86: 505-513, 2000.
- [30] R. Lidl and H. Niederreiter, Finite Fields, Cambridge Univ. Press, Cambridge, 1997.
- [31] C. Li, S. Lin and L. Qu, On the Covering Structures of Two Classes of Linear Codes From Perfect Nonlinear Functions, IEEE Trans. Inf. Theory, 55(1):70-82, 2009.
- [32] S. Mesnager and A. Sinak, Several classes of minimal linear codes with few weights from weakly regular plateaued functions. IEEE Trans. Inf. Theory, 66(4): 2296-2310, 2020.
- [33] S. Mesnager, F. Ozbudak, A. Sinak, Linear codes from weakly regular plateaued functions and their secret sharing schemes. Des. Codes Cryptogr. 87(2-3): 463-480, 2019.
- [34] S. Mesnager, Linear codes with few weights from weakly regular bent functions based on a generic construction. Cryptogr. Commun. 9(1): 71-84, 2017.
- [35] A. Munemasa and V. D. Tonchev, A new quasi-symmetric 2-(56,16,6) design obtained from codes, Discrete Math. 284: 231-234, 2004.
- [36] C. Tang, C. Xiang and K. Feng, Linear codes with few weights from inhomogeneous quadratic functions. Des. Codes Cryptogr. 83(3): 691-714,
- [37] C. Tang, C. Ding and M. Xiong, Steiner systems  $S(2,4,\frac{3^m-1}{2})$  and 2-designs from ternary linear codes of length  $\frac{3^m-1}{2}$ . Des. Codes Cryptogr. 87(12): 2793-2811 (2019)
- [38] C.Tang, C. Ding and M. Xiong, Codes, Differentially  $\delta$ -Uniform Functions, and  $t$ -Designs. IEEE Trans. Inf. Theory, 66(6): 3691-3703, 2020.
- [39] C. Tang, N. Li, Y. Qi, Z. Zhou, T. Helleseeth, Linear codes with two or three weights from weakly regular bent functions. IEEE Trans. Inf. Theory, 62(3): 1166-1176, 2016.
- [40] V. D. Tonchev, Codes and designs, In Handbook of Coding Theory, vol. II, V. S. Pless and W. C. Huffman, eds., Elsevier, Amsterdam, 1229-1268, 1998.
- [41] V. D. Tonchev, Codes, In Handbook of Combinatorial Designs, 2nd edition, C. J. Colbourn and J. H. Dinitz, eds., CRC Press, New York, 677-701, 2007.
- [42] Y. Wu, N. Li and X. Zeng, Linear Codes From Perfect Nonlinear Functions Over Finite Fields. IEEE Trans. Communications 68(1): 3-11, 2020.
- [43] J. Yuan, C. Carlet and C. Ding, The weight distribution of a class of linear codes from perfect nonlinear functions. IEEE Trans. Inf. Theory, 52(2): 712-717, 2006.
- [44] Z. Zha, G. Kyureghyan and X. Wang, Perfect nonlinear binomials and their semifields, Finite Fields Appl. 15(2): 125-133, 2009.
- [45] Z. Zhou, C. Ding, J. Luo and et al., A family of five-weight cyclic codes and their weight enumerators. IEEE Trans. Inf. Theory, 59(10): 6674-6682, 2013.
- [46] Z. Zhou and C. Ding, Seven classes of three-weight cyclic codes. IEEE Trans. Communications, 61(10): 4120-4126, 2013.