

A Sharp Blockwise Tensor Perturbation Bound for Orthogonal Iteration

Yuetian Luo

*Department of Statistics
University of Wisconsin-Madison
Madison, WI 53706, USA*

YLUO86@WISC.EDU

Garvesh Raskutti

*Department of Statistics
University of Wisconsin-Madison
Madison, WI 53706, USA*

RASKUTTI@STAT.WISC.EDU

Ming Yuan

*Department of Statistics
Columbia University
New York, NY 10027, USA*

MING.YUAN@COLUMBIA.EDU

Anru R. Zhang

*Department of Statistics
University of Wisconsin-Madison
Madison, WI 53706, USA*

ANRUZHANG@STAT.WISC.EDU

Editor: Animashree Anandkumar

Abstract

In this paper, we develop novel perturbation bounds for the higher-order orthogonal iteration (HOOI) (De Lathauwer et al., 2000a). Under mild regularity conditions, we establish blockwise tensor perturbation bounds for HOOI with guarantees for both tensor reconstruction in Hilbert-Schmidt norm $\|\widehat{\mathcal{T}} - \mathcal{T}\|_{\text{HS}}$ and mode- k singular subspace estimation in Schatten- q norm $\|\sin \Theta(\widehat{\mathbf{U}}_k, \mathbf{U}_k)\|_q$ for any $q \geq 1$. We show the upper bounds of mode- k singular subspace estimation are unilateral and converge linearly to a quantity characterized by blockwise errors of the perturbation and signal strength. For the tensor reconstruction error bound, we express the bound through a simple quantity ξ , which depends only on perturbation and the multilinear rank of the underlying signal. Rate matching deterministic lower bound for tensor reconstruction, which demonstrates the optimality of HOOI, is also provided. Furthermore, we prove that one-step HOOI (i.e., HOOI with only a single iteration) is also optimal in terms of tensor reconstruction and can be used to lower the computational cost. The perturbation results are also extended to the case that only partial modes of \mathcal{T} have low-rank structure. We support our theoretical results by extensive numerical studies. Finally, we apply the novel perturbation bounds of HOOI on two applications, tensor denoising and tensor co-clustering, from machine learning and statistics, which demonstrates the superiority of the new perturbation results.

Keywords: Tensor, Higher-order orthogonal iteration, Perturbation bounds, Minimax optimality

1. Introduction

The past decades have seen a large body of work on tensors or multiway arrays (Kolda and Bader, 2009; Sidiropoulos et al., 2017; Cichocki et al., 2015; Kroonenberg, 2008). Tensors arise in numerous applications involving multiway data (e.g., brain imaging (Zhou et al., 2013), hyperspectral imaging (Li and Li, 2010), recommender system design (Bi et al., 2018)). In addition, various methods have been developed and applied to fundamental tensor problems such as tensor completion (Yuan and Zhang, 2016; Xia et al., 2020; Yuan and Zhang, 2017; Zhang, 2019), tensor regression (Zhang et al., 2020; Zhou et al., 2013; Raskutti et al., 2019; Chen et al., 2019), tensor PCA/SVD (Zhang and Xia, 2018; Richard and Montanari, 2014; Liu et al., 2017), generalized tensor learning (Han et al., 2020). In many other problems where the observations are not necessarily tensors, tensor parameters arise, such as topic and latent variable models (Anandkumar et al., 2014a), additive index models (Balasubramanian et al., 2018), high-order interaction pursuit (Hao et al., 2020). We refer readers to recent survey papers (Kolda and Bader, 2009; Sidiropoulos et al., 2017; Cichocki et al., 2015).

Among these methods, tensor decomposition is one of the most important and a flurry of research have been devoted to it. Tensor decomposition plays a similar role to singular value decomposition (SVD) or eigendecomposition for matrices which is of fundamental importance throughout a wide range of fields including computer science, applied mathematics, machine learning, statistics, signal processing, etc. For matrices, truncated SVD achieves the best low-rank approximation in terms of any unitarily invariant norm by the well-known Eckart-Young-Mirsky theorem (Eckart and Young, 1936; Mirsky, 1960; Golub et al., 1987) and more importantly it is computationally efficient. Despite the well-established theory for low-rank decomposition of matrices, tensors present unique challenges. First there are several notions of low-rankness in tensors, moreover it has been shown that computing various best low-rank approximations of a given tensor is NP hard in general (Hillar and Lim, 2013).

Fortunately, many computationally efficient algorithms have been proposed to approximate the best low-rank tensor decomposition (Kroonenberg and De Leeuw, 1980; De Lathauwer et al., 2000a; Elden and Savas, 2009; Ishteva et al., 2011, 2009; Savas and Lim, 2010; De Lathauwer et al., 2000b; Vannieuwenhoven et al., 2012). One popular choice is the higher-order orthogonal iteration (HOOI) proposed in De Lathauwer et al. (2000a). HOOI is based on alternating least-squares. It can be seen as a “spectral” algorithm for tensors, generalizations of the 2D-PCA (Sheehan and Saad, 2007), and the power iteration refinement of HOSVD (De Lathauwer et al., 2000b) and sequential HOSVD (Vannieuwenhoven et al., 2012). Convergence properties of HOOI have been studied in Zhang and Golub (2001); Wang and Chu (2014); Uschmajew (2015); Xu (2018); Zhang and Xia (2018).

In addition to computing low-rank approximations of matrices and tensors, there is the more nuanced question of computing low-rank approximations under noise perturbation and determining how the perturbation impacts the quality of the decomposition. For matrices, perturbation theory is well studied and a number of results has been established (Bhatia, 2013; Stewart, 1990). However, perturbation theory for tensors is still in its infancy. It is difficult to extend results for matrices to tensors due to the complexity of tensor algebra and the fact many well established theories and concepts in matrices such as SVD

or eigendecomposition do not exist or are not easy to use for tensors. There are several attempts in the literature and most of them require the noise tensor to be random (Richard and Montanari, 2014; Zhang and Xia, 2018; Liu et al., 2017; Zhang and Han, 2019; Zheng and Tomioka, 2015; Xia and Zhou, 2019). In this paper, we move one step forward in this direction and provide the first general perturbation bounds of HOOI for tensors under the signal-plus-noise model

$$\tilde{\mathcal{T}} = \mathcal{T} + \mathcal{Z} \quad (1)$$

without putting any structural assumption on the perturbation noise \mathcal{Z} . Like the classical Wedin’s perturbation theory for matrices, we provide perturbation guarantees of estimated mode- k singular subspace in tensors. In addition, we also provide the perturbation bounds for tensor reconstruction. By providing the deterministic rate matching lower bound, we can further show HOOI with good initialization is optimal for tensor reconstruction.

1.1 Problem Statement

Formally, this paper considers the tensor perturbation model (1), where \mathcal{T} is the low-rank order- d signal tensor and \mathcal{Z} is the perturbation tensor with the same dimension as \mathcal{T} . Two popular choices of low-rankness in tensors are canonical polyadic (CP) low-rank and multilinear/Tucker low-rank and each of them has their respective benefits (see the surveys Kolda and Bader (2009); Sidiropoulos et al. (2017); Cichocki et al. (2015); Grasedyck et al. (2013)). The CP low-rank decomposition which approximates the original tensor by a sum of rank-1 outer products gives a compact and unique (under certain conditions) representation and multilinear/Tucker low-rank decomposition generally finds a better fit for the data by estimating the subspace of each mode. Since any CP low-rank tensor can be written as a multilinear low-rank tensor with a diagonal core tensor, we focus on the latter (Tucker rank) in this paper. Specifically, we assume \mathcal{T} admits the following low multilinear rank (Tucker) decomposition:

$$\mathcal{T} = \mathcal{S} \times_{\Omega_1} \mathbf{U}_1 \times \cdots \times_{\Omega_m} \mathbf{U}_m. \quad (2)$$

Here, \mathcal{S} is an order- d core tensor; $\{\mathbf{U}_i\}_{i=1}^m$ are group- i singular subspaces; $\{\Omega_i\}_{i=1}^m$ are symmetric index groups which will be introduced next; and “ \times_{Ω_i} ” is the tensor matrix product along modes in Ω_i . The formal definition of Tucker decomposition and tensor matrix product will be given in equations (5), (6) and (7) in Section 2.

The symmetric index groups $\{\Omega_i\}_{i=1}^m$ represent the specific symmetry structure of \mathcal{T} and satisfy $\Omega_i \subseteq [d], \Omega_i \neq \emptyset, \Omega_i \cap \Omega_j = \emptyset$ for $i \neq j$, and $\bigcup_{i=1}^m \Omega_i = [d]$, where $[d] := \{1, \dots, d\}$. It means by fixing indices outside the group and any permutation of indices within the group does not change the value of tensor corresponding to those indices. For example, if $\Omega_1 = \{1, 2, \dots, k\}$, then fixing coordinates in $\{\Omega_i\}_{i=2}^m$ i.e., coordinates $k+1, \dots, d$ and for any permutation δ of $[k]$, we have $\mathcal{T}_{[i_{\delta(1)}, \dots, i_{\delta(k)}, i_{k+1}, \dots, i_d]} = \mathcal{T}_{[i_1, \dots, i_d]}$. In addition to \mathcal{T} , we also assume \mathcal{S} and \mathcal{Z} have the same symmetric structure characterized by $\{\Omega_i\}_{i=1}^m$. The symmetric index groups have two extreme cases:

- Asymmetric: $\Omega_i = \{i\}$ for $i = 1, \dots, d$,
- Supersymmetric: $\Omega_1 = \{1, 2, \dots, d\}$.

To simplify the notation, if mode $j \in \Omega_i$, then we denote $j' := i$ as the group index of mode j . For symmetric group i , the dimension of \mathcal{T} in this group is denoted as p_i and its mode rank is denoted as r_i . The rigorous definition of mode rank is postponed to Section 2. Also throughout the paper, for $i \in [m]$, let $\bar{i} = \min\{j : j \in \Omega_i\}$ be the smallest index in Ω_i , and denote $\check{\Omega}_i := \Omega_i \setminus \{\bar{i}\}$, $\underline{\Omega}_i := \bigcup_{j=1}^{i-1} \Omega_j$, $\bar{\Omega}_i := \bigcup_{j=i+1}^d \Omega_j$ with $\underline{\Omega}_1 = \bar{\Omega}_d = \emptyset$. Finally, we define the signal strength of \mathcal{T} as $\lambda = \min_{i=1, \dots, m} \sigma_{r_i}(\mathcal{M}_{\bar{i}}(\mathcal{T}))$, the smallest singular values of matricization of \mathcal{T} in modes $\{\bar{i}\}_{i=1}^m$. Here for any matrix \mathbf{D} , $\sigma_i(\mathbf{D})$ denotes the i th largest singular values of \mathbf{D} and $\mathcal{M}_{\bar{i}}(\mathcal{T})$ represents the matricization of tensor \mathcal{T} along mode \bar{i} and its formal definition will be given in (4) in Section 2.

In summary, for the dimensions of the perturbation model (1), we have $\tilde{\mathcal{T}}, \mathcal{T}, \mathcal{Z} \in \mathbb{R}^{p_{1'} \times \dots \times p_{d'}}$ with symmetric index groups $\{\Omega_i\}_{i=1}^m$. The HOOI algorithm we study is provided in Algorithm 1. It is worth noting the original HOOI algorithm in De Lathauwer et al. (2000a) mainly focuses on asymmetric tensor decomposition and we generalize it to accommodate arbitrary symmetric structures of \mathcal{T} characterized by $\{\Omega_i\}_{i=1}^m$. In addition, in the literature (De Lathauwer et al., 2000a; Kolda and Bader, 2009), HOOI often refers to the overall procedure including both the initialization of $\tilde{\mathbf{U}}_i^{(0)}$ by HOSVD and the orthogonal iteration updates as detailed in Algorithm 1. We shall point out that this paper studies the orthogonal iteration with any initializers satisfying some mild conditions. Our results accommodate different types of initialization scheme and can be applied to a wide range of scenarios (see Remark 2). We conclude the algorithm by remarking that the matricization mode we choose in group i to perform SVD in (3) does not matter due to symmetry. For simplicity, we choose the smallest index in group i .

Algorithm 1 Higher-Order Orthogonal Iteration for Tensor Decomposition

Input: $\tilde{\mathcal{T}} \in \mathbb{R}^{p_{1'} \times \dots \times p_{d'}}$, symmetric index groups $\{\Omega_i\}_{i=1}^m$, initialization $\{\tilde{\mathbf{U}}_i^{(0)}\}_{i=1}^m$ with $\tilde{\mathbf{U}}_i^{(0)} \in \mathbb{O}_{p_i, r_i}$ having orthonormal columns, maximum number of iterations t_{\max} .

Output: $\{\hat{\mathbf{U}}_i\}_{i=1}^m, \hat{\mathcal{T}}$.

1: For $t = 1, \dots, t_{\max}$, do

(a) For $i = 1$ to m , update

$$\begin{aligned} \tilde{\mathbf{U}}_i^{(t+1)} = \text{SVD}_{r_i} & \left(\mathcal{M}_{\bar{i}}(\tilde{\mathcal{T}} \times_{\Omega_1} (\tilde{\mathbf{U}}_1^{(t+1)})^\top \times \dots \right. \\ & \left. \times_{\Omega_{i-1}} (\tilde{\mathbf{U}}_{i-1}^{(t+1)})^\top \times_{\check{\Omega}_i} (\tilde{\mathbf{U}}_i^{(t)})^\top \times \dots \times_{\Omega_d} (\tilde{\mathbf{U}}_d^{(t)})^\top \right), \end{aligned} \quad (3)$$

where for any matrix \mathbf{D} , $\text{SVD}_r(\mathbf{D})$ computes the subspace composed of the leading r left singular vectors of \mathbf{D} .

2: Let $\hat{\mathbf{U}}_i = \tilde{\mathbf{U}}_i^{(t_{\max})}$ for $i = 1, \dots, m$ and compute

$$\hat{\mathcal{T}} = \tilde{\mathcal{T}} \times_{\Omega_1} P_{\hat{\mathbf{U}}_1} \times \dots \times_{\Omega_m} P_{\hat{\mathbf{U}}_m}.$$

1.2 Our Contributions

Under the perturbation model (1) and the HOOI Algorithm 1, we make the following major contributions to the tensor perturbation theory of low-rank tensor decomposition based on HOOI.

- We provide the first sharp blockwise perturbation bounds of HOOI for tensors with guarantees for both the estimated mode- k singular subspace and the tensor reconstruction induced by low-rank approximation of the noise corrupted tensor. Specifically, in Theorem 3, we show that under suitable initialization $\{\tilde{\mathbf{U}}_i^{(0)}\}_{i=1}^m$ in Algorithm 1, the upper bound of $\max_i \left\| \sin \Theta \left(\hat{\mathbf{U}}_i, \mathbf{U}_i \right) \right\|_q$ converges linearly with respect to the iteration number to a fixed quantity characterized by \mathcal{Z} and the signal strength. In addition, a practically useful tensor reconstruction error bound in Hilbert-Schmidt norm is provided. Surprisingly, we found the upper bound of tensor reconstruction is free of the “condition number” of the signal tensor and can be expressed by a unified simple quantity ξ characterized only by the noise tensor \mathcal{Z} and the underlying multilinear rank of \mathcal{T} . ξ is closely related to the Gaussian width (Gordon, 1988), a common measure for the complexity of a given set, and its formal definition and more explanations will be given in Sections 2.1 and 6.
- In addition, we also generalize the main results to the case that only a subset of modes of \mathcal{T} have low-rank structure.
- Furthermore, we provide a deterministic minimax lower bound for the tensor reconstruction error under perturbation model (1) in Theorem 2. The lower bound matches the perturbation upper bound in Theorem 3 when the tensor order d is fixed, which demonstrates the optimality of HOOI for tensor reconstruction.
- In addition, by combining Theorems 2 and 3, we prove that the tensor reconstruction error rate of HOOI with only one iteration is also optimal and further iterations improve the coefficient in front of the error rate ξ . This suggests that in some applications where running full HOOI is too expensive and prohibitive compared to truncated HOSVD (De Lathauwer et al., 2000b) or sequentially truncated HOSVD (Vannieuwenhoven et al., 2012), we can just run HOOI for one iteration to obtain an optimal (up to constant) reconstruction. Details are provided in Remark 6 and numerical comparison is given in Section 7.2.
- In addition, we apply the new perturbation bounds of HOOI in two modern applications, tensor denoising and tensor co-clustering, from machine learning and statistics. Based on our perturbation results, we can easily recover the results of tensor denoising in current literature with a much shorter proof and provide the first guarantee of HOOI on tensor cocluster recovery that improves the state-of-the-art results.
- Finally, we perform extensive numerical studies to support our perturbation bounds and do a comparison with other existing low-rank tensor decomposition algorithms.

1.3 Related Literature

In this section, we give a brief overview of the literature on Tucker decomposition of tensors, matrix/tensor perturbation theory related to this article. Tensor decomposition has become one of the most important topics in the literature on tensors (Kolda and Bader, 2009; Sidiropoulos et al., 2017; Cichocki et al., 2015). An analogy of matrix SVD for Tucker decomposition of tensors, which is today commonly called high-order singular value decomposition (HOSVD), was first proposed in Tucker (1966) and then popularized by De Lathauwer et al. (2000b). However, unlike the truncated matrix SVD, truncated HOSVD (T-HOSVD) can provide a reasonable but not necessarily optimal low-multilinear-rank approximation for a given tensor. It has been shown that computing the best low multilinear rank approximation is NP hard in general (Hillar and Lim, 2013). On the other hand, various computationally efficient algorithms have been proposed to obtain better approximations than HOSVD (De Lathauwer et al., 2000a; Elden and Savas, 2009; Ishteva et al., 2011, 2009; Savas and Lim, 2010). Among them, the higher-order orthogonal iteration (HOOI) in De Lathauwer et al. (2000a) has become a popular choice in literature. HOOI has been applied to problems including but not limited to tensor PCA/tensor SVD (Richard and Montanari, 2014; Zhang and Xia, 2018), tensor completion (Xia et al., 2020), tensor regression (Zhang et al., 2020), hypergraph community recovery (Ke et al., 2019; Jing et al., 2020), independent component analysis (De Lathauwer and Vandewalle, 2004), tensor clustering (Luo and Zhang, 2020). Many variants of HOOI such as sparse high-order singular value decomposition (STAT-SVD) for tensors (Zhang and Han, 2019), regularized HOOI (Ke et al., 2019; Jing et al., 2020), generalized higher-order orthogonal iteration (gHOI) (Liu et al., 2014) have been proposed. Nowadays, HOOI has become a prevalent choice to compute the low-rank tensor approximation in many applications and been coded in common tensor software such as Matlab “Tensor Toolbox” (Bader and Kolda, 2012), “Tensorlab” (Sorber et al., 2014) and R “rTensor” package (Li et al., 2018). Moreover, it has been regarded as the gold standard guideline for comparison when developing even faster randomized or memory-efficient algorithms for low-rank tensor approximation (Sun et al., 2020; Malik and Becker, 2018; Tsourakakis, 2010; Kolda and Sun, 2008).

Perturbation theory is a long-existing field in mathematics. In particular, the perturbation theory on matrices has attracted much attention. The original work in matrix dated back to Weyl, Davis-Kahan and Wedin (Davis and Kahan, 1970; Wedin, 1972; Weyl, 1912; Stewart, 1998) (see Bhatia (2013); Stewart (1990) for an overview of classical perturbation results and historical development) and recently it has been further developed in Yu et al. (2014); Cai and Zhang (2018); Cape et al. (2019b). In addition, various generalizations and extensions have been made in different settings including random perturbation (Vu, 2011; Shabalin and Nobel, 2013; O’Rourke et al., 2018; Wang, 2015; Benaych-Georges and Nadakuditi, 2011; Abbe et al., 2020; Koltchinskii and Xia, 2016; Benaych-Georges and Nadakuditi, 2012; Cape et al., 2019a; Chen et al., 2018), structured perturbation (Fan et al., 2018; Stewart, 2006) and many others (Eldridge et al., 2017). Also the perturbation theory for matrices has been widely applied to a number of applications such as community detection (Rohe et al., 2011; Chaudhuri et al., 2012; Chin et al., 2015; Sussman et al., 2012; Cape et al., 2019b), covariance matrix estimation (Fan et al., 2018; Cape et al., 2019b), matrix denoising (Cai and Zhang, 2018), matrix completion (Cai et al., 2016), etc.

Studying perturbation theory for low-rank tensor approximation is much harder than matrices and there are only a few attempts in literature. Perturbation results for the best low multilinear rank approximation have been developed in De Lathauwer (2004); Elden and Savas (2011). These results are promising but less practically useful due to the computational hardness of the best multilinear rank approximation (Hillar and Lim, 2013). Hence more attention are given to the perturbation results for polynomial-time low-rank tensor approximation algorithms. A line of work has been done when the perturbation is random (Richard and Montanari, 2014; Zhang and Xia, 2018; Hopkins et al., 2015; Liu et al., 2017; Zhang and Han, 2019; Zheng and Tomioka, 2015; Xia and Zhou, 2019) and various perturbation results for robust tensor decomposition with sparse noise have been studied in Goldfarb and Qin (2014); Gu et al. (2014); Shah et al. (2015); Anandkumar et al. (2016). Perturbation bounds for orthogonally decomposable tensors have been studied in Mu et al. (2017); Auddy and Yuan (2020). In addition, Anandkumar et al. (2014a,b) provided perturbation guarantees for power iteration algorithm for symmetric orthogonal and non-orthogonal CP low-rank decomposition. However, we are not aware of any perturbation result for polynomial time algorithms under partial symmetric multilinear low-rank setting. In this paper, we make the first attempt in this direction and provide the first perturbation bounds of HOOI for tensors with guarantees for both singular subspaces and tensor reconstruction in the general setting. It is worth mentioning that the reconstruction error bound of this paper is often significantly better than the simple estimator and truncated HOSVD without power iteration. This is fundamentally different from the perturbation results in Anandkumar et al. (2014a). See further discussions in Remarks 5 and 8.

We end this section by remarking that in most situations there is a trade-off about the quality of low-rank tensor decomposition and computational cost of the algorithm. For example, computing truncated HOSVD and sequentially truncated HOSVD (ST-HOSVD) (Vannieuwenhoven et al., 2012) may be much faster than iterative algorithms such as HOOI, (quasi-)Newton-Grassmann method (Elden and Savas, 2009; Savas and Lim, 2010), geometric Newton method (Ishteva et al., 2009) and Riemannian trust region scheme (Ishteva et al., 2011) in the large scale settings. On the other hand, these iterative algorithms achieve higher accuracy. In the perturbation model (1), we show HOOI could achieve optimal tensor reconstruction error, which is not true for HOSVD and ST-HOSVD in general.

1.4 Organization of the Paper

The remainder of the article is organized as follows. In Section 2, after a brief introduction of notation and preliminaries, we define various blockwise errors of \mathcal{Z} as the key quantities in our perturbation bounds. We illustrate our main perturbation theorem in asymmetric order-3 case in Section 3 and at the end of the same section, we provide the deterministic lower bound for tensor reconstruction. In Section 4, we provide the perturbation bounds of HOOI applying on a corrupted general partial symmetric order- d tensor. In Section 5, we discuss the tensor perturbation bounds when the target tensor has the low-rank structure only along a subset of modes. In Section 6, we apply our perturbation bounds to two applications, tensor denoising and tensor co-clustering. In Section 7, we corroborate our theoretical results by extensive numerical studies. Conclusion and discussions are provided in Section 8. Due to space constraints, all technical proofs are postponed to the Appendix.

2. Notation and Preliminaries

The following notation will be used throughout this article. For any non-negative integer n , let $[n] = \{1, \dots, n\}$. Lowercase letters (e.g., a, b), lowercase boldface letters (e.g., \mathbf{u}, \mathbf{v}), uppercase boldface letters (e.g., \mathbf{U}, \mathbf{V}), and boldface calligraphic letters (e.g., \mathcal{T}, \mathcal{Z}) are used to denote scalars, vectors, matrices, and order-3-or-higher tensors, respectively. For any two series of numbers, say $\{a_n\}$ and $\{b_n\}$, denote $a \asymp b$ or $a = O(b)$ if $ca_n \leq b_n \leq Ca_n$ or $a_n \leq Cb_n$ for some uniform constants $c, C > 0$. For any matrix $\mathbf{D} \in \mathbb{R}^{m \times n}$ with singular value decomposition $\sum_{i=1}^{m \wedge n} \sigma_i(\mathbf{D}) \mathbf{u}_i \mathbf{v}_i^\top$ ($\sigma_1(\mathbf{D}) \geq \dots \geq \sigma_{m \vee n}(\mathbf{D})$), let $\mathbf{D}_{\max(r)} = \sum_{i=1}^r \sigma_i(\mathbf{D}) \mathbf{u}_i \mathbf{v}_i^\top$ be the leading rank- r SVD approximation of \mathbf{D} and $\mathbf{D}_{\max(-r)} = \sum_{i=r+1}^{m \wedge n} \sigma_i(\mathbf{D}) \mathbf{u}_i \mathbf{v}_i^\top$ be its complement. We also denote $\text{SVD}_r(\mathbf{D}) := [\mathbf{u}_1 \ \dots \ \mathbf{u}_r]$ as the subspace composed of the leading r left singular vectors of \mathbf{D} . The Schatten- q norm of matrix \mathbf{D} for $q \geq 1$ is defined as $\|\mathbf{D}\|_q := (\sum_{i=1}^{m \wedge n} \sigma_i^q(\mathbf{D}))^{1/q}$. Frobenius norm $\|\cdot\|_F$ and spectral norm $\|\cdot\|$ of a matrix are special cases of Schatten- q norm with $q = 2$ and $q = \infty$. In addition, \mathbf{I}_r represents the r -by- r identity matrix. Let $\mathbb{O}_{p,r} = \{\mathbf{U} : \mathbf{U}^\top \mathbf{U} = \mathbf{I}_r\}$ be the set of all p -by- r matrices with orthonormal columns. For any $\mathbf{U} \in \mathbb{O}_{p,r}$, $P_{\mathbf{U}} = \mathbf{U} \mathbf{U}^\top$ represents the projection matrix onto the column span of \mathbf{U} ; we also use $\mathbf{U}_\perp \in \mathbb{O}_{p,p-r}$ to represent the orthonormal complement of \mathbf{U} . We use bracket subscripts to denote sub-matrices. For example, $\mathbf{D}_{[i_1, i_2]}$ is the entry of \mathbf{D} on the i_1 -th row and i_2 -th column; $\mathbf{D}_{[(r+1):m, :]}$ contains the $(r+1)$ -th to the m -th rows of \mathbf{D} . For any matrices $\mathbf{U} \in \mathbb{R}^{p_1 \times p_2}$ and $\mathbf{V} \in \mathbb{R}^{m_1 \times m_2}$, let

$$\mathbf{U} \otimes \mathbf{V} = \begin{bmatrix} \mathbf{U}_{[1,1]} \cdot \mathbf{V} & \cdots & \mathbf{U}_{[1,p_2]} \cdot \mathbf{V} \\ \vdots & & \vdots \\ \mathbf{U}_{[p_1,1]} \cdot \mathbf{V} & \cdots & \mathbf{U}_{[p_1,p_2]} \cdot \mathbf{V} \end{bmatrix} \in \mathbb{R}^{(p_1 m_1) \times (p_2 m_2)}$$

be the Kronecker product of \mathbf{U} and \mathbf{V} .

For any order- d tensor $\mathcal{T} \in \mathbb{R}^{p_1 \times \dots \times p_d}$, let $\mathcal{M}_k(\cdot)$ be the matricization operation that unfolds or flattens the order- d tensor $\mathcal{T} \in \mathbb{R}^{p_1 \times \dots \times p_d}$ along mode k into the matrix $\mathcal{M}_k(\mathcal{T}) \in \mathbb{R}^{p_k \times p_{-k}}$ and here $p_{-k} := \prod_{j \neq k} p_j$. Specifically, the mode- k matricization of \mathcal{T} is formally defined as

$$\mathcal{M}_k(\mathcal{T}) \in \mathbb{R}^{p_k \times p_{-k}}, \quad (\mathcal{M}_k(\mathcal{T}))_{[i_k, j]} = \mathcal{T}_{[i_1, \dots, i_d]}, \quad j = 1 + \sum_{\substack{l=1 \\ l \neq k}}^d \left\{ (i_l - 1) \prod_{\substack{m=1 \\ m \neq k}}^{l-1} p_m \right\} \quad (4)$$

for any $1 \leq i_l \leq p_l, l = 1, \dots, d$. Also see (Kolda and Bader, 2009, Section 2.4) for more discussion on tensor matricizations. Given two tensors $\mathcal{T}_1, \mathcal{T}_2 \in \mathbb{R}^{p_1 \times \dots \times p_d}$, define their inner product as $\langle \mathcal{T}_1, \mathcal{T}_2 \rangle = \sum_{i_1, \dots, i_d} \langle \mathcal{T}_1_{[i_1, \dots, i_d]}, \mathcal{T}_2_{[i_1, \dots, i_d]} \rangle$. The Hilbert-Schmidt norm of \mathcal{T} is defined as $\|\mathcal{T}\|_{\text{HS}} = (\langle \mathcal{T}, \mathcal{T} \rangle)^{1/2}$. The multilinear rank of a tensor \mathcal{T} , $\text{rank}(\mathcal{T})$, is defined as a d -tuple (r_1, \dots, r_d) , where $r_k = \text{rank}(\mathcal{M}_k(\mathcal{T}))$ is the mode- k rank. For any multilinear rank- (r_1, \dots, r_d) tensor \mathcal{T} , it has Tucker decomposition (Tucker, 1966):

$$\mathcal{T} = \llbracket \mathcal{S}; \mathbf{U}_1, \dots, \mathbf{U}_d \rrbracket := \mathcal{S} \times_1 \mathbf{U}_1 \times \dots \times_d \mathbf{U}_d, \quad (5)$$

where $\mathcal{S} \in \mathbb{R}^{r_1 \times \dots \times r_d}$ is the core tensor and $\mathbf{U}_k = \text{SVD}_{r_k}(\mathcal{M}_k(\mathcal{T}))$ is the mode- k singular subspace. Here, the mode- k product of $\mathcal{T} \in \mathbb{R}^{p_1 \times \dots \times p_d}$ with a matrix $\mathbf{U} \in \mathbb{R}^{p_k \times r_k}$ is denoted

by $\mathcal{T} \times_k \mathbf{U}^\top$ and is of size $p_1 \times \cdots \times p_{k-1} \times r_k \times p_{k+1} \times \cdots \times p_d$, such that

$$(\mathcal{T} \times_k \mathbf{U}^\top)_{[i_1, \dots, i_{k-1}, j, i_{k+1}, \dots, i_d]} = \sum_{i_k=1}^{p_k} \mathcal{T}_{[i_1, i_2, \dots, i_d]} \mathbf{U}_{[i_k, j]}. \quad (6)$$

Given $S = \{i_1, \dots, i_d\}$, it is convenient to denote the product of \mathcal{T} along the modes indexed by S with the same matrix \mathbf{U} and with different matrices $\{\mathbf{U}_i\}$ respectively as

$$\mathcal{T} \times_S \mathbf{U} := \mathcal{T} \times_{i_1} \mathbf{U} \times \cdots \times_{i_d} \mathbf{U}, \quad \mathcal{T} \times_{i \in S} \mathbf{U}_i := \mathcal{T} \times_{i_1} \mathbf{U}_{i_1} \times \cdots \times_{i_d} \mathbf{U}_{i_d}. \quad (7)$$

Given $\mathcal{T} = \llbracket \mathcal{S}; \mathbf{U}_1, \dots, \mathbf{U}_d \rrbracket$, the following relationship of tensor matricization is used frequently in the proof:

$$\mathcal{M}_k(\mathcal{S} \times_1 \mathbf{U}_1 \times_2 \cdots \times_d \mathbf{U}_d) = \mathbf{U}_k \mathcal{M}_k(\mathcal{S})(\mathbf{U}_d^\top \otimes \cdots \otimes \mathbf{U}_{k+1}^\top \otimes \mathbf{U}_{k-1}^\top \otimes \cdots \otimes \mathbf{U}_1^\top). \quad (8)$$

We refer the readers to Kolda (2001); Kolda and Bader (2009) for a more comprehensive survey on tensor algebra.

Finally, we use $\sin \Theta$ distance to measure the difference between two p -by- r column orthogonal matrices $\widehat{\mathbf{U}}$ and \mathbf{U} . Suppose the singular values of $\widehat{\mathbf{U}}^\top \mathbf{U}$ are $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_r \geq 0$. Then $\Theta(\widehat{\mathbf{U}}, \mathbf{U})$ is defined as

$$\Theta(\widehat{\mathbf{U}}, \mathbf{U}) = \text{diag}(\cos^{-1}(\sigma_1), \cos^{-1}(\sigma_2), \dots, \cos^{-1}(\sigma_r)).$$

Common properties of $\sin \Theta$ distance can be found in (Cai and Zhang, 2018, Lemma 1) and (Luo et al., 2020, Lemma 6).

2.1 Blockwise Errors of \mathcal{Z}

In this subsection, we introduce key quantities appearing in the perturbation bounds that characterize the blockwise errors of \mathcal{Z} . For simplicity, we consider order-3 tensors and $\Omega_1 = \{1\}, \Omega_2 = \{2\}, \Omega_3 = \{3\}$ perturbation setting for illustration.

Define the blockwise errors of \mathcal{Z} that characterize the tensor perturbation:

$$\begin{aligned} \tau_1 &= \max_{k \in [3]} \tau_{1k}, \quad \tau_{1k} = \left\| (\mathcal{M}_k(\mathcal{Z} \times_{k+1} \mathbf{U}_{k+1}^\top \times_{k+2} \mathbf{U}_{k+2}^\top))_{\max(r_k)} \right\|_q, \quad k = 1, 2, 3; \\ \tau_2 &= \max_{k \in [3]} \left\{ \max_{\substack{\mathbf{V} \in \mathbb{R}^{(p_{k+1}-r_{k+1}) \times r_{k+1}} \\ \|\mathbf{V}\|_q \leq 1}} \left\| (\mathcal{M}_k(\mathcal{Z} \times_{k+1} (\mathbf{U}_{k+1\perp} \mathbf{V})^\top \times_{k+2} \mathbf{U}_{k+2}^\top))_{\max(r_k)} \right\|_q, \right. \\ &\quad \left. \max_{\substack{\mathbf{V} \in \mathbb{R}^{(p_{k+2}-r_{k+2}) \times r_{k+2}} \\ \|\mathbf{V}\|_q \leq 1}} \left\| (\mathcal{M}_k(\mathcal{Z} \times_{k+1} \mathbf{U}_{k+1}^\top \times_{k+2} (\mathbf{U}_{k+2\perp} \mathbf{V})^\top))_{\max(r_k)} \right\|_q \right\}; \\ \tau_3 &= \max_{k \in [3]} \max_{\substack{\mathbf{V} \in \mathbb{R}^{(p_{k+1}-r_{k+1}) \times r_{k+1}} \\ \mathbf{V}' \in \mathbb{R}^{(p_{k+2}-r_{k+2}) \times r_{k+2}} \\ \|\mathbf{V}\|_q \leq 1, \|\mathbf{V}'\|_q \leq 1}} \left\| (\mathcal{M}_k(\mathcal{Z} \times_{k+1} (\mathbf{U}_{k+1\perp} \mathbf{V})^\top \times_{k+2} (\mathbf{U}_{k+2\perp} \mathbf{V}')^\top))_{\max(r_k)} \right\|_q. \end{aligned} \quad (9)$$

Here all mode indices $(\cdot)_k$ of an order-3 tensor are in the sense of modulo-3, e.g., $r_1 = r_4$, $p_2 = p_5$.




Figure 1: Illustration of $\tau_{11}, \tau_{12}, \tau_{13}$. Here, we assume $\mathbf{U}_k^\top = [\mathbf{I}_{r_k} \mathbf{0}_{r_k \times (p_k - r_k)}]$, $k = 1, 2, 3$, for a better visualization. The red, green, blue blocks represent the corresponding blockwise errors $\tau_{11}, \tau_{12}, \tau_{13}$ in \mathcal{Z} .

τ_1, τ_2, τ_3 represent the maximum of blockwise errors of \mathcal{Z} in the projection spaces expanded by $\mathbf{U}_1, \mathbf{U}_2, \mathbf{U}_3$ and their complements. For example, in Figure 1 we illustrate the blockwise errors characterized by $\tau_{11}, \tau_{12}, \tau_{13}$. τ_2, τ_3 characterize blockwise errors of \mathcal{Z} in a similar fashion but with more complicated projections.

These blockwise errors of \mathcal{Z} are in fact a generalization of error terms in matrix perturbation theory. In the matrix setting $\tilde{\mathbf{T}} = \mathbf{T} + \mathbf{Z}$, let $\hat{\mathbf{U}}, \hat{\mathbf{V}}$ and \mathbf{U}, \mathbf{V} be leading left and right singular vectors of $\tilde{\mathbf{T}}$ and \mathbf{T} , respectively. Then by Wedin's perturbation theory, the upper bounds of $\|\sin \Theta(\hat{\mathbf{U}}, \mathbf{U})\|$ and $\|\sin \Theta(\hat{\mathbf{V}}, \mathbf{V})\|$ involve $\|\mathbf{Z}\hat{\mathbf{V}}\|$ and $\|\hat{\mathbf{U}}^\top \mathbf{Z}\|$, which are also blockwise errors of \mathbf{Z} .

Next, we introduce a simple quantity ξ that characterizes the error bound for tensor reconstruction. In this order-3 asymmetric setting, ξ is defined as

$$\xi := \sup_{\|\mathcal{Y}\|_{\text{HS}} \leq 1, \text{rank}(\mathcal{Y}) \leq (r_1, r_2, r_3)} \langle \mathcal{Z}, \mathcal{Y} \rangle. \quad (10)$$

In the following Lemma 1, we give another equivalent characterization of ξ .

Lemma 1 (Equivalent Characterizations of ξ)

$$\begin{aligned} \xi &:= \sup_{\|\mathcal{Y}\|_{\text{HS}} \leq 1, \text{rank}(\mathcal{Y}) \leq (r_1, r_2, r_3)} \langle \mathcal{Z}, \mathcal{Y} \rangle \\ &= \sup_{\mathbf{U}_i \in \mathbb{O}_{p_i, r_i}, 1 \leq i \leq 3} \|\mathcal{Z} \times_1 \mathbf{U}_1^\top \times \mathbf{U}_2^\top \times_d \mathbf{U}_3^\top\|_{\text{HS}}. \end{aligned}$$

By Lemma 1, ξ measures the maximum magnitude of the projection of \mathcal{Z} onto low rank subspaces in Hilbert-Schmidt norm. Another interpretation of ξ is from Gaussian width (Gordon, 1988), which will be discussed in Section 6.

Although the exact values of τ_j and ξ may be hard to compute in general, it is often practical to provide probabilistic bounds when we impose distributional assumptions on \mathcal{Z} . For example, if \mathcal{Z} is a random tensor with i.i.d. standard normal entries and consider $r_1 = r_2 = r_3 = r, p_1 = p_2 = p_3 = p, q = \infty$, then by random matrix theory (Vershynin,

2010), we can show that with high probability $\tau_{1i} \asymp (\sqrt{p} + r)$, $\tau_2 \asymp \sqrt{pr}$, $\tau_3 \asymp \sqrt{pr}$ and $\xi \asymp (r^{\frac{3}{2}} + \sqrt{pr})$. More details about the upper bounds for τ_j and ξ can be found in tensor denoising and tensor co-clustering applications in Section 6.

3. Illustration of Perturbation Bounds for $d = 3$ Asymmetric Case

In this section, we present our main results in the $d = 3$ asymmetric case to better illustrate the main ideas in this paper. The specialized HOOI algorithm for the $d = 3$ asymmetric case is given in Algorithm 2 and its guarantee is provided in Theorem 1.

Algorithm 2 Higher-Order Orthogonal Iteration for Tensor Decomposition ($d = 3$)

Input: $\tilde{\mathcal{T}} \in \mathbb{R}^{p_1 \times p_2 \times p_3}$, initialization $\{\tilde{\mathbf{U}}_i^{(0)}\}_{i=1}^3$ with $\tilde{\mathbf{U}}_i^{(0)} \in \mathbb{O}_{p_i, r_i}$, maximum number of iterations t_{\max} .

Output: $\{\hat{\mathbf{U}}_i\}_{i=1}^3, \hat{\mathcal{T}}$.

1: For $t = 1, \dots, t_{\max}$, do

$$\begin{aligned}\tilde{\mathbf{U}}_1^{(t+1)} &= \text{SVD}_{r_1} \left(\mathcal{M}_1 \left(\tilde{\mathcal{T}} \times_2 (\tilde{\mathbf{U}}_2^{(t)})^\top \times_3 (\tilde{\mathbf{U}}_3^{(t)})^\top \right) \right), \\ \tilde{\mathbf{U}}_2^{(t+1)} &= \text{SVD}_{r_2} \left(\mathcal{M}_2 \left(\tilde{\mathcal{T}} \times_1 (\tilde{\mathbf{U}}_1^{(t+1)})^\top \times_3 (\tilde{\mathbf{U}}_3^{(t)})^\top \right) \right), \\ \tilde{\mathbf{U}}_3^{(t+1)} &= \text{SVD}_{r_3} \left(\mathcal{M}_3 \left(\tilde{\mathcal{T}} \times_1 (\tilde{\mathbf{U}}_1^{(t+1)})^\top \times_2 (\tilde{\mathbf{U}}_2^{(t+1)})^\top \right) \right).\end{aligned}$$

2: Let $\hat{\mathbf{U}}_i = \tilde{\mathbf{U}}_i^{(t_{\max})}$ for $i = 1, 2, 3$ and compute

$$\hat{\mathcal{T}} = \tilde{\mathcal{T}} \times_1 P_{\hat{\mathbf{U}}_1} \times_2 P_{\hat{\mathbf{U}}_2} \times_3 P_{\hat{\mathbf{U}}_3}.$$

Theorem 1 (Tensor Perturbation Bounds for HOOI ($d = 3$)) *Consider the perturbation model (1) with $\tilde{\mathcal{T}}, \mathcal{T}, \mathcal{Z} \in \mathbb{R}^{p_1 \times p_2 \times p_3}$. Suppose $q \geq 1$. Define the block-wise errors as in (9) (10) and denote the initialization errors of $\{\tilde{\mathbf{U}}_k^{(0)}\}_{k=1}^3$ as $\tilde{e}_0 := \max_{k=1,2,3} \|\tilde{\mathbf{U}}_{k\perp}^{(0)\top} \mathbf{U}_k\|$, $e_0 := \max_{k=1,2,3} \|\tilde{\mathbf{U}}_{k\perp}^{(0)\top} \mathbf{U}_k\|_q$. Assume the initialization error and the signal strength satisfy*

$$\tilde{e}_0 \leq \sqrt{2}/2 \text{ and } \lambda \geq (20 + 28\sqrt{2})\xi. \quad (11)$$

Let $\tilde{\mathcal{T}}^{(t)} := \tilde{\mathcal{T}} \times_1 P_{\tilde{\mathbf{U}}_1^{(t)}} \times_2 P_{\tilde{\mathbf{U}}_2^{(t)}} \times_3 P_{\tilde{\mathbf{U}}_3^{(t)}}$ be the estimator of \mathcal{T} after t steps in Algorithm 1.

Then with inputs $\tilde{\mathcal{T}}, \{\tilde{\mathbf{U}}_i^{(0)}\}_{i=1}^3$, the mode- k singular subspace updates in Algorithm 2 after t iterations satisfy

$$\max_{k=1,2,3} \left\| \sin \Theta \left(\tilde{\mathbf{U}}_k^{(t)}, \mathbf{U}_k \right) \right\|_q \leq \frac{8\tau_1}{\lambda} + \frac{e_0}{2^t}, \quad (12)$$

and the t -step tensor estimation satisfies

$$\left\| \tilde{\mathcal{T}}^{(t)} - \mathcal{T} \right\|_{\text{HS}} \leq \left(1 + \frac{6}{1 - \left(8\frac{\tau_1}{\lambda} + \frac{\tilde{e}_0}{2^{t-1}} \right)^2} \right) \xi. \quad (13)$$

Moreover, when $t_{\max} \geq C \log(e_0 \lambda / \tau_1) \vee 1$ for some $C > 0$, the outputs of the estimated mode- k singular subspaces of Algorithm 2 satisfy

$$\max_{k=1,2,3} \left\| \sin \Theta \left(\hat{\mathbf{U}}_k, \mathbf{U}_k \right) \right\|_q \leq \frac{9\tau_1}{\lambda}, \quad (14)$$

$$\left\| \sin \Theta \left(\hat{\mathbf{U}}_k, \mathbf{U}_k \right) \right\|_q = \left\| \hat{\mathbf{U}}_{k\perp}^\top \mathbf{U}_k \right\|_q \leq 4 \left(\frac{\tau_{1k}}{\lambda} + \frac{18\tau_1\tau_2}{\lambda^2} + \frac{81\tau_1^2\tau_3}{\lambda^3} \right), \quad k = 1, 2, 3, \quad (15)$$

and the output of tensor reconstruction $\hat{\mathcal{T}}$ satisfies

$$\left\| \hat{\mathcal{T}} - \mathcal{T} \right\|_{\text{HS}} \leq \left\| [\mathcal{Z}; \hat{\mathbf{U}}_1^\top, \hat{\mathbf{U}}_2^\top, \hat{\mathbf{U}}_3^\top] \right\|_{\text{HS}} + \sum_{k=1}^3 \left\| \hat{\mathbf{U}}_{k\perp}^\top \mathcal{M}_k(\mathcal{T}) \right\|_F \leq 13\xi. \quad (16)$$

Remark 1 (Noise Tolerance and Least Singular Value λ) Our theory relies on a lower bound assumption of the least singular value: $\lambda \geq C\xi$, which is in the same vein as the classical matrix perturbation theory (Davis and Kahan, 1970; Wedin, 1972). Moreover, in the existing results on perturbation analysis for Canonical-Polyadic (CP) decomposition, e.g., Theorem 5.1 of Anandkumar et al. (2014a) and Theorem 1 of Anandkumar et al. (2014b), one assumes $\lambda \geq C\|\mathcal{Z}\|$. Since $\|\mathcal{Z}\| = \sup_{\mathbf{y}: \|\mathbf{y}\|_{\text{HS}} \leq 1, \text{rank}(\mathbf{y})=(1, \dots, 1)} \langle \mathcal{Z}, \mathbf{y} \rangle$, ξ defined in (10) can be seen as a counterpart of $\|\mathcal{Z}\|$ in Tucker decomposition.

Remark 2 (Initialization) In Theorem 1, we assume the initialization $\{\tilde{\mathbf{U}}_i^{(0)}\}_{i=1}^3$ is warm in the sense that the maximum error \tilde{e}_0 is upper bounded by a constant. The constant $\sqrt{2}/2$ in this upper bound is chosen for convenience and can be replaced by any fixed constant less than 1. Our perturbation bound applies to HOOI with any initialization as long as this condition holds, although the original HOOI algorithm was proposed with the initialization scheme named HOSVD, i.e., $\hat{\mathbf{U}}_k^{(0)} = \text{SVD}_{r_k} \left(\mathcal{M}_k(\tilde{\mathcal{T}}) \right)$. Next, we briefly discuss two specific initialization schemes: HOSVD for tensor PCA/SVD (Richard and Montanari, 2014; Zhang and Xia, 2018) and diagonal-deletion SVD for tensor completion (Xia et al., 2020). For convenience of presentation, we focus on the setting $p_1 = p_2 = p_3 = p$.

- (Tensor Denoising) Suppose we observe a tensor $\tilde{\mathcal{T}} = \mathcal{T} + \mathcal{Z} \in \mathbb{R}^{p \times p \times p}$ and aim to recover \mathcal{T} from $\tilde{\mathcal{T}}$. To this end, we can apply HOOI by inputting $\tilde{\mathcal{T}}$. When \mathcal{Z} has i.i.d. $N(0, \sigma^2)$ entries, Theorem 1 in Zhang and Xia (2018) showed if one initializes by HOSVD, as long as $\lambda \geq Cp^{3/4}$, the initialization condition $\left\| \sin \Theta \left(\tilde{\mathbf{U}}_k^{(0)}, \mathbf{U}_k \right) \right\| \leq \frac{\sqrt{2}}{2}$ holds with high probability. Zhang and Xia (2018) also showed the signal strength requirement $\lambda \geq Cp^{3/4}$ is essential, which means HOSVD is a proper initialization in the tensor denoising model.
- (Tensor Completion) Suppose we observe a set of entries, selected uniformly at random and indexed by Ω , from a noisy tensor $\bar{\mathcal{T}} = \mathcal{T} + \mathcal{Z}$. Denote $\bar{\mathcal{T}}_\Omega$ as

$$(\bar{\mathcal{T}}_\Omega)_{[i_1, \dots, i_d]} = \begin{cases} \bar{\mathcal{T}}_{[i_1, \dots, i_d]}, & (i_1, \dots, i_d) \in \Omega; \\ 0 & \text{otherwise.} \end{cases}$$

Suppose \mathcal{Z} has i.i.d. $N(0, \sigma^2)$ entries. Then, it is easy to check that $\tilde{\mathcal{T}}_\Omega/\rho$ is an unbiased estimator of \mathcal{T} , where ρ is the sampling ratio. Xia et al. (2020) proposed to apply HOOI on $\tilde{\mathcal{T}} := \tilde{\mathcal{T}}_\Omega/\rho$ to estimate \mathcal{T} . They proposed to set $\tilde{\mathbf{U}}_k^{(0)}$ as the leading r_k singular vectors of $\mathcal{M}_k(\tilde{\mathcal{T}})\mathcal{M}_k(\tilde{\mathcal{T}})^\top$ with diagonal deletion (i.e., zero the diagonal values of $\mathcal{M}_k(\tilde{\mathcal{T}})\mathcal{M}_k(\tilde{\mathcal{T}})^\top$) and showed that $\|\sin \Theta(\tilde{\mathbf{U}}_k^{(0)}, \mathbf{U}_k)\| \leq \frac{\sqrt{2}}{2}$ holds with high probability when $|\Omega| \geq Cp^{3/2}$, where $|\Omega|$ is the cardinality of Ω . At the same time, they proved that HOSVD requires $|\Omega| \geq Cp^2$ to achieve the same initialization performance and may not be an ideal initialization scheme for tensor completion.

In addition, the random initialization is also widely considered in the literature. For example, Anandkumar et al. (2014a) proposed to pick the best one among many random trials. It can be proved that if the number of random trials is large enough (usually polynomial in the dimension), one can find a trial such that the initialization is good enough (Anandkumar et al., 2014a).

Remark 3 (Mode- k Singular Subspace Linear Convergence Property) The upper bound in (12) includes two parts: a fixed quantity that represents the intrinsic estimation error, and another quantity that decays linearly to 0 with respect to iteration index t . The linear convergence of HOOI was observed in Ishteva et al. (2011), while Theorem 1 gives a rigorous proof for it. Note that HOOI can be viewed as a special alternating minimization method, which was shown to have asymptotic linear convergence rate in solving nonlinear least squares problems (Ruhe and Wedin, 1980). This fact also sheds light on the linear convergence of HOOI.

Remark 4 (Unilateral Perturbation Bounds for Mode- k Singular Subspace)

Our tensor perturbation bounds on singular subspace share the same spirit as the unilateral perturbation bounds on singular subspaces of matrix SVD in Cai and Zhang (2018). Consider the matrix perturbation setting mentioned in Section 2.1 with the additional assumption that \mathbf{T} is rank- r and has SVD $\mathbf{U}\Sigma\mathbf{V}^\top$. Cai and Zhang (2018) showed that the upper bound of $\|\sin \Theta(\hat{\mathbf{U}}, \mathbf{U})\|$ can be written as $\frac{a_1}{\lambda} + \frac{a_2}{\lambda^2}$, which can be interpreted as the sum of first and second order perturbations. In Theorem 1, the upper bound of $\left\|\sin \Theta\left(\hat{\mathbf{U}}_k, \mathbf{U}_k\right)\right\|_q$ can be also written as $\sum_{i=1}^3 \frac{a_i}{\lambda^i}$ which can be interpreted as summation of the first, second, and third order perturbations. This phenomenon also generalizes to order- d case in Theorem 3.

Due to the unilateral property, when the tensor dimension of each mode is at different order, the estimation error rate of singular subspace in each mode can vary significantly. For example in the tensor denoising setting, $\tilde{\mathcal{T}} = \mathcal{T} + \mathcal{Z}$ where $\mathcal{T} \in \mathbb{R}^{p_1 \times p_2 \times p_3}$ is a multilinear rank- (r_1, r_2, r_3) tensor and \mathcal{Z} is a random tensor with i.i.d. standard normal entries. Let $r_{\max} = \max_i r_i$ and suppose $p_1 \ll p_2 \ll p_3$, $r_{\max} \ll p_1^{1/2}$. Consider $q = \infty$, then by random matrix theory (Vershynin, 2010), we can show $\tau_{1i} \leq C\sqrt{p_i}$, $\tau_2 \leq C(\sqrt{p_3 r_{\max}})$ and $\tau_3 \leq C(\sqrt{p_3 r_{\max}})$ with high probability. Thus when $\lambda \geq Cp_3\sqrt{\frac{r_{\max}}{p_1}}$, Theorem 1 immediately implies, with high probability

$$\left\|\sin \Theta\left(\hat{\mathbf{U}}_k, \mathbf{U}_k\right)\right\| \leq C\frac{\sqrt{p_k}}{\lambda}, \quad k = 1, 2, 3$$

for some $C > 0$. So we can see the perturbation of $\widehat{\mathbf{U}}_k$ depends on p_k . Also as λ decreases, for different k , different order perturbations in (15) could dominate in the perturbation bound of $\widehat{\mathbf{U}}_k$. For example, when $\lambda \asymp Cp_3\sqrt{\frac{r_{\max}}{p_2}}$, Theorem 1 yields

$$\begin{aligned} \left\| \sin \Theta \left(\widehat{\mathbf{U}}_1, \mathbf{U}_1 \right) \right\| &\leq C \frac{\sqrt{p_3^2 r_{\max}}}{\lambda^2}, \\ \left\| \sin \Theta \left(\widehat{\mathbf{U}}_2, \mathbf{U}_2 \right) \right\| &\leq C \frac{\sqrt{p_2}}{\lambda} + C' \frac{\sqrt{p_3^2 r_{\max}}}{\lambda^2}, \\ \left\| \sin \Theta \left(\widehat{\mathbf{U}}_3, \mathbf{U}_3 \right) \right\| &\leq C \frac{\sqrt{p_3}}{\lambda} \end{aligned}$$

for constants $C, C' > 0$. More details about the application of HOOI perturbation bounds in order- d tensor denoising and numerical studies for this unilateral property of singular subspace perturbation can be found in Sections 6.1 and 7.1, respectively.

Remark 5 (Comparing Perturbation Bounds of truncated HOSVD and HOOI)

It is worth mentioning that the power iteration in Algorithm 1 plays an important role for refining tensor reconstruction. Without power iteration, the estimator $\widehat{\mathcal{T}}^{\text{T-HOSVD}} = \widetilde{\mathcal{T}} \times_1 P_{\widetilde{\mathbf{U}}_1^{(0)}} \times_2 P_{\widetilde{\mathbf{U}}_2^{(0)}} \times_3 P_{\widetilde{\mathbf{U}}_3^{(0)}}$ with $\widetilde{\mathbf{U}}_i^{(0)} = \text{SVD}_{r_i}(\mathcal{M}_i(\widetilde{\mathcal{T}}))$ is called truncated HOSVD (T-HOSVD) in the literature (De Lathauwer et al., 2000b). It is not hard to show $\|\widehat{\mathcal{T}}^{\text{T-HOSVD}} - \mathcal{T}\|_{\text{HS}} \leq C\|\mathcal{Z}\|_{\text{HS}}$ for some $C > 0$. Since $\|\mathcal{Z}\|_{\text{HS}} = \sup_{\|\mathcal{Y}\|_{\text{HS}} \leq 1} \langle \mathcal{Z}, \mathcal{Y} \rangle$ and may be much larger than ξ , we can see the power iteration can greatly improve the accuracy for tensor reconstruction, and this echos the findings in literature in tensor denoising setting (Zhang and Xia, 2018).

The following lemma provides an alternative way to bound $\left\| \left[\mathcal{Z}; \widehat{\mathbf{U}}_1^\top, \widehat{\mathbf{U}}_2^\top, \widehat{\mathbf{U}}_3^\top \right] \right\|_{\text{HS}}$ appearing in the reconstruction error bound (16).

Lemma 2 Suppose $\mathcal{Z} \in \mathbb{R}^{p_1 \times \dots \times p_d}$ is a general order- d tensor and $\mathbf{U}_k, \widehat{\mathbf{U}}_k \in \mathbb{O}_{p_k, r_k}$ are general matrices with orthonormal columns. For any subset $\Omega \subseteq \{1, \dots, d\}$, we further define projections of \mathcal{Z} on Ω as follows,

$$\theta_\Omega = \left\| \mathcal{Z} \times_{k \in \Omega} \mathbf{U}_k^\top \times_{k \in \Omega^c} \mathbf{U}_{k\perp}^\top \right\|_{\text{HS}}.$$

Then,

$$\left\| \left[\mathcal{Z}; \widehat{\mathbf{U}}_1^\top, \widehat{\mathbf{U}}_2^\top, \dots, \widehat{\mathbf{U}}_d^\top \right] \right\|_{\text{HS}} \leq \sum_{\Omega \subseteq \{1, \dots, d\}} \theta_\Omega \prod_{k \in \Omega^c} \left\| \sin \Theta(\widehat{\mathbf{U}}_k, \mathbf{U}_k) \right\|.$$

We end this section by introducing a deterministic rate matching lower bound for tensor reconstruction. Since the statement of the lower bound is relative simple, we state it in general order- d setting. In particular, we consider the following class of $(\mathcal{T}, \mathcal{Z})$ pairs of $p_1 \times \dots \times p_d$ tensors and perturbations,

$$\mathcal{F}_r(\xi) = \left\{ (\mathcal{T}, \mathcal{Z}) : \text{rank}(\mathcal{T}) = (r_1, \dots, r_d) \leq \mathbf{r}, \sup_{\substack{\|\mathcal{Y}\|_{\text{HS}} \leq 1, \\ \text{rank}(\mathcal{Y}) \leq (r_1, \dots, r_d)}} \langle \mathcal{Z}, \mathcal{Y} \rangle \leq \xi \right\},$$

here $\mathbf{r} = (r, \dots, r)$ and the comparison is entrywise.

Theorem 2 (Tensor Reconstruction Lower Bound under Perturbation)

Consider perturbation model (1), we have the following deterministic lower bound for reconstructing \mathcal{T} ,

$$\inf_{\hat{\mathcal{T}}} \sup_{(\mathcal{T}, \mathcal{Z}) \in \mathcal{F}_r(\xi)} \|\hat{\mathcal{T}} - \mathcal{T}\|_{\text{HS}} \geq \frac{\sqrt{2}}{2} \xi.$$

Remark 6 (Optimality of HOOI and one-step HOOI for Tensor Reconstruction)

When tensor order d is fixed, combining Theorem 1 and 2, we have shown that HOOI with good initialization is optimal for tensor reconstruction in the class $(\mathcal{T}, \mathcal{Z}) \in \mathcal{F}_r(\xi)$. At the same time, from (13), we see the error rate of tensor reconstruction is optimal even after one iteration of HOOI i.e., $t_{\max} = 1$ and more iterations can improve the coefficient in front of ξ . This suggests that in some applications where running HOOI until convergence is prohibitive, we can just run it for one iteration to get a fairly good reconstruction. See more in Section 7.2 about a numerical comparison of HOOI and one-step HOOI.

Apart from the optimality of our perturbation bound in tensor reconstruction, it is also interesting to study whether the perturbation bounds in (14), (15) for singular subspaces are optimal or not and we leave it as an interesting future work.

4. A Blockwise Perturbation Bound of Higher-order Orthogonal Iteration for Tensor Decomposition

In this section, we present the main results of perturbation bounds of HOOI given in Algorithm 1. In contrast with Theorem 1, Theorem 3 in this section covers the general order- d perturbation setting with $\tilde{\mathcal{T}}$ having symmetric index groups $\{\Omega_i\}_{i=1}^m$. Before stating the theorem, we first define the blockwise errors of \mathcal{Z} in this general setting. Let $\mathcal{S}_i^{(-\bar{k})} := \{S \subseteq [d] \setminus \{\bar{k}\} : |S| = i\}$ be the collection of all possible index sets with i elements from $[d] \setminus \{\bar{k}\}$ and $\mathcal{S}_0^{(-\bar{k})} := \emptyset$. For $S \in \mathcal{S}_i^{(-\bar{k})}$, we let $S^c = ([d] \setminus \{\bar{k}\}) \setminus S$. Now we define the blockwise errors of \mathcal{Z} as

$$\begin{aligned} \tau_1 &= \max_{k=1, \dots, m} \tau_{1k}, \quad \tau_{1k} = \left\| \left(\mathcal{M}_{\bar{k}}(\mathcal{Z} \times_{i \neq \bar{k}} \mathbf{U}_{i'}^\top) \right)_{\max(r_k)} \right\|_q, \quad k = 1, \dots, m; \\ \tau_j &= \max_{k \in [m]} \left\{ \max_{S \in \mathcal{S}_{j-1}^{(-\bar{k})}} \sup_{\substack{\mathbf{V}_{i'} \in \mathbb{R}^{(p_{i'} - r_{i'}) \times r_{i'}} \\ \|\mathbf{V}_{i'}\|_q \leq 1, i \in S}} \left\| \left(\mathcal{M}_{\bar{k}}(\mathcal{Z} \times_{i \in S} (\mathbf{U}_{i' \perp} \mathbf{V}_{i'})^\top \times_{i \in S^c} \mathbf{U}_{i'}^\top) \right)_{\max(r_k)} \right\|_q \right\}, \quad (17) \\ &\text{for } j = 2, \dots, m. \end{aligned}$$

Finally, ξ in this setting is defined as

$$\xi := \sup_{\|\mathcal{Y}\|_{\text{HS}} \leq 1, \text{rank}(\mathcal{Y}) \leq (r_{1'}, \dots, r_{d'})} \langle \mathcal{Z}, \mathcal{Y} \rangle. \quad (18)$$

Theorem 3 (General Perturbation Bounds for Tensor Power Iteration)

Consider the perturbation model (1) with $\tilde{\mathcal{T}}, \mathcal{T}, \mathcal{Z} \in \mathbb{R}^{p_{1'} \times \dots \times p_{d'}}$, symmetric index groups $(\Omega_1, \dots, \Omega_m)$ and blockwise errors in (17) (18). Suppose $q \geq 1$. Denote the initialization errors of $\{\tilde{\mathbf{U}}_k^{(0)}\}_{k=1}^m$ as $\tilde{e}_0 := \max_{k=1, \dots, m} \|\tilde{\mathbf{U}}_{k \perp}^{(0)\top} \mathbf{U}_k\|$, $e_0 := \max_{k=1, \dots, m} \|\tilde{\mathbf{U}}_{k \perp}^{(0)\top} \mathbf{U}_k\|_q$.

Assume the initialization error and the signal strength satisfy

$$\tilde{e}_0 \leq \frac{\sqrt{2}}{2} \quad \text{and} \quad \lambda \geq 2^{\frac{d+4}{2}} \left(1 + \frac{\sqrt{2}}{2}\right)^d \xi. \quad (19)$$

Let $\tilde{\mathcal{T}}^{(t)} := \tilde{\mathcal{T}} \times_{\Omega_1} P_{\tilde{\mathbf{U}}_1^{(t)}} \times \cdots \times_{\Omega_m} P_{\tilde{\mathbf{U}}_m^{(t)}}$ be the estimator of \mathcal{T} after t steps in Algorithm 1. Then with inputs $\tilde{\mathcal{T}}$, $\{\tilde{\mathbf{U}}_k^{(0)}\}_{k=1}^m$, $\{\Omega_i\}_{i=1}^m$, the mode- k singular subspace updates in Algorithm 1 after t iterations satisfy

$$\max_{k \in [m]} \left\| \sin \Theta \left(\tilde{\mathbf{U}}_k^{(t)}, \mathbf{U}_k \right) \right\|_q \leq 2^{\frac{d+3}{2}} \frac{\tau_1}{\lambda} + \frac{e_0}{2^t}, \quad (20)$$

and the t -step tensor estimation satisfies

$$\left\| \tilde{\mathcal{T}}^{(t)} - \mathcal{T} \right\|_{\text{HS}} \leq \left(1 + 2d \left(1 - \left(2^{\frac{d+3}{2}} \frac{\tau_1}{\lambda} + \frac{\tilde{e}_0}{2^{t-1}} \right)^2 \right)^{-\frac{d-1}{2}} \right) \xi. \quad (21)$$

Moreover, when $t_{\max} \geq C \log(e_0 \lambda / \tau_1) \vee 1$ for some $C > 0$, the outputs of estimated mode- k singular subspace of Algorithm 1 satisfy

$$\max_{k \in [m]} \left\| \sin \Theta \left(\hat{\mathbf{U}}_k, \mathbf{U}_k \right) \right\|_q \leq \left(2^{\frac{d+3}{2}} + 1 \right) \frac{\tau_1}{\lambda},$$

$$\left\| \sin \Theta \left(\hat{\mathbf{U}}_k, \mathbf{U}_k \right) \right\|_q \leq \frac{2}{(1 - c^*(\tau_1, \lambda, d))^{\frac{d-1}{2}}} \left(\frac{\tau_{1k}}{\lambda} + \sum_{j=1}^{d-1} \frac{\binom{d-1}{j} \left(2^{\frac{d+3}{2}} + 1 \right)^j \tau_1^j \tau_{j+1}}{\lambda^{j+1}} \right). \quad (22)$$

for $k = 1, \dots, m$ where $c^*(\tau_1, \lambda, d) := \frac{\left(2^{\frac{d+3}{2}} + 2 \right)^2 \tau_1^2}{\lambda^2} \leq \frac{1}{2}$, and the output of tensor reconstruction $\hat{\mathcal{T}}$ satisfies

$$\begin{aligned} \left\| \hat{\mathcal{T}} - \mathcal{T} \right\|_{\text{HS}} &\leq \left\| \mathcal{Z} \times_{\Omega_1} \hat{\mathbf{U}}_1^\top \times \cdots \times_{\Omega_m} \hat{\mathbf{U}}_m^\top \right\|_{\text{HS}} + \sum_{k=1}^m |\Omega_k| \left\| \hat{\mathbf{U}}_{k\perp}^\top \mathcal{M}_k(\mathcal{T}) \right\|_F \\ &\leq \left(1 + 2d (1 - c^*(\tau_1, \lambda, d))^{-\frac{d-1}{2}} \right) \xi. \end{aligned} \quad (23)$$

Remark 7 (Size of $c^*(\tau_1, \lambda, d)$) It is easy to check $c^*(\tau_1, \lambda, d) \leq \frac{1}{2}$ based on $\tau_1 \leq \xi$ and the requirement of the signal strength λ . So we have $(1 - c^*(\tau_1, \lambda, d))^{-\frac{d-1}{2}} \leq 2^{\frac{d-1}{2}}$ in the upper bounds of $\left\| \sin \Theta \left(\hat{\mathbf{U}}_k, \mathbf{U}_k \right) \right\|_q$ and $\left\| \hat{\mathcal{T}} - \mathcal{T} \right\|_{\text{HS}}$. However, in many practical applications, such as tensor denoising to be introduced in Section 6.1, tensor order d is fixed and $\lambda \gg (2^{\frac{d+3}{2}} + 2)\tau_1$. In this case $c^*(\tau_1, \lambda, d)$ could be much smaller than $\frac{1}{2}$ and the scale of $(1 - c^*(\tau_1, \lambda, d))^{-\frac{d-1}{2}}$ can be very close to 1.

Remark 8 (Comparison with Anandkumar et al. (2014a)) Compared with the perturbation bounds of power iteration for supersymmetric CP-low-rank decomposition (Anandkumar et al., 2014a, Theorem 5.1), our Theorem 3 covers more general symmetric and partial symmetric multilinear low-rank decomposition settings. Also in Theorem 5.1 of Anandkumar et al. (2014a), the tensor reconstruction error bound of power iteration is given in terms of tensor spectral norm, which does not improve upon the guarantee by the trivial estimator $\tilde{\mathcal{T}}$. On the other hand, the tensor reconstruction error of $\hat{\mathcal{T}}$ in Theorem 3 is given in Hilbert-Schmidt norm and can be significantly better than the guarantee for $\tilde{\mathcal{T}}$ as $\|\mathcal{Z}\|_{\text{HS}} \gg \xi$ in most of the applications.

Remark 9 (Dependence on Tensor Order d) We note that in Theorem 3, the constants in our condition (19) and perturbation bounds (20) and (21) scales exponentially w.r.t. the tensor order d . We think this exponential dependence on d is not sharp. In fact, in Theorem 1 of the recent work Luo and Zhang (2021), they show the dependence on d in (19) and (21) can be reduced to $\text{poly}(d)$.

Remark 10 (A Proof Sketch of Theorem 3) We provide a sketch on how to prove (20) and (21). The rest of the results (22) and (23) follow easily from (20), (21) by plugging in $t_{\max} \geq \log(e_0\lambda/\tau_1) \vee 1$. The idea is to develop the recursive error bounds of $\tilde{\mathbf{U}}_k^{(t+1)}$, i.e., the estimate of \mathbf{U}_k at iteration $t+1$, based on the error bound of $\tilde{\mathbf{U}}_k^{(t)}$, i.e., the estimate at iteration t . The argument can be divided into three steps. It is worth mentioning that all three steps involves complex tensor algebra and this makes the proof even more difficult.

First, we denote

$$\begin{aligned} \tilde{e}_t &= \max_k \tilde{e}_{t,k}, & \tilde{e}_{t,k} &= \left\| (\tilde{\mathbf{U}}_{k\perp}^{(t)})^\top \mathbf{U}_k \right\|, \\ e_t &= \max_k e_{t,k}, & e_{t,k} &= \left\| (\tilde{\mathbf{U}}_{k\perp}^{(t)})^\top \mathbf{U}_k \right\|_q, \quad k = 1, \dots, m; t = 0, 1, \dots \end{aligned}$$

Step 1. In HOOI procedure, the update for the mode- k singular subspace satisfies

$$\begin{aligned} \tilde{\mathbf{U}}_k^{(t+1)} &= \text{SVD}_{r_k} \left(\mathcal{M}_{\bar{k}} \left(\mathcal{T} \times_{i \in \underline{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} \times_{\check{\Omega}_k} \tilde{\mathbf{U}}_k^{(t)\top} \times_{i \in \bar{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t)\top} \right) \right. \\ &\quad \left. + \mathcal{M}_{\bar{k}} \left(\mathcal{Z} \times_{i \in \underline{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} \times_{\check{\Omega}_k} \tilde{\mathbf{U}}_k^{(t)\top} \times_{i \in \bar{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t)\top} \right) \right), \end{aligned}$$

here $\underline{\Omega}_i := \bigcup_{j=1}^{i-1} \Omega_j$, $\bar{\Omega}_i := \bigcup_{j=i+1}^d \Omega_j$. To give an upper bound for $e_{t+1,k}$, we aim to give an upper bound for

$$\left\| \left(\mathcal{M}_{\bar{k}} \left(\mathcal{Z} \times_{i \in \underline{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} \times_{\check{\Omega}_k} \tilde{\mathbf{U}}_k^{(t)\top} \times_{i \in \bar{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t)\top} \right) \right)_{\max(r_k)} \right\|_q \quad (24)$$

by using τ_1, \dots, τ_m , e_t, e_{t+1} . The main idea to bound (24) is to introduce $\mathbf{I} = P_{\mathbf{U}_{i'}} + P_{\mathbf{U}_{i'\perp}}$ in each mode multiplication, expand the mode products, then write the whole term into summation of many small terms.

Step 2. After getting an upper bound for (24), we use induction to prove the following claim,

$$e_t \leq 2^{(d+3)/2} \tau_1 / \lambda + e_0 / 2^t, \quad \tilde{e}_t \leq 2^{(d+3)/2} \tau_1 / \lambda + \tilde{e}_0 / 2^t; \quad t = 0, 1, \dots$$

One technical difficulty is to deal with the sequential updating of singular subspaces in HOOI and we use the induction idea again to tackle it. Tools we use in this step include the singular subspace bound in (Luo et al., 2020, Theorem 5).

Step 3. The final and most challenging step involves upper bounding the tensor reconstruction error $\|\tilde{\mathcal{T}} \times_{\Omega_1} P_{\hat{\mathbf{U}}_1} \times \cdots \times_{\Omega_m} P_{\hat{\mathbf{U}}_m} - \mathcal{T}\|_{\text{HS}}$ by the unified quantity ξ . By decomposing \mathcal{T} onto the estimated singular subspaces, we can show that

$$\begin{aligned} & \left\| \tilde{\mathcal{T}} \times_{\Omega_1} P_{\hat{\mathbf{U}}_1} \times \cdots \times_{\Omega_m} P_{\hat{\mathbf{U}}_m} - \mathcal{T} \right\|_{\text{HS}} \\ & \leq \left\| \mathcal{Z} \times_{\Omega_1} P_{\hat{\mathbf{U}}_1} \times \cdots \times_{\Omega_m} P_{\hat{\mathbf{U}}_m} \right\|_{\text{HS}} + \sum_{k=1}^d \left\| \hat{\mathbf{U}}_{k'}^\top \mathcal{M}_k(\mathcal{T}) \right\|_F. \end{aligned}$$

By definition, $\left\| \mathcal{Z} \times_{\Omega_1} P_{\hat{\mathbf{U}}_1} \times \cdots \times_{\Omega_m} P_{\hat{\mathbf{U}}_m} \right\|_{\text{HS}} \leq \xi$. We further show

$$\left\| \hat{\mathbf{U}}_{k'}^\top \mathcal{M}_k(\mathcal{T}) \right\|_F \stackrel{(a)}{\leq} C(\tau_1, d, \lambda) \xi.$$

Here $C(\tau_1, d, \lambda)$ is a quantity that depends on τ_1, d, λ . The main challenge to prove (a) is that $\hat{\mathbf{U}}_{k'}$ is not the left singular subspace of $\mathcal{M}_k(\tilde{\mathcal{T}})$. So to leverage the SVD property of $\hat{\mathbf{U}}_{k'}^\perp$, we have to project $\tilde{\mathcal{T}}$ onto $\tilde{\mathbf{U}}_i^{(t_{\max})}$ and $\tilde{\mathbf{U}}_i^{(t_{\max}-1)}$, then use the subspace perturbation bounds established before.

Note that Theorem 3 covers the general situation where \mathcal{T} may have partial symmetric modes. We provide a corollary for the common asymmetric case, i.e., $\Omega_i = \{i\}, i = 1, \dots, d$ in the Appendix.

5. Perturbation Bounds of Power Iteration for Tensors with Partial Low Multilinear Rank Structure

In some applications, e.g., multilayer network analysis (Lei et al., 2019), the tensor $\tilde{\mathcal{T}}$ only has low-rank structure on a subset of modes. Both the tensor power iteration algorithm and our perturbation theory can be generalized to such cases. For a better illustration, we present the modified algorithm and theory for order-3 tensor perturbation with mode 1 to be dense. Specifically, we consider

$$\tilde{\mathcal{T}} = \mathcal{T} + \mathcal{Z} \in \mathbb{R}^{p_1 \times p_2 \times p_3}, \tag{25}$$

where \mathcal{T} is the signal tensor and \mathcal{Z} is the noise. We assume \mathcal{T} is low-rank on modes 2 and 3, i.e., $\mathcal{T} = \mathcal{S} \times_2 \mathbf{U}_2 \times_3 \mathbf{U}_3$, where $\mathcal{S} \in \mathbb{R}^{p_1 \times r_2 \times r_3}$ is the core tensor and $\mathbf{U}_i \in \mathbb{O}_{p_i, r_i}$ for $i = 2, 3$. In this setting, we consider the modified tensor power iteration algorithm for low-rank tensor decomposition in Algorithm 3.

Algorithm 3 Power Iteration for Tensor Decomposition in Partial Multilinear Low-Rank Setting (25)

Input: $\tilde{\mathcal{T}} \in \mathbb{R}^{p_1 \times p_2 \times p_3}$, initialization $\{\tilde{\mathbf{U}}_i^{(0)} \in \mathbb{O}_{p_i, r_i}\}_{i=2,3}^3$, maximum number of iterations t_{\max} .

Output: $\{\hat{\mathbf{U}}_i\}_{i=2,3}^3, \hat{\mathcal{T}}$.

1: For $t = 1, \dots, t_{\max}$, do

$$\begin{aligned}\tilde{\mathbf{U}}_2^{(t+1)} &= \text{SVD}_{r_2} \left(\mathcal{M}_2 \left(\tilde{\mathcal{T}} \times_3 (\tilde{\mathbf{U}}_3^{(t)})^\top \right) \right) \\ \tilde{\mathbf{U}}_3^{(t+1)} &= \text{SVD}_{r_3} \left(\mathcal{M}_3 \left(\tilde{\mathcal{T}} \times_2 (\tilde{\mathbf{U}}_2^{(t+1)})^\top \right) \right).\end{aligned}$$

2: Let $\hat{\mathbf{U}}_i = \tilde{\mathbf{U}}_i^{(t_{\max})}$ for $i = 2, 3$ and compute

$$\hat{\mathcal{T}} = \tilde{\mathcal{T}} \times_2 P_{\hat{\mathbf{U}}_2} \times_3 P_{\hat{\mathbf{U}}_3}.$$

In the setting (25), we can define the blockwise errors of \mathcal{Z} as follows:

$$\begin{aligned}\tau_1 &= \max_{k=2,3} \tau_{1k}, \quad \tau_{12} = \left\| \left(\mathcal{M}_2(\mathcal{Z} \times_3 \mathbf{U}_3^\top) \right)_{\max(r_2)} \right\|_q, \quad \tau_{13} = \left\| \left(\mathcal{M}_3(\mathcal{Z} \times_2 \mathbf{U}_2^\top) \right)_{\max(r_3)} \right\|_q \\ \tau_2 &= \max \left\{ \max_{\substack{\mathbf{V} \in \mathbb{R}^{(p_3-r_3) \times r_3} \\ \|\mathbf{V}\|_q \leq 1}} \left\| \left(\mathcal{M}_2(\mathcal{Z} \times_3 (\mathbf{U}_{3\perp} \mathbf{V})^\top) \right)_{\max(r_2)} \right\|_q, \right. \\ &\quad \left. \max_{\substack{\mathbf{V} \in \mathbb{R}^{(p_2-r_2) \times r_2} \\ \|\mathbf{V}\|_q \leq 1}} \left\| \left(\mathcal{M}_3(\mathcal{Z} \times_2 (\mathbf{U}_{2\perp} \mathbf{V})^\top) \right)_{\max(r_3)} \right\|_q \right\}; \\ \xi &= \sup_{\text{rank}(\mathcal{Y}) \leq (p_1, r_2, r_3), \|\mathcal{Y}\|_{\text{HS}} \leq 1} \langle \mathcal{Z}, \mathcal{Y} \rangle.\end{aligned} \tag{26}$$

We have the following perturbation bounds for the outputs of Algorithm 3.

Theorem 4 (Tensor Perturbation Bounds with Partial Multilinear Low-Rank)

Consider the perturbation model (25) with $\tilde{\mathcal{T}}, \mathcal{T}, \mathcal{Z} \in \mathbb{R}^{p_1 \times p_2 \times p_3}$. Suppose $q \geq 1$. Define the blockwise errors as in (26) and denote the initialization errors of $\{\tilde{\mathbf{U}}_k^{(0)}\}_{k=2,3}^3$ as $\tilde{e}_0 := \max_{k=2,3} \|\tilde{\mathbf{U}}_{k\perp}^{(0)\top} \mathbf{U}_k\|$, $e_0 := \max_{k=2,3} \|\tilde{\mathbf{U}}_{k\perp}^{(0)\top} \mathbf{U}_k\|_q$. Assume the initialization error and the signal strength satisfy

$$\tilde{e}_0 \leq \sqrt{2}/2 \text{ and } \lambda \geq 16\xi. \tag{27}$$

Then with inputs $\tilde{\mathcal{T}}, \{\tilde{\mathbf{U}}_i^{(0)}\}_{i=2,3}^3$, the mode- k singular subspace updates in Algorithm 3 after t iterations satisfy

$$\max_{k=2,3} \left\| \sin \Theta \left(\tilde{\mathbf{U}}_k^{(t)}, \mathbf{U}_k \right) \right\|_q \leq \frac{4\sqrt{2}\tau_1}{\lambda} + \frac{e_0}{2^t}. \tag{28}$$

Moreover, when $t_{\max} \geq C \log(e_0 \lambda / \tau_1) \vee 1$ for some $C > 0$, the outputs of estimated mode- k singular subspace of Algorithm 3 satisfy

$$\max_{k=2,3} \left\| \sin \Theta \left(\hat{\mathbf{U}}_k, \mathbf{U}_k \right) \right\|_q \leq \frac{(4\sqrt{2} + 1)\tau_1}{\lambda},$$

$$\left\| \sin \Theta \left(\hat{\mathbf{U}}_k, \mathbf{U}_k \right) \right\|_q = \left\| \hat{\mathbf{U}}_{k\perp}^\top \mathbf{U}_k \right\|_q \leq 2\sqrt{2} \left(\frac{\tau_{1k}}{\lambda} + \frac{(4\sqrt{2} + 1)\tau_1\tau_2}{\lambda^2} \right), \quad k = 2, 3. \quad (29)$$

The output of tensor reconstruction $\hat{\mathcal{T}}$ satisfies

$$\left\| \hat{\mathcal{T}} - \mathcal{T} \right\|_{\text{HS}} \leq \left\| \mathcal{Z} \times_2 P_{\hat{\mathbf{U}}_2} \times_3 P_{\hat{\mathbf{U}}_3} \right\|_{\text{HS}} + \sum_{k=2}^3 \left\| \hat{\mathbf{U}}_{k\perp}^\top \mathcal{M}_k(\mathcal{T}) \right\|_F \leq (4\sqrt{2} + 1)\xi.$$

The proof of Theorem 4 follows the proof of Theorem 3 easily. For simplicity, we omit it here.

6. Implications in Statistics and Machine Learning

In this section, we consider a couple of applications of the HOOI perturbation bounds we developed in statistics and machine learning. Specifically, here we consider the perturbation model (1) and assume $\mathcal{Z}_{i_1, \dots, i_d}$'s are independent, mean-zero σ -subgaussian, where $\sigma > 0$ is the subgaussianity parameter. More precisely,

$$\mathbb{E} \exp(\lambda \mathcal{Z}_{i_1, \dots, i_d}) \leq \exp(C\lambda^2 \sigma^2), \text{ for all } (i_1, \dots, i_d) \in [p_1] \times \dots \times [p_d] \text{ and all } \lambda \in \mathbb{R},$$

where $C > 0$ is some absolute constant. For convenience, we let $p_{\max} = \max_i p_i, p_{\min} = \min_i p_i, r_{\max} = \max_i r_i, r_{\min} = \min_i r_i$.

In this setting, the quantity ξ is in fact closely related to the Gaussian width (Gordon, 1988) studied in literature that measures the size or complexity of a given set. Recall the Gaussian width of a set $\mathcal{S} \subset \mathbb{R}^{p_1 \times \dots \times p_d}$ is defined to be

$$w(\mathcal{S}) := \mathbb{E} \left(\sup_{\mathcal{Y} \in \mathcal{S}} \langle \mathcal{B}, \mathcal{Y} \rangle \right),$$

where $\mathcal{B} \in \mathbb{R}^{p_1 \times \dots \times p_d}$ is a tensor whose entries are independent $N(0, 1)$ random variables. In view of $w(\mathcal{S})$, we can regard ξ as the Gaussian width with no expectation and $\mathcal{S} = \{\mathcal{Y} : \|\mathcal{Y}\|_{\text{HS}} \leq 1, \text{rank}(\mathcal{Y}) \leq (r_1, \dots, r_d)\}$. It can be shown in the case that \mathcal{Z} has i.i.d. $N(0, 1)$ entries, ξ and $w(\mathcal{S})$ are the same up to constant with high probability (Raskutti et al., 2019).

In the following subsections, we consider two particular structures of \mathcal{T} , one is pure low multilinear rank structure, namely tensor denoising/tensor PCA/tensor SVD studied in literature (Richard and Montanari, 2014; Zhang and Xia, 2018; Perry et al., 2020; Hopkins et al., 2015) and another one is tensor co-clustering/block structure (Chi et al., 2020; Wang and Zeng, 2019).

6.1 HOOI for Tensor Denoising

In tensor denoising, we assume \mathcal{T} has the following structure,

$$\mathcal{T} = \mathcal{S} \times_1 \mathbf{U}_1 \times \dots \times_d \mathbf{U}_d, \quad (30)$$

where $\mathcal{S} \in \mathbb{R}^{r_1 \times \dots \times r_d}$ is the core tensor and $\{\mathbf{U}_i \in \mathbb{O}_{p_i, r_i}\}_{i=1}^d$ are loading matrices. With the established tensor perturbation bounds, we can establish the following theoretical guarantee for the performance of HOOI on tensor denoising with a very short proof.

Theorem 5 (Tensor Denoising: General Order d) Consider the tensor denoising problem “(1) + (30)” and Algorithm 1 with inputs $\tilde{\mathcal{T}}$, $\Omega_i = \{i\}$, initialization $\{\tilde{\mathbf{U}}_i^{(0)}\}_{i=1}^d$ and $t_{\max} = C \left(\log \frac{\lambda/\sigma}{\sqrt{p_{\max}}} \vee 1 \right)$ for some $C > 0$, where $\lambda = \min_k \sigma_{r_k}(\mathcal{M}_k(\mathcal{S}))$ is the minimal singular value of each matricization of \mathcal{S} . Assume $r_{\max} \leq p_{\min}^{1/(d-1)}$ and $\max_i \|\tilde{\mathbf{U}}_{i\perp}^{(0)\top} \mathbf{U}_i\| \leq \sqrt{2}/2$. Then if $\lambda/\sigma \geq 2^{(d+4)/2} (1 + \sqrt{2}/2)^d \sqrt{p_{\max} r_{\max}}$, with probability at least $1 - \exp(-cp_{\min})$, the output $\hat{\mathbf{U}}_k, \hat{\mathcal{T}}$ satisfy

$$\left\| \sin \Theta \left(\hat{\mathbf{U}}_k, \mathbf{U}_k \right) \right\| \leq C \left(1 - \frac{c}{r_{\max}} \right)^{-(d-1)/2} \left(\frac{\sqrt{p_k}}{\lambda/\sigma} + \frac{\sqrt{p_{\max}^2 r_{\max}}}{(\lambda/\sigma)^2} \right)$$

and

$$\|\hat{\mathcal{T}} - \mathcal{T}\|_{\text{HS}} \leq C \left(1 + 2d \left(1 - \frac{c}{r_{\max}} \right)^{-\frac{d-1}{2}} \right) \sigma \sqrt{\sum_{i=1}^d p_i r_i}, \quad (31)$$

for some constants $c, C > 0$.

When d is a constant, the upper bound for tensor reconstruction error matches the lower bound in (Zhang and Xia, 2018, Theorem 3), which shows HOOI achieves the optimal tensor reconstruction error in the tensor denoising problem.

6.2 Tensor Co-clustering/Block Model

Co-clustering is among the most important unsupervised learning methods that reveals the checkbox-like association pattern in data. A number of algorithms have been proposed (Wang et al., 2015; Wu et al., 2016; Kolda and Sun, 2008; Papalexakis et al., 2012; Sun et al., 2009; Jegelka et al., 2009) for tensor co-clustering in the literature, however most of the work does not provide statistical guarantees for recovering the underlying co-clustering structure. Very recently, Chi et al. (2020) and Wang and Zeng (2019) studied the performance of co-clustering estimation and cocluster recovery based on convex relaxation and combinatorial search algorithms. By using the tools of perturbation bounds of HOOI given in Section 4, we are able to provide the first guarantee for co-clustering estimation and cocluster recovery based on computational efficient HOOI algorithm. Compared to the convex relaxation approach (Chi et al., 2020), HOOI has a better guarantee for tensor reconstruction and it also gives guarantee for cocluster membership recovery. Specifically, in the tensor co-clustering/block model, we assume \mathcal{T} has the following structure,

$$\mathcal{T} = \mathcal{B} \times_1 \mathbf{\Pi}_1 \times \cdots \times_d \mathbf{\Pi}_d, \quad (32)$$

where $\mathbf{\Pi}_i \in \mathbb{M}_{p_i, r_i}$ and \mathbb{M}_{p_i, r_i} is the collection of all $p_i \times r_i$ membership matrices with each row has exactly one 1 and $(r_i - 1)$ 0's. For any $\mathbf{\Pi}_i$, the cocluster membership of node k is denoted by $g_i^{(k)} \in [r_i]$, which satisfies $(\mathbf{\Pi}_i)_{[k, g_i^{(k)}]} = 1$. Let $G_i^{(j)} \equiv G_i^{(j)}(\mathbf{\Pi}_i) = \{k \in [p_i] : g_i^{(k)} = j\}$ be the set of i th mode node indices that belongs to cocluster j and $p_i^{(j)} = |G_i^{(j)}|$ for all $j \in [r_i]$. For simplicity, we assume the cocluster sizes for each cluster are on the same order for every mode, i.e.,

$$p_i^{(1)} \asymp p_i^{(2)} \asymp \cdots \asymp p_i^{(r_i)} \asymp \frac{p_i}{r_i}, \quad \text{for } i = 1, \dots, d. \quad (A1)$$

We consider two cocluster membership recovery error metrics:

1. Let $\|\mathbf{M}\|_0$ be the number of nonzero entries in matrix \mathbf{M} . Suppose \mathbf{E}_{r_i} is the set of all $r_i \times r_i$ permutation matrices. Define

$$err(\hat{\mathbf{\Pi}}_i, \mathbf{\Pi}_i) = \frac{1}{p_i} \min_{\mathbf{J} \in \mathbf{E}_{r_i}} \|\hat{\mathbf{\Pi}}_i \mathbf{J} - \mathbf{\Pi}_i\|_0 \quad (33)$$

as the misclassification rate of $\hat{\mathbf{\Pi}}_i$.

2. Define

$$\widetilde{err}(\hat{\mathbf{\Pi}}_i, \mathbf{\Pi}_i) = \min_{\mathbf{J} \in \mathbf{E}_{r_i}} \max_{1 \leq j \leq r_i} \frac{1}{p_i^{(j)}} \|(\hat{\mathbf{\Pi}}_i \mathbf{J})_{[G_i^{(j)}, :]} - (\mathbf{\Pi}_i)_{[G_i^{(j)}, :]}\|_0.$$

Intuitively speaking, $\widetilde{err}(\hat{\mathbf{\Pi}}_i, \mathbf{\Pi}_i)$ measures the worst relative misclassification rates over all communities.

It is easy to check that $0 \leq err(\hat{\mathbf{\Pi}}_i, \mathbf{\Pi}_i) \leq \widetilde{err}(\hat{\mathbf{\Pi}}_i, \mathbf{\Pi}_i) \leq 2$.

The following lemma gives a Tucker decomposition of \mathcal{T} in the tensor block model (32). This decomposition bridges the tensor co-clustering model and the tensor signal-plus-noise model, which explains why HOOI would work for tensor co-clustering.

Lemma 3 *Suppose \mathcal{T} has the tensor co-clustering/block structure (32), where \mathcal{B} is a multilinear rank- (r_1, \dots, r_d) tensor. Assume $\mathcal{S} \times_1 \mathbf{V}_1 \times \dots \times_d \mathbf{V}_d$ with $\mathbf{V}_i \in \mathbb{O}_{r_i, r_i}$ is the Tucker decomposition of $\mathcal{B} \times_1 (\mathbf{\Pi}_1^\top \mathbf{\Pi}_1)^{\frac{1}{2}} \times \dots \times (\mathbf{\Pi}_d^\top \mathbf{\Pi}_d)^{\frac{1}{2}}$. Then*

$$\mathcal{T} = \mathcal{S} \times_1 \mathbf{U}_1 \times \dots \times_d \mathbf{U}_d$$

with $\mathbf{U}_i = \mathbf{\Pi}_i (\mathbf{\Pi}_i^\top \mathbf{\Pi}_i)^{-\frac{1}{2}} \mathbf{V}_i \in \mathbb{O}_{p_i, r_i}$ for $i = 1, \dots, d$.

The following Theorem 6 gives the theoretical guarantee on the performance of Algorithm 4 for tensor reconstruction and cocluster membership recovery.

Algorithm 4 HOOI for Tensor Co-clustering/Block Model

Input: Tensor $\tilde{\mathcal{T}} \in \mathbb{R}^{p_1 \times \dots \times p_d}$, indices group $\Omega_i = \{i\}$, initialization $\tilde{\mathbf{U}}_i^{(0)} \in \mathbb{R}^{p_i \times r_i}$ for $i = 1, \dots, d$, maximum number of iterations t_{\max} ,

Output: $\hat{\mathbf{\Pi}}_i \in \mathbb{M}_{p_i, r_i}, i = 1, \dots, d$ and $\hat{\mathcal{T}}$.

- 1: Apply Algorithm 1 with input $\tilde{\mathcal{T}}, \{\Omega_i\}_{i=1}^d, \{\tilde{\mathbf{U}}_i^{(0)}\}_{i=1}^d$, maximum number of iterations t_{\max} and get outputs $\{\hat{\mathbf{U}}_i\}_{i=1}^d$ and $\hat{\mathcal{T}}$.
- 2: For each mode i , apply ϵ -approximation K-means (Kumar et al., 2004) on $\hat{\mathbf{U}}_i$, i.e., compute $\hat{\mathbf{\Pi}}_i \in \mathbb{M}_{p_i, r_i}, \hat{\mathbf{X}}_i \in \mathbb{R}^{r_i \times r_i}$ such that

$$\|\hat{\mathbf{\Pi}}_i \hat{\mathbf{X}}_i - \hat{\mathbf{U}}_i\|_F^2 \leq (1 + \epsilon) \min_{\mathbf{\Pi} \in \mathbb{M}_{p_i, r_i}, \mathbf{X} \in \mathbb{R}^{r_i \times r_i}} \|\mathbf{\Pi} \mathbf{X} - \hat{\mathbf{U}}_i\|_F^2. \quad (34)$$

Theorem 6 (HOOI for Tensor Co-clustering/Block Model) Consider the tensor co-clustering/block model “(1) + (32)” and the Algorithm 4 with inputs $\tilde{\mathcal{T}}$, initializations $\{\tilde{\mathbf{U}}_i^{(0)}\}_{i=1}^d$ and $t_{\max} = C \left(\log \left(\frac{\lambda/\sigma}{\sqrt{p_{\max}}} \sqrt{\frac{\prod_{i=1}^d p_i}{\prod_{i=1}^d r_i}} \right) \vee 1 \right)$ for some $C > 0$, where $\lambda = \min_k \sigma_{r_k}(\mathcal{M}_k(\mathcal{B}))$ is the minimal singular value at each matricization of the core tensor parameter \mathcal{B} . Assume $r_{\max} \leq p_{\min}^{1/(d-1)}$, $\max_i \|\tilde{\mathbf{U}}_{i\perp}^{(0)\top} \mathbf{U}_i\| \leq \frac{\sqrt{2}}{2}$, and (A1) holds. Then if

$$\lambda/\sigma \geq C \sqrt{\frac{p_{\max}^2 r_{\max} \prod_{i=1}^d r_i}{p_{\min} \prod_{i=1}^d p_i}},$$

for sufficiently large constant $C > 0$, with probability at least $1 - \exp(-cp_{\min})$ for some $c > 0$, $\hat{\mathbf{U}}_k, \hat{\mathcal{T}}$ satisfy

$$\|\sin \Theta(\hat{\mathbf{U}}_k, \mathbf{U}_k)\| \leq C(d) \frac{\sqrt{p_k}}{\lambda/\sigma} \sqrt{\frac{\prod_{i=1}^d r_i}{\prod_{i=1}^d p_i}}, \quad k = 1, \dots, d$$

$$\|\hat{\mathcal{T}} - \mathcal{T}\|_{\text{HS}} \leq C(d) \sigma \sqrt{\sum_{i=1}^d p_i r_i};$$

and we also have the following upper bound on cocluster recovery error,

$$\begin{aligned} \text{err}(\hat{\mathbf{\Pi}}_i, \mathbf{\Pi}_i) &\leq C_1(d, \epsilon) \frac{p'_{i, \max} r_i \prod_{i=1}^d r_i}{(\lambda/\sigma)^2 \prod_{i=1}^d p_i}, \\ \tilde{\text{err}}(\hat{\mathbf{\Pi}}_i, \mathbf{\Pi}_i) &\leq C_2(d, \epsilon) \frac{p_i r_i \prod_{i=1}^d r_i}{(\lambda/\sigma)^2 \prod_{i=1}^d p_i}. \end{aligned}$$

Here $C(d), C_1(d, \epsilon), C_2(d, \epsilon) > 0$ are some constants depending only on d and ϵ , $p'_{i, \max}$ is the second largest cocluster size at mode i .

Note that our cocluster recovery guarantee is new for polynomial-time algorithms. When $p = p_1 = \dots = p_d$, the best polynomial time algorithm guarantee for tensor reconstruction is p^{d-1} in Chi et al. (2020) and our result can be significantly better.

7. Numerical Studies

In this section, we first provide numerical studies to support the main theoretical results in Section 4 and then compare HOOI with other existing algorithms for tensor decomposition in applications of tensor denoising and tensor co-clustering. Throughout the simulation, we consider order-3 tensor perturbation setting $\tilde{\mathcal{T}} = \mathcal{T} + \mathcal{Z}$ with \mathcal{Z} being the noise tensor with i.i.d. $N(0, \sigma^2)$ entries. Without particular specification, we set $p = p_1 = p_2 = p_3, r = r_1 = r_2 = r_3$. The error metrics we consider for tensor reconstruction and mode- k singular subspace estimation are root mean square error (RMSE) $\|\hat{\mathcal{T}} - \mathcal{T}\|_{\text{HS}}$ and $\|\sin \Theta(\hat{\mathbf{U}}_k, \mathbf{U}_k)\|$, respectively. All simulations are repeated 100 times and the average statistics are reported.




Figure 2: HOOI with good initialization. (a) Tensor reconstruction error $\|\hat{\mathcal{T}} - \mathcal{T}\|_{\text{HS}}$ for $p \in \{20, 30, \dots, 100\}$, $r = 5$, $\sigma \in \{1, 2, 3, 4\}$ and $\lambda = 5\sqrt{pr}\sigma$; (b) Mode- k singular subspace estimation with and without rescaling under $p_1 = 10, p_2 = 100, p_3 = 500$, $r \in \{3, 5\}$, $\sigma = 1$ and $\lambda = \alpha \cdot p_3 \frac{\sqrt{r_1}}{\sqrt{p_1}}$ with varying α

7.1 Perturbation Bounds of HOOI with good initialization

In this simulation, we study the perturbation bounds of HOOI with randomly generated good initialization. Let $\mathcal{T} = \mathcal{S} \times_1 \mathbf{U}_1 \times_2 \mathbf{U}_2 \times_3 \mathbf{U}_3$, where $\mathbf{U}_i \in \mathbb{R}^{p_i \times r}$ is generated uniformly at random from $\mathbb{O}_{p_i, r}$ and $\mathcal{S} \in \mathbb{R}^{r \times r \times r}$ is a diagonal tensor with diagonal values $\{i\lambda\}_{i=1}^r$. The initializations of \mathbf{U}_i of Algorithm 1 are $\tilde{\mathbf{U}}_i^{(0)} = \frac{1}{\sqrt{2}}\mathbf{U}_i + \frac{1}{\sqrt{2}}\mathbf{U}'_i$, where $\mathbf{U}'_i = \mathbf{U}_{i\perp}\mathbf{O}$ for some random orthogonal matrix $\mathbf{O} \in \mathbb{O}_{p_i-r, r}$. It is easy to check that $\|\sin \Theta(\mathbf{U}_i, \tilde{\mathbf{U}}_i^{(0)})\| = \frac{\sqrt{2}}{2}$ for $i = 1, 2, 3$.

First for tensor reconstruction, let $p \in \{20, 30, \dots, 100\}$, $r = 5$, $\sigma \in \{1, 2, 3, 4\}$ and $\lambda = 5\sqrt{pr}\sigma$. We can check that with high probability, $\|\mathcal{Z}\|_{\text{HS}} \leq Cp^{\frac{3}{2}}\sigma$ and $\xi \leq C\sqrt{pr}\sigma$ for some $C > 0$ following the same proof as Theorem 5. In Figure 2(a), the RMSE of tensor reconstruction of HOOI is presented. We find as the perturbation results in Section 4 suggest, $\|\hat{\mathcal{T}} - \mathcal{T}\|_{\text{HS}}$ can be much smaller than $\|\mathcal{Z}\|_{\text{HS}}$. This demonstrates the superior performance of the HOOI estimator compared to the trivial estimator $\hat{\mathcal{T}}$. At the same time, the RMSE for tensor reconstruction increases as p and σ become bigger and this matches our theoretical findings in Theorem 3 that the error bound of HOOI for $\|\hat{\mathcal{T}} - \mathcal{T}\|_{\text{HS}}$ is $O(\xi)$, which increases as p, σ increase.

Next we demonstrate the unilateral perturbation bounds for mode- k singular subspace estimation. Specifically, we consider $p_1 = 10, p_2 = 100, p_3 = 500$, $r \in \{3, 5\}$, $\sigma = 1$ and $\lambda = \alpha \cdot p_3 \frac{\sqrt{r}}{\sqrt{p_1}}$ with varying α . The errors of the mode-1, mode-2, mode-3 estimated singular subspaces with and without rescaling are provided in Figure 2(b). We can see from Figure 2(b) left panel the errors of estimated singular subspaces converge to different values depending on the corresponding mode size p_i , and a further rescaling of estimation error by $\sqrt{p_i}$ makes them roughly on the same level (see Figure 2(b) right panel). This matches the unilateral property of the singular subspace perturbation results in Remark 4 that when $\lambda = O(p_3 \frac{\sqrt{r}}{\sqrt{p_1}})$, $\|\sin \Theta(\hat{\mathbf{U}}_k, \mathbf{U}_k)\| \leq C \frac{\sqrt{p_i}}{\lambda/\sigma}$ for some $C > 0$, and this upper bound increases linearly with respect to $\sqrt{p_i}$.

	HOOI	O-HOOI	ST-HOSVD	T-HOSVD
Complexity	$O(p^{d+1} + t_{\max} dr p^d)$	$O(p^{d+1} + dr p^d)$	$O(p^{d+1})$	$O(dp^{d+1})$

Table 1: Time Complexity of HOOI, O-HOOI, ST-HOSVD and T-HOSVD under setting $p_1 = \dots = p_d = p, r_1 = \dots = r_d = r, r \ll p$. HOOI and O-HOOI is initialized by ST-HOSVD.

7.2 Comparison of HOOI with other Algorithms

In this section, we do a comparison of HOOI with truncated HOSVD (T-HOSVD) (De Lathauwer et al., 2000b) and sequentially truncated HOSVD (ST-HOSVD) (Vannieuwenhoven et al., 2012) in the tensor denoising and tensor co-clustering applications. We also include one-step HOOI (O-HOOI), since it might be useful as a surrogate of HOOI in large scale tensor decomposition settings as we mentioned in Remark 6. The initialization we consider for HOOI and O-HOOI are ST-HOSVD with natural truncation order, i.e., $\tilde{\mathbf{U}}_i^{(0)} = \text{SVD}_r(\mathcal{M}_i(\tilde{\mathcal{T}} \times_{j < i} \tilde{\mathbf{U}}_j^{(0)}))$. In Table 1, we give the time complexity of HOOI, O-HOOI, ST-HOSVD and T-HOSVD. We can see that as long as $dr \leq p$, a common case in practice, the time complexity of O-HOOI and ST-HOSVD are on the same order, and they could be faster than full HOOI and HOSVD in general.

In tensor denoising, the generating process of \mathcal{T} is the same as before. Let $p = 100$, $r = 5$, $\sigma \in \{1, 2\}$, $\lambda = \alpha \cdot p^{\frac{3}{4}} \sigma$ with varying α . The comparison of these algorithms for tensor reconstruction and singular subspace estimation are given in Figure 3. First, we find that HOOI is best in both tensor reconstruction and singular subspace estimation among four algorithms. Meanwhile, O-HOOI is slightly worse than HOOI for small α and has very close performance with HOOI when α is relative large, which suggests that in some computationally heavy applications, we can just run HOOI for one iteration to achieve reasonable estimation. Part of this phenomenon can be explained by the one-iteration optimality of HOOI for tensor reconstruction as we discussed in Remark 6. On the other hand, HOOI and O-HOOI are often much better than T-HOSVD and ST-HOSVD for both tensor reconstruction and mode- k singular subspace estimation within a wide range of settings.

Finally, we study the performance of the HOOI-based Algorithm 4 in tensor co-clustering recovery and do a comparison of it with T-HOSVD, ST-HOSVD, and O-HOOI based clustering algorithms. In this simulation, we generate $\mathcal{T} = \mathcal{B} \times_1 \mathbf{\Pi}_1 \times_2 \mathbf{\Pi}_2 \times_3 \mathbf{\Pi}_3$ such that $\{\mathbf{\Pi}_i\}_{i=1}^3$ have balanced cluster size and $\mathcal{B} = \frac{\mathcal{B}_0}{\min_i \sigma_r(\mathcal{M}_i(\mathcal{B}_0))} \lambda$ with $\mathcal{B}_0 \stackrel{i.i.d.}{\sim} N(0, 1)$. The error metric we consider is the average cocluster membership misclassification error rate in (33). The performance of Algorithm 4 under $p \in \{50, 80\}$, $r \in \{3, 5, 8\}$, $\sigma = 1$, $\lambda = \alpha \cdot \frac{r^{3/2}}{p^{3/4}} \sigma$ is presented in Figure 4(a). We can see the misclassification error decreases as the signal strength increases and cocluster number decreases. The comparison of Algorithm 4 and T-HOSVD, ST-HOSVD, O-HOOI based spectral clustering is given in Figure 4(b) under the same setting with $r = 5$. Again, HOOI-based algorithm has the best performance in cocluster recovery. O-HOOI and ST-HOSVD perform similarly here and both of them are much better than T-HOSVD.




Figure 3: Comparison of HOOI, one-step HOOI (O-HOOI), truncated HOSVD (T-HOSVD), sequentially truncated HOSVD (ST-HOSVD) in tensor denoising under $p = 100$, $r = 5$, $\sigma \in \{1, 2\}$, $\lambda = \alpha \cdot p^{\frac{3}{4}}\sigma$ with $\alpha \in [1, 4]$. (a) Tensor reconstruction; (b) Averaged singular subspace estimation




Figure 4: Tensor cocluster recovery under $\sigma = 1$, $\lambda = \alpha \cdot \frac{r^{3/2}}{p^{3/4}}\sigma$ with varying α . (a) HOOI on tensor cocluster recovery under $p \in \{50, 80\}$, $r \in \{3, 5, 8\}$. (b) Comparison of HOOI, one-step HOOI (O-HOOI), truncated HOSVD (T-HOSVD), sequentially truncated HOSVD (ST-HOSVD) in cocluster recovery under $p = 80$, $r = 5$.

8. Conclusion and Discussion

In this paper, we provide the first sharp blockwise perturbation bounds of HOOI for tensors with guarantees for both tensor reconstruction and mode- k singular subspace estimation. Furthermore, we show both HOOI and one-step HOOI with good initialization is optimal in terms of tensor reconstruction by providing rate matching lower bound. Finally, we support our theoretical results with extensive numerical studies and apply them to tensor denoising and tensor co-clustering applications. Apart from the applications mentioned above, the main perturbation results can be applied to many other applications where tensor “spectral method” HOOI is applicable, such as tensor completion (Yuan and Zhang, 2016, 2017; Xia et al., 2020; Xia and Yuan, 2017), hypergraphic stochastic block model (Ghoshdastidar and Dukkipati, 2014, 2017; Ke et al., 2019; Chien et al., 2019; Ahn et al., 2018; Kim et al., 2018), multilayer network (Lei et al., 2019; Jing et al., 2020), MPCA (Lu et al., 2008), latent variable model (Anandkumar et al., 2014a), etc. In tensor completion (Xia et al., 2020) and many other applications, more specialized initializers can achieve better performance than HOSVD – the classic initializers for HOOI in the literature (De Lathauwer et al., 2000a). Our tensor perturbation bounds still apply to these cases as our theoretical analysis admits all initializers satisfying certain mild conditions.

At the same time, due to the NP hardness of computing many tensor quantities (Hillar and Lim, 2013), the Alternating Least Square (ALS) and Power iteration have been the “workhorse” algorithms in computing low-rank tensor approximation and solving many other tensor problems (Kolda and Bader, 2009). Our induction proof idea in Theorem 3 could also shed light on how to analyze other iterative ALS/Power iteration procedures for tensor problems (Zhou et al., 2013; Lu et al., 2008; Wang and Zeng, 2019; Xu et al., 2005; Yan et al., 2005, 2006; Lee and Wang, 2020).

The convergence rate of HOOI is another important topic related to the results in this paper. Savas and Lim (2010); Elden and Savas (2009) observed that HOOI converge fast when the target tensor has fast-decaying multilinear singular values (in other words, the tensor is approximately Tucker low-rank). On the other hand, HOOI may converge slowly when the target tensor is high-rank or approximately sparse (Savas and Lim, 2010; Elden and Savas, 2009). In this paper, we studied how HOOI converges when the target tensor is approximately Tucker low-rank. It is interesting to further explore how HOOI converges in the less ideal settings, such as for the high-rank or approximately sparse tensors.

In addition to the perturbation results for HOOI, it is also interesting to develop perturbation results for the second-order algorithms, such as (quasi-)Newton-Grassmann method (Elden and Savas, 2009; Savas and Lim, 2010), geometric Newton method (Ishteva et al., 2009), Riemannian trust region scheme (Ishteva et al., 2011) since these second-order methods may take much fewer iterations than HOOI to converge. Also, this paper mainly focuses on Tucker format of tensor decomposition. Although Tucker format has many advantages, in ultra higher-order tensor problems, the storage cost of the core tensor in Tucker format scales exponentially with respect to the tensor order and it is more desirable to consider other low-rank tensor decomposition formats, such as the hierarchical Tucker decomposition (Ballani and Grasedyck, 2013; Grasedyck, 2010; Hackbusch and Kühn, 2009) and Tensor-Train decomposition (Oseledets, 2011; Oseledets and Tyrtshnikov, 2009; Zhou et al., 2020).

It is interesting to develop the perturbation bounds for algorithms on Hierarchical Tucker or Tensor Train tensor decompositions.

Acknowledgments

The authors would like to thank the Action Editor Animashree Anandkumar and the anonymous referees for constructive suggestions. Y.L. would like to thank Rungang Han for helpful discussions during the project. The research of Y. L. and A. R. Zhang was in part by NSF Grants CAREER-1944904, NSF DMS-1811868, NSF DMS-2023239, NIH Grant R01 GM131399, and Wisconsin Alumni Research Foundation (WARF); the research of G. R. was supported in part by NSF Grant DMS-2015285; the research of M. Y. was supported in part by NSF Grant DMS-1811767.

Appendix A. Tensor Perturbation Bounds for HOOI in Asymmetric Case

In this section, we present a Corollary of Theorem 3 in the case where $\Omega_i = \{i\}$, $i = 1, \dots, d$, which appears frequently in practice.

Corollary 1 (Tensor Perturbation Bounds for HOOI in Asymmetric Case)

Consider the perturbation model (1) with $\tilde{\mathcal{T}}, \mathcal{T}, \mathcal{Z} \in \mathbb{R}^{p_1 \times \dots \times p_d}$ and $\Omega_i = \{i\}$, $i = 1, \dots, d$. Define $\mathcal{S}_i^{(-k)} := \{S \subseteq [d] \setminus \{k\} : |S| = i\}$ as the set of all possible index sets with i elements from $[d] \setminus \{k\}$ and $\mathcal{S}_0^{(-k)} := \emptyset$. For $S \in \mathcal{S}_i^{(-k)}$, let $S^c = ([d] \setminus \{k\}) \setminus S$. Now we define the blockwise errors as

$$\begin{aligned} \tau_1 &= \max_{k=1, \dots, d} \tau_{1k}, \quad \tau_{1k} = \left\| \left(\mathcal{M}_k(\mathcal{Z} \times_{i \neq k} \mathbf{U}_i^\top) \right)_{\max(r_k)} \right\|_q, \quad k = 1, \dots, d; \\ \tau_j &= \max_{k=1, \dots, d} \left\{ \max_{S \in \mathcal{S}_{j-1}^{(-k)}} \sup_{\substack{\mathbf{V}_i \in \mathbb{R}^{(p_i - r_i) \times r_i} \\ \|\mathbf{V}_i\|_q \leq 1, i \in S}} \left\| \left(\mathcal{M}_k(\mathcal{Z} \times_{i \in S} (\mathbf{U}_{i \perp} \mathbf{V}_i)^\top \times_{i \in S^c} \mathbf{U}_i^\top) \right)_{\max(r_k)} \right\|_q \right\}, \end{aligned}$$

for $j = 2, \dots, d$;

Denote the initialization errors of $\{\tilde{\mathbf{U}}_k^{(0)}\}_{k=1}^d$ as $\tilde{e}_0 := \max_{k=1, \dots, d} \|\tilde{\mathbf{U}}_{k \perp}^{(0)\top} \mathbf{U}_k\|$, $e_0 := \max_{k=1, \dots, d} \|\tilde{\mathbf{U}}_{k \perp}^{(0)\top} \mathbf{U}_k\|_q$. Assume the initialization error and the signal strength satisfy

$$\tilde{e}_0 \leq \frac{\sqrt{2}}{2} \quad \text{and} \quad \lambda \geq 2^{\frac{d+4}{2}} \left(1 + \frac{\sqrt{2}}{2} \right)^d \xi,$$

where $\xi := \sup_{\|\mathcal{Y}\|_{\text{HS}} \leq 1, \text{rank}(\mathcal{Y}) \leq (r_1, \dots, r_d)} \langle \mathcal{Z}, \mathcal{Y} \rangle$.

Then with inputs $\tilde{\mathcal{T}}, \{\tilde{\mathbf{U}}_k^{(0)}\}_{k=1}^d, \{\Omega_i\}_{i=1}^d$, the estimated mode- k singular subspaces updates in Algorithm 1 after t iterations satisfy

$$\max_{k \in [d]} \left\| \sin \Theta \left(\tilde{\mathbf{U}}_k^{(t)}, \mathbf{U}_k \right) \right\|_q \leq 2^{\frac{d+3}{2}} \frac{\tau_1}{\lambda} + \frac{e_0}{2^t}.$$

Moreover, when $t_{\max} \geq \log(e_0 \lambda / \tau_1) \vee 1$, the outputs of estimated mode- k singular subspaces of Algorithm 1 satisfy

$$\max_{k \in [d]} \left\| \sin \Theta \left(\hat{\mathbf{U}}_k, \mathbf{U}_k \right) \right\|_q \leq \left(2^{\frac{d+3}{2}} + 1 \right) \frac{\tau_1}{\lambda},$$

$$\left\| \sin \Theta \left(\widehat{\mathbf{U}}_k, \mathbf{U}_k \right) \right\|_q \leq \frac{2}{(1 - c^*(\tau_1, \lambda, d))^{\frac{d-1}{2}}} \left(\frac{\tau_{1k}}{\lambda} + \sum_{j=1}^{d-1} \frac{\binom{d-1}{j} \left(2^{\frac{d+3}{2}} + 1 \right)^j \tau_1^j \tau_{j+1}}{\lambda^{j+1}} \right),$$

for $k = 1, \dots, d$, where $c^*(\tau_1, \lambda, d) = \left(2^{\frac{d+3}{2}} + 2 \right)^2 \frac{\tau_1^2}{\lambda^2} \leq \frac{1}{2}$, and the output of tensor reconstruction $\widehat{\mathcal{T}}$ satisfies

$$\begin{aligned} \left\| \widehat{\mathcal{T}} - \mathcal{T} \right\|_{\text{HS}} &\leq \left\| \llbracket \mathcal{Z}; \widehat{\mathbf{U}}_1^\top, \dots, \widehat{\mathbf{U}}_d^\top \rrbracket \right\|_{\text{HS}} + \sum_{k=1}^d \left\| \widehat{\mathbf{U}}_{k\perp}^\top \mathcal{M}_k(\mathcal{T}) \right\|_F \\ &\leq \left(1 + 2d(1 - c^*(\tau_1, \lambda, d))^{-\frac{d-1}{2}} \right) \xi. \end{aligned}$$

Appendix B. Additional Proofs

B.1 Proof of Lemma 1

For generality, here we present the proof for order d case. First we show the first equivalent characterization.

$$\begin{aligned} \sup_{\|\mathbf{y}\|_{\text{HS}} \leq 1, \text{rank}(\mathbf{y}) \leq (r_1, \dots, r_d)} \langle \mathcal{Z}, \mathbf{y} \rangle &\geq \sup_{\|\mathbf{s}\|_{\text{HS}} \leq 1, \mathbf{U}_i \in \mathbb{O}_{p_i, r_i}} \langle \mathcal{Z}, \mathbf{s} \times_1 \mathbf{U}_1 \times \dots \times_d \mathbf{U}_d \rangle \\ &= \sup_{\|\mathbf{s}\|_{\text{HS}} \leq 1, \mathbf{U}_i \in \mathbb{O}_{p_i, r_i}} \langle \mathcal{Z} \times_1 \mathbf{U}_1^\top \times \dots \times_d \mathbf{U}_d^\top, \mathbf{s} \rangle \\ &= \sup_{\mathbf{U}_i \in \mathbb{O}_{p_i, r_i}} \left\| \mathcal{Z} \times_1 \mathbf{U}_1^\top \times \dots \times_d \mathbf{U}_d^\top \right\|_{\text{HS}}. \end{aligned}$$

On the other hand by Theorem 2 of De Lathauwer et al. (2000b), we have

$$\mathbf{y} = \left(\mathbf{y} \times_1 \widetilde{\mathbf{U}}_1^\top \times \dots \times \widetilde{\mathbf{U}}_d^\top \right) \times_1 \widetilde{\mathbf{U}}_1 \times \dots \times_d \widetilde{\mathbf{U}}_d,$$

where $\widetilde{\mathbf{U}}_i \in \mathbb{R}^{p_i \times r_i}$ is the left singular space of $\mathcal{M}_i(\mathbf{y})$. Since $\|\mathbf{y} \times_1 \widetilde{\mathbf{U}}_1^\top \times \dots \times \widetilde{\mathbf{U}}_d^\top\|_{\text{HS}} \leq 1$,

$$\sup_{\|\mathbf{y}\|_{\text{HS}} \leq 1, \text{rank}(\mathbf{y}) \leq (r_1, \dots, r_d)} \langle \mathcal{Z}, \mathbf{y} \rangle \leq \sup_{\|\mathbf{s}\|_{\text{HS}} \leq 1, \mathbf{U}_i \in \mathbb{O}_{p_i, r_i}} \langle \mathcal{Z}, \mathbf{s} \times_1 \mathbf{U}_1 \times \dots \times_d \mathbf{U}_d \rangle.$$

So, we have proved the lemma. \blacksquare

B.2 Proof of Lemma 2

This proof idea of this lemma is to project \mathcal{Z} onto orthogonal subspaces \mathbf{U}_k and $\mathbf{U}_{k\perp}$ at each modes.

$$\begin{aligned} &\mathcal{Z} \times_1 \widehat{\mathbf{U}}_1^\top \times \dots \times \widehat{\mathbf{U}}_d^\top \\ &= (\mathcal{Z} \times_1 (P_{\mathbf{U}_1} + P_{\mathbf{U}_{1\perp}}) \times \dots \times_d (P_{\mathbf{U}_d} + P_{\mathbf{U}_{d\perp}})) \times_1 \widehat{\mathbf{U}}_1^\top \times \dots \times_d \widehat{\mathbf{U}}_d^\top \\ &= \left(\sum_{\Omega \subseteq [d]} \mathcal{Z} \times_{k \in \Omega} P_{\mathbf{U}_k} \times_{k \in \Omega^c} P_{\mathbf{U}_{k\perp}} \right) \times_1 \widehat{\mathbf{U}}_1^\top \times \dots \times_d \widehat{\mathbf{U}}_d^\top \\ &= \sum_{\Omega \subseteq [d]} \mathcal{Z} \times_{k \in \Omega} \widetilde{\mathbf{U}}_k^\top P_{\mathbf{U}_k} \times_{k \in \Omega^c} \widetilde{\mathbf{U}}_k^\top P_{\mathbf{U}_{k\perp}}. \end{aligned}$$

So by triangle inequality, we have

$$\begin{aligned}
 \left\| [\mathcal{Z}; \widehat{\mathbf{U}}_1^\top, \dots, \widehat{\mathbf{U}}_d^\top] \right\|_{\text{HS}} &\leq \sum_{\Omega \subseteq [d]} \left\| \mathcal{Z} \times_{k \in \Omega} \widehat{\mathbf{U}}_k^\top P_{\mathbf{U}_k} \times_{k \in \Omega^c} \widehat{\mathbf{U}}_k^\top P_{\mathbf{U}_{k\perp}} \right\|_{\text{HS}} \\
 &\leq \sum_{\Omega \subseteq [d]} \left\| \mathcal{Z} \times_{k \in \Omega} \mathbf{U}_k^\top \times_{k \in \Omega^c} \mathbf{U}_{k\perp}^\top \right\|_{\text{HS}} \prod_{k \in \Omega} \|\widehat{\mathbf{U}}_k^\top \mathbf{U}_k\| \prod_{k \in \Omega^c} \|\widehat{\mathbf{U}}_k^\top \mathbf{U}_{k\perp}\| \\
 &\leq \sum_{\Omega \subseteq [d]} \left\| \mathcal{Z} \times_{k \in \Omega} \mathbf{U}_k^\top \times_{k \in \Omega^c} \mathbf{U}_{k\perp}^\top \right\|_{\text{HS}} \prod_{k \in \Omega^c} \|\sin \Theta(\widehat{\mathbf{U}}_k, \mathbf{U}_k)\| \\
 &= \sum_{\Omega \subseteq \{1, \dots, d\}} \theta_\Omega \prod_{k \in \Omega^c} \|\sin \Theta(\widehat{\mathbf{U}}_k, \mathbf{U}_k)\|.
 \end{aligned}$$

Here the second inequality is due to the fact that $\|\mathcal{Z} \times_i \mathbf{A}\mathbf{B}\|_{\text{HS}} = \|\mathbf{A}\mathbf{B}\mathcal{M}_i(\mathcal{Z})\|_{\text{HS}} \leq \|\mathbf{A}\| \|\mathbf{B}\mathcal{M}_i(\mathcal{Z})\|_{\text{HS}} = \|\mathbf{A}\| \|\mathcal{Z} \times_i \mathbf{B}\|_{\text{HS}}$ and we apply iteratively for each mode with $\mathbf{A} = \widehat{\mathbf{U}}_k^\top \mathbf{U}_k$ (or $\widehat{\mathbf{U}}_k^\top \mathbf{U}_{k\perp}$) and $\mathbf{B} = \mathbf{U}_k^\top$ (or $\mathbf{U}_{k\perp}^\top$). The third inequality is due to that $\|\widehat{\mathbf{U}}_k^\top \mathbf{U}_k\| \leq 1$ and $\|\widehat{\mathbf{U}}_k^\top \mathbf{U}_{k\perp}\| = \|\sin \Theta(\widehat{\mathbf{U}}_k, \mathbf{U}_k)\|$. ■

B.3 Proof of Theorem 2

The proof is done by construction. Let's denote $\mathcal{I}_r \in (\mathbb{R}^r)^{\otimes d}$ as the order- d identity tensor with entries (i, i, \dots, i) to be 1 and others are 0. We construct

$$\mathcal{Z}_1 = \frac{\xi}{\sqrt{r}} \mathcal{I}_r \times_1 \begin{pmatrix} \mathbf{0}_{r \times r} \\ \mathbf{I}_r \\ \mathbf{0}_{(p_1-2r) \times r} \end{pmatrix} \times \dots \times_d \begin{pmatrix} \mathbf{0}_{r \times r} \\ \mathbf{I}_r \\ \mathbf{0}_{(p_d-2r) \times r} \end{pmatrix},$$

where $\mathbf{0}_{m \times n}$ denotes a $m \times n$ matrix with all entries to be 0.

It is easy to check that $\sup_{\|\mathcal{Y}\|_{\text{HS}} \leq 1, \text{rank}(\mathcal{Y}) \leq (r, \dots, r)} \langle \mathcal{Z}_1, \mathcal{Y} \rangle \leq \|\mathcal{Z}_1\|_{\text{HS}} = \xi$. Similarly we construct

$$\mathcal{T}_1 = \frac{\xi}{\sqrt{r}} \mathcal{I}_r \times_1 \begin{pmatrix} \mathbf{I}_r \\ \mathbf{0}_{r \times r} \\ \mathbf{0}_{(p_1-2r) \times r} \end{pmatrix} \times \dots \times_d \begin{pmatrix} \mathbf{I}_r \\ \mathbf{0}_{r \times r} \\ \mathbf{0}_{(p_d-2r) \times r} \end{pmatrix}.$$

Also we let $\mathcal{Z}_2 = \mathcal{T}_1$ and $\mathcal{T}_2 = \mathcal{Z}_1$, and it is easy to check $(\mathcal{T}_1, \mathcal{Z}_1), (\mathcal{T}_2, \mathcal{Z}_2) \in \mathcal{F}_r(\xi)$. At the same time, we have $\mathcal{Z}_1 + \mathcal{T}_1 = \mathcal{T}_2 + \mathcal{Z}_2$. Thus

$$\begin{aligned}
 \inf_{\widehat{\mathcal{T}}} \sup_{(\mathcal{T}, \mathcal{Z}) \in \mathcal{F}_r(\xi)} \|\widehat{\mathcal{T}} - \mathcal{T}\|_{\text{HS}} &\geq \inf_{\widehat{\mathcal{T}}} \max \left\{ \|\widehat{\mathcal{T}} - \mathcal{T}_1\|_{\text{HS}}, \|\widehat{\mathcal{T}} - \mathcal{T}_2\|_{\text{HS}} \right\} \\
 &\geq \inf_{\widehat{\mathcal{T}}} \frac{1}{2} \left(\|\widehat{\mathcal{T}} - \mathcal{T}_1\|_{\text{HS}} + \|\widehat{\mathcal{T}} - \mathcal{T}_2\|_{\text{HS}} \right) \\
 &\geq \frac{1}{2} \|\mathcal{T}_1 - \mathcal{T}_2\|_{\text{HS}} = \frac{\sqrt{2}}{2} \xi.
 \end{aligned}$$

■

B.4 Proof of Theorem 3

The proof is long and nontrivial. The main idea of the proof is to develop the recursive error bound of $\tilde{\mathbf{U}}_k^{(t+1)}$ i.e., the estimate of \mathbf{U}_k at iteration $t+1$, based on the error bound of $\tilde{\mathbf{U}}_k^{(t)}$, i.e., the estimate at iteration t . The outline of the proof is the following: after a briefly introduction of notations, the main proof could be divided into three steps.

- Step 1: Recall in HOOI procedure, the update for the mode- k singular subspace satisfies

$$\begin{aligned} \tilde{\mathbf{U}}_k^{(t+1)} = & \text{SVD}_{r_k} \left(\mathcal{M}_{\bar{k}} \left(\mathcal{T} \times_{i \in \underline{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} \times_{\check{\Omega}_k} \tilde{\mathbf{U}}_k^{(t)\top} \times_{i \in \bar{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t)\top} \right) \right. \\ & \left. + \mathcal{M}_{\bar{k}} \left(\mathcal{Z} \times_{i \in \underline{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} \times_{\check{\Omega}_k} \tilde{\mathbf{U}}_k^{(t)\top} \times_{i \in \bar{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t)\top} \right) \right). \end{aligned}$$

To give an upper bound for $e_{t+1,k}$, we aim to give an upper bound for

$$\left\| \left(\mathcal{M}_{\bar{k}} \left(\mathcal{Z} \times_{i \in \underline{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} \times_{\check{\Omega}_k} \tilde{\mathbf{U}}_k^{(t)\top} \times_{i \in \bar{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t)\top} \right) \right)_{\max(r_k)} \right\|_q$$

by using $\tau_1, \dots, \tau_m, e_t, e_{t+1}$ in this step..

- Step 2: After getting an upper bound for (24), in this step we use induction to prove the following claim,

$$\begin{aligned} e_t & \leq 2^{(d+3)/2} \tau_1 / \lambda + e_0 / 2^t; t = 0, 1, \dots \\ \tilde{e}_t & \leq 2^{(d+3)/2} \tau_1 / \lambda + \tilde{e}_0 / 2^t; t = 0, 1, \dots \end{aligned}$$

- Step 3: Derive the error bound for $\|\hat{\mathcal{T}} - \mathcal{T}\|_{\text{HS}}$ by the unified quantity ξ .

For convenience, in this proof we denote

$$\mathbf{T}_k = \mathcal{M}_{\bar{k}}(\mathcal{T}), \quad \tilde{\mathbf{T}}_k = \mathcal{M}_{\bar{k}}(\tilde{\mathcal{T}}), \quad \mathbf{Z}_k = \mathcal{M}_{\bar{k}}(\mathcal{Z}), \quad k = 1, \dots, m.$$

Suppose

$$\begin{aligned} e_t & = \max_k e_{t,k}, \quad e_{t,k} = \left\| (\tilde{\mathbf{U}}_{k\perp}^{(t)})^\top \mathbf{U}_k \right\|_q, \quad k = 1, \dots, m; t = 0, 1, \dots \\ \tilde{e}_t & = \max_k \tilde{e}_{t,k}, \quad \tilde{e}_{t,k} = \left\| (\tilde{\tilde{\mathbf{U}}}_{k\perp}^{(t)})^\top \mathbf{U}_k \right\|, \quad k = 1, \dots, m; t = 0, 1, \dots \end{aligned} \tag{35}$$

Step 1. Recall the procedure of HOOI that

$$\begin{aligned} \tilde{\mathbf{U}}_k^{(t+1)} = & \text{SVD}_{r_k} \left(\mathcal{M}_{\bar{k}} \left(\tilde{\mathcal{T}} \times_{i \in \underline{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} \times_{i \in \check{\Omega}_k} \tilde{\mathbf{U}}_k^{(t)\top} \times_{i \in \bar{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t)\top} \right) \right) \\ = & \text{SVD}_{r_k} \left(\mathcal{M}_{\bar{k}} \left(\mathcal{T} \times_{i \in \underline{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} \times_{\check{\Omega}_k} \tilde{\mathbf{U}}_k^{(t)\top} \times_{i \in \bar{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t)\top} \right) \right. \\ & \left. + \mathcal{M}_{\bar{k}} \left(\mathcal{Z} \times_{i \in \underline{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} \times_{\check{\Omega}_k} \tilde{\mathbf{U}}_k^{(t)\top} \times_{i \in \bar{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t)\top} \right) \right). \end{aligned} \tag{36}$$

Notice that $\text{rank} \left(\mathcal{M}_{\bar{k}} \left(\mathcal{T} \times_{i \in \Omega_k} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} \times_{\check{\Omega}_k} \tilde{\mathbf{U}}_k^{(t)\top} \times_{i \in \bar{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t)\top} \right) \right) \leq r_k$, to apply Theorem 5 in Luo et al. (2020), the key is to give an upper bound for

$$\left\| \left(\mathcal{M}_{\bar{k}} \left(\mathcal{Z} \times_{i \in \Omega_k} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} \times_{\check{\Omega}_k} \tilde{\mathbf{U}}_k^{(t)\top} \times_{i \in \bar{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t)\top} \right) \right)_{\max(r_k)} \right\|_q.$$

To simplify the notation, for $S_j \in \mathcal{S}_j^{(-\bar{k})}$, we let $S_{j1} = S_j \cap \Omega_k, S_{j2} = S_j \cap (\check{\Omega}_k \cup \bar{\Omega}_k)$ and $S_{j1}^c = S_j^c \cap \Omega_k, S_{j2}^c = S_j^c \cap (\check{\Omega}_k \cup \bar{\Omega}_k)$. First we can introduce $\mathbf{I} = (P_{\mathbf{U}_{i'}} + P_{\mathbf{U}_{i'\perp}})$ in the expression,

$$\begin{aligned} & \left\| \left(\mathcal{M}_{\bar{k}} \left(\mathcal{Z} \times_{i \in \Omega_k} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} \times_{\check{\Omega}_k} \tilde{\mathbf{U}}_k^{(t)\top} \times_{i \in \bar{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t)\top} \right) \right)_{\max(r_k)} \right\|_q \\ &= \left\| \left(\mathcal{M}_{\bar{k}} \left(\mathcal{Z} \times_{i \in \Omega_k} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} (P_{\mathbf{U}_{i'}} + P_{\mathbf{U}_{i'\perp}}) \times_{\check{\Omega}_k} \tilde{\mathbf{U}}_k^{(t)\top} (P_{\mathbf{U}_k} + P_{\mathbf{U}_{k\perp}}) \right. \right. \right. \\ & \quad \left. \left. \left. \times_{i \in \bar{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t)\top} (P_{\mathbf{U}_{i'}} + P_{\mathbf{U}_{i'\perp}}) \right) \right)_{\max(r_k)} \right\|_q. \end{aligned} \quad (37)$$

Then

$$\begin{aligned} (37) & \stackrel{(a)}{\leq} \left\| \left(\mathcal{M}_{\bar{k}} \left(\mathcal{Z} \times_{i \in \Omega_k} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} P_{\mathbf{U}_{i'}} \times_{\check{\Omega}_k} \tilde{\mathbf{U}}_k^{(t)\top} P_{\mathbf{U}_k} \times_{i \in \bar{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t)\top} P_{\mathbf{U}_{i'}} \right) \right)_{\max(r_k)} \right\|_q \\ & + \sum_{S_1 \in \mathcal{S}_1^{(-\bar{k})}} \left\| \left(\mathcal{M}_{\bar{k}} \left(\mathcal{Z} \times_{i \in S_{11}} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} P_{\mathbf{U}_{i'\perp}} \times_{i \in S_{11}^c} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} P_{\mathbf{U}_{i'}} \right. \right. \right. \\ & \quad \left. \left. \left. \times_{i \in S_{12}} \tilde{\mathbf{U}}_{i'}^{(t)\top} P_{\mathbf{U}_{i'\perp}} \times_{i \in S_{12}^c} \tilde{\mathbf{U}}_{i'}^{(t)\top} P_{\mathbf{U}_{i'}} \right) \right)_{\max(r_k)} \right\|_q + \dots \\ & + \sum_{S_j \in \mathcal{S}_j^{(-\bar{k})}} \left\| \left(\mathcal{M}_{\bar{k}} \left(\mathcal{Z} \times_{i \in S_{j1}} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} P_{\mathbf{U}_{i'\perp}} \times_{i \in S_{j1}^c} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} P_{\mathbf{U}_{i'}} \right. \right. \right. \\ & \quad \left. \left. \left. \times_{i \in S_{j2}} \tilde{\mathbf{U}}_{i'}^{(t)\top} P_{\mathbf{U}_{i'\perp}} \times_{i \in S_{j2}^c} \tilde{\mathbf{U}}_{i'}^{(t)\top} P_{\mathbf{U}_{i'}} \right) \right)_{\max(r_k)} \right\|_q + \dots \quad (38) \\ & + \left\| \left(\mathcal{M}_{\bar{k}} \left(\mathcal{Z} \times_{i \in \Omega_k} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} P_{\mathbf{U}_{i'\perp}} \times_{\check{\Omega}_k} \tilde{\mathbf{U}}_k^{(t)\top} P_{\mathbf{U}_{k\perp}} \times_{i \in \bar{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t)\top} P_{\mathbf{U}_{i'\perp}} \right) \right)_{\max(r_k)} \right\|_q \\ &= \left\| \left(\mathcal{M}_{\bar{k}} \left(\mathcal{Z} \times_{i \in \Omega_k} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} P_{\mathbf{U}_{i'}} \times_{\check{\Omega}_k} \tilde{\mathbf{U}}_k^{(t)\top} P_{\mathbf{U}_k} \times_{i \in \bar{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t)\top} P_{\mathbf{U}_{i'}} \right) \right)_{\max(r_k)} \right\|_q \\ & + \sum_{j=1}^{d-1} \sum_{S_j \in \mathcal{S}_j^{(-\bar{k})}} \left\| \left(\mathcal{M}_{\bar{k}} \left(\mathcal{Z} \times_{i \in S_{j1}} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} P_{\mathbf{U}_{i'\perp}} \times_{i \in S_{j1}^c} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} P_{\mathbf{U}_{i'}} \right. \right. \right. \\ & \quad \left. \left. \left. \times_{i \in S_{j2}} \tilde{\mathbf{U}}_{i'}^{(t)\top} P_{\mathbf{U}_{i'\perp}} \times_{i \in S_{j2}^c} \tilde{\mathbf{U}}_{i'}^{(t)\top} P_{\mathbf{U}_{i'}} \right) \right)_{\max(r_k)} \right\|_q. \end{aligned}$$

Here (a) is due to the triangle inequality for truncated Schatten- q norm given in Lemma 4 of Luo et al. (2020). The right hand side of (38) can be divided into the sum of d groups and the value of j^{th} group is denoted as G_j where

$$G_0 := \left\| \left(\mathcal{M}_{\bar{k}} \left(\mathcal{Z} \times_{i \in \Omega_k} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} P_{\mathbf{U}_{i'}} \times_{\check{\Omega}_k} \tilde{\mathbf{U}}_k^{(t)\top} P_{\mathbf{U}_k} \times_{i \in \bar{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t)\top} P_{\mathbf{U}_{i'}} \right) \right)_{\max(r_k)} \right\|_q$$

and for $1 \leq j \leq d-1$, define G_j to be,

$$G_j = \sum_{S_j \in \mathcal{S}_j^{(-\bar{k})}} \left\| \left(\mathcal{M}_{\bar{k}} \left(\mathbf{Z} \times_{i \in S_{j1}} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} P_{\mathbf{U}_{i'\perp}} \times_{i \in S_{j1}^c} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} P_{\mathbf{U}_{i'}} \right. \right. \right. \\ \left. \left. \left. \times_{i \in S_{j2}} \tilde{\mathbf{U}}_{i'}^{(t)\top} P_{\mathbf{U}_{i'\perp}} \times_{i \in S_{j2}^c} \tilde{\mathbf{U}}_{i'}^{(t)\top} P_{\mathbf{U}_{i'}} \right) \right)_{\max(r_k)} \right\|_q.$$

Next we are going to upper bound G_j ($0 \leq j \leq d-1$).

- Upper Bound of G_0 .

$$\begin{aligned} G_0 &= \left\| \left(\mathcal{M}_{\bar{k}} \left(\mathbf{Z} \times_{i \in \underline{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} P_{\mathbf{U}_{i'}} \times_{\check{\Omega}_k} \tilde{\mathbf{U}}_k^{(t)\top} P_{\mathbf{U}_k} \times_{i \in \bar{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t)\top} P_{\mathbf{U}_{i'}} \right) \right)_{\max(r_k)} \right\|_q \\ &= \left\| \left(\mathcal{M}_{\bar{k}} \left(\left(\mathbf{Z} \times_{i \neq \bar{k}} \mathbf{U}_{i'}^\top \right) \times_{i \in \underline{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} \mathbf{U}_{i'} \times_{\check{\Omega}_k} \tilde{\mathbf{U}}_k^{(t)\top} \mathbf{U}_k \right. \right. \\ &\quad \left. \left. \times_{i \in \bar{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t)\top} \mathbf{U}_{i'} \right) \right)_{\max(r_k)} \right\|_q \\ &\leq \left\| \left(\mathcal{M}_{\bar{k}} \left(\mathbf{Z} \times_{i \neq \bar{k}} \mathbf{U}_{i'}^\top \right) \right)_{\max(r_k)} \right\|_q = \tau_{1k}. \end{aligned}$$

Here the inequality is due to the fact $\|\tilde{\mathbf{U}}_{i'}^{(t)\top} \mathbf{U}_{i'}\| \leq 1$ for any t and the following fact: for any matrix \mathbf{A} s.t. $\|\mathbf{A}\| \leq 1$,

$$\sigma_k(\mathcal{M}_i(\mathbf{Z} \times_j \mathbf{A})) = \sigma_k(\mathcal{M}_i(\mathbf{Z}) \cdot \mathbf{A}^\top) \leq \sigma_k(\mathcal{M}_i(\mathbf{Z})) \|\mathbf{A}\| \leq \sigma_k(\mathcal{M}_i(\mathbf{Z})), \quad (39)$$

where the first inequality is due to Lemma 7 of Luo et al. (2020).

- Upper Bound of G_j ($1 \leq j \leq d-1$).

$$\begin{aligned} G_j &\leq \binom{d-1}{j} \times \max_{S_j \in \mathcal{S}_j^{(-\bar{k})}} \\ &\left\| \left(\mathcal{M}_{\bar{k}} \left(\mathbf{Z} \times_{i \in S_{j1}} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} P_{\mathbf{U}_{i'\perp}} \times_{i \in S_{j1}^c} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} P_{\mathbf{U}_{i'}} \times_{i \in S_{j2}} \tilde{\mathbf{U}}_{i'}^{(t)\top} P_{\mathbf{U}_{i'\perp}} \right. \right. \\ &\quad \left. \left. \times_{i \in S_{j2}^c} \tilde{\mathbf{U}}_{i'}^{(t)\top} P_{\mathbf{U}_{i'}} \right) \right)_{\max(r_k)} \right\|_q \\ &\stackrel{(a)}{\leq} \binom{d-1}{j} \max_{S_j \in \mathcal{S}_j^{(-\bar{k})}} \left(\prod_{i \in S_{j1}} \|\tilde{\mathbf{U}}_{i'}^{(t+1)\top} \mathbf{U}_{i'\perp}\|_q \prod_{i \in S_{j2}} \|\tilde{\mathbf{U}}_{i'}^{(t)\top} \mathbf{U}_{i'\perp}\|_q \right) \times \\ &\left\| \left(\mathcal{M}_{\bar{k}} \left(\mathbf{Z} \times_{i \in S_{j1}} \frac{\tilde{\mathbf{U}}_{i'}^{(t+1)\top} P_{\mathbf{U}_{i'\perp}}}{\|\tilde{\mathbf{U}}_{i'}^{(t+1)\top} P_{\mathbf{U}_{i'\perp}}\|_q} \times_{i \in S_{j2}} \frac{\tilde{\mathbf{U}}_{i'}^{(t)\top} P_{\mathbf{U}_{i'\perp}}}{\|\tilde{\mathbf{U}}_{i'}^{(t)\top} P_{\mathbf{U}_{i'\perp}}\|_q} \times_{i \in S_j^c} \mathbf{U}_{i'}^\top \right) \right)_{\max(r_k)} \right\|_q, \end{aligned} \quad (40)$$

where (a) is due to the fact that $\|\tilde{\mathbf{U}}_i^{(t)\top} \mathbf{U}_{i'}\| \leq 1$ for any t . By simplifying (40), we get

$$\begin{aligned}
 G_j &\leq \binom{d-1}{j} \max_{S_j \in \mathcal{S}_j^{(-\bar{k})}} \left(\prod_{i \in S_{j1}} \|\tilde{\mathbf{U}}_{i'}^{(t+1)\top} \mathbf{U}_{i'\perp}\|_q \prod_{i \in S_{j2}} \|\tilde{\mathbf{U}}_{i'}^{(t)\top} \mathbf{U}_{i'\perp}\|_q \right) \\
 &\quad \times \left\| \left(\mathcal{M}_{\bar{k}} \left(\mathbf{Z} \times_{i \in S_{j1}} \frac{\tilde{\mathbf{U}}_{i'}^{(t+1)\top} \mathbf{U}_{i'\perp}}{\|\tilde{\mathbf{U}}_{i'}^{(t+1)\top} \mathbf{U}_{i'\perp}\|_q} \mathbf{U}_{i'\perp}^\top \right. \right. \right. \\
 &\quad \left. \left. \left. \times_{i \in S_{j2}} \frac{\tilde{\mathbf{U}}_{i'}^{(t)\top} \mathbf{U}_{i'\perp}}{\|\tilde{\mathbf{U}}_{i'}^{(t)\top} \mathbf{U}_{i'\perp}\|_q} \mathbf{U}_{i'\perp}^\top \times_{i \in S_j^c} \mathbf{U}_{i'}^\top \right) \right)_{\max(r_k)} \right\|_q \quad (41) \\
 &\stackrel{(b)}{\leq} \binom{d-1}{j} \max_{S_j \in \mathcal{S}_j^{(-\bar{k})}} \left(\prod_{i \in S_{j1}} \|\tilde{\mathbf{U}}_{i'}^{(t+1)\top} \mathbf{U}_{i'\perp}\|_q \prod_{i \in S_{j2}} \|\tilde{\mathbf{U}}_{i'}^{(t)\top} \mathbf{U}_{i'\perp}\|_q \right) \times \tau_{j+1} \\
 &\leq \binom{d-1}{j} \max_{S_j \in \mathcal{S}_j^{(-\bar{k})}} (e_t)^{|S_{j2}|} (e_{t+1})^{|S_{j1}|} \tau_{j+1}
 \end{aligned}$$

where (b) is due to the fact that $\left\| \frac{\tilde{\mathbf{U}}_{i'}^{(t+1)\top} \mathbf{U}_{i'\perp}}{\|\tilde{\mathbf{U}}_{i'}^{(t+1)\top} \mathbf{U}_{i'\perp}\|_q} \right\| \leq 1$ and the definition of τ_{j+1} .

So in summary, plug (41) into (38), now we have the following bound

$$\begin{aligned}
 &\left\| \left(\mathcal{M}_{\bar{k}} \left(\mathbf{Z} \times_{i \in \Omega_k} \tilde{\mathbf{U}}_{i'}^{(t+1)\top} \times_{\check{\Omega}_k} \tilde{\mathbf{U}}_k^{(t)\top} \times_{i \in \bar{\Omega}_k} \tilde{\mathbf{U}}_{i'}^{(t)\top} \right) \right)_{\max(r_k)} \right\|_q \\
 &\leq \tau_{1k} + \sum_{j=1}^{d-1} \binom{d-1}{j} \max_{S_j \in \mathcal{S}_j^{(-\bar{k})}} (e_t)^{|S_{j2}|} (e_{t+1})^{|S_{j1}|} \tau_{j+1}. \quad (42)
 \end{aligned}$$

Step 2. In this step we want to use induction to prove the following claims,

$$e_t \leq 2^{\frac{d+3}{2}} \frac{\tau_1}{\lambda} + e_0/2^t \quad \text{and} \quad \tilde{e}_t \leq 2^{\frac{d+3}{2}} \frac{\tau_1}{\lambda} + \tilde{e}_0/2^t; \quad t = 0, 1, \dots \quad (43)$$

Since the proof of two statements in (43) are similar, we mainly focus on the proof of the first statement. Claim (43) clearly holds if $t = 0$. Assume the first claim of (43) holds for t and next we show it also holds for $t + 1$.

Let's first show the upper bound of e_t can be used to upper bound $e_{t+1,1}$. Notice

$$\begin{aligned}
 2 \left\| \left(\mathcal{M}_{\bar{1}} \left(\mathbf{Z} \times_{i \neq \bar{1}} \tilde{\mathbf{U}}_{i'}^{(t)\top} \right) \right)_{\max(r_1)} \right\|_q &\stackrel{(a)}{\geq} \left\| \tilde{\mathbf{U}}_{1\perp}^{(t+1)\top} \mathbf{U}_1 \right\|_q \sigma_{r_1} \left(\mathcal{M}_{\bar{1}} \left(\mathcal{T} \times_{i \neq \bar{1}} \tilde{\mathbf{U}}_{i'}^{(t)\top} \right) \right) \\
 &\stackrel{(b)}{\geq} (e_{t+1,1}) \sigma_{r_1} \left(\mathbf{T}_1 \left(\otimes_{i \neq \bar{1}} \mathbf{U}_{i'} \right) \cdot \left(\otimes_{i \neq \bar{1}} \mathbf{U}_{i'}^\top \tilde{\mathbf{U}}_{i'}^{(t)} \right) \right) \\
 &\stackrel{(c)}{\geq} (e_{t+1,1}) \lambda \left(\prod_{i \neq \bar{1}} \sigma_{\min} \left(\mathbf{U}_{i'}^\top \tilde{\mathbf{U}}_{i'}^{(t)} \right) \right) \\
 &\stackrel{(d)}{\geq} (e_{t+1,1}) \lambda (1 - \tilde{e}_t^2)^{\frac{d-1}{2}}. \quad (44)
 \end{aligned}$$

Here (a) is due to Theorem 5 in Luo et al. (2020) and the fact the left singular space of $\mathcal{M}_{\bar{1}} \left(\mathcal{T} \times_{i \neq \bar{1}} \tilde{\mathbf{U}}_{i'}^{(t)\top} \right)$ is \mathbf{U}_1 and its rank is less than r_1 , (b) is due to the fact that the right singular space of \mathbf{T}_1 is $\otimes_{i \neq \bar{1}} \mathbf{U}_{i'}$, equation (8), (c) is by Lemma 7 of Luo et al. (2020) and (d) is due to the fact $\sigma_{\min} \left(\mathbf{U}_{i'}^\top \tilde{\mathbf{U}}_i^{(t)} \right) = \sqrt{1 - \|\mathbf{U}_{i'}^\top \tilde{\mathbf{U}}_i^{(t)}\|^2} \geq \sqrt{1 - \tilde{e}_t^2}$ by Lemma 1 of Cai and Zhang (2018).

Plugging the upper bound of $\left\| \left(\mathcal{M}_{\bar{1}} \left(\mathcal{Z} \times_{i \neq \bar{1}} \tilde{\mathbf{U}}_{i'}^{(t)\top} \right) \right)_{\max(r_1)} \right\|_q$ in (42) into (44), we get

$$e_{t+1,1} \leq 2 \frac{\tau_{11} + \sum_{j=1}^{d-1} \binom{d-1}{j} (e_t)^j \tau_{j+1}}{\lambda(1 - \tilde{e}_t^2)^{\frac{d-1}{2}}}. \quad (45)$$

Next we show under condition (19), $e_{t+1,1} \leq 2^{\frac{d+3}{2}} \frac{\tau_1}{\lambda} + e_0/2^t$, i.e., the upper bound of e_t can be used to upper bound $e_{t+1,1}$. First, under (19), we have

$$\lambda \geq 2^{\frac{d+6}{2}} \tau_1 \vee \left(\sum_{j=1}^{d-1} 2^{\frac{d+4-j}{2}} \binom{d-1}{j} \tau_{j+1} \right), \quad (46)$$

due to the fact that $\xi \geq \tau_j$ for $j = 1, \dots, d$. And under the assumption of λ , we have $\tilde{e}_t \leq \sqrt{2}/2$. So by (45), we have

$$\begin{aligned} \left(\frac{1}{2} \right)^{\frac{d-1}{2}} e_{t+1,1} &\leq 2 \frac{\tau_{11} + \sum_{j=1}^{d-1} \binom{d-1}{j} (e_t)^j \tau_{j+1}}{\lambda} \\ &\leq 2 \left(\frac{\tau_{11}}{\lambda} + \frac{\sum_{j=1}^{d-1} \binom{d-1}{j} (\frac{\sqrt{2}}{2})^{j-1} \tau_{j+1}}{\lambda} e_t \right) \\ &\stackrel{(46)}{\leq} 2 \left(\frac{\tau_{11}}{\lambda} + 2^{-\frac{d+3}{2}} e_t \right) \end{aligned} \quad (47)$$

Since (43) holds for t and plug in the upper bound of e_t in (47), multiply $2^{\frac{d-1}{2}}$ at both side of (47), we get

$$e_{t+1,1} \leq 2^{\frac{d+3}{2}} \frac{\tau_1}{\lambda} + \frac{e_0}{2^{t+1}} \leq 2^{\frac{d+3}{2}} \frac{\tau_1}{\lambda} + \frac{e_0}{2^t}.$$

So the upper bound of e_t can be used to bound $e_{t+1,1}$ and by doing similar analysis for all modes, we conclude that the upper bound for e_t also holds for e_{t+1} , i.e., we have

$$e_{t+1} \leq 2^{\frac{d+3}{2}} \frac{\tau_1}{\lambda} + \frac{e_0}{2^t}. \quad (48)$$

Now we can show the first statement in (43) also holds for $t+1$ given it holds for t . With (48), when we do the similar analysis of (44) for other modes, we can use the same upper bound of e_t to bound e_{t+1} . So repeat (44), for any $1 \leq k \leq m$, we have

$$e_{t+1,k} (1 - \tilde{e}_t^2)^{\frac{d-1}{2}} \leq 2 \frac{\tau_{1k} + \sum_{j=1}^{d-1} \binom{d-1}{j} (e_t)^j \tau_{j+1}}{\lambda}. \quad (49)$$

Since it is true for every k , we obtain the following recursive inequality for e_{t+1} ,

$$e_{t+1} \left(\frac{1}{2}\right)^{\frac{d-1}{2}} \leq 2 \frac{\tau_1 + \sum_{j=1}^{d-1} \binom{d-1}{j} (e_t)^j \tau_{j+1}}{\lambda}. \quad (50)$$

By applying the same argument as in (47), we can show that

$$e_{t+1} \leq 2^{\frac{d+3}{2}} \frac{\tau_1}{\lambda} + \frac{e_0}{2^{t+1}},$$

given (43) holds for e_t . Similarly we can show $\tilde{e}_{t+1} \leq 2^{\frac{d+3}{2}} \frac{\tau_1}{\lambda} + \frac{\tilde{e}_0}{2^{t+1}}$

Thus, (43) holds for all t . Given $t_{\max} \geq \log(e_0 \lambda / \tau_1) \vee 1$, we get the upper bounds for e_t and \tilde{e}_t :

$$e_t \leq \left(2^{\frac{d+3}{2}} + 1\right) \frac{\tau_1}{\lambda}, \quad \tilde{e}_t \leq \left(2^{\frac{d+3}{2}} + 1\right) \frac{\tau_1}{\lambda} \quad (51)$$

Plug (50) and (46) into the upper bound (49), we get

$$e_{t+1,k} \leq 2 \left(1 - \frac{\left(2^{\frac{d+3}{2}} + 1\right)^2 \tau_1^2}{\lambda^2}\right)^{-\frac{d-1}{2}} \left(\frac{\tau_{1k}}{\lambda} + \sum_{j=1}^{d-1} \frac{\binom{d-1}{j} \left(2^{\frac{d+3}{2}} + 1\right)^j \tau_1^j \tau_{j+1}}{\lambda^{j+1}}\right),$$

for $k = 1, \dots, m$.

Finally the perturbation of signal subspaces follows by observing that $e_{t,k} := \left\| \left(\tilde{\mathbf{U}}_{k\perp}^{(t)}\right)^\top \mathbf{U}_k \right\|_q = \left\| \sin \Theta \left(\tilde{\mathbf{U}}_k^{(t)}, \mathbf{U}_k\right) \right\|_q$ due to Lemma 6 of Luo et al. (2020).

Step 3. In this step, we are going to give an upper bound for the tensor reconstruction error for $\left\| \tilde{\mathcal{T}} \times_{\Omega_1} P_{\hat{\mathbf{U}}_1} \times \dots \times_{\Omega_m} P_{\hat{\mathbf{U}}_m} - \mathcal{T} \right\|_{\text{HS}}$.

First, notice the following decomposition

$$\begin{aligned} \mathcal{T} &= \mathcal{T} \times_1 \left(P_{\hat{\mathbf{U}}_{1'}} + P_{\hat{\mathbf{U}}_{1'\perp}}\right) \times_2 \left(P_{\hat{\mathbf{U}}_{2'}} + P_{\hat{\mathbf{U}}_{2'\perp}}\right) \times \dots \times_d \left(P_{\hat{\mathbf{U}}_{d'}} + P_{\hat{\mathbf{U}}_{d'\perp}}\right) \\ &= \mathcal{T} \times_1 P_{\hat{\mathbf{U}}_{1'}} \times \dots \times_d P_{\hat{\mathbf{U}}_{d'}} + \mathcal{T} \times_1 P_{\hat{\mathbf{U}}_{1'\perp}} \times_2 P_{\hat{\mathbf{U}}_{2'}} \times \dots \times_d P_{\hat{\mathbf{U}}_{d'}} \\ &\quad + \mathcal{T} \times_1 \mathbf{I}_{p_{1'}} \times_2 P_{\hat{\mathbf{U}}_{2'\perp}} \times \dots \times_d P_{\hat{\mathbf{U}}_{d'}} + \mathcal{T} \times_1 \mathbf{I}_{p_{1'}} \times_2 \mathbf{I}_{p_{2'}} \times \dots \times_d P_{\hat{\mathbf{U}}_{d'\perp}} \\ &\quad + \dots + \mathcal{T} \times_{i \leq d-1} \mathbf{I}_{p_{i'}} \times_d P_{\hat{\mathbf{U}}_{d'\perp}} \\ &= \mathcal{T} \times_{\Omega_1} P_{\hat{\mathbf{U}}_1} \times \dots \times_{\Omega_m} P_{\hat{\mathbf{U}}_m} + \sum_{k=1}^d \mathcal{T} \times_{i < k} \mathbf{I}_{p_{i'}} \times_k P_{\hat{\mathbf{U}}_{k'\perp}} \times_{i > k} P_{\hat{\mathbf{U}}_{i'}}. \end{aligned}$$

Thus,

$$\begin{aligned} &\left\| \tilde{\mathcal{T}} \times_{\Omega_1} P_{\hat{\mathbf{U}}_1} \times \dots \times_{\Omega_m} P_{\hat{\mathbf{U}}_m} - \mathcal{T} \right\|_{\text{HS}} \\ &\leq \left\| (\tilde{\mathcal{T}} - \mathcal{T}) \times_{\Omega_1} P_{\hat{\mathbf{U}}_1} \times \dots \times_{\Omega_m} P_{\hat{\mathbf{U}}_m} \right\|_{\text{HS}} + \sum_{k=1}^d \left\| \mathcal{T} \times_{i < k} \mathbf{I}_{p_{i'}} \times_k P_{\hat{\mathbf{U}}_{k'\perp}} \times_{i > k} P_{\hat{\mathbf{U}}_{i'}} \right\|_{\text{HS}} \quad (52) \\ &\leq \left\| \mathcal{Z} \times_{\Omega_1} P_{\hat{\mathbf{U}}_1} \times \dots \times_{\Omega_m} P_{\hat{\mathbf{U}}_m} \right\|_{\text{HS}} + \sum_{k=1}^d \left\| \hat{\mathbf{U}}_{k'\perp}^\top \mathcal{M}_k(\mathcal{T}) \right\|_F. \end{aligned}$$

Notice that for $k_1, k_2 \in \Omega_k$, $\mathcal{M}_{k_1}(\mathcal{T}) = \mathcal{M}_{k_2}(\mathcal{T})P_1$ where P_1 is a permutation matrix. So $\sum_{k=1}^d \left\| \widehat{\mathbf{U}}_{k'\perp}^\top \mathcal{M}_k(\mathcal{T}) \right\|_F = \sum_{k=1}^m |\Omega_k| \left\| \widehat{\mathbf{U}}_{k\perp}^\top \mathbf{T}_k \right\|_F$.

$$\begin{aligned}
 & \left\| \widehat{\mathbf{U}}_{k\perp}^\top \mathbf{T}_k \right\|_F \left(1 - \frac{\left(2^{\frac{d+3}{2}} + 2 \right)^2 \tau_1^2}{\lambda^2} \right)^{\frac{d-1}{2}} \\
 & \stackrel{(a)}{\leq} \left\| \widehat{\mathbf{U}}_{k\perp}^\top \mathbf{T}_k \right\|_F \sigma_{\min} \left(\left(\otimes_{i \in \Omega_k} \mathbf{U}_{i'}^\top \widetilde{\mathbf{U}}_i^{(t_{\max})} \otimes_{i \in \check{\Omega}_k \cup \bar{\Omega}_k} \mathbf{U}_{i'}^\top \widetilde{\mathbf{U}}_{i'}^{(t_{\max}-1)} \right) \right) \\
 & \stackrel{(b)}{=} \left\| \widehat{\mathbf{U}}_{k\perp}^\top \mathbf{T}_k \otimes_{i \neq \bar{k}} \mathbf{U}_{i'} \right\|_F \sigma_{\min} \left(\left(\otimes_{i \in \Omega_k} \mathbf{U}_{i'}^\top \widetilde{\mathbf{U}}_i^{(t_{\max})} \otimes_{i \in \check{\Omega}_k \cup \bar{\Omega}_k} \mathbf{U}_{i'}^\top \widetilde{\mathbf{U}}_{i'}^{(t_{\max}-1)} \right) \right) \\
 & \stackrel{(c)}{\leq} \left\| \widehat{\mathbf{U}}_{k\perp}^\top \mathbf{T}_k \left(\otimes_{i \neq \bar{k}} \mathbf{U}_{i'} \right) \left(\otimes_{i \neq \bar{k}} \mathbf{U}_{i'} \right)^\top \left(\otimes_{i \in \Omega_k} \widetilde{\mathbf{U}}_{i'}^{(t_{\max})} \otimes_{i \in \check{\Omega}_k \cup \bar{\Omega}_k} \widetilde{\mathbf{U}}_{i'}^{(t_{\max}-1)} \right) \right\|_F \\
 & \stackrel{(d)}{\leq} 2 \left\| \left(\mathcal{M}_{\bar{k}} \left(\mathbf{Z} \times_{i \in \Omega_k} \widetilde{\mathbf{U}}_{i'}^{(t_{\max})\top} \times_{i \in \check{\Omega}_k \cup \bar{\Omega}_k} \widetilde{\mathbf{U}}_{i'}^{(t_{\max}-1)\top} \right) \right)_{\max(r_k)} \right\|_F \stackrel{(e)}{\leq} 2\xi,
 \end{aligned} \tag{53}$$

where $\xi := \sup_{\|\mathbf{Y}\|_{\text{HS}} \leq 1, \text{rank}(\mathbf{Y}) \leq (r_1, \dots, r_{d'})} \langle \mathbf{Z}, \mathbf{Y} \rangle$. (a) is due to the fact $\sigma_{\min} \left(\mathbf{U}_{i'}^\top \widetilde{\mathbf{U}}_i^{(t)} \right) =$

$\sqrt{1 - \|\mathbf{U}_{i'\perp}^\top \widetilde{\mathbf{U}}_i^{(t)}\|^2} \geq \sqrt{1 - \tilde{e}_t^2}$ and $\max(\tilde{e}_{t_{\max}}, \tilde{e}_{t_{\max}-1}) \leq \frac{(2^{\frac{d+3}{2}} + 2)\tau_1}{\lambda}$; (b) is due to the fact that the right singular space of \mathbf{T}_k is $\otimes_{i \neq \bar{k}} \mathbf{U}_{i'}$; (c) is due to Lemma 7 of Luo et al. (2020), equation (8) and properties of Kronecker product, (d) is due to Theorem 2 in Luo et al. (2020) and

$$\text{rank} \left(\widehat{\mathbf{U}}_{k\perp}^\top \mathbf{T}_k \left(\otimes_{i \in \Omega_k} \widetilde{\mathbf{U}}_{i'}^{(t_{\max})} \otimes_{i \in \check{\Omega}_k \cup \bar{\Omega}_k} \widetilde{\mathbf{U}}_{i'}^{(t_{\max}-1)} \right) \right) \leq r_k;$$

and the last inequality (e) is due to the fact that

$$\begin{aligned}
 & \left\| \left(\mathcal{M}_{\bar{k}} \left(\mathbf{Z} \times_{i \in \Omega_k} \widetilde{\mathbf{U}}_{i'}^{(t_{\max})\top} \times_{i \in \check{\Omega}_k \cup \bar{\Omega}_k} \widetilde{\mathbf{U}}_{i'}^{(t_{\max}-1)\top} \right) \right)_{\max(r_k)} \right\|_F \\
 & = \sup_{\|\mathbf{X}\|_F \leq 1, \text{rank}(\mathbf{X}) \leq r_k} \left\langle \mathcal{M}_{\bar{k}} \left(\mathbf{Z} \times_{i \in \Omega_k} \widetilde{\mathbf{U}}_{i'}^{(t_{\max})\top} \times_{i \in \check{\Omega}_k \cup \bar{\Omega}_k} \widetilde{\mathbf{U}}_{i'}^{(t_{\max}-1)\top} \right), \mathbf{X} \right\rangle \leq \xi,
 \end{aligned}$$

for $k = 1, \dots, d$ and the equality is due to Lemma 2 of Luo et al. (2020).

Combined with the fact $\left\| \mathbf{Z} \times_{\Omega_1} P_{\widehat{\mathbf{U}}_1} \times \dots \times_{\Omega_m} P_{\widehat{\mathbf{U}}_m} \right\|_{\text{HS}} \leq \xi$ and (53) and plug them into (52), we have

$$\begin{aligned}
 & \left\| \widetilde{\mathcal{T}} \times_{\Omega_1} P_{\widehat{\mathbf{U}}_1} \times \dots \times_{\Omega_m} P_{\widehat{\mathbf{U}}_m} - \mathcal{T} \right\|_{\text{HS}} \\
 & \leq \left\| \mathbf{Z} \times_{\Omega_1} P_{\widehat{\mathbf{U}}_1} \times \dots \times_{\Omega_m} P_{\widehat{\mathbf{U}}_m} \right\|_{\text{HS}} + \sum_{k=1}^d \left\| \widehat{\mathbf{U}}_{k'\perp}^\top \mathcal{M}_k(\mathcal{T}) \right\|_F \\
 & = \left\| \mathbf{Z} \times_{\Omega_1} \widehat{\mathbf{U}}_1^\top \times \dots \times_{\Omega_m} \widehat{\mathbf{U}}_m^\top \right\|_{\text{HS}} + \sum_{k=1}^m |\Omega_k| \left\| \widehat{\mathbf{U}}_{k\perp}^\top \mathcal{M}_k(\mathcal{T}) \right\|_F \\
 & \leq \left(1 + 2d \left(1 - \frac{\left(2^{\frac{d+3}{2}} + 2 \right)^2 \tau_1^2}{\lambda^2} \right)^{-\frac{d-1}{2}} \right) \xi.
 \end{aligned} \tag{54}$$

Finally, notice that (21) follows exact the same proof as Step 3 except that in (53), we need to replace $\tilde{\mathbf{U}}_i^{(t_{\max})}, \tilde{\mathbf{U}}_i^{(t_{\max}-1)}$ with $\tilde{\mathbf{U}}_i^{(t)}, \tilde{\mathbf{U}}_i^{(t-1)}$ and use $\max(\tilde{e}_t, \tilde{e}_{t-1}) \leq 2^{\frac{d+3}{2}} \frac{\tau_1}{\lambda} + \frac{\tilde{e}_0}{2^{t-1}}$. Therefore, we have finished the proof of this theorem. \blacksquare

B.5 Proof of Theorem 5

In this setting, we consider applying Corollary 1. To apply Corollary 1, we only need to compute τ_{1i}, ξ .

Consider the case $q = \infty$ and let's first bound τ_{1k} . Notice

$$\mathcal{M}_k(\mathcal{Z} \times_{i \neq k} \mathbf{U}_i^\top) = \mathcal{M}_k(\mathcal{Z}) \otimes_{i \neq k} \mathbf{U}_i.$$

Each row of $\mathcal{M}_k(\mathcal{Z}) \otimes_{i \neq k} \mathbf{U}_i$ is independent multivariate Gaussian with covariance matrix $\sigma \mathbf{I}_{r-k}$ where $r-k = \prod_{i \neq k} r_i$. By random matrix theory (Vershynin, 2010), we have

$$\mathbb{P}(\|\mathcal{M}_k(\mathcal{Z} \times_{i \neq k} \mathbf{U}_i^\top)\|/\sigma \leq \sqrt{p_k} + \sqrt{r-k}) \geq 1 - c \exp(-(p_k + r-k)/2).$$

Since $r_{\max} \leq p_{\min}^{\frac{1}{d-1}}$, we have $\tau_{1k} \leq C\sigma\sqrt{p_k}$ with probability at least $1 - c \exp(-(p_k + r-k)/2)$.

Similarly by Lemma 5 of Zhang and Xia (2018), we have

$$\xi \leq C\sigma \sqrt{\sum_{i=1}^d p_i r_i}$$

w.p. at least $1 - \exp(-cp_{\min})$.

As we mentioned in Remark 7, $\tau_j \leq \xi$. So the results follows by plugging the bound of τ_j, ξ into Corollary 1 and noticing that

$$\frac{\sum_{j=1}^{d-1} \binom{d-1}{j} \left(\left(2^{\frac{d+3}{2}} + 1 \right) \tau_1 / \lambda \right)^j \tau_{j+1}}{\lambda} \leq C \frac{\tau_1 \max_j \tau_j}{\lambda^2},$$

under the assumption of the signal to ratio λ/σ . \blacksquare

B.6 Proof of Lemma 3

The proof of this Lemma is straight forward.

$$\begin{aligned} \mathcal{T} &= \mathcal{B} \times_1 \mathbf{\Pi}_1 \times \cdots \times_d \mathbf{\Pi}_d \\ &= \mathcal{B} \times_1 \mathbf{\Pi}_1 (\mathbf{\Pi}_1^\top \mathbf{\Pi}_1)^{-\frac{1}{2}} (\mathbf{\Pi}_1^\top \mathbf{\Pi}_1)^{\frac{1}{2}} \times \cdots \times_d \mathbf{\Pi}_d (\mathbf{\Pi}_d^\top \mathbf{\Pi}_d)^{-\frac{1}{2}} (\mathbf{\Pi}_d^\top \mathbf{\Pi}_d)^{\frac{1}{2}} \\ &= \left(\mathcal{B} \times_1 (\mathbf{\Pi}_1^\top \mathbf{\Pi}_1)^{\frac{1}{2}} \times \cdots \times_d (\mathbf{\Pi}_d^\top \mathbf{\Pi}_d)^{\frac{1}{2}} \right) \times_1 \mathbf{\Pi}_1 (\mathbf{\Pi}_1^\top \mathbf{\Pi}_1)^{-\frac{1}{2}} \times \cdots \times_d \mathbf{\Pi}_d (\mathbf{\Pi}_d^\top \mathbf{\Pi}_d)^{-\frac{1}{2}} \\ &= \mathcal{S} \times_1 \mathbf{\Pi}_1 (\mathbf{\Pi}_1^\top \mathbf{\Pi}_1)^{-\frac{1}{2}} \mathbf{V}_1 \times \cdots \times_d \mathbf{\Pi}_d (\mathbf{\Pi}_d^\top \mathbf{\Pi}_d)^{-\frac{1}{2}} \mathbf{V}_d, \end{aligned}$$

where the last inequality comes from the assumption about the decomposition of $\mathcal{B} \times_1 (\mathbf{\Pi}_1^\top \mathbf{\Pi}_1)^{\frac{1}{2}} \times \cdots \times_d (\mathbf{\Pi}_d^\top \mathbf{\Pi}_d)^{\frac{1}{2}}$. \blacksquare

B.7 Proof of Theorem 6

First by the same argument in the proof of Theorem 5, with probability at least $1 - \exp(-cp_{\min})$, $\tau_{1k} \leq C\sigma\sqrt{p_k}$ and $\xi \leq C\sigma\sqrt{\sum_{i=1}^d p_i r_i}$.

Under the assumption (A1), we have $\lambda(\mathcal{S}) := \min_i \sigma_{r_i}(\mathcal{M}_i(\mathcal{S})) \geq C\lambda\sqrt{\frac{\prod_{i=1}^d p_i}{\prod_{i=1}^d r_i}} \geq C\sqrt{\frac{p_{\max}^2 r_{\max}}{p_{\min}}}$ by the definition of \mathcal{S} .

Notice that under the signal strength in Theorem 6, the first order perturbation error in $\|\sin \Theta(\hat{\mathbf{U}}_i, \mathbf{U}_i)\|$ dominates as we discussed in Remark 4. So from Corollary 1 we have

$$\|\sin \Theta(\hat{\mathbf{U}}_i, \mathbf{U}_i)\| \leq C \frac{\sqrt{p_k}}{\lambda/\sigma} \sqrt{\frac{\prod_{i=1}^d r_i}{\prod_{i=1}^d p_i}},$$

and the bound for $\|\hat{\mathcal{T}} - \mathcal{T}\|_{\text{HS}}$.

For the cocluster membership recovery, first we have

$$\|\sin \Theta(\hat{\mathbf{U}}_i, \mathbf{U}_i)\|_F \leq \sqrt{r_i} \|\sin \Theta(\hat{\mathbf{U}}_i, \mathbf{U}_i)\| \leq C \frac{\sqrt{p_i r_i}}{\lambda/\sigma} \sqrt{\frac{\prod_{i=1}^d r_i}{\prod_{i=1}^d p_i}}. \quad (55)$$

The following proof is relative standard for proving misclassification error based on singular subspace estimation. Without loss of generality, we focus on the mode-1 clustering analysis. Take $\delta_k = \sqrt{\frac{1}{p_1^{(k)}} + \frac{1}{\max\{p_1^{(l)} : l \neq k\}}}$, and let $S_k = \{i \in G_1^{(k)} : \|(\hat{\mathbf{\Pi}}_1 \hat{\mathbf{X}}_1)_{[i,:]} - (\mathbf{U}_1)_{[i,:]}\| \geq \frac{\delta_k}{2}\}$ where \mathbf{U}_1 is defined in Lemma 3. Here we can imagine S_k as the set of nodes in cluster k of mode 1 that may not be correctly clustered. By Lemma 5.3 of Lei and Rinaldo (2015) and (55), we have

$$\sum_{k=1}^{r_1} |S_k| \delta_k^2 \leq C \frac{p_i r_i}{(\lambda/\sigma)^2} \frac{\prod_{i=1}^d r_i}{\prod_{i=1}^d p_i}.$$

Moreover, under the signal strength condition in Theorem 6, we have that there exists a permutation matrix such that after permutation, the nodes in $G_1 = \bigcup_{k=1}^{r_1} (G_1^{(k)} \setminus S_k)$ can be perfectly recovered based on Lemma 5.3 of Lei and Rinaldo (2015). So to analyze the misclassification error, we just need to consider about S_k .

$$\widetilde{err}(\hat{\mathbf{\Pi}}_1, \mathbf{\Pi}_1) \leq \max_{1 \leq k \leq r_1} \frac{|S_k|}{p_1^{(k)}} \leq \sum_{1 \leq k \leq r_1} \frac{|S_k|}{p_1^{(k)}} \stackrel{(a)}{\leq} \sum_{k=1}^{r_1} |S_k| \delta_k^2 \leq C \frac{p_1 r_1}{(\lambda/\sigma)^2} \frac{\prod_{i=1}^d r_i}{\prod_{i=1}^d p_i},$$

where (a) is because of the choice of δ_k and

$$err(\hat{\mathbf{\Pi}}_1, \mathbf{\Pi}_1) \leq \sum_{1 \leq k \leq r_1} \frac{|S_k|}{p_1} \leq C \frac{p_{1,\max}' r_1}{(\lambda/\sigma)^2} \frac{\prod_{i=1}^d r_i}{\prod_{i=1}^d p_i}.$$

The proof for other modes are similar and this finishes the proof of this theorem. \blacksquare

References

- Emmanuel Abbe, Jianqing Fan, Kaizheng Wang, Yiqiao Zhong, et al. Entrywise eigenvector analysis of random matrices with low expected rank. *Annals of Statistics*, 48(3):1452–1474, 2020.
- Kwangjun Ahn, Kangwook Lee, and Changho Suh. Hypergraph spectral clustering in the weighted stochastic block model. *IEEE Journal of Selected Topics in Signal Processing*, 12(5):959–974, 2018.
- Anima Anandkumar, Prateek Jain, Yang Shi, and Uma Naresh Niranjan. Tensor vs. matrix methods: Robust tensor decomposition under block sparse perturbations. In *Artificial Intelligence and Statistics*, pages 268–276, 2016.
- Animashree Anandkumar, Rong Ge, Daniel Hsu, Sham M Kakade, and Matus Telgarsky. Tensor decompositions for learning latent variable models. *Journal of Machine Learning Research*, 15:2773–2832, 2014a.
- Animashree Anandkumar, Rong Ge, and Majid Janzamin. Guaranteed non-orthogonal tensor decomposition via alternating rank-1 updates. *arXiv preprint arXiv:1402.5180*, 2014b.
- Arnab Auddy and Ming Yuan. Perturbation bounds for orthogonally decomposable tensors and their applications in high dimensional data analysis. *arXiv preprint arXiv:2007.09024*, 2020.
- Brett W Bader and Tamara G Kolda. Matlab tensor toolbox version 2.5. *Available online*, January, 7, 2012.
- Krishnakumar Balasubramanian, Jianqing Fan, and Zhuoran Yang. Tensor methods for additive index models under discordance and heterogeneity. *arXiv preprint arXiv:1807.06693*, 2018.
- Jonas Ballani and Lars Grasedyck. A projection method to solve linear systems in tensor format. *Numerical linear algebra with applications*, 20(1):27–43, 2013.
- Florent Benaych-Georges and Raj Rao Nadakuditi. The eigenvalues and eigenvectors of finite, low rank perturbations of large random matrices. *Advances in Mathematics*, 227(1):494–521, 2011.
- Florent Benaych-Georges and Raj Rao Nadakuditi. The singular values and vectors of low rank perturbations of large rectangular random matrices. *Journal of Multivariate Analysis*, 111:120–135, 2012.
- Rajendra Bhatia. *Matrix analysis*, volume 169. Springer Science & Business Media, 2013.
- Xuan Bi, Annie Qu, and Xiaotong Shen. Multilayer tensor factorization with applications to recommender systems. *The Annals of Statistics*, 46(6B):3308–3333, 2018.

- T Tony Cai and Anru Zhang. Rate-optimal perturbation bounds for singular subspaces with applications to high-dimensional statistics. *The Annals of Statistics*, 46(1):60–89, 2018.
- Tianxi Cai, T. Tony Cai, and Anru Zhang. Structured matrix completion with applications to genomic data integration. *Journal of the American Statistical Association*, 111(514):621–633, 2016.
- Joshua Cape, Minh Tang, and Carey E Priebe. Signal-plus-noise matrix models: eigenvector deviations and fluctuations. *Biometrika*, 106(1):243–250, 2019a.
- Joshua Cape, Minh Tang, and Carey E Priebe. The two-to-infinity norm and singular subspace geometry with applications to high-dimensional statistics. *The Annals of Statistics*, 47(5):2405–2439, 2019b.
- Kamalika Chaudhuri, Fan Chung, and Alexander Tsiatas. Spectral clustering of graphs with general degrees in the extended planted partition model. In *Conference on Learning Theory*, pages 35–1, 2012.
- Han Chen, Garvesh Raskutti, and Ming Yuan. Non-convex projected gradient descent for generalized low-rank tensor regression. *The Journal of Machine Learning Research*, 20(1):172–208, 2019.
- Yuxin Chen, Chen Cheng, and Jianqing Fan. Asymmetry helps: Eigenvalue and eigenvector analyses of asymmetrically perturbed low-rank matrices. *arXiv preprint arXiv:1811.12804*, 2018.
- Eric C Chi, Brian J Gaines, Will Wei Sun, Hua Zhou, and Jian Yang. Provable convex co-clustering of tensors. *Journal of Machine Learning Research*, 21(214):1–58, 2020.
- I Chien, Chung-Yi Lin, and I-Hsiang Wang. On the minimax misclassification ratio of hypergraph community detection. *IEEE Transactions on Information Theory*, 2019.
- Peter Chin, Anup Rao, and Van Vu. Stochastic block model and community detection in sparse graphs: A spectral algorithm with optimal rate of recovery. In *Conference on Learning Theory*, pages 391–423, 2015.
- Andrzej Cichocki, Danilo Mandic, Lieven De Lathauwer, Guoxu Zhou, Qibin Zhao, Cesar Caiafa, and Huy Anh Phan. Tensor decompositions for signal processing applications: From two-way to multiway component analysis. *IEEE Signal Processing Magazine*, 32(2):145–163, 2015.
- Chandler Davis and William Morton Kahan. The rotation of eigenvectors by a perturbation. iii. *SIAM Journal on Numerical Analysis*, 7(1):1–46, 1970.
- Lieven De Lathauwer. First-order perturbation analysis of the best rank-(r_1, r_2, r_3) approximation in multilinear algebra. *Journal of Chemometrics: A Journal of the Chemometrics Society*, 18(1):2–11, 2004.

- Lieven De Lathauwer and Joos Vandewalle. Dimensionality reduction in higher-order signal processing and rank- (r_1, r_2, r_n) reduction in multilinear algebra. *Linear Algebra and its Applications*, 391:31–55, 2004.
- Lieven De Lathauwer, Bart De Moor, and Joos Vandewalle. On the best rank-1 and rank- (r_1, r_2, \dots, r_n) approximation of higher-order tensors. *SIAM journal on Matrix Analysis and Applications*, 21(4):1324–1342, 2000a.
- Lieven De Lathauwer, Bart De Moor, and Joos Vandewalle. A multilinear singular value decomposition. *SIAM journal on Matrix Analysis and Applications*, 21(4):1253–1278, 2000b.
- Carl Eckart and Gale Young. The approximation of one matrix by another of lower rank. *Psychometrika*, 1(3):211–218, 1936.
- Lars Elden and Berkant Savas. A newton–grassmann method for computing the best multilinear rank- (r_1, r_2, r_3) approximation of a tensor. *SIAM Journal on Matrix Analysis and Applications*, 31(2):248–271, 2009.
- Lars Elden and Berkant Savas. Perturbation theory and optimality conditions for the best multilinear rank approximation of a tensor. *SIAM journal on matrix analysis and applications*, 32(4):1422–1450, 2011.
- Justin Eldridge, Mikhail Belkin, and Yusu Wang. Unperturbed: spectral analysis beyond davis-kahan. *arXiv preprint arXiv:1706.06516*, 2017.
- Jianqing Fan, Weichen Wang, and Yiqiao Zhong. An l_∞ eigenvector perturbation bound and its application to robust covariance estimation. *Journal of Machine Learning Research*, 18(207):1–42, 2018.
- Debarghya Ghoshdastidar and Ambedkar Dukkipati. Consistency of spectral partitioning of uniform hypergraphs under planted partition model. In *Advances in Neural Information Processing Systems*, pages 397–405, 2014.
- Debarghya Ghoshdastidar and Ambedkar Dukkipati. Consistency of spectral hypergraph partitioning under planted partition model. *The Annals of Statistics*, 45(1):289–315, 2017.
- Donald Goldfarb and Zhiwei Qin. Robust low-rank tensor recovery: Models and algorithms. *SIAM Journal on Matrix Analysis and Applications*, 35(1):225–253, 2014.
- Gene H Golub, Alan Hoffman, and Gilbert W Stewart. A generalization of the eckart–young–mirsky matrix approximation theorem. *Linear Algebra and its applications*, 88:317–327, 1987.
- Yehoram Gordon. On milman’s inequality and random subspaces which escape through a mesh in r^n . In *Geometric aspects of functional analysis*, pages 84–106. Springer, 1988.
- Lars Grasedyck. Hierarchical singular value decomposition of tensors. *SIAM Journal on Matrix Analysis and Applications*, 31(4):2029–2054, 2010.

- Lars Grasedyck, Daniel Kressner, and Christine Tobler. A literature survey of low-rank tensor approximation techniques. *GAMM-Mitteilungen*, 36(1):53–78, 2013.
- Quanquan Gu, Huan Gui, and Jiawei Han. Robust tensor decomposition with gross corruption. In *Advances in Neural Information Processing Systems*, pages 1422–1430, 2014.
- Wolfgang Hackbusch and Stefan Kühn. A new scheme for the tensor representation. *Journal of Fourier analysis and applications*, 15(5):706–722, 2009.
- Rungang Han, Rebecca Willett, and Anru Zhang. An optimal statistical and computational framework for generalized tensor estimation. *arXiv preprint arXiv:2002.11255*, 2020.
- Botao Hao, Anru Zhang, and Guang Cheng. Sparse and low-rank tensor estimation via cubic sketchings. *IEEE Transactions on Information Theory*, 2020.
- Christopher J Hillar and Lek-Heng Lim. Most tensor problems are np-hard. *Journal of the ACM (JACM)*, 60(6):45, 2013.
- Samuel B Hopkins, Jonathan Shi, and David Steurer. Tensor principal component analysis via sum-of-square proofs. In *Conference on Learning Theory*, pages 956–1006, 2015.
- Mariya Ishteva, Lieven De Lathauwer, P-A Absil, and Sabine Van Huffel. Differential-geometric newton method for the best rank-(r1, r2,r3) approximation of tensors. *Numerical Algorithms*, 51(2):179–194, 2009.
- Mariya Ishteva, P-A Absil, Sabine Van Huffel, and Lieven De Lathauwer. Best low multi-linear rank approximation of higher-order tensors, based on the riemannian trust-region scheme. *SIAM Journal on Matrix Analysis and Applications*, 32(1):115–135, 2011.
- Stefanie Jegelka, Suvrit Sra, and Arindam Banerjee. Approximation algorithms for tensor clustering. In *International Conference on Algorithmic Learning Theory*, pages 368–383. Springer, 2009.
- Bing-Yi Jing, Ting Li, Zhongyuan Lyu, and Dong Xia. Community detection on mixture multi-layer networks via regularized tensor decomposition. *arXiv preprint arXiv:2002.04457*, 2020.
- Zheng Tracy Ke, Feng Shi, and Dong Xia. Community detection for hypergraph networks via regularized tensor power iteration. *arXiv preprint arXiv:1909.06503*, 2019.
- Chiheon Kim, Afonso S Bandeira, and Michel X Goemans. Stochastic block model for hypergraphs: Statistical limits and a semidefinite programming approach. *arXiv preprint arXiv:1807.02884*, 2018.
- Tamara G Kolda. Orthogonal tensor decompositions. *SIAM Journal on Matrix Analysis and Applications*, 23(1):243–255, 2001.
- Tamara G Kolda and Brett W Bader. Tensor decompositions and applications. *SIAM review*, 51(3):455–500, 2009.

- Tamara G Kolda and Jimeng Sun. Scalable tensor decompositions for multi-aspect data mining. In *2008 Eighth IEEE international conference on data mining*, pages 363–372. IEEE, 2008.
- Vladimir Koltchinskii and Dong Xia. Perturbation of linear forms of singular vectors under gaussian noise. *High Dimensional Probability VII*, pages 397–423, 2016.
- Pieter M Kroonenberg. *Applied multiway data analysis*, volume 702. John Wiley & Sons, 2008.
- Pieter M Kroonenberg and Jan De Leeuw. Principal component analysis of three-mode data by means of alternating least squares algorithms. *Psychometrika*, 45(1):69–97, 1980.
- Amit Kumar, Yogish Sabharwal, and Sandeep Sen. A simple linear time $(1+\epsilon)$ -approximation algorithm for k-means clustering in any dimensions. In *45th Annual IEEE Symposium on Foundations of Computer Science*, pages 454–462. IEEE, 2004.
- Chanwoo Lee and Miaoyan Wang. Tensor denoising and completion based on ordinal observations. *arXiv preprint arXiv:2002.06524*, 2020.
- Jing Lei and Alessandro Rinaldo. Consistency of spectral clustering in stochastic block models. *The Annals of Statistics*, 43(1):215–237, 2015.
- Jing Lei, Kehui Chen, and Brian Lynch. Consistent community detection in multi-layer network data. *Biometrika*, 2019.
- James Li, Jacob Bien, and Martin T Wells. rtensor: An r package for multidimensional array (tensor) unfolding, multiplication, and decomposition. *Journal of Statistical Software*, 87(10):1–31, 2018.
- Nan Li and Baoxin Li. Tensor completion for on-board compression of hyperspectral images. In *2010 IEEE International Conference on Image Processing*, pages 517–520. IEEE, 2010.
- Tianqi Liu, Ming Yuan, and Hongyu Zhao. Characterizing spatiotemporal transcriptome of human brain via low rank tensor decomposition. *arXiv preprint arXiv:1702.07449*, 2017.
- Yuanyuan Liu, Fanhua Shang, Wei Fan, James Cheng, and Hong Cheng. Generalized higher-order orthogonal iteration for tensor decomposition and completion. In *Advances in Neural Information Processing Systems*, pages 1763–1771, 2014.
- Haiping Lu, Konstantinos N Plataniotis, and Anastasios N Venetsanopoulos. MPCA: Multilinear principal component analysis of tensor objects. *IEEE transactions on Neural Networks*, 19(1):18–39, 2008.
- Yuetian Luo and Anru R Zhang. Tensor clustering with planted structures: Statistical optimality and computational limits. *arXiv preprint arXiv:2005.10743*, 2020.
- Yuetian Luo and Anru R Zhang. Low-rank tensor estimation via riemannian gauss-newton: Statistical optimality and second-order convergence. *arXiv preprint arXiv:2104.12031*, 2021.

- Yuetian Luo, Rungang Han, and Anru R Zhang. A Schatten- q matrix perturbation analysis via perturbation projection error bound. *arXiv preprint arXiv:2008.01312*, 2020.
- Osman Asif Malik and Stephen Becker. Low-rank Tucker decomposition of large tensors using tensorsketch. In *Advances in Neural Information Processing Systems*, pages 10096–10106, 2018.
- Leon Mirsky. Symmetric gauge functions and unitarily invariant norms. *The quarterly journal of mathematics*, 11(1):50–59, 1960.
- Cun Mu, Daniel Hsu, and Donald Goldfarb. Greedy approaches to symmetric orthogonal tensor decomposition. *SIAM Journal on Matrix Analysis and Applications*, 38(4):1210–1226, 2017.
- Sean O’Rourke, Van Vu, and Ke Wang. Random perturbation of low rank matrices: Improving classical bounds. *Linear Algebra and its Applications*, 540:26–59, 2018.
- Ivan V Oseledets. Tensor-train decomposition. *SIAM Journal on Scientific Computing*, 33(5):2295–2317, 2011.
- Ivan V Oseledets and Eugene E Tyrtysnikov. Breaking the curse of dimensionality, or how to use SVD in many dimensions. *SIAM Journal on Scientific Computing*, 31(5):3744–3759, 2009.
- Evangelos E Papalexakis, Nicholas D Sidiropoulos, and Rasmus Bro. From k-means to higher-way co-clustering: Multilinear decomposition with sparse latent factors. *IEEE transactions on signal processing*, 61(2):493–506, 2012.
- Amelia Perry, Alexander S Wein, Afonso S Bandeira, et al. Statistical limits of spiked tensor models. In *Annales de l’Institut Henri Poincaré, Probabilités et Statistiques*, volume 56, pages 230–264. Institut Henri Poincaré, 2020.
- Garvesh Raskutti, Ming Yuan, and Han Chen. Convex regularization for high-dimensional multiresponse tensor regression. *The Annals of Statistics*, 47(3):1554–1584, 2019.
- Emile Richard and Andrea Montanari. A statistical model for tensor PCA. In *Advances in Neural Information Processing Systems*, pages 2897–2905, 2014.
- Karl Rohe, Sourav Chatterjee, and Bin Yu. Spectral clustering and the high-dimensional stochastic blockmodel. *The Annals of Statistics*, 39(4):1878–1915, 2011.
- Axel Ruhe and Per Ake Wedin. Algorithms for separable nonlinear least squares problems. *SIAM review*, 22(3):318–337, 1980.
- Berkant Savas and Lek-Heng Lim. Quasi-Newton methods on Grassmannians and multilinear approximations of tensors. *SIAM Journal on Scientific Computing*, 32(6):3352–3393, 2010.
- Andrey A Shabalin and Andrew B Nobel. Reconstruction of a low-rank matrix in the presence of Gaussian noise. *Journal of Multivariate Analysis*, 118:67–76, 2013.

- Parikshit Shah, Nikhil Rao, and Gongguo Tang. Sparse and low-rank tensor decomposition. In *Advances in Neural Information Processing Systems*, pages 2548–2556, 2015.
- Bernard N Sheehan and Yousef Saad. Higher order orthogonal iteration of tensors (hooi) and its relation to pca and glam. In *Proceedings of the 2007 SIAM International Conference on Data Mining*, pages 355–365. SIAM, 2007.
- Nicholas D Sidiropoulos, Lieven De Lathauwer, Xiao Fu, Kejun Huang, Evangelos E Papalexakis, and Christos Faloutsos. Tensor decomposition for signal processing and machine learning. *IEEE Transactions on Signal Processing*, 65(13):3551–3582, 2017.
- Laurent Sorber, Marc Van Barel, and Lieven De Lathauwer. Tensorlab v2. 0. *Available online, January*, 2014.
- Gilbert W Stewart. Matrix perturbation theory. 1990.
- Gilbert W Stewart. Perturbation theory for the singular value decomposition. Technical report, 1998.
- Michael Stewart. Perturbation of the svd in the presence of small singular values. *Linear algebra and its applications*, 419(1):53–77, 2006.
- Jimeng Sun, Spiros Papadimitriou, Ching-Yung Lin, Nan Cao, Shixia Liu, and Weihong Qian. Multivis: Content-based social network exploration through multi-way visual analysis. In *Proceedings of the 2009 SIAM International Conference on Data Mining*, pages 1064–1075. SIAM, 2009.
- Yiming Sun, Yang Guo, Charlene Luo, Joel Tropp, and Madeleine Udell. Low-rank tucker approximation of a tensor from streaming data. *SIAM Journal on Mathematics of Data Science*, 2(4):1123–1150, 2020.
- Daniel L Sussman, Minh Tang, Donniell E Fishkind, and Carey E Priebe. A consistent adjacency spectral embedding for stochastic blockmodel graphs. *Journal of the American Statistical Association*, 107(499):1119–1128, 2012.
- Charalampos E Tsourakakis. Mach: Fast randomized tensor decompositions. In *Proceedings of the 2010 SIAM International Conference on Data Mining*, pages 689–700. SIAM, 2010.
- Ledyard R Tucker. Some mathematical notes on three-mode factor analysis. *Psychometrika*, 31(3):279–311, 1966.
- Andre Uschmajew. A new convergence proof for the higher-order power method and generalizations. *Pacific Journal Of Optimization*, 11(ARTICLE):309–321, 2015.
- Nick Vannieuwenhoven, Raf Vandebril, and Karl Meerbergen. A new truncation strategy for the higher-order singular value decomposition. *SIAM Journal on Scientific Computing*, 34(2):A1027–A1052, 2012.
- Roman Vershynin. Introduction to the non-asymptotic analysis of random matrices. *arXiv preprint arXiv:1011.3027*, 2010.

- Van Vu. Singular vectors under random perturbation. *Random Structures & Algorithms*, 39(4):526–538, 2011.
- Chang-Dong Wang, Jian-Huang Lai, and S Yu Philip. Multi-view clustering based on belief propagation. *IEEE Transactions on Knowledge and Data Engineering*, 28(4):1007–1021, 2015.
- Liqi Wang and Moody T Chu. On the global convergence of the alternating least squares method for rank-one approximation to generic tensors. *SIAM Journal on Matrix Analysis and Applications*, 35(3):1058–1072, 2014.
- Miaoyan Wang and Yuchen Zeng. Multiway clustering via tensor block models. In *Advances in Neural Information Processing Systems*, pages 713–723, 2019.
- Rongrong Wang. Singular vector perturbation under gaussian noise. *SIAM Journal on Matrix Analysis and Applications*, 36(1):158–177, 2015.
- Per-Ake Wedin. Perturbation bounds in connection with singular value decomposition. *BIT Numerical Mathematics*, 12(1):99–111, 1972.
- Hermann Weyl. Das asymptotische verteilungsgesetz der eigenwerte linearer partieller differentialgleichungen (mit einer anwendung auf die theorie der hohlraumstrahlung). *Mathematische Annalen*, 71(4):441–479, 1912.
- Tao Wu, Austin R Benson, and David F Gleich. General tensor spectral co-clustering for higher-order data. In *Advances in Neural Information Processing Systems*, pages 2559–2567, 2016.
- Dong Xia and Ming Yuan. On polynomial time methods for exact low-rank tensor completion. *Foundations of Computational Mathematics*, pages 1–49, 2017.
- Dong Xia and Fan Zhou. The sup-norm perturbation of hosvd and low rank tensor denoising. *Journal of Machine Learning Research*, 20(61):1–42, 2019.
- Dong Xia, Ming Yuan, and Cun-Hui Zhang. Statistically optimal and computationally efficient low rank tensor completion from noisy entries. *The Annals of Statistics*, to appear, 2020.
- Dong Xu, Shuicheng Yan, Lei Zhang, Hong-Jiang Zhang, Zhengkai Liu, and Heung-Yeung Shum. Concurrent subspaces analysis. In *2005 IEEE Computer Society Conference on Computer Vision and Pattern Recognition (CVPR’05)*, volume 2, pages 203–208. IEEE, 2005.
- Yangyang Xu. On the convergence of higher-order orthogonal iteration. *Linear and Multilinear Algebra*, 66(11):2247–2265, 2018.
- Shuicheng Yan, Dong Xu, Qiang Yang, Lei Zhang, Xiaoou Tang, and Hong-Jiang Zhang. Discriminant analysis with tensor representation. In *2005 IEEE Computer Society Conference on Computer Vision and Pattern Recognition (CVPR’05)*, volume 1, pages 526–532. IEEE, 2005.

- Shuicheng Yan, Dong Xu, Qiang Yang, Lei Zhang, Xiaoou Tang, and Hong-Jiang Zhang. Multilinear discriminant analysis for face recognition. *IEEE Transactions on Image Processing*, 16(1):212–220, 2006.
- Yi Yu, Tengyao Wang, and Richard J Samworth. A useful variant of the davis–kahan theorem for statisticians. *Biometrika*, 102(2):315–323, 2014.
- Ming Yuan and Cun-Hui Zhang. On tensor completion via nuclear norm minimization. *Foundations of Computational Mathematics*, 16(4):1031–1068, 2016.
- Ming Yuan and Cun-Hui Zhang. Incoherent tensor norms and their applications in higher order tensor completion. *IEEE Transactions on Information Theory*, 63(10):6753–6766, 2017.
- Anru Zhang. Cross: Efficient low-rank tensor completion. *The Annals of Statistics*, 47(2):936–964, 2019.
- Anru Zhang and Rungang Han. Optimal sparse singular value decomposition for high-dimensional high-order data. *Journal of the American Statistical Association*, 114(528):1708–1725, 2019.
- Anru Zhang and Dong Xia. Tensor svd: Statistical and computational limits. *IEEE Transactions on Information Theory*, 64(11):7311–7338, 2018.
- Anru R Zhang, Yuetian Luo, Garvesh Raskutti, and Ming Yuan. Islet: Fast and optimal low-rank tensor regression via importance sketching. *SIAM Journal on Mathematics of Data Science*, 2(2):444–479, 2020.
- Tong Zhang and Gene H Golub. Rank-one approximation to high order tensors. *SIAM Journal on Matrix Analysis and Applications*, 23(2):534–550, 2001.
- Qinqing Zheng and Ryota Tomioka. Interpolating convex and non-convex tensor decompositions via the subspace norm. In *Advances in Neural Information Processing Systems*, pages 3106–3113, 2015.
- Hua Zhou, Lexin Li, and Hongtu Zhu. Tensor regression with applications in neuroimaging data analysis. *Journal of the American Statistical Association*, 108(502):540–552, 2013.
- Yuchen Zhou, Anru R Zhang, Lili Zheng, and Yazhen Wang. Optimal high-order tensor svd via tensor-train orthogonal iteration. *arXiv preprint arXiv:2010.02482*, 2020.