

# On the invertibility in periodic ARFIMA models

AMINE AMIMOUR<sup>1</sup>

and

KARIMA BELAIDE<sup>2</sup>

Department of Mathematics  
Applied Mathematics Laboratory  
University of Bejaia, Bejaia. Algeria

[amineamimour@gmail.com](mailto:amineamimour@gmail.com)<sup>1</sup>

[k\\_tim2002@yahoo.fr](mailto:k_tim2002@yahoo.fr)<sup>2</sup>

## Abstract

The present paper, characterizes the invertibility and causality conditions of a periodic ARFIMA (PARFIMA) models. We first, discuss the conditions in the multivariate case, by considering the corresponding p-variate stationary ARFIMA models. Second, we construct the conditions using the univariate case and we deduce a new infinite autoregressive representation for the PARFIMA model, the results are investigated through a simulation study.

**Keywords:** PARFIMA. Invertibility condition. Fractionally process. Simulation. Long range dependence. Long memory.

## 1 Introduction and notations

A long memory processes have long been studied in the literature. (See, e.g. Granger and Joyeux [6] and Hosking [7] for early work, Baillie [3] for more background). Hosking [7] proposed a purely fractionally differenced autoregressive moving average process  $(X_t)_{t \in \mathbb{Z}}$  denoted by ARFIMA(0,  $d$ , 0) or ARFIMA for short, given by the equation

$$(1 - B)^d X_t = \varepsilon_t, \quad (1.1)$$

where  $d$  is a real number which denote the lag of the ARFIMA process,  $\varepsilon_t$  is a sequence of independent and identically distributed (i.i.d) random variables with mean zero and finite variance  $\sigma^2$ , and  $B$  is the back shift operator, such that  $X_{t-j} = B^j X_t$ . The invertibility concept

of time series models, loosely says that the ARFIMA model is invertible when we are able to express the noise process  $\varepsilon_t$  as a convergent series of the observations  $X_t$ . The characteristic equation for a ARFIMA process is

$$\varepsilon_t = \sum_{j=0}^{\infty} \pi_j X_{t-j}, \quad (1.2)$$

where

$$\pi_j = \frac{\Gamma(-d+j)}{\Gamma(-d)\Gamma(j+1)}.$$

$\Gamma(\cdot)$  is a gamma function and it is given by

$$\Gamma(z) = \begin{cases} \int_0^{\infty} s^{z-1} e^{-s} ds, & \text{si } z > 0 \\ \infty, & \text{si } z = 0 \end{cases},$$

if  $z < 0$ ,  $\Gamma(z)$  is defined by the recurrence formula  $z\Gamma(z) = \Gamma(z+1)$ . In order for a ARFIMA process to be invertible, the infinite sum of the coefficients  $\pi_j$  must be absolutely summable. However, when the lag  $d > -\frac{1}{2}$ , the infinite sum of the coefficients  $\pi_j$  is absolutely summable. Odaki[9] noticed that the ARFIMA process is invertible even when  $-1 < d$ . These conditions concerning the univariate case. Chung [4] used a Vector ARFIMA model to define the invertibility condition of multivariate p-dimensional stationary process in the sense of Hosking [7]. Recently, the invertibility condition for the bivariate and multivariate case is achieved by Kechagias and Pipiras [8], they extended the definition of the bivariate LRD of Robinson [10], to the multivariate case, concluding that the trigonometric power-law coefficients, can be used to construct new univariate and multivariate causal LRD series.

Since a several time series encountered in practice exhibit periodic autocorrelation structure, a fact that cannot be explained, by the classical seasonal models or by the time series models with parameters invariants in time. In fact, Franses and Ooms [5] considered the PARFIMA models defined by

$$X_t = (1 - B)^{-d_s} \varepsilon_t, \quad (1.3)$$

where,  $d_s$  is the fractional parameter which can vary with season  $s = 1, 2, 3, 4$ , to study quarterly inflation in the UK, allowing the degree of fractional integration to vary with season, the series analyzed in this study concern the quarterly inflation rate verified over a 33-year period, after the presentation of the correlogram, they concluded that the autocorrelations do not decrease rapidly, which indicates long-memory behavior in the series. Moreover, if the transformation of the series be performed by classical differentiation in the sense of Box Jenkins, the autocorrelations indicate that the differentiated series can be overdifferentiated, because the sum of the autocorrelations will be close to  $-0.5$ , then the observations suggest the utility possible of ARFIMA models. Therefore, for a more appropriate modeling, which takes into account seasonal variation they have proposed to merge the presence of long memory and the periodic dynamics, into a new model. The key result of these authors is that this model compared with the periodic short-memory models and long-memory models with constant coefficients, not only

provides a plausible informative description of the periodic series, but also, can be useful for out-of-sample forecasts. Amimour and Belaïde [2] studied recently, the probabilistic property of PARFIMA models, in the sense that the long memory parameter varies periodically in time such that  $d_{t+p} = d_t$  with period  $p \in \mathbb{N}^*$ ;  $p > 1$ , denoted by PtvARFIMA and has the following stochastic equation

$$(1 - B)^{d_i} X_{i+pm} = \varepsilon_{i+pm}, \quad (1.4)$$

where for all  $t \in \mathbb{Z}$ , there exists  $i = \{1, \dots, p\}$ ,  $m \in \mathbb{Z}$ , such that  $t = i + pm$  and the variance is periodic in  $t$  such that  $\sigma_{t+pm}^2 = \sigma_t^2$ , the authors constructed also a local asymptotic normality property for this model see [1]. Indeed, the PARFIMA models have received little attention from time series analysts due to these complexities, one of the main difficulties is to achieving an invertibility condition. Noting that the works cited above based only on the sufficient condition of invertibility and causality, that is

when  $d_i > 0$ , the process (1.4) is invertible and has an infinite autoregressive representation as follows

$$\varepsilon_{i+pm} = (1 - B)^{d_i} X_{i+pm} = \sum_{j=0}^{\infty} \pi_j^i X_{i+pm-j}, \quad (1.5)$$

where

$$\pi_j^i = \frac{\Gamma(j-d_i)}{\Gamma(j+1)\Gamma(-d_i)},$$

when  $d_i < \frac{1}{2}$ , the process (1.4) is causal and has an infinite moving-average representation as follows

$$X_{i+pm} = (1 - B)^{-d_i} \varepsilon_{i+pm} = \sum_{j=0}^{\infty} \psi_j^i \varepsilon_{i+pm-j}, \quad (1.6)$$

where

$$\psi_j^i = \frac{\Gamma(j+d_i)}{\Gamma(j+1)\Gamma(d_i)}.$$

$$\psi_j^i \sim v_i j^{d_i-1}, v_i > 0, \text{ as } j \rightarrow \infty. \quad (1.7)$$

See propositions (2.1) and (2.2) in Amimour and Belaïde [2]. For instance, precise information on the invertibility conditions of a PtvARFIMA model is important to circumscribe a model's parameter space. Moreover, verifying invertibility in the necessary and sufficient conditions

sense, is required for statistical inference, in this article we address the above topic. To find an invertibility condition of PtvARFIMA.

The work is organized as follows. We first discuss in section 2 some characteristics of our model using his associated stationary multivariate process. In section 3, we focus on the constructing of new infinite autoregressive representation of our model. We finish this work by the simulation for illustrating the results in section 4.

## 2 PtvARFIMA and p-variate stationary ARFIMA models

The PtvARFIMA model is first introduced by Amimour and Belaïde [2], this model extends the ARFIMA model from Hosking [7]. PtvARFIMA time series exhibit long range dependence (LRD), and the periodic phenomenon, the periodic covariance function  $\gamma_X^i(h)$  echoes LRD and decays periodically as the lag  $h$  increases, the asymptotic behavior of this function is given by the following explicit form, as  $h \rightarrow \infty$

$$\gamma_X^i(h) \simeq \begin{cases} \gamma_X^1(h) \simeq \begin{cases} \sigma_1^2 \frac{\Gamma(1-d_1-d_2)}{\Gamma(d_2)\Gamma(1-d_2)}(h)^{d_1+d_2-1}, & \text{if } h \equiv 1[p], \\ \sigma_1^2 \frac{\Gamma(1-d_1-d_3)}{\Gamma(d_3)\Gamma(1-d_3)}(h)^{d_1+d_3-1}, & \text{if } h \equiv 2[p], \\ \cdot \\ \cdot \\ \sigma_1^2 \frac{\Gamma(1-2d_1)}{\Gamma(d_1)\Gamma(1-d_1)}(h)^{2d_1-1}, & \text{if } h \equiv 0[p]. \end{cases} \\ \gamma_X^2(h) \simeq \begin{cases} \sigma_2^2 \frac{\Gamma(1-d_2-d_3)}{\Gamma(d_3)\Gamma(1-d_3)}(h)^{d_2+d_3-1}, & \text{if } h \equiv 1[p], \\ \sigma_2^2 \frac{\Gamma(1-d_2-d_4)}{\Gamma(d_4)\Gamma(1-d_4)}(h)^{d_2+d_4-1}, & \text{if } h \equiv 2[p], \\ \cdot \\ \cdot \\ \sigma_2^2 \frac{\Gamma(1-2d_2)}{\Gamma(d_2)\Gamma(1-d_2)}(h)^{2d_2-1}, & \text{if } h \equiv 0[p]. \end{cases} \\ \cdot \\ \cdot \\ \cdot \\ \gamma_X^p(h) \simeq \begin{cases} \sigma_p^2 \frac{\Gamma(1-d_p-d_1)}{\Gamma(d_1)\Gamma(1-d_1)}(h)^{d_p+d_1-1}, & \text{if } h \equiv 1[p], \\ \sigma_p^2 \frac{\Gamma(1-d_p-d_2)}{\Gamma(d_2)\Gamma(1-d_2)}(h)^{d_p+d_2-1}, & \text{if } h \equiv 2[p], \\ \cdot \\ \cdot \\ \sigma_p^2 \frac{\Gamma(1-2d_p)}{\Gamma(d_p)\Gamma(1-d_p)}(h)^{2d_p-1}, & \text{if } h \equiv 0[p]. \end{cases} \end{cases}$$

In this section, we will focus on the PtvARFIMA model as a multivariate stationary ARFIMA time series, that is,  $\mathbb{R}^p$ -valued time series  $X_m = (X_{1+pm}, \dots, X_{p+pm})^T$ , where  $T$  denotes the transpose.

**Proposition 2.1.** *A PtvARFIMA model defined in (1.4) is a  $p$ -variate second order stationary long memory time series.*

*Proof.* Suppose that for  $a > 0$ , a diagonal matrix  $N = \text{diag}(n_1, \dots, n_p)$ , we write  $a^N = \text{diag}(a^{n_1}, \dots, a^{n_p})$ . The periodic covariance function  $\gamma_X^i(h)$  previously given, can be expressed as a matrix function of a second order stationary series  $X = \{X_m\}_{m \in \mathbb{Z}}$ , given by

$$\gamma_X^{i,k}(h) \simeq \sigma_i^2 h^{D-\frac{1}{2}I} R h^{D-\frac{1}{2}I} = (\sigma_i^2 R^{i,k} h^{d_i+d_k-1}), \quad i, k = 1, \dots, p, \quad \text{as } h \rightarrow \infty, \quad (2.1)$$

where  $D = \text{diag}(d_1, \dots, d_p)$ , and  $R^{i,k} > 0$  for all  $i, k = 1, \dots, p$ .  $\square$

**Proposition 2.2.** *The causal representation of the  $p$ -variate ARFIMA time series, associated with the PtvARFIMA model with period  $p$  is*

$$X_m = \sum_{j=0}^{\infty} \Upsilon_j \varepsilon_{m-j}, \quad (2.2)$$

and

$$R^{i,k} = \frac{\Gamma(d_i)\Gamma(d_k)}{\Gamma(d_i+d_k)} \left( v_i v_k \frac{\sin(\pi d_k)}{\sin(\pi(d_i+d_k))} \right), \quad (2.3)$$

where  $\Upsilon_j$  is a diagonal matrix  $\text{diag}(\psi_j^1, \dots, \psi_j^p)$  and  $\varepsilon_m$  is an  $\mathbb{R}^p$ -valued white noise  $(\varepsilon_{1+pm}, \dots, \varepsilon_{p+pm})^T$ , satisfying  $\mathbb{E}[\varepsilon_m] = 0$  and  $\mathbb{E}[\varepsilon_m \varepsilon_m'] = \sigma_i^2 I$ .

*Proof.* Let us calculate the autocovariance function  $\gamma_X^{i,k}(h)$

$$\begin{aligned} \gamma_X^{i,k}(h) &\simeq \sigma_i^2 \sum_{j=0}^{\infty} v_i j^{d_i-1} v_k (j+h)^{d_k-1} \\ &\simeq \sigma_i^2 \sum_{j=0}^{\infty} v_i v_k j^{d_i-1} (j+h)^{d_k-1} \\ &\simeq \sigma_i^2 \sum_{j=0}^{\infty} v_i v_k j^{d_i-1} (j+h)^{d_k-1} \\ &\simeq \sigma_i^2 \sum_{j=0}^{\infty} v_i v_k j^{d_i-1} (j+h)^{d_k-1} = \sum_{j=0}^{\infty} v_i v_k h^{d_i-1} \left(\frac{j}{h}\right)^{d_i-1} h^{d_k-1} \left(\frac{j}{h} + 1\right)^{d_k-1}, \end{aligned}$$

from (Appendix A, in Kechagias and Pipiras [8])

$$\gamma_X^{i,k}(h) \simeq \sigma_i^2 h^{d_i+d_k+1} v_i v_k \int_1^{\infty} z^{d_i-1} (z+1)^{d_k-1} dz \sim R^{i,k} h^{d_i+d_k-1},$$

where

$$R^{i,k} = v_i v_k \frac{\Gamma(d_i)\Gamma(1-d_i-d_k)}{\Gamma(1-d_k)},$$

and using the identity

$$\Gamma(z)\Gamma(1-z) = \frac{\pi}{\sin(\pi z)}, \quad 0 < z < 1,$$

we deduce the expression of  $R^{i,k}$ .  $\square$

### 3 A new infinite autoregressive representation for Pt-vARFIMA

**Theorem 3.1.** Let  $\{X_{i+pm}\}_{m \in \mathbb{Z}}$  be the PARFIMA(0,  $d_i$ , 0) process, given by the expression (1.4), then

(i) When  $-\frac{1}{2} < d_i < \frac{3}{2}$ , or  $|d_i| < 1$ .  $\{X_{i+pm}\}_{m \in \mathbb{Z}}$  is a invertible process with infinite autoregressive representation given by

$$\varepsilon_{i+pm} = \sum_{j=0}^{\infty} \Pi_j(i) X_{i+pm-j}. \quad (3.1)$$

The coefficients  $\Pi_j(i)$   $j \geq 1$  in (3.1) satisfy the following relation

$$\Pi_j(i) = -(\psi_j(i) + \sum_{l=1}^{j-1} \Pi_l(i) \psi_{j-l}(i+k)), \quad (3.2)$$

where the weights  $\psi_j(i)$   $j \geq 0$  are given in (1.6) and  $l \equiv k[p]$ ,  $\Pi_0(i) = 1$ , with  $\sum_{j=0}^{\infty} |\Pi_j(i)| < \infty$ .

(ii) When  $-\frac{3}{2} < d_i < \frac{1}{2}$ , or  $|d_i| < 1$ .  $\{X_{i+pm}\}_{m \in \mathbb{Z}}$  is a causal process with infinite moving average representation given by

$$X_{i+pm} = \sum_{j=0}^{\infty} \Psi_j(i) \varepsilon_{i+pm-j}. \quad (3.3)$$

The coefficients  $\Psi_j(i)$   $j \geq 1$  in (3.3) satisfy the following relation

$$\Psi_j(i) = -(\pi_j(i) + \sum_{l=1}^{j-1} \Psi_l(i) \pi_{j-l}(i+k)), \quad (3.4)$$

where the weights  $\pi_j(i)$   $j \geq 0$  are given in (1.5) and  $l \equiv k[p]$ ,  $\Pi_0(i) = 1$ , with  $\sum_{j=0}^{\infty} |\Psi_j(i)| < \infty$ .

*Proof.* For simplicity of notation, we use  $\psi_j(i) = \psi_j^i$  and  $\pi_j(i) = \pi_j^i$ .

(i) We have

$$X_{i+pm} = \sum_{j=0}^{\infty} \psi_j(i) \varepsilon_{i+pm-j}, \quad (3.5)$$

we can develop the series as follows

$$\begin{aligned} X_{i+pm} &= \varepsilon_{i+pm} + \psi_1(i) \varepsilon_{i+pm-1} + \psi_2(i) \varepsilon_{i+pm-2} + \dots \\ &= \varepsilon_{i+pm} \\ &\quad + \psi_1(i) \overbrace{[\varepsilon_{i+pm-1} + \psi_1(i+1) \varepsilon_{i+pm-2} + \psi_2(i+1) \varepsilon_{i+pm-3} \dots]}^{X_{i+pm-1}} \\ &\quad - \psi_1(i) [\psi_1(i+1) \varepsilon_{i+pm-2} + \psi_2(i+1) \varepsilon_{i+pm-3} \dots] \\ &\quad + \psi_2(i) \varepsilon_{i+pm-2} + \psi_3(i) \varepsilon_{i+pm-3} \dots \end{aligned}$$

$$\begin{aligned}
X_{i+pm} - \psi_1(i)X_{i+pm-1} &= \varepsilon_{i+pm} \\
&+ [\psi_2(i) - \psi_1(i)\psi_1(i+1)] \overbrace{[\varepsilon_{i+pm-2} + \psi_1(i+2)\varepsilon_{i+pm-3} + \dots]}^{X_{i+pm-2}} \\
&- [\psi_2(i) - \psi_1(i)\psi_1(i+1)] [\psi_1(i+2)\varepsilon_{i+pm-3} \dots] \\
&+ [\psi_3(i) - \psi_1(i)\psi_2(i+1)] \varepsilon_{i+pm-3} + \dots
\end{aligned}$$

$$\begin{aligned}
X_{i+pm} - \overbrace{\psi_1(i)}^{\phi_1(i)} X_{i+pm-1} &= \varepsilon_{i+pm} \\
&+ \overbrace{[\psi_2(i) - \psi_1(i)\psi_1(i+1)]}^{\phi_2(i)} \overbrace{[\varepsilon_{i+pm-2} + \psi_1(i+2)\varepsilon_{i+pm-3} + \dots]}^{X_{i+pm-2}} \\
&- [\psi_2(i) - \psi_1(i)\psi_1(i+1)] [\psi_1(i+2)\varepsilon_{i+pm-3} \dots] \\
&+ [\psi_3(i) - \psi_1(i)\psi_2(i+1)] \varepsilon_{i+pm-3} + \dots
\end{aligned}$$

$$X_{i+pm} - \phi_1(i)X_{i+pm-1} - \phi_2(i)X_{i+pm-2} \dots = \varepsilon_{i+pm}, \quad (3.6)$$

such that

$$\phi_j(i) = \psi_j(i) - \sum_{l=1}^{j-1} \phi_l(i)\psi_{j-l}(i+k), \quad (3.7)$$

with  $l \equiv k[p]$ . Next it is easy to show that  $\Pi_j(i) = -\phi_j(i)$ . This result is illustrated by the simulation in the next section.

(ii) The proof is similar to (i) but  $d_i$  must be replaced by  $-d_i$ .  $\square$

## 4 Simulation

In this section, we provide some simple examples, to illustrate the above theoretical results in the invertible case, as well as the non-invertible case for  $p = 2$ . More precisely, we show that the absolute sum of coefficients  $\Pi_j(i)$  is finite for  $-\frac{1}{2} < d_i < \frac{3}{2}$  or  $|d_i| < 1$  in the first case and infinite for  $d_1 < -\frac{1}{2}$  and  $d_2 < \frac{3}{2}$ ,  $d_1 > -\frac{1}{2}$  and  $d_2 > \frac{3}{2}$  or  $|d_i| > 1$  in the second case, for all  $N$  such that  $j = 0, 1, \dots, N$ . The sum of  $\Pi_j(i)$  is summable absolutely if the difference between the previous and the present value of  $\Pi_j(i)$ , e.g.  $(|\Pi_N(i) - \Pi_{N+1}(i)|)$ , will be very low or negligible. Numerical values of  $|\Pi_N(i) - \Pi_{N+1}(i)|$  are presented in table (1) for the invertible case. Table (2) for non invertible case, we find that the results indicated in the tables reflect the theoretical framework.

For  $-\frac{1}{2} < d_i < \frac{3}{2}$  or  $|d_i| < 1$  (see table 1) it is easy to see that the sum of  $|\Pi_j(i)|$  converges, i.e the sum of  $|\Pi_j(i)|$  stabilizes quickly in this case, since the quantity  $|\Pi_N(i) - \Pi_{N+1}(i)|$  decrease when  $N$  increase and have inconsiderable values. Contrarily, if the values of  $d_i$  are outside the intervals cited (see table 2), this demonstrates that the sum diverges even for large values of

$N$	$d_1$	$d_2$	$d_1$	$d_2$	$d_1$	$d_2$	$d_1$	$d_2$	$d_1$	$d_2$
	0.15	0.8	1.49	-0.49	0.75	0.2	0.9	0.09	-0.2	-0.7
10	$1.976584 \times 10^{-03}$		4.02289467		0.0128260513		$6.675804 \times 10^{-03}$		$2.775860 \times -04$	
25	$7.551323 \times 10^{-04}$		1.04204572		0.0016859353		$6.473788 \times 10^{-04}$		$1.642120 \times -04$	
50	$2.060515 \times 10^{-04}$		0.71213528		0.0007002542		$2.067307 \times 10^{-04}$		$5.866394 \times -05$	
75	$1.214109 \times 10^{-04}$		0.50327845		0.0003220433		$8.734869 \times 10^{-05}$		$4.469138 \times -05$	
100	$7.671041 \times 10^{-05}$		0.09702500		0.0001664161		$4.752873 \times 10^{-05}$		$3.821700 \times -05$	
$N$	$d_1$	$d_2$	$d_1$	$d_2$	$d_1$	$d_2$	$d_1$	$d_2$	$d_1$	$d_2$
	-0.6	-0.3	-0.9	-0.09	-0.5	-0.4	-0.49	-0.9	0.49	0.09
10	0.0252137		4.0228946		0.01134118		$2.696864 \times 10^{-03}$		0.01195167	
25	0.0092933		1.0420457		0.00540992		$4.487907 \times 10^{-04}$		0.00190099	
50	0.0061061		0.7121352		0.00242034		$9.400265 \times 10^{-05}$		0.00097871	
75	0.0040766		0.5032784		0.00173045		$4.431224 \times 10^{-05}$		0.00047477	
100	0.0032788		0.0970250		0.00131972		$2.515928 \times 10^{-05}$		0.00032591	

Table 1: Values of  $(|\Pi_N(i) - \Pi_{N+1}(i)|)$  for  $-\frac{1}{2} < d_i < \frac{3}{2}$  or  $|d_i| < 1$  and various  $N$ 

$N$	$d_1$	$d_2$	$d_1$	$d_2$	$d_1$	$d_2$
	-0.6	1.49	-0.4	1.65	-1.2	-1.4
10	1.4451672		0.7828075		0.2448915	
25	11.2747847		4.0371864		0.4049227	
50	21.7021959		3.6322384		0.4341756	
75	332.0970990		27.3838075		0.5108372	
100	1686.6896433		68.7898114		0.5445771	

Table 2: Values of  $(|\Pi_N(i) - \Pi_{N+1}(i)|)$  for  $d_1 < -\frac{1}{2}$  and  $d_2 < \frac{3}{2}$ ,  $d_1 > -\frac{1}{2}$  and  $d_2 > \frac{3}{2}$  or  $|d_i| > 1$  and various  $N$



$N$ , in this case the quantity  $|\Pi_N(i) - \Pi_{N+1}(i)|$  is considerable and sometimes increases when  $N$  increase.

## 5 Conclusion

In this article, we have discussed in one hand the p-variate stationary ARFIMA models associated with the PARFIMA model considered here. We have represented the model as a causal representation, showing that the asymptotic behavior of the periodic autocovariance function can be obtained using the multivariate moving average representation. On the other hand, we have established a new infinite autoregressive representation for the PARFIMA model and illustrated the results via a simulation study.

## References

- [1] A. AMIMOUR, K. BELAIDE, Local asymptotic normality for a periodically time varying long memory parameter. *Communications in Statistics - Theory and Methods*, doi : 10.1080/03610926.2020.1784435, (2020).
- [2] A. AMIMOUR, K. BELAIDE, A long memory time series with a periodic degree of fractional differencing. *arXiv preprint arXiv: 2008.01939*, (2020).
- [3] R.T. BAILLIE, Long memory processes and fractional integration in econometrics. *Journal of econometrics*, **73** (1996), pages 5–59.
- [4] CF. CHUNG, Sample means, Sample autocovariances, and linear regression of stationary multivariate long memory processes. *Econometric Theory*, **18** (2002), pages 51–78.
- [5] P.H. FRANSES, M. OOMS, A periodic long-memory model for quarterly UK inflation. *International Journal of Forecasting* , **13** (1997), pages 117–126.
- [6] C.W.J. GRANGER, R. JOYEUX, An introduction to long memory time series models and fractional differencing. *J. Time Ser. Anal.*, **1** (1980), pages 15–29.
- [7] J.R.M. HOSKING, Fractional differencing. *Biometrika.*, **68** (1981), pages 165–176.
- [8] S. KECHAGIAS, V. PIPIRAS, Definitions and representations of multivariate long-range dependent time series. *J. Time Ser. Anal.*, **36** (2015), pages 1–25.
- [9] M. ODAKI, On the invertibility of fractionally differenced ARIMA processes . *Biometrika* **80** (1993), pages 703–709.
- [10] P.M ROBINSON, Multiple local whittle estimation in stationary systems. *The annals of statistics*, **36** (2008), pages 2508–2530.