

# Minimax estimation of norms of a probability density: II. Rate-optimal estimation procedures

A. Goldenshluger <sup>\*</sup> and O. V. Lepski <sup>†</sup>

*Department of Statistics  
University of Haifa  
Mount Carmel  
Haifa 31905, Israel  
e-mail: [goldensh@stat.haifa.ac.il](mailto:goldensh@stat.haifa.ac.il)*

*Institut de Mathématique de Marseille  
Aix-Marseille Université  
39, rue F. Joliot-Curie  
13453 Marseille, France  
e-mail: [oleg.lepski@univ-amu.fr](mailto:oleg.lepski@univ-amu.fr)*

**Abstract:** In this paper we develop rate-optimal estimation procedures in the problem of estimating the  $\mathbb{L}_p$ -norm,  $p \in (0, \infty)$  of a probability density from independent observations. The density is assumed to be defined on  $\mathbb{R}^d$ ,  $d \geq 1$  and to belong to a ball in the anisotropic Nikolskii space. We adopt the minimax approach and construct rate-optimal estimators in the case of integer  $p \geq 2$ . We demonstrate that, depending on parameters of Nikolskii's class and the norm index  $p$ , the risk asymptotics ranges from inconsistency to  $\sqrt{n}$ -estimation. The results in this paper complement the minimax lower bounds derived in the companion paper [Goldenshluger and Lepski \(2020\)](#).

**AMS 2000 subject classifications:** 62G05, 62G20.

**Keywords and phrases:** density estimation, minimax risk,  $\mathbb{L}_p$ -norm,  $U$ -statistics, anisotropic Nikolskii class.

## 1. Introduction

Suppose that we observe i.i.d. random vectors  $X_i \in \mathbb{R}^d$ ,  $i = 1, \dots, n$ , with common probability density  $f$ . Let  $p > 1$  be a given number. We are interested in estimating the  $\mathbb{L}_p$ -norm of  $f$ ,

$$\|f\|_p := \left[ \int_{\mathbb{R}^d} |f(x)|^p dx \right]^{1/p},$$

from observation  $X^{(n)} = (X_1, \dots, X_n)$ . By estimator of  $\|f\|_p$  we mean any  $X^{(n)}$ -measurable map  $\tilde{N} : \mathbb{R}^n \rightarrow \mathbb{R}$ . Accuracy of an estimator  $\tilde{N}$  is measured by the quadratic risk

$$\mathcal{R}_n[\tilde{N}, f] := \left( \mathbb{E}_f [\tilde{N} - \|f\|_p]^2 \right)^{1/2},$$

where  $\mathbb{E}_f$  denotes expectation with respect to the probability measure  $\mathbb{P}_f$  of the observations  $X^{(n)} = (X_1, \dots, X_n)$ .

We adopt minimax approach to measuring estimation accuracy. Let  $\mathfrak{F}$  denote the set of all probability densities defined on  $\mathbb{R}^d$ . The maximal risk of an estimator  $\tilde{N}$  on the set  $\mathbb{F} \subset \mathfrak{F}$  is

<sup>\*</sup>Supported by the ISF grant No. 361/15.

<sup>†</sup>This work has been carried out in the framework of the Labex Archimède (ANR-11-LABX-0033) and of the A\*MIDEX project (ANR-11-IDEX-0001-02), funded by the "Investissements d'Avenir" French Government program managed by the French National Research Agency (ANR).

defined by

$$\mathcal{R}_n[\tilde{N}, \mathbb{F}] := \sup_{f \in \mathbb{F}} \mathcal{R}_n[\tilde{N}, f],$$

and the minimax risk is

$$\mathcal{R}_n[\mathbb{F}] := \inf_{\tilde{N}} \mathcal{R}_n[\tilde{N}; \mathbb{F}],$$

where inf is taken over all possible estimators.

In the companion paper [Goldenshluger and Lepski \(2020\)](#) (referred to hereafter as Part I), we derived lower bounds on the minimax risk over functional classes  $\mathbb{F} = \mathbb{N}_{\vec{r}, d}(\vec{\beta}, \vec{L}) \cap \mathbb{B}_q(Q)$ , where  $\mathbb{N}_{\vec{r}, d}(\vec{\beta}, \vec{L})$  denotes anisotropic Nikolskii's class [see Definition 1 below], and  $\mathbb{B}_q(Q) := \{f : \|f\|_q \leq Q\}$  is the ball in  $\mathbb{L}_q(\mathbb{R}^d)$  of radius  $Q$ . Specifically, we found the sequence  $\phi_n$  completely determined by the  $\vec{\beta}, \vec{L}, \vec{r}, q, p$  and  $n$  such that

$$\mathcal{R}_n[\mathbb{N}_{\vec{r}, d}(\vec{\beta}, \vec{L}) \cap \mathbb{B}_q(Q)] \gtrsim \phi_n, \quad n \rightarrow \infty.$$

The goal of the present paper is to develop a rate-optimal estimator, say,  $\hat{N}$ , such that for any given  $\vec{\beta}, \vec{r}, \vec{L}, q, Q$

$$\limsup_{n \rightarrow \infty} \phi_n^{-1} \mathcal{R}_n[\hat{N}, \mathbb{N}_{\vec{r}, d}(\vec{\beta}, \vec{L}) \cap \mathbb{B}_q(Q)] < \infty.$$

We provide explicit construction of such an estimator for integer values of  $p \geq 2$ .

The problem of estimating nonlinear functionals of a probability density has been studied in the literature: we refer to Part I for background and pointers to the relevant literature. Here we restrict ourselves with brief reminder of main definitions and results obtained in Part I.

We begin with the definition of the anisotropic Nikolskii's classes. Let  $(e_1, \dots, e_d)$  denote the canonical basis of  $\mathbb{R}^d$ . For a function  $G : \mathbb{R}^d \rightarrow \mathbb{R}^1$  and real number  $u \in \mathbb{R}$  the first order difference operator with step size  $u$  in direction of the variable  $x_j$  is defined by  $\Delta_{u, j} G(x) = G(x + ue_j) - G(x)$ ,  $j = 1, \dots, d$ . By induction, the  $k$ -th order difference operator is

$$\Delta_{u, j}^k G(x) = \Delta_{u, j} \Delta_{u, j}^{k-1} G(x) = \sum_{l=1}^k (-1)^{l+k} \binom{k}{l} \Delta_{ul, j} G(x).$$

**Definition 1.** For given vectors  $\vec{\beta} = (\beta_1, \dots, \beta_d) \in (0, \infty)^d$ ,  $\vec{r} = (r_1, \dots, r_d) \in [1, \infty]^d$ , and  $\vec{L} = (L_1, \dots, L_d) \in (0, \infty)^d$  a function  $G : \mathbb{R}^d \rightarrow \mathbb{R}^1$  is said to belong to anisotropic Nikolskii's class  $\mathbb{N}_{\vec{r}, d}(\vec{\beta}, \vec{L})$  if  $\|G\|_{r_j} \leq L_j$  for all  $j = 1, \dots, d$  and there exist natural numbers  $k_j > \beta_j$  such that

$$\|\Delta_{u, j}^{k_j} G\|_{r_j} \leq L_j |u|^{\beta_j}, \quad \forall u \in \mathbb{R}, \quad \forall j = 1, \dots, d.$$

Important quantities that are related to Nikolskii's classes and determine asymptotics of the minimax risk are the following:

$$\frac{1}{\beta} := \sum_{j=1}^d \frac{1}{\beta_j}, \quad \frac{1}{\omega} := \sum_{j=1}^d \frac{1}{\beta_j r_j}, \quad L_\beta := \prod_{j=1}^d L_j^{\frac{1}{\beta_j}},$$

$$\tau(s) := 1 - \frac{1}{\omega} + \frac{1}{\beta s}, \quad s \in [1, \infty].$$

It is worth mentioning that  $\tau(\cdot)$  appears in embedding theorems for Nikolskii's spaces; see Section 5.1.3 for details.

In Part I we have established the lower bound (1) on the minimax risk with sequence  $\phi_n$  defined by:

$$\theta = \begin{cases} \frac{1}{\tau(1)}, & \tau(p) \geq 1; \\ \frac{1/p-1/q}{1-1/q-(1-1/p)\tau(q)}, & \tau(p) < 1, \tau(q) < 0; \\ \frac{\tau(p)}{\tau(1)}, & \tau(p) < 1, \tau(q) \geq 0; \end{cases}$$

$$\phi_n = L_{\beta}^{\frac{1-1/p}{\tau(1)}} n^{-\theta^*}, \quad \theta^* = 2^{-1} \wedge \theta; \quad (1.1)$$

see Theorem 1 of Part I. Our goal is to develop a rate-optimal estimator whose risk converges to zero at the rate  $\phi_n$ .

## 2. Estimator construction

Assuming that  $p$  is integer let us first discuss the problem of estimating a closely related functional  $\|f\|_p^p$ .

Let  $K : [-1, 1]^d \rightarrow \mathbb{R}$  be a given function (kernel), and let  $h = (h_1, \dots, h_d) \in (0, 1]^d$  be a given vector (bandwidth). Let

$$K_h(x) := (1/V_h)K(x/h), \quad V_h := \prod_{k=1}^d h_k,$$

where here in all what follows  $y/x$  denotes the coordinate-wise division for  $x, y \in \mathbb{R}^d$ . Define

$$S_h(x) := \int K_h(x-y)f(y)dy, \quad B_h(x) := S_h(x) - f(x). \quad (2.1)$$

Obviously,  $B_h(x)$  is the bias of the kernel density estimator of  $f(x)$  associated with kernel  $K$  and bandwidth  $h$ .

The construction of our estimator for  $\|f\|_p^p$  is based on a simple observation formulated below as Lemma 1.

**Lemma 1.** *For any  $p \in \mathbb{N}^*, p \geq 2$ ,  $f \in \mathfrak{F}$  and  $h \in (0, \infty)^d$  one has*

$$\begin{aligned} \|f\|_p^p &= (1-p) \int S_h^p(x)dx + p \int S_h^{p-1}(x)f(x)dx \\ &\quad + \sum_{j=2}^p \binom{p}{j} (-1)^j \int [S_h(x)]^{p-j} B_h^j(x)dx. \end{aligned} \quad (2.2)$$

The proof is elementary and given in Appendix. It does not require any assumption on the kernel  $K$  except of existence of the integrals on the right hand side of (2.2).

Let  $\hat{T}_{1,h}$  and  $\hat{T}_{2,h}$  be estimators of

$$T_p^{(1)}(f) := \int S_h^p(x)dx, \quad T_p^{(2)}(f) := \int S_h^{p-1}(x)f(x)dx,$$

respectively. Then we estimate  $\|f\|_p^p$  by

$$\hat{T}_h = (1-p)\hat{T}_{1,h} + p\hat{T}_{2,h}.$$

Note that if  $\hat{T}_{1,h}$  and  $\hat{T}_{2,h}$  are unbiased estimators of  $T_p^{(1)}(f)$  and  $T_p^{(2)}(f)$  then, in view of Lemma 1, the bias of  $\hat{T}_h$  in estimation of  $\|f\|_p^p$  is

$$\sum_{j=2}^p \binom{p}{j} (-1)^j \int [S_h(x)]^{p-j} B_h^j(x) dx.$$

The last quantity can be efficiently bounded from above via norms of the bias  $B_h(\cdot)$  and the underlying density  $f$ .

The natural unbiased estimators for  $T_p^{(1)}(f)$  and  $T_p^{(2)}(f)$  are based on the U–statistics:

$$\begin{aligned} \hat{T}_{1,h} &:= \frac{1}{\binom{n}{p}} \sum_{i_1, \dots, i_p} U_h^{(1)}(X_{i_1}, \dots, X_{i_p}), \\ \hat{T}_{2,h} &:= \frac{1}{\binom{n}{p}} \sum_{i_1, \dots, i_p} U_h^{(2)}(X_{i_1}, \dots, X_{i_p}), \end{aligned}$$

where the summations are taken over all possible combinations of  $p$  distinct elements  $\{i_1, \dots, i_p\}$  of  $\{1, \dots, n\}$ , and

$$\begin{aligned} U_h^{(1)}(x_1, \dots, x_p) &:= \int K_h(y - x_1) \cdots K_h(y - x_p) dy, \\ U_h^{(2)}(x_1, \dots, x_p) &:= \frac{1}{p} \sum_{i=1}^p \prod_{\substack{j=1 \\ j \neq i}}^p K_h(x_j - x_i). \end{aligned}$$

It is worth mentioning that not only there is the explicit formula for the bias of  $\hat{T}_h$ , but also its variance admits a rather simple analytical bound. The following result states an upper bound on the variance of  $\hat{T}_h$ ; the proof is given in Appendix.

**Lemma 2.** *Let  $K$  be symmetric bounded function supported on  $[-1, 1]^d$ . Then for all  $p \in \mathbb{N}^*$ ,  $p \geq 2$ ,  $f \in \mathfrak{F}$  and  $h \in (0, \infty)^d$  one has*

$$\text{var}_f [\hat{T}_h] \leq C \|K\|_\infty^{2p} \sum_{k=1}^p \|f\|_{2^{p-k}}^{2p-k} (n^k V_h^{k-1})^{-1},$$

where  $C$  is a constant depending on  $p$  and  $d$  only.

If  $\hat{T}_h$  is a "reasonable" estimator of  $\|f\|_p^p$  then it seems natural to define

$$\hat{N}_h := |\hat{T}_h|^{1/p} \tag{2.3}$$

as an estimator for  $\|f\|_p$ .

### 3. Main result

In this section we demonstrate that the estimator  $\hat{N}_h$  with properly chosen bandwidth  $h$  is a rate–optimal estimator for  $\|f\|_p$ , provided that

$$\vec{r} \in [1, p]^d \cup [p, \infty]^d, \quad p \in \mathbb{N}^*, \quad p \geq 2,$$

i.e.,  $r_j \leq p$  for all  $j = 1, \dots, d$ , or  $r_j \geq p$  for all  $j = 1, \dots, d$ . The proof is based on the derivation of tight uniform upper bounds on the bias and the variance of  $\hat{T}_h$  over anisotropic Nikol'skii's classes. To get such bounds for the bias term we use a special construction of kernel  $K$  [see, e.g. Goldenshluger and Lepski (2014)].

For a given positive integer  $\ell \in \mathbb{N}^*$  and function  $\mathcal{K} : \mathbb{R} \rightarrow \mathbb{R}$  supported on  $[-1, 1]$  define

$$\mathcal{K}_\ell(y) = \sum_{i=1}^{\ell} \binom{\ell}{i} (-1)^{i+1} i^{-1} \mathcal{K}(y/i).$$

**Assumption 1.**  $\mathcal{K}$  is symmetric,  $\int_{\mathbb{R}} \mathcal{K}(y) dy = 1$ ,  $\|\mathcal{K}\|_\infty < \infty$ , and let

$$K(x) = \prod_{j=1}^d \mathcal{K}_\ell(x_j), \quad \forall x \in \mathbb{R}^d.$$

Let us introduce the following notation. For  $j = 1, \dots, d$  let

$$\varkappa_j := \begin{cases} \frac{\beta_j \tau(p)}{\tau(r_j)}, & r_j \leq p, \tau(q) > 0; \\ \beta_j, & \text{otherwise,} \end{cases} \quad p_j := \begin{cases} \frac{2(1-1/p)}{1-1/r_j}, & r_j \geq p; \\ 2, & r_j \leq p, \tau(q) > 0; \\ \frac{2(1/p-1/q)}{1/r_j-1/q}, & r_j < p, \tau(q) \leq 0, \end{cases} \quad (3.1)$$

and let

$$\frac{1}{v} := \sum_{j=1}^d \frac{1}{p_j \varkappa_j}.$$

Define  $\mathfrak{h} = (\mathfrak{h}_1, \dots, \mathfrak{h}_d)$  by

$$\mathfrak{h}_j := L_j^{-1/\varkappa_j} (\mathfrak{L} n^{-1})^{\frac{2}{\varkappa_j p_j [1+2(1-1/p)/v]}}, \quad (3.2)$$

where  $\mathfrak{L}$  is a constant that is completely determined by the class parameters  $\vec{\beta}, \vec{L}, \vec{r}$  and  $p$  (a cumbersome but explicit expression for  $\mathfrak{L}$  is given in Section 5.3.3).

The main result of this paper is given in the next theorem.

**Theorem.** Let  $p \in \mathbb{N}^*, p \geq 2$ ,  $Q > 0$ ,  $\vec{\beta} \in (0, \infty)^d$ ,  $\vec{L} \in (0, \infty)^d$ ,  $\vec{r} \in [1, p]^d \cup [p, \infty]^d$  and  $q \geq 2p - 1$  be fixed. Let Assumption 1 hold with  $\ell > \max_{j=1, \dots, d} \beta_j$ . Let  $\hat{N}_{\mathfrak{h}}$  be the estimator (2.3) associated with bandwidth  $\mathfrak{h}$  defined in (3.2); then

$$\limsup_{n \rightarrow \infty} \phi_n^{-1} \mathcal{R}_n[\hat{N}_{\mathfrak{h}}, \mathbb{N}_{\vec{r}, d}(\vec{\beta}, \vec{L}) \cap \mathbb{B}_q(Q)] < \infty,$$

where  $\phi_n$  is given in (1.1).

**Remark 1.**

(i) Combining the result of the theorem with that of Theorem 1 in Part I we conclude that the suggested estimator  $\hat{N}_{\mathfrak{h}}$  is rate-optimal. Thus, the problem of constructing a rate-optimal estimator is solved completely if  $r_j \leq p$  for all  $j = 1, \dots, d$ , or  $r_j \geq p$  for all  $j = 1, \dots, d$ . The general setting when values of the coordinates of  $\vec{r}$  may be arbitrary with respect to  $p$  remains an open problem. We conjecture that in the general setting a rate-optimal estimator does not belong to the family  $\{\hat{N}_h, h \in \mathbb{R}^d\}$ , and a different estimation procedure has to be developed. Another

intriguing question is whether condition  $q \geq 2p-1$  is necessary in order to guarantee the obtained estimation accuracy?

(ii) It is surprising that the  $\mathbb{L}_p$ -norm can be estimated with parametric rate  $n^{-1/2}$ . To the best of our knowledge this phenomenon has not been observed in the literature. Note that the parametric regime is possible only if  $\tau(q) > 0$ . Indeed,

$$\frac{1/p - 1/q}{1 - 1/q - (1 - 1/p)\tau(q)} \geq \frac{1}{2} \Leftrightarrow \frac{2}{p} - 1 - \frac{1}{q} \geq -\left(1 - \frac{1}{p}\right)\tau(q),$$

and the last inequality is impossible if  $\tau(q) \leq 0$  because  $p \geq 2$ .

(iii) A particularly simple description of the minimax rate of convergence  $\phi_n$  is obtained in the specific case  $p = 2$  and  $q = \infty$ . Here  $\theta^* = (\max\{\tau(1), 2\})^{-1}$  if  $\vec{r} \in [2, \infty]^d$ , and

$$\theta^* = \begin{cases} \frac{1}{2}, & \omega > 1; \\ \frac{1}{1+1/\omega}, & \omega \leq 1, \end{cases}$$

if  $\vec{r} \in [1, 2]^d$ . As we see, the regime corresponding to the exponent  $\tau(p)/\tau(1)$  does not appear if  $p = 2, q = \infty$ .

## 4. Discussion

In this section we compare and contrast our results with other results in the literature. Our discussion is restricted to the problem of estimating the  $\mathbb{L}_p$ -norm with integer  $p$  only. The fundamental differences between our results and other results are mainly due to the following two reasons.

- *Statistical model.* We argue that when estimating the  $\mathbb{L}_p$ -norm estimation in the density model, much better accuracy can be achieved than in the regression/Gaussian white noise model.
- *Domain of observations.* We also explain why estimation accuracy is completely different for compactly supported densities  $f$ , and densities  $f$  supported on the entire space  $\mathbb{R}^d$ .

### 4.1. Norm estimation in density and Gaussian white noise models

To the best of our knowledge, in the context of the Gaussian white noise model, estimation of the  $\mathbb{L}_p$ -norm was studied only in the one-dimensional case,  $d = 1$ . So we focus on the one-dimensional Nikolskii class that is denoted  $\mathbb{N}_{r,1}(\beta, \mathbf{L})$ ,  $\beta > 0$ ,  $\mathbf{L} > 0$ , and  $\mathbf{r} \in [1, \infty]$ .

Let  $W$  be the standard Wiener process and assume that we observe the trajectory of the process

$$X_n(t) = \int_0^t f(u)du + n^{-1/2}W(t), \quad t \in [0, 1]. \quad (4.1)$$

The estimation of  $\|f\|_p$ ,  $1 \leq p < \infty$  in the Gaussian white noise model (4.1) was initiated in Lepski et al. (1999) under assumption that  $f$  belongs to Hölder's functional class, i.e.,  $f \in \mathbb{N}_{\infty,1}(\beta, \mathbf{L})$ . The recent paper Han et al. (2019) considers the same problem for Nikolskii's classes with  $\mathbf{r} \in [p, \infty)$ . In the Gaussian white noise model (4.1) even in the case of integer  $p$  there is a big difference between the minimax rates of convergence obtained for odd and even values of

$p$ . Such phenomenon does not occur in the density model because density is a positive function. Thus in the subsequent discussion of the results in the Gaussian white noise model we restrict ourselves with the case of even  $p$ .

According to the results obtained in Lepski et al. (1999) and Han et al. (2019) in the case of even  $p$ , the minimax rate of convergence (expressed in our notation) is given by  $\varphi_n = n^{-\frac{1}{1+\tau(1)}}$ . Note that in both papers  $\mathbf{r} \geq p$  and, therefore,

$$\varphi_n = n^{-\frac{1}{1+\tau(1)}} \gg n^{-\frac{1}{\tau(1)}} = \phi_n,$$

where  $\phi_n$  is defined in (1.1). Also, since  $\tau(1) > 1$  for any  $\mathbf{r} \neq 1$  we conclude that the parametric regime is impossible in the model (4.1). Before providing the explanations why the results discussed above are so different, let us discuss another result.

Lepski and Spokoiny (1999) studied a hypothesis testing problem in the model (4.1) when the set of alternatives consists of functions from  $\mathbb{N}_{\mathbf{r},1}(\beta, \mathbf{L})$ ,  $\mathbf{r} \in [1, 2)$  separated away from zero in the  $\mathbb{L}_2$ -norm. It is well known that this problem is equivalent to the problem of estimating the  $\mathbb{L}_2$ -norm, and the minimax rate of testing coincides with the rate of estimation. The minimax rate found in Lepski and Spokoiny (1999) is  $\varphi_n = n^{-\frac{1+\tau(2)}{2[1+\tau(1)]}}$  under assumption  $\tau(\infty) > 0$ . Noting that  $\tau(\infty) > 0$  implies  $q = \infty$  and comparing this result with the result obtained in this paper for the case  $p = 2$ ,  $\mathbf{r} \in [1, 2)$  and  $\tau(\infty) > 0$  we conclude

$$\varphi_n = n^{-\frac{1+\tau(2)}{2[1+\tau(1)]}} \gg n^{-1/2} = \phi_n.$$

The last inequality follows from the fact that  $\tau(1) > \tau(2)$ . Thus we again conclude that the parametric regime is impossible in the model (4.1). Another interesting feature should be mentioned: the approach used in Lepski and Spokoiny (1999) is based on rather sophisticated pointwise bandwidth selection scheme while in our estimation procedure the bandwidth is a fixed vector.

The explanation why estimation accuracy in the density model is better than in the Gaussian white noise model (4.1) is rather simple. In fact, the maximal value of the risk in the density model is attained on the densities with small  $\mathbb{L}_p$ -norm. Analysis of our estimation strategy shows that in the density model the variance of the corresponding  $U$ -statistics is proportional to the  $\mathbb{L}_p$ -norm of the density (see Lemma 2 and Proposition 2 for details). The smaller this norm, the smaller the stochastic error of the estimation procedure, and careful bandwidth selection employed in our procedure takes into account this fact and improves considerably the estimation accuracy. In contrast, in the Gaussian white noise model (4.1) the stochastic error of the estimation procedure is independent of the signal  $f$  and of the value of its  $\mathbb{L}_p$ -norm.

#### 4.2. Norm estimation for compactly and non-compactly supported densities

We start with the following simple observation. If a probability density  $f$  is defined on a compact set  $\mathcal{I} \subset \mathbb{R}^d$  then necessarily  $\|f\|_p \geq |\mathcal{I}|^{-1/p} > 0$  for any  $p \in (1, \infty]$ . This fact reduces estimation of  $\|f\|_p$  for compactly supported densities to the problem of estimating  $\|f\|_p^p$ , which is a much smoother functional. Indeed, let  $\tilde{N}$  be an estimator of  $\|f\|_p^p$ . Then, for any density  $f$  and any  $p \in \mathbb{N}^*$ ,  $p \geq 2$

$$\left| |\tilde{N}|^{1/p} - \|f\|_p \right| \leq \|f\|_p^{-(p-1)} \left| |\tilde{N}| - \|f\|_p^p \right| \leq |\mathcal{I}|^{1-1/p} |\tilde{N} - \|f\|_p^p|.$$

The problem of estimating  $\|f\|_p^p$  has been considered by many authors starting from the seminal paper Bickel and Ritov (1988); see, for instance, Birgé and Massart (1995), Kerkycharian and Picard (1996), Laurent (1997), Cai and Low (2005), Tchetgen et al. (2008) among others. It is worth

noting that majority of the papers deals with compactly supported densities belonging to a semi-isotropic functional class, that is  $r_l = \mathbf{r}$  for any  $l = 1, \dots, d$ . Below we discuss these results and present them in a unified way.

Let  $p \in \mathbb{N}^*, p \geq 2$  be fixed and let  $\mathbb{N}_{\mathbf{r},d}(\vec{\beta}, \vec{L}, \mathcal{I})$  denote a semi-isotropic Nikolskii class of densities supported on  $\mathcal{I}$ . Assume that  $\mathbf{r} \geq p$ . Then the minimax rate of convergence in estimating  $\|f\|_p^p$  on this functional class, and, therefore, of  $\|f\|_p$  as well, is given by

$$\varphi_n = n^{-\left(\frac{4}{4+1/\beta} \wedge \frac{1}{2}\right)}.$$

Hence, taking into account that  $\tau(1) = 1 - 1/(\mathbf{r}\beta) + 1/\beta \leq 1 + 1/(2\beta)$  for any  $\mathbf{r} \geq p \geq 2$  we conclude that

$$\varphi_n = n^{-\left(\frac{4}{4+1/\beta} \wedge \frac{1}{2}\right)} \ll n^{-\left(\frac{1}{1+1/2\beta} \wedge \frac{1}{2}\right)} = \phi_n.$$

In particular, the parametric regime in estimating compactly supported densities is possible if and only if  $\beta \geq 1/4$  which is a less restrictive condition than  $\tau(1) \leq 2$ .

## 5. Proofs

### 5.1. Preliminaries

In this section we collect known facts from functional analysis as well as some recent results related to anisotropic Nikolskii's spaces. These results will be used in the subsequent proofs.

#### 5.1.1. Strong maximal operator

For locally integrable function  $f$  on  $\mathbb{R}^d$  the strong Hardy-Littlewood maximal operator of  $f$  is defined by

$$\mathfrak{M}[f](x) = \sup \left\{ \frac{1}{|I|} \int_I f(y) dy : x \in I \right\},$$

where the supremum is taken over all rectangles  $I$  with edges parallel to the coordinate axes and containing point  $x$ ; here  $|\cdot|$  stands for the Lebesgue measure. In view of the Lebesgue differentiation theorem

$$f(x) \leq \mathfrak{M}[f](x) \quad a.e. \tag{5.1}$$

Moreover, if  $f \in \mathbb{L}_p(\mathbb{R}^d)$  then

$$\|\mathfrak{M}[f]\|_p \leq c_0 \|f\|_p, \quad 1 < p \leq \infty \tag{5.2}$$

with constant  $c_0$  depending on  $p$  and  $d$  only; see, e.g., [Guzman \(1975\)](#).

#### 5.1.2. Some useful inequalities

For the ease of reference we recall some well known inequalities that are routinely used in the sequel. These results can be found, e.g., in [Folland \(1999\)](#).

- *Interpolation inequality.* Let  $1 \leq s_0 < s < s_1 \leq \infty$ . If  $f \in \mathbb{L}_{s_0}(\mathbb{R}^d) \cap \mathbb{L}_{s_1}(\mathbb{R}^d)$  then  $f \in \mathbb{L}_s(\mathbb{R}^d)$  and

$$\|f\|_s \leq (\|f\|_{s_0})^{\frac{(s_1-s)s_0}{(s_1-s_0)s}} (\|f\|_{s_1})^{\frac{(s-s_0)s_1}{(s_1-s_0)s}}.$$

- *Young's inequality (general form).* Let  $1 \leq p, q, r \leq \infty$  and  $1 + 1/r = 1/p + 1/q$ . If  $f \in \mathbb{L}_p(\mathbb{R}^d)$  and  $g \in \mathbb{L}_q(\mathbb{R}^d)$  then  $f * g \in \mathbb{L}_r(\mathbb{R}^d)$  and

$$\|f * g\|_r \leq \|f\|_p \|g\|_q.$$



### 5.1.3. Some facts related to Nikolskii's classes

Let  $s > 1$ , and let the functional class  $\mathbb{N}_{\vec{r},d}(\vec{\beta}, \vec{L})$  be fixed. Define for any  $j = 1, \dots, d$

$$\gamma_j(s) := \begin{cases} \beta_j \tau(s) / \tau(r_j), & r_j < s; \\ \beta_j, & r_j \geq s, \end{cases}$$

and let

$$s^* := s \vee \left[ \max_{j=1, \dots, d} r_j \right], \quad \vec{s} := (r_1 \vee s, \dots, r_d \vee s).$$

Recall that the bias  $B_h(\cdot) = B_h(\cdot, f)$  of a kernel density estimator associated with kernel  $K$  is defined in (2.1). The following result has been proved in Goldenshluger and Lepski (2014).

**Lemma 3.** *Let Assumption 1 hold with  $\ell > \max_{j=1, \dots, d} \beta_j$ . Then  $B_h(x)$  admits for any  $x \in \mathbb{R}^d$  the representation  $B_h(x) = \sum_{j=1}^d B_h^{(j)}(x)$  with functions  $B_h^{(j)}$  satisfying the following inequalities. There exist  $C_1 > 0$  and  $C_2 > 0$  independent of  $\vec{L}$  such that for any  $f \in \mathbb{N}_{\vec{r},d}(\vec{\beta}, \vec{L})$  and  $h \in (0, \infty)^d$*

$$\|B_h^{(j)}\|_{r_j} \leq C_1 L_j h_j^{\beta_j}, \quad \forall j = 1, \dots, d. \quad (5.3)$$

Moreover, for any  $s > 1$  satisfying  $\tau(s^*) > 0$  one has

$$\|B_h^{(j)}\|_{s_j} \leq C_2 L_j h_j^{\gamma_j(s)}, \quad \forall j = 1, \dots, d. \quad (5.4)$$

Finally, for any  $r \in [1, \infty]$  and  $R > 0$  there exists  $C_3 > 0$  such that for any  $f \in \mathbb{B}_r(R)$  and any  $h \in (0, \infty)^d$

$$\|B_h^{(j)}\|_r \leq C_3, \quad \forall j = 1, \dots, d. \quad (5.5)$$

The next lemma presents an inequality between different norms of a function belonging to anisotropic Nikolskii's class. The proof of this lemma is given in Appendix.

**Lemma 4.** *Let  $1 \leq p < \infty$  be fixed. For any  $s \in (1, \infty]$  satisfying  $s \geq p \vee [\max_{j=1, \dots, d} r_j]$  and  $\tau(s) > 0$  there exists constant  $C > 0$  independent of  $\vec{L}$  such that for any  $f \in \mathbb{N}_{\vec{r},d}(\vec{\beta}, \vec{L})$*

$$\|f\|_s \leq C (L_{\gamma(s)})^{\frac{(1/p-1/s)\tau(s)}{\tau(p)}} (\|f\|_p)^{\frac{\tau(s)}{\tau(p)}}, \quad L_{\gamma(s)} := \prod_{j=1}^d L_j^{\frac{1}{\gamma_j(s)}}.$$

## 5.2. Reduction to the risk of $\hat{T}_h$

We will analyze the risk of the proposed estimator  $\hat{N}_h$  using two different upper bounds that relate the risk of  $\hat{N}_h$  to the risk of  $\hat{T}_h$  via elementary inequalities:

$$\begin{aligned} \mathbb{E}_f |\hat{N}_h - \|f\|_p|^2 &= \mathbb{E}_f |\hat{T}_h^{1/p} - \|f\|_p|^2 \leq \mathbb{E}_f |\hat{T}_h - \|f\|_p^p|^{2/p} \\ &\leq \left[ \mathbb{E}_f (\hat{T}_h - \|f\|_p^p)^2 \right]^{1/p}; \\ \mathbb{E}_f |\hat{N}_h - \|f\|_p|^2 &= \mathbb{E}_f \left| \frac{\hat{N}_h^p - \|f\|_p^p}{\sum_{i=0}^{p-1} \hat{N}_h^i \|f\|_p^{p-1+i}} \right|^2 \leq \frac{\mathbb{E}_f (\hat{T}_h - \|f\|_p^p)^2}{\|f\|_p^{2p-2}}. \end{aligned}$$

Thus, for any underlying density  $f$  and any  $p \in \mathbb{N}^*$ ,  $p \geq 2$

$$\mathbb{E}_f |\hat{N}_h - \|f\|_p|^2 \leq \min \left\{ \left[ \mathbb{E}_f (\hat{T}_h - \|f\|_p^p)^2 \right]^{1/p}, \frac{\mathbb{E}_f (\hat{T}_h - \|f\|_p^p)^2}{\|f\|_p^{2p-2}} \right\}. \quad (5.6)$$

### 5.3. Proof of Theorem

The proof is divided in three steps. First we establish upper bounds (uniform over  $\mathbb{N}_{\vec{r},d}(\vec{\beta}, \vec{L}) \cap \mathbb{B}_q(Q)$ ) on the bias and the variance of the estimator  $\hat{T}_h$  for any value of  $h$ . Then using the derived upper bounds on the risk of  $\hat{T}_h$  and the risk reduction argument given in Section 5.2, we complete the proof.

#### 5.3.1. Bound for the bias

The next proposition states the upper bound on the bias of the estimator  $\hat{T}_h$ .

**Proposition 1.** *Let  $\vec{r} \in [1, p]^d \cup [p, \infty]^d$ . Then there exists  $C > 0$  independent of  $\vec{L}$  such that for any  $f \in \mathbb{N}_{\vec{r},d}(\vec{\beta}, \vec{L}) \cap \mathbb{B}_q(Q)$  and any  $h \in (0, \infty)^d$*

$$\mathbb{E}_f |\hat{T}_h - \|f\|_p^p| \leq C \|f\|_p^{p-2} \sum_{j=1}^d L_j^{p_j} h_j^{p_j \varkappa_j},$$

where  $p_j$  and  $\varkappa_j$  are defined in (3.1).

**Proof** Our first objective is to prove that

$$|\mathbb{E}_f(\hat{T}_h) - \|f\|_p^p| \leq c_1 \|f\|_p^{p-2} \|B_h\|_p^2, \quad \forall h \in (0, \infty)^d. \quad (5.7)$$

If  $p = 2$  then, by Lemma 1, (5.7) holds as equality with  $c_1 = 1$ . If  $p \geq 3$  then we have in view of Lemma 1

$$|\mathbb{E}_f(\hat{T}_h) - \|f\|_p^p| \leq c_2 \sum_{k=2}^p \int |B_h(x)|^k |S_h(x)|^{p-k} dx. \quad (5.8)$$

Note that for any  $h \in (0, \infty)^d$  in view of (5.1)

$$|B_h(x)| \leq |S_h(x)| + f(x) = \|K\|_1 \mathfrak{M}[f](x) + f(x) \leq c_3 \mathfrak{M}[f](x) \quad a.e.$$

It yields together with (5.8)

$$|\mathbb{E}_f(\hat{T}_h) - \|f\|_p^p| \leq c_4 \int |B_h(x)|^2 (\mathfrak{M}[f](x))^{p-2} dx.$$

Hence, applying Hölder's inequality with exponents  $p/2$  and  $p/(p-2)$  and (5.2) we obtain

$$|\mathbb{E}_f(\hat{T}_h) - \|f\|_p^p| \leq c_4 \|\mathfrak{M}[f]\|_p^{p-2} \|B_h\|_p^2 \leq c_5 \|f\|_p^{p-2} \|B_h\|_p^2.$$

This completes the proof of (5.7). We deduce from (5.7) and Lemma 3

$$|\mathbb{E}_f \hat{T}_h - \|f\|_p^p| \leq c_6 \|f\|_p^{p-2} \sum_{j=1}^d \|B_h^{(j)}\|_p^2, \quad \forall h \in (0, \infty)^d. \quad (5.9)$$

Consider now separately three cases. Let  $\vec{r} \in [p, \infty]^d$ . Then, for any  $j = 1, \dots, d$ , applying the interpolation inequality with  $s_0 = 1$ ,  $s = p$  and  $s_1 = r_j$ , and (5.3) and (5.5) of Lemma 3 we obtain

$$\|B_h^{(j)}\|_p^2 \leq c_7 \|B_h^{(j)}\|_{r_j}^{\frac{2(1-1/p)}{1-1/r_j}} \leq c_8 (L_j h_j^{\beta_j})^{p_j} = c_8 L_j^{p_j} h_j^{\varkappa_j p_j}, \quad (5.10)$$

for any  $f \in \mathbb{N}_{\vec{r},d}(\vec{\beta}, \vec{L})$ .

Let now  $\vec{r} \in [1, p]^d$  and  $\tau(q) > 0$ , and recall that in this case  $\tau(p) > 0$  because  $q > p$ . Applying (5.4) of Lemma 3 with  $s = p$  and  $s_j = p, j = 1, \dots, d$ , we obtain

$$\|B_h^{(j)}\|_p^2 \leq c_9 (L_j h_j^{\gamma_j^{(p)}})^2 = c_9 L_j^{p_j} h_j^{\alpha_j p_j}, \quad (5.11)$$

for any  $f \in \mathbb{N}_{\vec{r},d}(\vec{\beta}, \vec{L})$ .

It remains to consider the case  $\vec{r} \in [1, p]^d$  and  $\tau(q) \leq 0$ . Applying the interpolation inequality with  $s_0 = r_j, s = p$  and  $s_1 = q$  we get for any  $j = 1, \dots, d$  and any  $f \in \mathbb{N}_{\vec{r},d}(\vec{\beta}, \vec{L}) \cap \mathbb{B}_q(Q)$

$$\|B_h^{(j)}\|_p^2 \leq (\|B_h^{(j)}\|_{r_j})^{\frac{2(1/p-1/q)}{1/r_j-1/q}} (\|B_h^{(j)}\|_q)^{\frac{2(1/r_j-1/p)}{1/r_j-1/q}} \leq c_{10} \|B_h^{(j)}\|_{r_j}^{p_j}.$$

To get the last inequality we have used (5.5) of Lemma 3 with  $r = q$  and  $R = Q$ . It yields together with (5.3) of Lemma 3 that

$$\|B_h^{(j)}\|_p^2 \leq c_{11} L_j^{p_j} h_j^{\alpha_j p_j}, \quad (5.12)$$

The required result follows from (5.9), (5.10), (5.11) and (5.12).  $\blacksquare$

### 5.3.2. Bound for the variance

Now we derive the upper bound on the variance of  $\hat{T}_h$ . Recall that  $q \geq 2p - 1$  and define

$$\mathcal{L} := \begin{cases} \max_{k=1, \dots, p} (L_{\gamma(2p-k)})^{\frac{(1-k/p)\tau(2p-k)}{\tau(p)}}, & \tau(q) > 0; \\ 1, & \tau(q) \leq 0, \end{cases}$$

**Proposition 2.** *Let  $\vec{\beta}, \vec{L}, \vec{r}, q > 2p - 1$  and  $Q > 0$  be fixed. Then there exists  $C > 0$  independent of  $\vec{L}$  such that for any  $f \in \mathbb{N}_{\vec{r},d}(\vec{\beta}, \vec{L}) \cap \mathbb{B}_q(Q)$*

$$\text{var}_f [\hat{T}_h] \leq C \mathcal{L} \begin{cases} \sum_{k=1}^p (\|f\|_p)^{\frac{(2p-k)\tau(2p-k)}{\tau(p)}} (n^k V_h^{k-1})^{-1}, & \tau(q) > 0 \\ \sum_{k=1}^p (\|f\|_p)^{\frac{(q-2p+k)p}{q-p}} (n^k V_h^{k-1})^{-1}, & \tau(q) \leq 0. \end{cases}$$

**Proof** In view of Lemma 2

$$\text{var}_f [\hat{T}_h] \leq c_1 \sum_{k=1}^p \|f\|_{2p-k}^{2p-k} (n^k V_h^{k-1})^{-1},$$

Assume first that  $\tau(q) > 0$ ; this also implies that  $\tau(r) > 0$  for any  $r \leq q$ . Applying Lemma 4 with  $s = 2p - k$  we get for any  $k \in \{1, \dots, p\}$

$$\|f\|_{2p-k}^{2p-k} \leq (L_{\gamma(2p-k)})^{\frac{(1-k/p)\tau(2p-k)}{\tau(p)}} (\|f\|_p)^{\frac{(2p-k)\tau(2p-k)}{\tau(p)}} \leq \mathcal{L} (\|f\|_p)^{\frac{(2p-k)\tau(2p-k)}{\tau(p)}}.$$

Now assume that  $\tau(q) \leq 0$ . Applying the interpolation inequality with  $s_0 = p, s = 2p - k$  and  $s_1 = q$  we get for any  $k \in \{1, \dots, p\}$

$$\|f\|_{2p-k}^{2p-k} \leq (\|f\|_p)^{\frac{(q-2p+k)p}{q-p}} (\|f\|_q)^{\frac{(p-k)q}{q-p}} \leq c_2 (\|f\|_p)^{\frac{(q-2p+k)p}{q-p}}.$$

This completes the proof.  $\blacksquare$

## 5.3.3. Completion of the theorem proof

Now we are in a position to complete the proof of the theorem.

Define

$$N := (\mathfrak{L}n^{-1})^{\frac{1}{1+2(1-1/p)/v}}, \quad \mathfrak{L} := \mathcal{L}^{1/p} L_{\varkappa}^{1-1/p}, \quad L_{\varkappa} := \prod_{j=1}^d L_j^{1/\varkappa_j},$$

and recall that  $\mathfrak{h} = (\mathfrak{h}_1, \dots, \mathfrak{h}_d)$  is defined as follows.

$$\mathfrak{h}_j := L_j^{-1/\varkappa_j} (\mathfrak{L}n^{-1})^{\frac{2}{\varkappa_j p_j [1+2(1-1/p)/v]}} = L_j^{-1/\varkappa_j} N^{\frac{2}{\varkappa_j p_j}}.$$

1<sup>0</sup>. Several remarks are in order. Direct computations show that

$$N^{p-2} \sum_{j=1}^d L_j^{p_j} \mathfrak{h}_j^{p_j \varkappa_j} = dN^p; \quad (5.13)$$

$$\mathcal{L}N^p [n^p (V_{\mathfrak{h}})^{p-1}]^{-1} = N^{2p}. \quad (5.14)$$

The following useful equality is deduced from (5.14):

$$[nV_{\mathfrak{h}}]^{-1} = [\mathcal{L}^{-1}nN^p]^{\frac{1}{p-1}} = [\mathcal{L}^{-1}L_{\varkappa}]^{1/p} N^{1-\frac{2}{pv}}. \quad (5.15)$$

Let  $R_n(f)$  denote the quadratic risk of the estimator  $\hat{T}_{\mathfrak{h}}$ , and let

$$\mathbb{F} := \mathbb{N}_{\vec{r},d}(\vec{\beta}, \vec{L}) \cap \mathbb{B}_q(Q) \cap \mathbb{B}_p(N), \quad \bar{\mathbb{F}} := [\mathbb{N}_{\vec{r},d}(\vec{\beta}, \vec{L}) \cap \mathbb{B}_q(Q)] \setminus \mathbb{F}$$

We obviously have

$$\mathcal{R}_n := \mathcal{R}_n[\hat{N}_{\mathfrak{h}}, \mathbb{N}_{\vec{r},d}(\vec{\beta}, \vec{L}) \cap \mathbb{B}_q(Q)] = \max \left( \sup_{f \in \mathbb{F}} \mathcal{R}_n[\hat{N}_{\mathfrak{h}}, f], \sup_{f \in \bar{\mathbb{F}}} \mathcal{R}_n[\hat{N}_{\mathfrak{h}}, f] \right)$$

and, we deduce from (5.6) that

$$\mathcal{R}_n^2 \leq \max \left( \sup_{f \in \bar{\mathbb{F}}} [R_n(f)]^{1/p}, \sup_{f \in \bar{\mathbb{F}}} \|f\|_p^{2-2p} R_n(f) \right)$$

2<sup>0</sup>. In view of Propositions 1 and 2, and by (5.13) and (5.15) we have that for any  $f \in \mathbb{F}$

$$\mathbb{E}_f |\hat{T}_{\mathfrak{h}} - \|f\|_p^p| \leq c_1 N^p; \quad (5.16)$$

$$\text{var}_f [\hat{T}_{\mathfrak{h}}] \leq \frac{C_1(\vec{L})}{n} \begin{cases} \sum_{k=1}^p N^{\frac{(2p-k)\tau(2p-k)}{\tau(p)} + (1-\frac{2}{pv})(k-1)}, & \tau(q) > 0 \\ \sum_{k=1}^p N^{\frac{(q-2p+k)p}{q-p} + (1-\frac{2}{pv})(k-1)}, & \tau(q) \leq 0. \end{cases}$$

Setting  $\theta = 1 - \frac{2}{pv} + b$ ,

$$a = \begin{cases} \frac{(2p-1)\tau(2p-1)}{\tau(p)}, & \tau(q) > 0; \\ \frac{(q-2p+1)p}{q-p}, & \tau(q) \leq 0, \end{cases} \quad b = \begin{cases} -\frac{\tau(\infty)}{\tau(p)}, & \tau(q) > 0; \\ \frac{p}{q-p}, & \tau(q) \leq 0, \end{cases}$$

we obtain for any  $f \in \mathbb{F}$

$$\text{var}_f [\hat{T}_h] \leq \frac{C_2(\vec{L})N^a}{n} \sum_{m=0}^{p-1} N^{(1-\frac{2}{pv}+b)m} = \frac{C_2(\vec{L})N^{a+[\theta(p-1)]_-}}{n} = C_3(\vec{L})N^{\mathfrak{z}},$$

where here and later  $y_- := \min(y, 0)$ , and  $\mathfrak{z} := a + 1 + \frac{2(p-1)}{pv} + [\theta(p-1)]_-$ . It yields together with (5.16)

$$\sup_{f \in \mathbb{F}} [R_n(f)]^{1/p} \leq C_4(\vec{L}) [N^2 + N^{\mathfrak{z}/p}]. \quad (5.17)$$

3<sup>0</sup>. Let us compute the quantity  $1/v$ . If  $\vec{r} \in [p, \infty]^d$  we have

$$\frac{1}{v} = \frac{p}{2(p-1)} \sum_{j=1}^d \frac{1-1/r_j}{\beta_j} = \frac{p(1/\beta - 1/\omega)}{2(p-1)}. \quad (5.18)$$

If  $\vec{r} \in [1, p]^d$  and  $\tau(q) > 0$  we have

$$\frac{1}{v} = \frac{1}{2\tau(p)} \sum_{j=1}^d \frac{\tau(r_j)}{\beta_j} = \frac{1}{2\tau(p)} \left( \frac{\tau(\infty)}{\beta} + \frac{1}{\omega\beta} \right) = \frac{1}{2\beta\tau(p)}. \quad (5.19)$$

Finally, if  $\vec{r} \in [1, p]^d$  and  $\tau(q) \leq 0$  we have

$$\frac{1}{v} = \frac{pq}{2(q-p)} \sum_{j=1}^d \frac{1/r_j - 1/q}{\beta_j} = \frac{pq}{2(q-p)} \left( \frac{1}{\omega} - \frac{1}{\beta q} \right). \quad (5.20)$$

It yields, in particular,

$$(p-1)\theta = \begin{cases} \frac{p-1}{p\beta\tau(p)} - \frac{1}{\beta} + \frac{1}{\omega}, & \vec{r} \in [p, \infty]^d; \\ 0, & \vec{r} \in [1, p]^d, \quad \tau(q) > 0; \\ \frac{(p-1)q\tau(q)}{q-p}, & \vec{r} \in [1, p]^d, \quad \tau(q) \leq 0. \end{cases}$$

Noting that  $\vec{r} \in [p, \infty]^d$  implies  $\tau(p) \geq 1$  we assert

$$\frac{p-1}{p\beta\tau(p)} - \frac{1}{\beta} + \frac{1}{\omega} \leq \frac{p-1}{p\beta} - \frac{1}{\beta} + \frac{1}{\omega} = \frac{1}{\omega} - \frac{1}{p\beta} = 1 - \tau(p) \leq 0.$$

Hence in all cases  $(p-1)\theta \leq 0$  and we conclude that

$$\begin{aligned} \mathfrak{z} &= a + 1 + \frac{2(p-1)}{pv} + \theta(p-1) = a + b(p-1) + p \\ &= p + \frac{(2p-1)\tau(\infty) + 1/\beta}{\tau(p)} - \frac{(p-1)\tau(\infty)}{\tau(p)} = 2p. \end{aligned}$$

Hence we deduce from (5.17) that

$$\sup_{f \in \mathbb{F}} [R_n(f)]^{1/p} \leq C_4(\vec{L})N^2. \quad (5.21)$$

4<sup>0</sup>. Note that for any  $f \in \bar{\mathbb{F}}$  one has in view of Proposition 1 and (5.13)

$$\|f\|_p^{2-2p} (\mathbb{E}_f |\hat{T}_h - \|f\|_p^p|)^2 \leq c_1 \|f\|_p^{-2} \left( \sum_{j=1}^d L_j^{p_j} h_j^{p_j \varkappa_j} \right)^2 \leq c_2 N^2. \quad (5.22)$$

Here we have used that  $N$  is small then  $n$  is large.

(a). If  $\tau(q) > 0$  then for any  $k \in \{1, \dots, p\}$

$$\begin{aligned} (2p-k)\tau(2p-k) - (2p-2)\tau(p) &= -[(k-2)\tau(\infty) + (p-2)/(p\beta)] \\ &= -(k-2)[\tau(\infty) + 1/(p\beta)] - (p-k)/(p\beta). \\ &= -(k-2)\tau(p) - (p-k)/(p\beta). \end{aligned}$$

Since  $\tau(p) > 0$ , for any  $k \in \{2, \dots, p\}$

$$\frac{(2p-k)\tau(2p-k)}{\tau(p)} - (2p-2) = -(k-2) - \frac{(p-k)}{p\beta\tau(p)} \leq 0.$$

Putting  $\alpha = 1 - \frac{(p-1)}{\tau(p)p\beta}$  we deduce from Proposition 2 for any  $f \in \bar{\mathbb{F}}$

$$\begin{aligned} \text{var}_f [\hat{T}_h] &\leq C\mathcal{L}n^{-1} \left[ N^{\alpha-} + \sum_{k=2}^p N^{-(k-2) - \frac{(p-k)}{p\beta\tau(p)}} (nV_h)^{1-k} \right] \\ &\leq C_4(\vec{L})n^{-1} \left[ N^{\alpha-} + \sum_{k=2}^p N^{-(k-2) - \frac{(p-k)}{p\beta\tau(p)} + (1 - \frac{2}{pv})(k-1)} \right] \\ &\leq C_5(\vec{L})n^{-1} [N^{\alpha-} + N^{\gamma+(p-2)\theta}], \end{aligned}$$

where we have put  $\gamma = 1 - \frac{2}{pv} - \frac{p-2}{p\beta\tau(p)}$  and used that  $\theta \leq 0$ . The last bound can be rewritten as

$$\text{var}_f [\hat{T}_h] \leq C_6(\vec{L}) [n^{-1}N^{\alpha-} + N^z], \quad (5.23)$$

where  $z = \gamma + 1 + \frac{2(p-1)}{pv} + (p-2)\theta$ . We have

$$z = p - \frac{p-2}{p\beta\tau(p)} + (p-2)b = p - \frac{p-2}{\tau(p)} [\tau(\infty) + 1/(p\beta)] = 2$$

and (5.23) becomes

$$\text{var}_f [\hat{T}_h] \leq C_7(\vec{L}) [n^{-1}N^\alpha + N^2], \quad \forall f \in \bar{\mathbb{F}}. \quad (5.24)$$

Thus, if  $\tau(q) > 0$  we conclude from (5.21), (5.22) and (5.24)

$$\mathcal{R}_n^2 \leq C_7(\vec{L}) [n^{-1}N^\alpha + N^2]. \quad (5.25)$$

Additionally we remark that if  $\alpha < 0$  then

$$n^{-1}N^{\alpha-} = C_8(\vec{L})N^{\alpha+1+2(1-1/p)/v}.$$

If  $\vec{r} \in [p, \infty]^d$  then  $\tau(p) > 1$ , and we get from (5.18)

$$\alpha + 1 + 2(1-1/p)/v = 2 - \frac{(p-1)}{\tau(p)p\beta} + \frac{1}{\beta} - \frac{1}{\omega} \geq 2 + \frac{1}{p\beta} - \frac{1}{\omega} \geq 2.$$

If  $\vec{r} \in [1, p]^d$  then in view of (5.19)

$$\alpha + 1 + 2(1 - 1/p)/v = 2 - \frac{(p-1)}{\tau(p)p\beta} + \frac{(p-1)}{p\beta\tau(p)} = 2.$$

Thus, (5.25) is equivalent to

$$\mathcal{R}_n^2 \leq C_9(\vec{L}) \begin{cases} n^{-1} \vee N^2, & \alpha \geq 0; \\ N^2, & \alpha < 0. \end{cases} \quad (5.26)$$

The definition of  $N$  implies that

$$N^2 \leq n^{-1} \Leftrightarrow 2(1 - 1/p)/v \leq 1 \Leftrightarrow 1 \geq \begin{cases} 1/\beta - 1/\omega, & \vec{r} \in [p, \infty]^d; \\ \frac{(p-1)}{p\beta\tau(p)}, & \vec{r} \in [1, p]^d. \end{cases}$$

In the case  $\vec{r} \in [1, p]^d$  it yields immediately that  $N^2 \leq n^{-1} \Leftrightarrow a \geq 0$ .

It remains to note that if  $\vec{r} \in [p, \infty]^d$  then

$$\alpha \geq 0 \Leftrightarrow 1 - \frac{(p-1)}{p\beta\tau(p)} = 1 - \frac{1}{\beta} + \frac{1}{\omega} + \left[ \frac{(p-1)}{p\beta} - \frac{(p-1)}{p\beta\tau(p)} \right] + \left[ \frac{1}{p\beta} - \frac{1}{\omega} \right] \geq 0.$$

Since in the considered case  $\tau(p) \geq 1 \Leftrightarrow 1/(p\beta) > 1/\omega$  we assert that

$$1 \geq 1/\beta - 1/\omega \Rightarrow a \geq 0.$$

Thus, we deduce from (5.26)

$$\mathcal{R}_n^2 \leq C_9(\vec{L}) \max \left[ n^{-1}, n^{-\frac{2}{1+2(1-1/p)/v}} \right]$$

and the assertion of the theorem in the case  $\tau(q) > 0$  follows from (5.18) and (5.19).

(b). Let  $\tau(q) \leq 0$ . For any  $k \in \{1, \dots, p\}$  one has

$$\frac{(q - 2p + k)p}{q - p} - 2(p - 1) = 2 - p + \frac{p(k - p)}{q - p} \leq 0$$

and, therefore, we have in view of Proposition 2 and (5.15) for any  $f \in \bar{\mathbb{F}}$

$$\begin{aligned} \text{var}_f[\hat{T}_h] &\leq C\mathcal{L}N^{2-p} \sum_{k=1}^p N^{\frac{(k-p)p}{q-p}} (n^k V_h^{k-1})^{-1} \\ &\leq C\mathcal{L}N^{\frac{2q-p-pq}{q-p}} n^{-1} \sum_{k=1}^p (N^{\frac{p}{q-p}} (nV_h)^{-1})^{k-1} \\ &= C_{10}(\vec{L}) N^{2 - \frac{q(p-1)\tau(q)}{q-p}} \sum_{k=1}^p N^{\frac{q\tau(q)(k-1)}{q-p}} = C_{10}(\vec{L}) N^2. \end{aligned}$$

To get the penultimate equality we have used (5.20). It yields together with (5.22)

$$\sup_{f \in \bar{\mathbb{F}}} \|f\|_p^{2-2p} R_n(f) \leq C_{11}(\vec{L}) N^2.$$

which together with (5.21) leads to

$$\mathcal{R}_n^2 \leq C_{11}(\vec{L}) N^2 = C_{11}(\vec{L}) n^{-\frac{2}{1+2(1-1/p)/v}} = C_{11}(\vec{L}) n^{-\frac{2(1/p-1/q)}{1-1/q-(1-1/p)\tau(q)}}.$$

This completes the theorem proof. ■

## 6. Appendix

**Proof of Lemma 1** Using the identity  $a^p = \sum_{j=0}^p \binom{p}{j} b^{p-j} (a-b)^j$  with  $a = f(x)$  and  $b = S_h(x)$  we obtain for all  $h \in (0, \infty)^d$  and  $x \in \mathbb{R}^d$

$$\begin{aligned} f^p(x) &= \sum_{j=0}^p \binom{p}{j} [S_h(x)]^{p-j} [f(x) - S_h(x)]^j dx \\ &= S_h^p(x)(1-p) + p[S_h(x)]^{p-1}f(x) + \sum_{j=2}^p \binom{p}{j} (-1)^j [S_h(x)]^{p-j} B_h^j(x). \end{aligned}$$

Integrating the last equality, we come to the statement of the lemma. ■

**Proof of Lemma 2** For any  $y \in \mathbb{R}^d$  let

$$I_h(y) := \otimes_{j=1}^d \{x \in \mathbb{R}^d : |x_j - y_j| \leq h_j\}.$$

1<sup>o</sup>. We start with bounding the variance of  $\hat{T}_{1,h}$ . Define

$$\begin{aligned} g_k(x_1, \dots, x_k) &:= \mathbb{E}_f [U_h^{(1)}(x_1, \dots, x_k, X_{k+1}, \dots, X_p)], \quad k = 1, \dots, p-1, \\ g_p(x_1, \dots, x_p) &:= U_h^{(1)}(x_1, \dots, x_p). \end{aligned}$$

Then the variance of  $\hat{T}_{1,h}$  is given by the following well known formula [see, e.g., [Serfling \(1980\)](#)]:

$$\text{var}_f [\hat{T}_{1,h}] = \frac{1}{\binom{n}{p}} \sum_{k=1}^p \binom{p}{k} \binom{n-p}{p-k} \zeta_k, \quad \zeta_k := \text{var}_f [g_k(X_1, \dots, X_k)].$$

We note that  $g_k(x_1, \dots, x_k)$ ,  $k = 1, \dots, p$  are symmetric functions. Observe that for  $k = 1, \dots, p-1$

$$\begin{aligned} |g_k(x_1, \dots, x_k)| &= |\mathbb{E}_f [g(x_1, \dots, x_k, X_{k+1}, \dots, X_p)]| \\ &\leq \int \prod_{i=1}^k |K_h(y - x_i)| \left[ \prod_{i=k+1}^p \int |K_h(y - x_i)| f(x_i) dx_i \right] dy \\ &\leq \|K\|_\infty^{p-k} \int \prod_{i=1}^k |K_h(y - x_i)| (\mathfrak{M}[f](y))^{p-k} dy. \end{aligned}$$



Then

$$\begin{aligned}
\zeta_k &\leq \mathbb{E}_f [g_k(X_1, \dots, X_k)]^2 \\
&\leq \|K\|_\infty^{2(p-k)} \iint (\mathfrak{M}[f](y))^{p-k} (\mathfrak{M}[f](z))^{p-k} \\
&\quad \times \left[ \prod_{i=1}^k \int K_h(y-x_i) K_h(z-x_i) f(x_i) dx_i \right] dy dz \\
&\leq \|K\|_\infty^{2p} V_h^{-k} \iint (\mathfrak{M}[f](y))^{p-k} (\mathfrak{M}[f](z))^{p-k} \mathbf{1}\{y-x \in I_{2h}(0)\} \\
&\quad \times \left[ \prod_{i=1}^k \int K_h(z-x_i) f(x_i) dx_i \right] dy dz \\
&\leq \|K\|_\infty^{2p} V_h^{-k} \iint (\mathfrak{M}[f](y))^{p-k} (\mathfrak{M}[f](z))^p \mathbf{1}\{y-x \in I_{2h}(0)\} dz dy \\
&\leq c_1 \|K\|_\infty^{2p} V_h^{-k+1} \int (\mathfrak{M}[f](y))^{p-k} \mathfrak{M}[\mathfrak{M}^p[f]](y) dy.
\end{aligned}$$

Furthermore, by the Hölder inequality and (5.2)

$$\begin{aligned}
&\int (\mathfrak{M}[f](y))^{p-k} \mathfrak{M}[\mathfrak{M}^p[f]](y) \leq \|\mathfrak{M}[f]\|_{2^{p-k}}^{p-k} \|\mathfrak{M}[\mathfrak{M}^p[f]]\|_{(2^{p-k})/p} \\
&\leq c_2 \|f\|_{2^{p-k}}^{p-k} \|\mathfrak{M}^p[f]\|_{(2^{p-k})/p} = c_2 \|f\|_{2^{p-k}}^{p-k} \|\mathfrak{M}[f]\|_{2^{p-k}}^p \leq c_3 \|f\|_{2^{p-k}}^{2p-k}.
\end{aligned} \tag{6.1}$$

Therefore we obtain for any  $k = 1, \dots, p-1$

$$\zeta_k \leq c_2 \|K\|_\infty^{2p} V_h^{-k+1} \int [f(x)]^{2p-k} dx.$$

In addition,

$$\begin{aligned}
\mathbb{E}_f [g_p(X_1, \dots, X_p)]^2 &= \iint \left[ \prod_{i=1}^p \int K_h(y-x_i) K_h(z-x_i) f(x_i) dx_i \right] dy dz \\
&\leq \frac{\|K\|_\infty^{2p}}{V_h^p} \iint \mathbf{1}\{y-z \in I_{2h}(0)\} (\mathfrak{M}[f](y))^p dy dz \leq \frac{c_3 \|K\|_\infty^{2p}}{V_h^{p-1}} \int [f(x)]^p dx.
\end{aligned}$$

Thus we obtain

$$\text{var}_f [\hat{T}_{1,h}] \leq C_1 \|K\|_\infty^{2p} \sum_{k=1}^p \frac{1}{n^k V_h^{k-1}} \int [f(x)]^{2p-k} dx. \tag{6.2}$$

<sup>20</sup>. Bounding the variance of  $\hat{T}_{2,h}$  goes along the same lines. Define

$$\begin{aligned}
g_k(x_1, \dots, x_k) &:= \mathbb{E}_f [U_h^{(2)}(x_1, \dots, x_k, X_{k+1}, \dots, X_p)], \quad k = 1, \dots, p-1, \\
g_p(x_1, \dots, x_p) &:= U_h^{(2)}(x_1, \dots, x_p).
\end{aligned}$$

We have for  $k = 1, \dots, p-1$

$$\begin{aligned} g_k(x_1, \dots, x_k) &= \mathbb{E}_f[U_h^{(2)}(x_1, \dots, x_k, X_{k+1}, \dots, X_p)] \\ &= \frac{1}{p} \sum_{i=1}^k \prod_{\substack{j=1 \\ j \neq i}}^k K_h(x_j - x_i) \left[ \prod_{j=k+1}^p \int K_h(x_j - x_i) f(x_j) dx_j \right] \\ &\quad + \frac{1}{p} \sum_{i=k+1}^p \int \prod_{j=1}^k K_h(x_j - x_i) \left[ \prod_{\substack{j=k+1 \\ j \neq i}}^p \int K_h(x_j - x_i) f(x_j) dx_j \right] f(x_i) dx_i. \end{aligned}$$

Therefore

$$\begin{aligned} |g_k(x_1, \dots, x_k)| &\leq \frac{\|K\|_\infty^{p-k}}{p} \sum_{i=1}^k (\mathfrak{M}[f](x_i))^{p-k} \prod_{\substack{j=1 \\ j \neq i}}^k |K_h(x_j - x_i)| \\ &\quad + \frac{\|K\|_\infty^{p-k-1}}{p} \sum_{i=k+1}^p \int \prod_{j=1}^k |K_h(x_j - x_i)| (\mathfrak{M}[f](x_i))^{p-k-1} f(x_i) dx_i \\ &:= g_k^{(1)}(x_1, \dots, x_k) + g_k^{(2)}(x_1, \dots, x_k). \end{aligned}$$

For the first term on the right hand side we obtain

$$\begin{aligned} \mathbb{E}_f |g_k^{(1)}(X_1, \dots, X_k)|^2 &\leq c_1 \|K\|_\infty^{2(p-k)} \int (\mathfrak{M}[f](x_i))^{2(p-k)} \\ &\quad \times \left[ \prod_{\substack{j=1 \\ j \neq i}}^k \int K_h^2(x_j - x_i) f(x_j) dx_j \right] f(x_i) dx_i \\ &\leq \frac{c_1 \|K\|_\infty^{2p-2}}{V_h^{k-1}} \int (\mathfrak{M}[f](x))^{2p-k-1} f(x) dx \\ &\leq \frac{c_1 \|K\|_\infty^{2p-2}}{V_h^{k-1}} \int (\mathfrak{M}[f](x))^{2p-k} dx \leq \frac{c_2 \|K\|_\infty^{2p-2}}{V_h^{k-1}} \int [f(x)]^{2p-k} dx. \end{aligned}$$

The expectation of the squared second term is bounded as follows:

$$\begin{aligned} \mathbb{E}_f |g_k^{(2)}(x_1, \dots, x_k)|^2 &\leq c_4 \|K\|_\infty^{2(p-k-1)} \iint \left\{ \prod_{j=1}^k \int K_h(x_j - y) K_h(x_j - z) f(x_j) dx_j \right\} \\ &\quad \times (\mathfrak{M}[f](y))^{p-k} (\mathfrak{M}[f](z))^{p-k} dy dz \\ &\leq \frac{c_3 \|K\|_\infty^{2p-2}}{V_h^{k-1}} \iint \mathbf{1}\{z - y \in I_{2h}(0)\} (\mathfrak{M}[f](y))^p (\mathfrak{M}[f](z))^{p-k} dz dy \\ &\leq \frac{c_4 \|K\|_\infty^{2p-2}}{V_h^{k-1}} \int (\mathfrak{M}[f](z))^{p-k} \mathfrak{M}[\mathfrak{M}^p[f]](z) dz \\ &\leq \frac{c_5 \|K\|_\infty^{2p-2}}{V_h^{k-1}} \int [f(z)]^{2p-k} dz, \end{aligned}$$

where we have used (6.1). Finally,

$$\begin{aligned} \mathbb{E}_f |g_p(X_1, \dots, X_p)|^2 &\leq c_6 \int \left[ \prod_{j=2}^p \int K_h^2(x_j - x) f(x_j) dx_j \right] f(x) dx \\ &\leq \frac{c_6 \|K\|_\infty^{2p-2}}{V_h^{p-1}} \int (\mathfrak{M}[f](x))^{p-1} f(x) dx \\ &\leq \frac{c_7 \|K\|_\infty^{2p-2}}{V_h^{p-1}} \int (\mathfrak{M}[f](x))^p dx \leq \frac{c_7 \|K\|_\infty^{2p-2}}{V_h^{p-1}} \int [f(x)]^p dx. \end{aligned}$$

Thus, we obtain

$$\text{var}_f [\hat{T}_{2,h}] \leq C_2 \|K\|_\infty^{2p-2} \sum_{k=1}^p \frac{1}{n^k V_h^{k-1}} \int [f(x)]^{2p-k} dx. \quad (6.3)$$

The assertion of the lemma follows now from (6.2) and (6.3).  $\blacksquare$

**Proof of Lemma 4** Let  $K$  be a kernel satisfying Assumption 1 with  $\ell \geq \max_{j=1, \dots, d} \beta_j$ . For any  $\eta = (\eta_1, \dots, \eta_d)$ ,  $\eta_j > 0$ ,  $j = 1, \dots, d$  we have in view of Lemma 3

$$\|f\|_s \leq \|B_\eta\|_s + \|S_\eta\|_s \leq \sum_{j=1}^d \|B_\eta^{(j)}\|_s + \|K_\eta * f\|_s.$$

By the general form of Young's inequality with  $1/q = 1 + 1/s - 1/p$

$$\|K_\eta * f\|_s \leq \|f\|_p \|K_\eta\|_q \leq c_1 (V_\eta)^{1/q-1} \|f\|_p = c_1 (V_\eta)^{1/s-1/p} \|f\|_p.$$

Furthermore, it follows from (5.4) of Lemma 3 with  $s^* = s$  and  $s_j = s$  that

$$\sum_{j=1}^d \|B_\eta^{(j)}\|_s \leq c_2 \sum_{j=1}^d L_j \eta_j^{\gamma_j(s)}, \quad \gamma_j(s) = \frac{\beta_j \tau(s)}{\tau(r_j)}.$$

Therefore, for any  $\eta_j > 0$ ,  $j = 1, \dots, d$

$$\|f\|_s \leq c_1 (V_\eta)^{1/s-1/p} \|f\|_p + c_2 \sum_{j=1}^d L_j \eta_j^{\gamma_j(s)}.$$

Putting  $1/\gamma = \sum_{j=1}^d 1/\gamma_j(s)$  and choosing  $\eta_1, \dots, \eta_d$  from the equality

$$(V_\eta)^{1/s-1/p} \|f\|_p = \sum_{j=1}^d L_j \eta_j^{\gamma_j(s)}$$

we come to the following bound

$$\|f\|_s \leq c_3 \left( \prod_{j=1}^d L_j^{\frac{1}{\gamma_j(s)}} \right)^{\frac{1/p-1/s}{1+\frac{1/p-1/s}{\gamma}}} \|f\|_p^{\frac{1}{1+\frac{1/p-1/s}{\gamma}}}.$$

Noting that

$$1 + \frac{1/p - 1/s}{\gamma} = 1 + \left[ \frac{1}{p} - \frac{1}{s} \right] \left[ \frac{\tau(\infty)}{\beta\tau(s)} + \frac{1}{\beta\omega\tau(s)} \right] = 1 + \frac{1/p - 1/s}{\beta\tau(s)} = \frac{\tau(p)}{\tau(s)}$$

we complete the proof. ■

## References

- BICKEL, R.J. and RITOV, Y. (1988). Estimating integrated squared density derivatives: sharp best order of convergence estimates. *Sankhya: The Indian Journal of Statistics* **50**, 381-393.
- BIRGÉ, L. and MASSART, P. (1995). Estimation of the integral functionals of a density. *Ann. Statist.* **23**, 11-29.
- CAI, T.T. and LOW, M.G. (2005). Nonquadratic estimators of a quadratic functional. *Ann. Statist.* **33**, 2930-2956.
- FOLLAND, G. (1999). *Real Analysis: Modern Techniques and Their Applications*. Second edition. John Wiley & Sons, New York.
- GOLDENSHLUGER, A. and LEPSKI, O.V. (2014). On adaptive minimax density estimation on  $\mathbb{R}^d$ . *Probab. Theory Related Fields* **159**, 479-543.
- GOLDENSHLUGER, A. and LEPSKI, O.V. (2020). Minimax estimation of norms of a probability density: I. Lower bounds. *Manuscript*.
- DE GUZMAN, M. (1975). *Differentiation of Integrals in  $R^n$* . Lecture Notes in Mathematics, Vol. 481. Springer-Verlag, Berlin-New York.
- HAN, Y., JIAO, J. and MUKHERJEE, R. (2019). On estimation of  $L_r$ -norms in Gaussian White Noise Models. *arXiv:1710.03863 [math.ST]*.
- KERKYACHARIAN, G. and PICARD, D. (1996). Estimating nonquadratic functionals of a density using Haar wavelets. *Ann. Statist.* **24**, 485-507.
- LEPSKI, O.V., NEMIROVSKI, A. and SPOKOINY, V. (1999). On estimation of the  $\mathbb{L}_r$ -norm of a regression function *Probab. Theory and Related Fields* **113**, 221-253.
- LEPSKI, O.V. and SPOKOINY, V. (1999). Minimax nonparametric hypothesis testing: the case of an inhomogeneous alternative. *Bernoulli* **5**, 333-358.
- LAURENT, B. (1997). Estimation of integral functionals of a density and its derivatives. *Bernoulli* **3**, 181-211.
- NIKOLSKII S.M. (1977). *Priblizhenie Funktsii Mnogikh Peremennykh i Teoremy Vlozheniya*. (in Russian). [*Approximation of functions of several variables and embedding theorems.*] 2nd ed., revised and supplemented. Nauka, Moscow.
- SERFLING, R. J. (1980). *Approximation Theorems of Mathematical Statistics*. John Wiley & Sons, Inc.
- TCHETGEN, E., LI, L., ROBINS, J. and VAN DER VAART, A. (2008). Minimax estimation of the integral of a power of a density. *Stat. Probab. Letters* **78**, 3307-3311.