

Sharp finite-sample concentration of independent variables

Akshay Balsubramani

AKSHAY7@GMAIL.COM

Abstract

We show an extension of Sanov’s theorem on large deviations, controlling the tail probabilities of i.i.d. random variables with matching concentration and anti-concentration bounds. This result has a general scope, applies to samples of any size, and has a short information-theoretic proof using elementary techniques.

Independently and identically distributed (i.i.d.) data are drawn from a distribution P . A central focus in statistics, machine learning, and probability is what can be gleaned about P from a sample of these data – how much data must be sampled for the empirical distribution of the data to concentrate near P ?

1. Setup

We describe distributions P and Q over a measurable space \mathcal{X} using the quantities of information theory. The entropy of a distribution P is $H(P) := \mathbb{E}_{x \sim P} \left[\ln \frac{1}{P(x)} \right]$. The relative entropy of Q with respect to P is $D(Q \parallel P) := \mathbb{E}_{x \sim Q} \left[\ln \frac{Q(x)}{P(x)} \right]$. The cross entropy of P with respect to Q is $H(Q, P) := \mathbb{E}_{x \sim Q} \left[\ln \frac{1}{P(x)} \right]$. The empirical measure of any sample $Z = (z_1, \dots, z_n) \in \mathcal{X}^n$ is $\hat{\mu}_Z$, a distribution over \mathcal{X} .¹ In our case, $Z \sim P^n$ is often an i.i.d. data sample – a random variable; as shorthand, the resulting (random) empirical measure over \mathcal{X} is written \hat{P}_n .

Concentration of the empirical distribution \hat{P}_n to P is addressed by Sanov’s theorem (Sanov, 1957), when we are looking to measure if \hat{P}_n falls in a set \mathcal{A} of distributions over \mathcal{X} . Sanov’s theorem bounds the probability of any such \mathcal{A} if \mathcal{A} is “nice” (convex), in the limit of infinite samples.

Theorem 1 (Sanov’s Theorem (Cover and Thomas, 2006)) *Fix a distribution P over \mathcal{X} and a convex set \mathcal{A} of such distributions. Then*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \ln \Pr \left(\hat{P}_n \in \mathcal{A} \right) = - \min_{Q \in \mathcal{A}} D(Q \parallel P) = -D(P_{\mathcal{A}}^* \parallel P)$$

where $P_{\mathcal{A}}^* \in \arg \min_{Q \in \mathcal{A}} D(Q \parallel P)$ is called the *I-projection* of P on \mathcal{A} .

The right-hand side is typically a finite negative quantity, so this precisely describes the rate of exponential decay of “tail” events which decay exponentially fast in n . For finite n , this is not directly useful, but a variety of concentration bounds (e.g. Chernoff-type

1. It assigns probability to any $E \subseteq \mathcal{X}$ of $\hat{\mu}_Z(E) := \frac{1}{n} \sum_{i=1}^n \mathbf{1}(z_i \in E)$.

bounds) can be derived by approximating the decay for extreme-enough tail events and finite n . These typically apply to special subcases of the setting of Theorem 1 only, and can be loose.

Linear families. For any real-valued functions f_1, f_2, \dots, f_k on \mathcal{X} and $\alpha_1, \alpha_2, \dots, \alpha_k \in \mathbb{R}$, the set $\{P : \mathbb{E}_{a \sim P} [f_i(a)] = \alpha_i, \forall i \in [k]\}$ is called a linear family of distributions. For any $Q \in \mathcal{A}$ if \mathcal{A} is convex,

$$D(Q \parallel P) \geq D(Q \parallel P_{\mathcal{A}}^*) + D(P_{\mathcal{A}}^* \parallel P) \quad (1)$$

with equality if \mathcal{A} is a linear family (Csiszár and Shields, 2004). This is the Pythagorean (in)equality for I-projections.

2. Main results

2.1 A tight characterization with the I-projection

We prove a tight general-purpose information-theoretic characterization of the probability of \mathcal{A} under the empirical measure \hat{P}_n . This is stated in terms of the conditional distribution $\mu_{\mathcal{A}}(y) := \Pr_{Z \sim P^n} (Z = y \mid \hat{\mu}_Z \in \mathcal{A})$, for $y \in \mathcal{X}^n$.

Theorem 2 *Fix a distribution P over \mathcal{X} and a convex set \mathcal{A} of such distributions. For any $\mathcal{B} \subseteq \mathcal{A}$,*

$$\frac{1}{n} \ln \Pr \left(\hat{P}_n \in \mathcal{B} \right) \leq -D(P_{\mathcal{A}}^* \parallel P) - \frac{1}{n} D(\mu_{\mathcal{B}} \parallel P_{\mathcal{A}}^{*n}) \quad (2)$$

If \mathcal{A} is a linear family, equality holds.

It is instructive to consider $\mathcal{B} = \mathcal{A}$. The upper bound of Theorem 2 resembles the asymptotic Sanov bound $-D(P_{\mathcal{A}}^* \parallel P)$ of Theorem 1. However, it is strengthened by a term $\frac{1}{n} D(\mu_{\mathcal{A}} \parallel P_{\mathcal{A}}^{*n})$, which vanishes as $n \rightarrow \infty$, by Sanov's theorem.

2.2 A more general extension

Instead of using I-projections, we can state a slightly more general version of Theorem 2. Define $\mu_{\mathcal{A}}$'s marginal $\omega_{\mathcal{A}}(u) := \Pr_{Z \sim P^n} (Z_i = u \mid \hat{\mu}_Z \in \mathcal{A})$ for $u \in \mathcal{X}$, along any coordinate $i = 1, \dots, n$.²

Theorem 3 *Fix any distribution P over \mathcal{X} and a set \mathcal{A} of such distributions. Then:*

$$\frac{1}{n} \ln \Pr \left(\hat{P}_n \in \mathcal{A} \right) = -D(\omega_{\mathcal{A}} \parallel P) - \frac{1}{n} D(\mu_{\mathcal{A}} \parallel \omega_{\mathcal{A}}^n) \quad (3)$$

Also,

$$1) \frac{1}{n} \ln \Pr \left(\hat{P}_n \in \mathcal{A} \right) \stackrel{(a)}{\leq} -D(\omega_{\mathcal{A}} \parallel P) \stackrel{(b)}{\leq} -D(P_{\mathcal{A}}^* \parallel P)$$

2. These are all the same, by exchangeability.

$$2) \frac{1}{n} \ln \Pr \left(\hat{P}_n \in \mathcal{A} \right) \stackrel{(a)}{\geq} -\mathrm{H}(\omega_{\mathcal{A}}, P) \stackrel{(b)}{\geq} -\max_{Q \in \mathcal{A}} \mathrm{H}(Q, P)$$

The inequalities (1b) and (2b) require \mathcal{A} to be a convex set.

The upper bound $-\mathrm{D}(\omega_{\mathcal{A}} \parallel P) = -\mathrm{H}(\omega_{\mathcal{A}}, P) + \mathrm{H}(\omega_{\mathcal{A}})$ and lower bound $-\mathrm{H}(\omega_{\mathcal{A}}, P)$ (part (2) of the theorem) can be compared:

$$\left| \frac{\text{upper bound}}{\text{lower bound}} - 1 \right| = \left| \frac{-\mathrm{H}(\omega_{\mathcal{A}}, P) + \mathrm{H}(\omega_{\mathcal{A}})}{-\mathrm{H}(\omega_{\mathcal{A}}, P)} - 1 \right| = \frac{\mathrm{H}(\omega_{\mathcal{A}})}{\mathrm{H}(\omega_{\mathcal{A}}, P)} \quad (4)$$

In situations where $\mathrm{H}(\omega_{\mathcal{A}}) \ll \mathrm{H}(\omega_{\mathcal{A}}, P)$ (\mathcal{A} represents a very atypical event under P , so that conditioning on \mathcal{A} significantly changes P), the bounds tend to match well.

For finite n , the slack in the upper bound is $-\frac{1}{n}\mathrm{D}(\mu_{\mathcal{A}} \parallel \omega_{\mathcal{A}}^n)$. This quantity $\mathrm{D}(\mu_{\mathcal{A}} \parallel \omega_{\mathcal{A}}^n)$ is called the *total correlation* between the n marginal distributions of $\mu_{\mathcal{A}}$. The total correlation, a multivariate generalization of the mutual information (Watanabe, 1960), is the information gained by knowing the joint distribution instead of knowing just the marginals. It decreases as n increases here because the marginals are nearly independent, to become negligible as $n \rightarrow \infty$ in Sanov's theorem.

3. Discussion

The proof techniques here are from Csiszár (Csiszár, 1984), underpinned by an information-theoretic identity (Csiszár (1984), Eq. 2.11). This expresses large deviation probabilities of an n -sample random variable in terms of the information geometry of its conditional distribution.

Lemma 4 *Fix a distribution P over \mathcal{X} and a set \mathcal{A} of distributions over \mathcal{X} . Then*

$$-\ln \Pr \left(\hat{P}_n \in \mathcal{A} \right) = \mathrm{D}(\mu_{\mathcal{A}} \parallel P^n) = n\mathrm{D}(\omega_{\mathcal{A}} \parallel P) + \mathrm{D}(\mu_{\mathcal{A}} \parallel \omega_{\mathcal{A}}^n)$$

Sanov's theorem has notably been bounded in a non-asymptotic form in different situations, including for finite discrete spaces \mathcal{X} (Cover and Thomas, 2006) and using vastly more advanced techniques (Dembo and Zeitouni (1998), Ex. 6.2.19). The distinctive additional features of this manuscript's result are the elementary proof and general scope. Extensions and applications are left to future work.

4. Proofs

Proof [Proof of Theorem 2] To prove (2),

$$\begin{aligned} \frac{1}{n} \ln \Pr \left(\hat{P}_n \in \mathcal{B} \right) &\stackrel{(a)}{=} -\mathrm{D}(\omega_{\mathcal{B}} \parallel P) - \frac{1}{n}\mathrm{D}(\mu_{\mathcal{B}} \parallel \omega_{\mathcal{B}}^n) \\ &\stackrel{(b)}{\leq} -\mathrm{D}(P_{\mathcal{A}}^* \parallel P) - \mathrm{D}(\omega_{\mathcal{B}} \parallel P_{\mathcal{A}}^*) - \frac{1}{n}\mathrm{D}(\mu_{\mathcal{B}} \parallel \omega_{\mathcal{B}}^n) \\ &\stackrel{(c)}{=} -\mathrm{D}(P_{\mathcal{A}}^* \parallel P) - \frac{1}{n}\mathrm{D}(\mu_{\mathcal{B}} \parallel P_{\mathcal{A}}^{*n}) \end{aligned}$$

where (a) invokes Lemma 4 with the set \mathcal{B} , (b) uses the Pythagorean inequality for I-projections (1) (since $\omega_{\mathcal{B}} \in \mathcal{B} \subseteq \mathcal{A}$), and (c) uses Lemma 5. \blacksquare

Proof [Proof of Lemma 4] First, note that for all $y \in \mathcal{X}^n$, by definition $\mu_{\mathcal{A}}(y) = \frac{P^n(y)}{\Pr(\hat{P}_n \in \mathcal{A})} \mathbf{1}(\hat{\mu}_y \in \mathcal{A})$. Therefore,

$$\begin{aligned} D(\mu_{\mathcal{A}} \parallel P^n) &= \mathbb{E}_{Z \sim \mu_{\mathcal{A}}} \left[\ln \frac{\mu_{\mathcal{A}}(Z)}{P^n(Z)} \right] = \frac{\mathbb{E}_{Z \sim P^n} \left[\ln \frac{\mu_{\mathcal{A}}(Z)}{P^n(Z)} \mathbf{1}(\hat{\mu}_Z \in \mathcal{A}) \right]}{\Pr(\hat{P}_n \in \mathcal{A})} \\ &= \frac{\mathbb{E}_{Z \sim P^n} \left[-\ln \Pr(\hat{P}_n \in \mathcal{A}) \mathbf{1}(\hat{\mu}_Z \in \mathcal{A}) \right]}{\Pr(\hat{P}_n \in \mathcal{A})} = -\ln \Pr(\hat{P}_n \in \mathcal{A}) \end{aligned} \quad (5)$$

Calling Lemma 5 (with $P, \mu_{\mathcal{A}}, \omega_{\mathcal{A}}$) shows that $D(\mu_{\mathcal{A}} \parallel P^n) - D(\mu_{\mathcal{A}} \parallel \omega_{\mathcal{A}}^n) = nD(\omega_{\mathcal{A}} \parallel P)$. Combining this with (5) gives the result. \blacksquare

Lemma 5 Fix a distribution P over \mathcal{X} and a set \mathcal{A} of distributions over \mathcal{X} , and define $\mu_{\mathcal{A}}, \omega_{\mathcal{A}}$ with respect to these. For any distribution Q over \mathcal{X} ,

$$D(\mu_{\mathcal{A}} \parallel Q^n) - D(\mu_{\mathcal{A}} \parallel \omega_{\mathcal{A}}^n) = nD(\omega_{\mathcal{A}} \parallel Q)$$

Proof [Proof of Lemma 5]

$$\begin{aligned} D(\mu_{\mathcal{A}} \parallel Q^n) - D(\mu_{\mathcal{A}} \parallel \omega_{\mathcal{A}}^n) &= \mathbb{E}_{Z \sim \mu_{\mathcal{A}}} \left[\ln \frac{\omega_{\mathcal{A}}^n(Z)}{Q^n(Z)} \right] = \sum_{i=1}^n \mathbb{E}_{Z \sim \mu_{\mathcal{A}}} \left[\ln \frac{\omega_{\mathcal{A}}(Z_i)}{Q(Z_i)} \right] \\ &\stackrel{(a)}{=} \sum_{i=1}^n \mathbb{E}_{u \sim \omega_{\mathcal{A}}} \left[\ln \frac{\omega_{\mathcal{A}}(u)}{Q(u)} \right] = \sum_{i=1}^n D(\omega_{\mathcal{A}} \parallel Q) = nD(\omega_{\mathcal{A}} \parallel Q) \end{aligned} \quad (6)$$

where (a) is because of the definition of the marginal distribution $\omega_{\mathcal{A}}$. \blacksquare

Proof [Proof of Theorem 3] Lemma 4 is equivalent to Eq. (3).

We get part (1a) by bounding Eq. (3) using the expression $D(\mu_{\mathcal{A}} \parallel \omega_{\mathcal{A}}^n) \geq 0$. Using the convexity of \mathcal{A} , and the fact that $\omega_{\mathcal{A}} \in \mathcal{A}$ (this can be proved³ from the definitions), and further applying $D(\omega_{\mathcal{A}} \parallel P) \geq \min_{Q \in \mathcal{A}} D(Q \parallel P)$ gives part (1b).

3. For any set $F \subseteq \mathcal{X}$, $\omega_{\mathcal{A}}$ is a linear combination of distributions in \mathcal{A} :

$$\begin{aligned} \omega_{\mathcal{A}}(F) &= \int_{Z \in \mathcal{X}^n} \mu_{\mathcal{A}}(Z) \mathbf{1}(Z_i \in F) = \int_{Z \in \mathcal{X}^n} \mu_{\mathcal{A}}(Z) \left(\frac{1}{n} \sum_{i=1}^n \mathbf{1}(Z_i \in F) \right) \\ &= \int_{Z \in \mathcal{X}^n} \mu_{\mathcal{A}}(Z) \hat{\mu}_Z(F) = \int_{\hat{\mu}_Z \in \mathcal{A}} \mu_{\mathcal{A}}(Z) \hat{\mu}_Z(F) \end{aligned}$$

where the last equality is because $\{Z : \hat{\mu}_Z \notin \mathcal{A}\}$ are given measure zero by $\mu_{\mathcal{A}}$. Since \mathcal{A} is convex, $\omega_{\mathcal{A}} \in \mathcal{A}$.

For part (2a), note that

$$\begin{aligned}
D(\mu_{\mathcal{A}} \parallel \omega_{\mathcal{A}}^n) &= H(\mu_{\mathcal{A}}, \omega_{\mathcal{A}}^n) - H(\mu_{\mathcal{A}}) = \mathbb{E}_{Z \sim \mu_{\mathcal{A}}} \left[-\ln \left(\prod_{i=1}^n \omega_{\mathcal{A}}(Z_i) \right) \right] - H(\mu_{\mathcal{A}}) \\
&= \sum_{i=1}^n \mathbb{E}_{Z \sim \mu_{\mathcal{A}}} [-\ln(\omega_{\mathcal{A}}(Z_i))] - H(\mu_{\mathcal{A}}) = \sum_{i=1}^n \mathbb{E}_{u \sim \omega_{\mathcal{A}}} [-\ln(\omega_{\mathcal{A}}(u))] - H(\mu_{\mathcal{A}}) \\
&= nH(\omega_{\mathcal{A}}) - H(\mu_{\mathcal{A}})
\end{aligned}$$

Using this on Eq. (3), $\frac{1}{n} \ln \Pr(\hat{P}_n \in \mathcal{A}) = -(D(\omega_{\mathcal{A}} \parallel P) + H(\omega_{\mathcal{A}})) + \frac{1}{n} H(\mu_{\mathcal{A}}) = -H(\omega_{\mathcal{A}}, P) + \frac{1}{n} H(\mu_{\mathcal{A}}) \geq -H(\omega_{\mathcal{A}}, P)$. This proves part (2a). Using the convexity of \mathcal{A} and $H(\omega_{\mathcal{A}}, P) \leq \max_{Q \in \mathcal{A}} H(Q, P)$ gives part (2b). ■

References

- Thomas M Cover and Joy A Thomas. *Elements of Information Theory*. Wiley Series in Telecommunications and Signal Processing, 2006.
- Imre Csiszár. Sanov property, generalized i -projection and a conditional limit theorem. *The Annals of Probability*, 12(3):768–793, 1984.
- Imre Csiszár and Paul C Shields. Information theory and statistics: a tutorial. *Communications and Information Theory*, 1(4):417–528, 2004.
- Amir Dembo and Ofer Zeitouni. *Large deviations techniques and applications*, volume 38. Springer Applications of Mathematics, Stochastic Modelling, and Applied Probability, 1998.
- Ivan Nikolaevich Sanov. On the probability of large deviations of random magnitudes. *Matematicheskii Sbornik*, 84(1):11–44, 1957.
- Satosi Watanabe. Information theoretical analysis of multivariate correlation. *IBM Journal of research and development*, 4(1):66–82, 1960.