

EXISTENCE OF DISCRETE EIGENVALUES FOR THE DIRICHLET LAPLACIAN IN A TWO-DIMENSIONAL TWISTED STRIP

RAFAEL T. AMORIM* and ALESSANDRA A. VERRI

November 23, 2021

Abstract

We study the spectrum of the Dirichlet Laplacian operator in twisted strips on ruled surfaces in any space dimension. It is shown that a suitable twisted effect can create discrete eigenvalues for the operator. In particular, we also study the case where the twisted effect “grows” at infinity while the width of the strip goes to zero. In this situation, we find an asymptotic behavior for the eigenvalues.

Mathematics Subject Classification (2020). Primary: 81Q10; Secondary: 35P15, 47F05, 58J50.

Keywords. Dirichlet Laplacian, unbounded strips, twisted strips, discrete eigenvalues.

1 Introduction

Let Ω be a strip on a surface in \mathbb{R}^d , $d \geq 2$, and denote by $-\Delta_\Omega^D$ the Dirichlet Laplacian operator in Ω . A problem extensively studied in the literature is to find spectral information of $-\Delta_\Omega^D$. If Ω is a bounded strip, it is known that the spectrum $\sigma(-\Delta_\Omega^D)$ is purely discrete. Otherwise, the existence of discrete eigenvalues is a non-trivial property and it depends on the geometry of Ω [1, 2, 4, 5, 10, 11, 13, 14, 16, 17, 18, 22, 23, 24, 27, 30].

At first, consider the following class of planar strips. Let $r : \mathbb{R} \rightarrow \mathbb{R}^2$ be a C^∞ curve parameterized by its arc-length s and denote by $k(s)$ its curvature at the point $r(s)$. Consider the case where Ω is obtained by moving a bounded segment (c, d) along r with respect to its normal vector field. In the pioneering paper [15], the authors studied the spectral problem of $-\Delta_\Omega^D$. In particular, they proved the existence of discrete spectrum for the operator on the conditions that $k(s) \neq 0$, for some $s \in \mathbb{R}$, and that $k(s)$ decays fast enough at infinity. In addition, the authors assumed some regularity for r and that $d - c$ is small enough. In subsequent studies, the results were improved and generalized [10, 19, 25]. In [19], the authors proved the existence of discrete eigenvalues without the restriction on the width of the strip. In [10] the authors estimated the number of discrete eigenvalues. They proved that this number is finite and bounded for a constant that does not depend on the width of Ω . In [25], the authors proved the existence of discrete spectrum on the conditions that r is of class C^2 , $k(s) \neq 0$, and $k(s) \rightarrow 0$, as $|s| \rightarrow \infty$; we also emphasize that the authors made an overview of some new and old results on spectral properties of $-\Delta_\Omega^D$, including other boundary conditions in $\partial\Omega$.

In [3], the authors introduced a new, two-dimensional model of strip to study the spectral problem of $-\Delta_\Omega^D$. In that work, Ω is a strip in \mathbb{R}^2 which is built by translating a segment oriented in a constant direction along an unbounded curve in the plane. The spectrum of the operator $-\Delta_\Omega^D$ was carefully studied and the model covers different effects: purely essential spectrum, discrete spectrum or a combination of both.

*The first author was supported by CNPq, Brazil (grant 141842/2019-9).

One can consider strips embedded in a Riemannian manifold instead of the Euclidean space. For example, suppose that Ω is a strip of constant width which is defined as a tubular neighborhood of an infinite curve in a two-dimensional Riemannian manifold. This situation was considered in [22]. In particular, the author proved that the discrete spectrum of $-\Delta_{\Omega}^D$ is nonempty for non-negatively curved strips.

Strips on ruled surfaces are also regions of great interest [23, 24, 28, 31]; roughly speaking, a ruled surface is generated by straight lines translating along a curve in the Euclidean space. Consider the case where Ω is a twisted strip composed of segments translated along the straight line in \mathbb{R}^3 with respect to a rotation angle. In [24], assuming that the twisted effect diverges at infinity, the authors studied the spectral problem of $-\Delta_{\Omega}^D$. In particular, they proved that this kind of geometry can create discrete eigenvalues. In a similar situation, in [31] was proved that the discrete spectrum of $-\Delta_{\Omega}^D$ is nonempty since the twisted effect “grows” at infinity while the width of Ω goes to zero.

Some results for strips on ruled surfaces can be extended to higher dimensions. In other words, consider Ω as a twisted and bent two-dimensional strip embedded in \mathbb{R}^d with $d \geq 3$. In [28], the spectrum of the Dirichlet Laplacian $-\Delta_{\Omega}^D$ was carefully studied. The authors also proved that, in the limit when the width of the strip tends to zero, the Dirichlet Laplacian converges in the norm resolvent sense to a one-dimensional Schrödinger operator whose potential depends on the deformations of twisting and bending. An interesting point is that the geometric construction of those strips were performed with a *relatively parallel adapted frame* instead of a Frenet frame. In fact, it is known that a Frenet frame of a curve does not need to exist. However, the authors proved that a relatively parallel adapted frame always exists for an arbitrary curve. The goal of this work is to find additional information about the spectrum of the Dirichlet Laplacian in this situation. In the next paragraphs, we present the formal construction of the strip and more details of the problem.

Let $\Gamma : \mathbb{R} \rightarrow \mathbb{R}^{n+1}$, $n \geq 1$, be a curve of class $C^{1,1}$ parameterized by its arc-length s , i.e., $|\Gamma'(s)| = 1$, for all $s \in \mathbb{R}$. The vector $T(s) := \Gamma'(s)$ denotes its unitary tangent vector at the point $\Gamma(s)$. Note that T is a locally Lipschitz continuous function which is differentiable almost everywhere in \mathbb{R} . The number $k(s) := |\Gamma''(s)|$, $s \in \mathbb{R}$, is called the curvature of Γ at the position $\Gamma(s)$. In Appendix A of [28], the authors proved the existence of a relatively parallel adapted frame for the curve Γ . More exactly, it was shown that there exist n almost-everywhere differentiable normal vector fields N_1, \dots, N_n so that

$$\begin{pmatrix} T \\ N_1 \\ \vdots \\ N_n \end{pmatrix}' = \begin{pmatrix} 0 & k_1 & \cdots & k_n \\ -k_1 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ -k_n & 0 & \cdots & 0 \end{pmatrix} \begin{pmatrix} T \\ N_1 \\ \vdots \\ N_n \end{pmatrix},$$

where $k_j : \mathbb{R} \rightarrow \mathbb{R}$, $j \in \{1, \dots, n\}$, are locally bounded functions. In particular, the vector (k_1, \dots, k_n) satisfies $k_1^2 + \dots + k_n^2 = k^2$.

Now, let $\Theta_j : \mathbb{R} \rightarrow \mathbb{R}$, $j \in \{1, \dots, n\}$, be functions of class $C^{0,1}$ so that

$$\Theta_1^2 + \dots + \Theta_n^2 = 1, \quad (1)$$

and define

$$N_{\Theta} := \Theta_1 N_1 + \dots + \Theta_n N_n. \quad (2)$$

Consider the region

$$\Omega_{\varepsilon} := \{\Gamma(s) + N_{\Theta}(s) \varepsilon t : (s, t) \in \mathbb{R} \times (-1, 1)\}; \quad (3)$$

Ω_{ε} is obtained by translating the segment $(-\varepsilon, \varepsilon)$ along Γ with respect to a normal field (2). Let $-\Delta_{\Omega_{\varepsilon}}^D$ be the Dirichlet Laplacian operator in Ω_{ε} . More precisely, $-\Delta_{\Omega_{\varepsilon}}^D$ is defined as the self-adjoint operator associated with the quadratic form

$$a_{\varepsilon}(\varphi) = \int_{\Omega_{\varepsilon}} |\nabla \varphi|^2 dx, \quad \text{dom } a_{\varepsilon} = H_0^1(\Omega_{\varepsilon}); \quad (4)$$

$\nabla\varphi$ denotes the gradient of φ corresponding to the metric induced by the immersion defined in (13), Section 2. For simplicity, we denote $-\Delta_\varepsilon := -\Delta_{\Omega_\varepsilon}^D$.

Define the function $\Theta : \mathbb{R} \rightarrow \mathbb{R}^n$,

$$\Theta(s) := (\Theta_1(s), \dots, \Theta_n(s)),$$

and write $|\Theta'(s)| := (\Theta_1'^2(s) + \dots + \Theta_n'^2(s))^{1/2}$. Note that $\Theta \in C^{0,1}(\mathbb{R}; \mathbb{R}^n)$. As in [28], Θ is called *twisting vector*; if $\Theta' = 0$, Ω_ε is called *untwisted* or *purely bent* strip; if $k \cdot \Theta := k_1\Theta_1 + \dots + k_n\Theta_n = 0$, Ω_ε is called *unbent* or *purely twisted* strip. Geometrically, interpreting Γ as a curve in Ω_ε , $k \cdot \Theta$ is the *geodesic curvature* of Γ ; $-|\Theta'|^2/(1+|\Theta'|^2\varepsilon^2t^2)^2$ is the *Gauss curvature* of Ω_ε . One can see [28] for a detailed geometric description of Ω_ε .

Let $(\pi/2\varepsilon)^2$ be the first eigenvalue of the Dirichlet Laplacian $-\Delta_{(-\varepsilon,\varepsilon)}^D$ in $L^2(-\varepsilon,\varepsilon)$. In [28], the authors presented a detailed study of the spectrum of $-\Delta_\varepsilon$. In particular, under the conditions $k \cdot \Theta, |\Theta'| \in L^\infty(\mathbb{R})$, $\varepsilon\|k \cdot \Theta\|_{L^\infty(\mathbb{R})} < 1$, and $(k \cdot \Theta)(s) \rightarrow 0$, $|\Theta'(s)| \rightarrow 0$, as $|s| \rightarrow \infty$, they proved that:

- (i) $\sigma_{ess}(-\Delta_\varepsilon) = [(\pi/2\varepsilon)^2, \infty)$;
- (ii) if $\Theta' = 0$ and $k \cdot \Theta \neq 0$, then the discrete spectrum of $-\Delta_\varepsilon$ is nonempty;
- (iii) if $k \cdot \Theta = 0$ and $\Theta' \neq 0$ with $\varepsilon\|\Theta'\|_{L^\infty(\mathbb{R})} \leq \sqrt{2}$, then the discrete spectrum of $-\Delta_\varepsilon$ is empty.

The conditions in (i) and (iii) show that a local twisted effect does not create discrete eigenvalues for $-\Delta_\varepsilon$. Then, a natural question is to know if some appropriated twisted effect can be used to create discrete eigenvalues for the operator. This problem is the subject of this work.

In the first part of this paper, we assume that

$$\Theta \in C^{1,1}(\mathbb{R}; \mathbb{R}^n), \quad (k \cdot \Theta)(s) = 0, \quad \text{and} \quad |\Theta'(s)| = \gamma - \beta(s), \quad \forall s \in \mathbb{R}, \quad (5)$$

where γ is a positive number, and $\beta : \mathbb{R} \rightarrow \mathbb{R}$ is a continuous, almost-everywhere differentiable function with compact support so that $\beta' \in L^\infty(\mathbb{R})$. Geometrically, the second and third conditions in (5) mean that Ω_ε is a purely twisted strip; the properties of the function $\beta(s)$ imply that the twisted effect locally slows down. In this situation, we will find information about the spectrum of $-\Delta_\varepsilon$. In particular, we will show that the conditions in (5) can create discrete eigenvalues for $-\Delta_\varepsilon$. Note that the condition $k \cdot \Theta = 0$ does not necessarily implies that Γ is a straight line.

At first, we study the essential spectrum of $-\Delta_\varepsilon$. The strategy is based on a direct integral decomposition of the operator; see Section 3. In particular, consider the one-dimensional operator

$$D_\varepsilon(0) := -\frac{\partial_t^2}{\varepsilon^2} + Y_\varepsilon^0(t), \quad \text{dom } D_\varepsilon(0) = H_0^1(-1,1) \cap H^2(-1,1), \quad (6)$$

where $\partial_t := \partial/\partial t$,

$$Y_\varepsilon^0(t) := -\frac{3\gamma^4\varepsilon^2t^2}{4h_\varepsilon^4(t)} + \frac{\gamma^2}{2h_\varepsilon^2(t)}, \quad h_\varepsilon(t) := \sqrt{1 + \gamma^2\varepsilon^2t^2}. \quad (7)$$

Since $Y_\varepsilon^0 \in C^\infty[-1,1]$, $D_\varepsilon(0)$ has compact resolvent. Denote by $\lambda_{\varepsilon,1}(0)$ its first eigenvalue and by $u_{\varepsilon,1}^0$ the corresponding orthonormal eigenfunction; $\lambda_{\varepsilon,1}(0)$ is simple. Take $\varepsilon_0 > 0$ so that

$$Y_\varepsilon^0 > 0, \quad \forall \varepsilon \in (0, \varepsilon_0). \quad (8)$$

Thus, for each $\varepsilon \in (0, \varepsilon_0)$, $u_{\varepsilon,1}^0$ can be chosen to be real and positive in $(-1,1)$; see, e.g., Chapter 6 of [12] for more details. The condition in (8) will be useful in the proof of Proposition 3 in Section 3.

Now, we have conditions to state the following result.

Theorem 1. *Assume the conditions (1), (5) and (8). Then,*

$$\sigma_{ess}(-\Delta_\varepsilon) = [\lambda_{\varepsilon,1}(0), \infty).$$

Theorem 1 is a consequence of Propositions 2 and 3 of Section 3. In that same section, Remark 4 gives an asymptotic behavior for the sequence $\{\lambda_{\varepsilon,1}(0)\}_\varepsilon$; in fact, $\varepsilon^2 \lambda_{\varepsilon,1}(0) \rightarrow (\pi/2)^2$, as $\varepsilon \rightarrow 0$.

The next results provide sufficient conditions to the existence of (discrete) eigenvalues for $-\Delta_\varepsilon$ below $\lambda_{\varepsilon,1}(0)$. Let $s_0 > 0$ be so that $\text{supp } \beta \subset [-s_0, s_0]$.

Theorem 2. *Assume the conditions (1), (5) and (8). If $\int_{-s_0}^{s_0} (|\Theta'(s)|^2 - \gamma^2) ds < 0$, then there exists $\varepsilon_1 > 0$ so that, for each $\varepsilon \in (0, \varepsilon_1)$,*

$$\inf \sigma(-\Delta_\varepsilon) < \lambda_{\varepsilon,1}(0),$$

i.e., $\sigma_{dis}(-\Delta_\varepsilon) \neq \emptyset$.

The assumption $\int_{-s_0}^{s_0} (|\Theta'(s)|^2 - \gamma^2) ds < 0$ in Theorem 2 is not a necessary condition to create discrete eigenvalues for $-\Delta_\varepsilon$. For example,

Theorem 3. *Assume the conditions (1), (5) and (8). If $\int_{-s_0}^{s_0} (|\Theta'(s)|^2 - \gamma^2) ds = 0$, then there exists $\varepsilon_2 > 0$ so that, for each $\varepsilon \in (0, \varepsilon_2)$,*

$$\inf \sigma(-\Delta_\varepsilon) < \lambda_{\varepsilon,1}(0),$$

i.e., $\sigma_{dis}(-\Delta_\varepsilon) \neq \emptyset$.

Theorems 2 and 3 show as an appropriated twisted effect can create discrete eigenvalues for $-\Delta_\varepsilon$; its proofs are presented in Section 4.

Now, the goal of the second part of this paper is to discuss a different situation where this phenomenon can also be obtained.

As already commented in the previous paragraphs, in [24, 31], the authors studied the Dirichlet Laplacian restricted in a two-dimensional twisted (straight) strip in \mathbb{R}^3 . We emphasize the paper [24] where the authors considered the region

$$\mathcal{S} := \{(s, t \cos \theta(s), t \sin \theta(s)) : (s, t) \in \mathbb{R} \times (a_1, a_2)\}, \quad (9)$$

where $\theta : \mathbb{R} \rightarrow \mathbb{R}$ is a (locally) Lipschitz continuous function and $a_1, a_2 \in \mathbb{R}$. One of the results of that work shows that the assumptions

$$\lim_{|s| \rightarrow \infty} |\theta'(s)| = \infty, \quad a_1 a_2 \leq 0, \quad (10)$$

ensure that the discrete spectrum of the Dirichlet Laplacian in \mathcal{S} is nonempty. As noted by the authors, an interesting geometric fact is that the conditions in (10) show that the two-dimensional strip \mathcal{S} looks at infinity like a three-dimensional tube of annular cross-section $\{x \in \mathbb{R}^2 : 0 < |x| < r\}$, where $r := \max\{|a_1|, |a_2|\}$. Inspired by [24], in [31] the author considered a twisted strip similar to that in (9). However, in that work, the twisted effect “grows” at infinity while the width of the strip goes to zero. Then, since the strip is thin enough, it was shown that the discrete spectrum of the Dirichlet Laplacian is nonempty and was found an asymptotic behavior for the eigenvalues. In the next paragraphs, we present an adaptation the results of [31] for the model of strips treated in this work.

In this new situation, for $n \geq 1$, assume that $\Theta : \mathbb{R} \rightarrow \mathbb{R}^n$ is a $C^{1,1}$ function, which satisfies (1), and

(I) $\lim_{|s| \rightarrow \infty} |\Theta'(s)| = \infty$;

(II) $|\Theta'|$ is decreasing in $(-\infty, 0)$ and increasing in $(0, \infty)$.

Fix a number $0 < a < 1/3$. For each $\varepsilon > 0$ small enough, let $\nu_1(\varepsilon) < 0$ and $\nu_2(\varepsilon) > 0$ so that

$$|\Theta'(\nu_i(\varepsilon))| = \frac{1}{\varepsilon^a}, \quad i \in \{1, 2\}. \quad (11)$$

Define $I_\varepsilon := (\nu_1(\varepsilon), \nu_2(\varepsilon))$ and let $\Theta_\varepsilon : \mathbb{R} \rightarrow \mathbb{R}^n$ be a function of class $C^{1,1}$ so that

(III) $\Theta_\varepsilon(s) = \Theta(s)$, for all $s \in I_\varepsilon$;

(IV) $|\Theta'_\varepsilon(s)| \leq |\Theta'(s)|$, for all $s \in \mathbb{R}$;

(V) $|\Theta'_\varepsilon(s)|$ is non-increasing in $(-\infty, 0)$ and non-decreasing in $(0, +\infty)$.

Now, assume that the sequence $\{\Theta_\varepsilon\}_\varepsilon$ satisfies

(VI) there exists $K > 0$ so that

$$|\Theta'_\varepsilon(s)| \leq \frac{K}{\varepsilon^a}, \quad |\Theta''_\varepsilon(s)| \leq \frac{K}{\varepsilon^b}, \quad |\Theta'''_\varepsilon(s)| \leq \frac{K}{\varepsilon^c}, \quad \forall s \in \mathbb{R},$$

for all $\varepsilon > 0$ small enough, where b, c are real numbers so that $b < 1$, $a + c < 2$;

(VII) $|\Theta'_\varepsilon(s)| \leq |\Theta'_{\varepsilon'}(s)|$, for all $s \in \mathbb{R}$, if $\varepsilon > \varepsilon'$.

Finally, we use the notation $\Theta_\varepsilon := (\Theta_\varepsilon^1, \dots, \Theta_\varepsilon^n)$, and we assume that

(VIII) $(\Theta_\varepsilon^1)^2 + \dots + (\Theta_\varepsilon^n)^2 = 1$.

For each $\varepsilon > 0$ small enough, let $\Gamma_\varepsilon : \mathbb{R} \rightarrow \mathbb{R}^{n+1}$ be a curve of class $C^{1,1}$ whose curvature k_ε satisfies

(IX) $\text{supp } k_\varepsilon \subset I_\varepsilon$, and $(k_\varepsilon \cdot \Theta_\varepsilon)(s) = 0$, for all $s \in \mathbb{R}$.

The normal vector fields of Γ_ε are denoted by $N_1^\varepsilon, \dots, N_n^\varepsilon$, and $N_{\Theta_\varepsilon}^\varepsilon := \Theta_\varepsilon^1 N_1^\varepsilon + \dots + \Theta_\varepsilon^n N_n^\varepsilon$.

Consider the strip

$$\tilde{\Omega}_\varepsilon := \{\Gamma_\varepsilon(s) + N_{\Theta_\varepsilon}^\varepsilon(s) \varepsilon t : (s, t) \in \mathbb{R} \times (-1, 1)\}.$$

Geometrically, $\tilde{\Omega}_\varepsilon$ is a locally twisting (and locally bending) strip for which the twisted effect “grows” at infinity while its width goes to zero.

Let $-\Delta_{\tilde{\Omega}_\varepsilon}^D$ be the Dirichlet Laplacian operator in $\tilde{\Omega}_\varepsilon$, i.e., the self-adjoint operator associated with the quadratic form

$$\tilde{a}_\varepsilon(\varphi) = \int_{\tilde{\Omega}_\varepsilon} |\nabla \varphi|^2 dx, \quad \text{dom } \tilde{a}_\varepsilon = H_0^1(\tilde{\Omega}_\varepsilon). \quad (12)$$

For simplicity, write $-\tilde{\Delta}_\varepsilon := -\Delta_{\tilde{\Omega}_\varepsilon}^D$.

Remark 1. Let T be a self-adjoint operator that is bounded from below. We denote by $\{\lambda_j(T)\}_{j \in \mathbb{N}}$ the non-decreasing sequence of numbers corresponding to the spectral problem of T according to the Min-Max Principle; see, for example, Theorem XIII.1 in [29].

Let $\Pi(\varepsilon)$ be the infimum of the essential spectrum of $-\tilde{\Delta}_\varepsilon$. Denote by $N(\varepsilon) \leq \infty$ the number of eigenvalues $\lambda_j(-\tilde{\Delta}_\varepsilon)$ of $-\tilde{\Delta}_\varepsilon$ below $\Pi(\varepsilon)$. Let $-\Delta_{\mathbb{R}}$ be the one-dimensional Laplacian and consider the self-adjoint operator $-\Delta_{\mathbb{R}} + (|\Theta'(s)|^2/2)\mathbf{1}$ acting in $L^2(\mathbb{R})$; $\mathbf{1}$ denotes the identity operator in $L^2(\mathbb{R})$. Due to the condition (I), this operator has purely discrete spectrum. In Section 5 of this work, we prove the following result.

Theorem 4. *Assume the conditions (I)-(IX). For $\varepsilon > 0$ small enough, the discrete spectrum $\sigma_{dis}(-\tilde{\Delta}_\varepsilon)$ is nonempty and $N(\varepsilon) \rightarrow \infty$, as $\varepsilon \rightarrow 0$. Furthermore, for each $j \in \mathbb{N}$,*

$$\lim_{\varepsilon \rightarrow 0} \left[\lambda_j(-\tilde{\Delta}_\varepsilon) - \left(\frac{\pi}{2\varepsilon} \right)^2 \right] = \lambda_j \left(-\Delta_{\mathbb{R}} + \frac{|\Theta'(s)|^2}{2} \mathbf{1} \right).$$

Estimating the number of discrete eigenvalues of the Dirichlet Laplacian operator in unbounded strips is also an interesting problem [6, 7, 8, 31]. In the case of curved strips in \mathbb{R}^2 , in [10], the authors found that this number is finite and bounded for a constant that does not depend on the width of the strip. Now, consider the “broken” strip defined by $\{(s, t) \in \mathbb{R}^2 : s \tan \theta < |t| < (s + \pi/\sin \theta) \tan \theta\}$, $\theta \in (0, \pi/2)$. In [6],

the authors ensured the existence of discrete eigenvalues for the Dirichlet Laplacian operator in that region. They proved that there are a finite number of them and this number tends to infinity as $\theta \rightarrow 0$.

As a consequence of Theorem 4, the number of discrete eigenvalues of $-\tilde{\Delta}_\varepsilon$ grows when the width of the strip goes to zero. In [31], a result similar to Theorem 4 was obtained as a consequence of a convergence in the norm resolvent sense of the operators associated to the problem. In this text, we present a simpler proof where the strategy is based on to find upper and lower bounds for the eigenvalues $\lambda_j(-\tilde{\Delta}_\varepsilon)$.

We finish this introduction with some remarks and examples of the model presented.

Remark 2. Theorem 4 shows that the locally bending effect imposed by (IX) does not affect the final result.

Example 1. A simple example of a family of curves $\{\Gamma_\varepsilon\}_\varepsilon$ satisfying (IX) is the following. Let Θ be a function satisfying (I), and let $\Gamma : \mathbb{R} \rightarrow \mathbb{R}^{n+1}$ be a curve of class $C^{2,1}$ whose curvature k has compact support and satisfies $(k \cdot \Theta)(s) = 0$, for all $s \in \mathbb{R}$. Define $\Gamma_\varepsilon := \Gamma$, for all $\varepsilon > 0$ small enough. Then, the condition (IX) is satisfied for all $\varepsilon > 0$ small enough.

Remark 3. In the case $n = 2$, conditions (1) and (I) imply that Θ can be written as $\Theta = (\cos(\psi), \sin(\psi))$, for some function $\psi \in C^{1,1}(\mathbb{R}; \mathbb{R})$ so that $|\psi'(s)| \rightarrow \infty$, as $|s| \rightarrow \infty$. In the case $n = 3$, due to the condition (1), Θ can be written as $\Theta = (\cos(\phi) \cos(\psi), \sin(\phi) \cos(\psi), \sin(\psi))$, for functions $\phi, \psi \in C^{1,1}(\mathbb{R}; \mathbb{R})$. Since $|\Theta'| = \sqrt{(\phi')^2 \cos^2(\psi) + (\psi')^2}$, the condition (I) is satisfied, for example, if $|\psi'(s)| \rightarrow \infty$, as $|s| \rightarrow \infty$, or, if $\psi(s) \in (c_1, c_2] \subset (0, \pi/2)$, for all $s \in \mathbb{R}$, and $|\phi'(s)| \rightarrow \infty$, as $|s| \rightarrow \infty$.

Example 2. Consider $\Theta : \mathbb{R} \rightarrow \mathbb{R}^2$ defined by $\Theta(s) = (\cos(s^2), \sin(s^2))$. Some calculations show that $|\Theta'(s)| = 2|s|$, $s \in \mathbb{R}$; Θ satisfies (I)-(II). Fix a number $0 < a < 1/3$. For each $\varepsilon > 0$ small enough, define the functions $\alpha_\varepsilon : \mathbb{R} \rightarrow \mathbb{R}$,

$$\alpha_\varepsilon(s) := \begin{cases} (-2\varepsilon^a s - 1)/\varepsilon^{2a}, & s \leq -1/\varepsilon^a, \\ s^2, & s \in (-1/\varepsilon^a, 1/\varepsilon^a), \\ (2\varepsilon^a s - 1)/\varepsilon^{2a}, & s \geq 1/\varepsilon^a, \end{cases}$$

and $\Theta_\varepsilon(s) := (\cos(\alpha_\varepsilon(s)), \sin(\alpha_\varepsilon(s)))$, $s \in \mathbb{R}$. Taking $b = 2a$ and $c = 3a$, the conditions in (VI) are satisfied. In fact, the sequence $\{\Theta_\varepsilon\}_\varepsilon$ satisfies the conditions (III)-(VIII).

Example 3. Consider the function $\Theta : \mathbb{R} \rightarrow \mathbb{R}^3$ defined by

$$\Theta(s) = \left(\cos(s^2) \cos\left(\frac{1}{1+s^2}\right), \sin(s^2) \cos\left(\frac{1}{1+s^2}\right), \sin\left(\frac{1}{1+s^2}\right) \right).$$

Note that the condition (1) is satisfied and the function

$$|\Theta'(s)| = \sqrt{4s^2 \cos^2\left(\frac{1}{1+s^2}\right) + \frac{4s^2}{(1+s^2)^4}}, \quad s \in \mathbb{R},$$

satisfies (I)-(II). Fix a number $0 < a < 1/3$. For each $\varepsilon > 0$ small enough, take $\nu(\varepsilon) > 0$ so that $|\Theta'(\pm\nu(\varepsilon))| = 1/\varepsilon^a$; then, there exists $K > 0$ so that $|\nu(\varepsilon)| \leq K/\varepsilon^a$, for all $\varepsilon > 0$ small enough. Let $\alpha_\varepsilon : \mathbb{R} \rightarrow \mathbb{R}$ be the function defined by

$$\alpha_\varepsilon(s) := \begin{cases} -2\nu(\varepsilon)s - \nu(\varepsilon)^2, & s \leq -\nu(\varepsilon), \\ s^2, & s \in (-\nu(\varepsilon), \nu(\varepsilon)), \\ 2\nu(\varepsilon)s - \nu(\varepsilon)^2, & s \geq \nu(\varepsilon); \end{cases}$$

note that $\alpha_\varepsilon \in C^{1,1}(\mathbb{R}; \mathbb{R})$. Now, define

$$\Theta_\varepsilon(s) := \left(\cos(\alpha_\varepsilon(s)) \cos\left(\frac{1}{1+s^2}\right), \sin(\alpha_\varepsilon(s)) \cos\left(\frac{1}{1+s^2}\right), \sin\left(\frac{1}{1+s^2}\right) \right), \quad s \in \mathbb{R}.$$

The sequence $\{\Theta_\varepsilon\}_\varepsilon$ satisfies the conditions (III)-(VII), where $b = 2a$ and $c = 3a$.

This paper is organized as follows. In Section 2 we present some details of the construction of the region in (3) and we make usual change of coordinates in the quadratic form in (4). Sections 3 and 4 are dedicated to study the essential and discrete spectrum of $-\Delta_\varepsilon$, respectively. In Section 5, we study the spectral problem of $-\tilde{\Delta}_\varepsilon$. In Appendix 6 are presented results that are useful in this text. Along the text, K is used to denote different constants.

2 Geometry of the region and change of coordinates

Recall the region Ω_ε given by (3) in the Introduction and the straight strip $\Lambda := \mathbb{R} \times (-1, 1)$. In this section we identify Ω_ε with the Riemannian manifold $(\Lambda, \mathcal{G}_\varepsilon)$, where \mathcal{G}_ε is given by (14), below. After that, we perform usual changes of coordinates in the quadratic form $a_\varepsilon(\varphi)$.

Consider the map

$$\begin{aligned} \mathcal{L}_\varepsilon : \mathbb{R}^2 &\longrightarrow \mathbb{R}^{n+1}, \\ (s, t) &\longmapsto \Gamma(s) + N_\Theta(s)\varepsilon t. \end{aligned} \quad (13)$$

We have $\Omega_\varepsilon = \mathcal{L}_\varepsilon(\Lambda)$. Define the metric $\mathcal{G}_\varepsilon := \nabla \mathcal{L}_\varepsilon \cdot (\nabla \mathcal{L}_\varepsilon)^\perp$. Some calculations show that

$$\mathcal{G}_\varepsilon = \begin{pmatrix} f_\varepsilon^2 & 0 \\ 0 & \varepsilon^2 \end{pmatrix}, \quad f_\varepsilon(s, t) := \sqrt{1 + |\Theta'(s)|^2 \varepsilon^2 t^2}. \quad (14)$$

Let \mathcal{J}_ε be the Jacobian matrix of \mathcal{L}_ε . One has $\det \mathcal{J}_\varepsilon = |\det \mathcal{G}_\varepsilon|^{1/2} = \varepsilon f_\varepsilon > 0$, for all $(s, t) \in \Lambda$. If Γ and Θ are smooth functions, the map $\mathcal{L}_\varepsilon : \Lambda \rightarrow \Omega_\varepsilon$ is a local smooth diffeomorphism. Then, Ω_ε can be identified with the Riemannian manifold $(\Lambda, \mathcal{G}_\varepsilon)$. However, as mentioned in the Introduction of this work, the assumptions about Γ and Θ are more general. At first, one has

Proposition 1. *Assume that $\Gamma \in C^{1,1}(\mathbb{R}; \mathbb{R}^{n+1})$ and $\Theta \in C^{0,1}(\mathbb{R}; \mathbb{R}^n)$. Then, the map $\mathcal{L}_\varepsilon : \Lambda \rightarrow \Omega_\varepsilon$ is a local $C^{0,1}$ -diffeomorphism.*

The proof of this result can be found in [28]. We emphasize that in that work does not necessarily $k \cdot \Theta = 0$, and Proposition 1 is proven under the additional assumptions that $k \cdot \Theta \in L^\infty(\mathbb{R})$ and $\varepsilon \|k \cdot \Theta\|_{L^\infty(\mathbb{R})} < 1$.

In particular, Proposition 1 ensures that \mathcal{L}_ε is a $C^{0,1}$ -immersion. In addition, assume that \mathcal{L}_ε is injective. Thus, the strip Ω_ε does not self-intersect and it is interpreted as an immersed submanifold in \mathbb{R}^{n+1} . As a consequence, $(\Lambda, \mathcal{G}_\varepsilon)$ is an abstract Riemannian manifold.

Now, we perform a change of coordinates so that the quadratic form $a_\varepsilon(\varphi)$ starts to act in the Hilbert space $L^2(\Lambda)$ (with the usual metric of \mathbb{R}^2) instead of $L^2(\Omega_\varepsilon)$. At first, consider the unitary operator

$$\begin{aligned} \mathcal{U}_\varepsilon : L^2(\Omega_\varepsilon) &\longrightarrow L^2(\Lambda, f_\varepsilon ds dt), \\ \psi &\longmapsto \varepsilon^{1/2} \psi \circ \mathcal{L}_\varepsilon, \end{aligned}$$

and define the quadratic form

$$\begin{aligned} b_\varepsilon(\psi) &:= a_\varepsilon(\mathcal{U}_\varepsilon^{-1} \psi) = \int_\Lambda \langle \nabla \psi, \mathcal{G}_\varepsilon^{-1} \nabla \psi \rangle f_\varepsilon ds dt \\ &= \int_\Lambda \frac{|\partial_s \psi|^2}{f_\varepsilon} ds dt + \frac{1}{\varepsilon^2} \int_\Lambda |\partial_t \psi|^2 f_\varepsilon ds dt, \end{aligned}$$

$\text{dom } b_\varepsilon := \mathcal{U}_\varepsilon(H_0^1(\Omega_\varepsilon))$; $\partial_t = \partial/\partial t$ and $\partial_s = \partial/\partial s$. Then, consider

$$\begin{aligned} \mathcal{V}_\varepsilon : L^2(\Lambda) &\longrightarrow L^2(\Lambda, f_\varepsilon ds dt), \\ \psi &\longmapsto f_\varepsilon^{-1/2} \psi, \end{aligned}$$

which is also a unitary operator, and, finally, define

$$c_\varepsilon(\psi) := b_\varepsilon(\mathcal{V}_\varepsilon\psi) = \int_\Lambda \frac{1}{f_\varepsilon^2} \left| \partial_s \psi - \frac{\partial_s f_\varepsilon}{2f_\varepsilon} \psi \right|^2 ds dt + \frac{1}{\varepsilon^2} \int_\Lambda |\partial_t \psi|^2 ds dt + \int_\Lambda V_\varepsilon |\psi|^2 ds dt,$$

where

$$V_\varepsilon(s, t) := -\frac{3|\Theta'(s)|^4 \varepsilon^2 t^2}{4f_\varepsilon^4(s, t)} + \frac{|\Theta'(s)|^2}{2f_\varepsilon^2(s, t)},$$

$\text{dom } c_\varepsilon = \mathcal{V}_\varepsilon^{-1}(\mathcal{U}_\varepsilon(H_0^1(\Omega_\varepsilon)))$. Due to the conditions in (5), one has $\text{dom } c_\varepsilon = H_0^1(\Lambda)$. Denote by C_ε the self-adjoint operator associated with the quadratic form $c_\varepsilon(\psi)$.

3 Essential spectrum

This section is dedicated to prove Theorem 1. Recall the functions h_ε and Y_ε^0 defined by (7) in the Introduction. Consider the quadratic form

$$d_\varepsilon(\psi) := \int_\Lambda \frac{|\partial_s \psi|^2}{h_\varepsilon^2} ds dt + \frac{1}{\varepsilon^2} \int_\Lambda |\partial_t \psi|^2 ds dt + \int_\Lambda Y_\varepsilon^0 |\psi|^2 ds dt, \quad \text{dom } d_\varepsilon := H_0^1(\Lambda).$$

Denote by D_ε the self-adjoint operator associated with $d_\varepsilon(\psi)$. We start with the following result.

Proposition 2. *Assume the conditions in (5). Then, $\sigma_{\text{ess}}(C_\varepsilon) = \sigma_{\text{ess}}(D_\varepsilon)$.*

The proof of this result is presented in Appendix 6. As a consequence, we start to study the essential spectrum of D_ε .

Let $\mathcal{F}_s : L^2(\Lambda) \rightarrow L^2(\Lambda)$ be the Fourier transform with respect to s . \mathcal{F}_s is a unitary operator and, for functions in $L^1(\Lambda)$, its action is given by

$$(\mathcal{F}_s \psi)(p, t) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-ips} \psi(s, t) ds.$$

Then, the operator $\hat{D}_\varepsilon := \mathcal{F}_s D_\varepsilon \mathcal{F}_s^{-1}$ admits the direct integral decomposition

$$\hat{D}_\varepsilon = \int_{\mathbb{R}}^{\oplus} D_\varepsilon(p) dp, \tag{15}$$

where, for each $p \in \mathbb{R}$, $D_\varepsilon(p)$ is the self-adjoint operator associated with the quadratic form

$$d_\varepsilon(p)(v) := \frac{1}{\varepsilon^2} \int_{-1}^1 |\partial_t v|^2 dt + \int_{-1}^1 Y_\varepsilon^p |v|^2 dt, \quad \text{dom } d_\varepsilon(p) = H_0^1(-1, 1),$$

where $Y_\varepsilon^p(t) := p^2/h_\varepsilon^2(t) + Y_\varepsilon^0(t)$. More precisely,

$$D_\varepsilon(p) = -\frac{\partial_t^2}{\varepsilon^2} + Y_\varepsilon^p(t), \quad \text{dom } D_\varepsilon(p) = H^2(-1, 1) \cap H_0^1(-1, 1);$$

the case $p = 0$ corresponds to the operator defined by (6) in the Introduction. Since $Y_\varepsilon^p \in C^\infty[-1, 1]$, each $D_\varepsilon(p)$ has compact resolvent. Denote by $\{\lambda_{\varepsilon, n}(p)\}_{n \in \mathbb{N}}$ the sequence of eigenvalues of $D_\varepsilon(p)$ and by $\{u_{\varepsilon, n}(p)\}_{n \in \mathbb{N}}$ the sequence of the corresponding normalized eigenfunctions, i.e.,

$$D_\varepsilon(p)u_{\varepsilon, n}(p) = \lambda_{\varepsilon, n}(p)u_{\varepsilon, n}(p), \quad n \in \mathbb{N}, \quad p \in \mathbb{R}.$$

Due to the decomposition in (15), we have

$$\sigma(D_\varepsilon) = \cup_{p \in \mathbb{R}} \sigma(D_\varepsilon(p)) = \cup_{n \in \mathbb{N}} \{\lambda_{\varepsilon, n}(p) : p \in \mathbb{R}\}. \tag{16}$$

In particular, denote $u_{\varepsilon, 1}^0 := u_{\varepsilon, 1}(0)$. Due to the condition (8) in the Introduction, $u_{\varepsilon, 1}^0$ can be chosen to be real and positive in $(-1, 1)$. This property will be used in the proof of Proposition 3 below.

Lemma 1. For each $n \in \mathbb{N}$, $\lambda_{\varepsilon,n}(\cdot)$ is a real analytic function in p and

$$\lim_{p \rightarrow \pm\infty} \lambda_{\varepsilon,n}(p) = \infty.$$

Proof. At first, note that $\text{dom } D_\varepsilon(p) = \text{dom } D_\varepsilon(0)$, for all $p \in \mathbb{R}$. One can write $D_\varepsilon(p) = D_\varepsilon(0) + p^2/h_\varepsilon^2$. Since $h_\varepsilon \geq 1$, it holds the estimate $\|(p^2/h_\varepsilon^2)v\| \leq p^2\|v\|$, for all $v \in \text{dom } D_\varepsilon(0)$, for all $p \in \mathbb{R}$. Then, p^2/h_ε^2 is $D_\varepsilon(0)$ -bounded with relative bound zero. Consequently, $\{D_\varepsilon(p) : p \in \mathbb{R}\}$ is a type A analytic family. By Theorem 3.9 in [21], $\lambda_{\varepsilon,n}(\cdot)$ is a real analytic function in p .

Now, for each $v \in \text{dom } d_\varepsilon(p)$, we have

$$d_\varepsilon(p)(v) = \frac{1}{\varepsilon^2} \int_{-1}^1 |\partial_t v|^2 dt + \int_{-1}^1 Y_\varepsilon^p |v|^2 dt \geq \left(\frac{p^2}{1 + \gamma^2 \varepsilon^2} - \varepsilon^2 \gamma^4 \right) \int_{-1}^1 |v|^2 dt.$$

As a consequence, for each $n \in \mathbb{N}$,

$$\lambda_{\varepsilon,n}(p) = d_\varepsilon(p)(u_{\varepsilon,n}(p)) \geq \frac{p^2}{1 + \gamma^2 \varepsilon^2} - \varepsilon^2 \gamma^4, \quad p \in \mathbb{R}.$$

Thus, we obtain the punctual limit $\lambda_{\varepsilon,n}(p) \rightarrow \infty$, as $p \rightarrow \pm\infty$. \square

Proposition 3. One has $\sigma(D_\varepsilon) = [\lambda_{\varepsilon,1}(0), \infty)$.

Proof. Since $\lambda_{\varepsilon,1}(p)$ is a real analytic function in p , by (16), we have $[\lambda_{\varepsilon,1}(0), \infty) \subset \sigma(D_\varepsilon)$. Now, we need to show that

$$(-\infty, \lambda_{\varepsilon,1}(0)) \cap \sigma(D_\varepsilon) = \emptyset. \quad (17)$$

Take $\psi \in C_0^\infty(\Lambda)$. Since $u_{\varepsilon,1}^0$ is positive, we can write $\psi(s, t) = \phi(s, t)u_{\varepsilon,1}^0(t)$, with $\phi \in C_0^\infty(\Lambda)$. Some calculations show that

$$\begin{aligned} d_\varepsilon(\psi) - \lambda_{\varepsilon,1}(0) \int_\Lambda |\psi|^2 ds dt &= \int_\Lambda \left(\frac{|\partial_s \phi|^2}{h_\varepsilon^2} + \frac{|\partial_t \phi|^2}{\varepsilon^2} \right) |u_{\varepsilon,1}^0|^2 ds dt + \frac{2}{\varepsilon^2} \mathcal{R} \int_\Lambda \bar{\phi} \partial_t \phi u_{\varepsilon,1}^0 \partial_t u_{\varepsilon,1}^0 ds dt \\ &\quad + \frac{1}{\varepsilon^2} \int_\Lambda |\phi|^2 |\partial_t u_{\varepsilon,1}^0|^2 ds dt + \int_\Lambda |\phi|^2 (Y_\varepsilon^0 |u_{\varepsilon,1}^0|^2 - \lambda_{\varepsilon,1}(0) |u_{\varepsilon,1}^0|^2) ds dt. \end{aligned}$$

An integration by parts, and since $D_\varepsilon(0)u_{\varepsilon,1}^0 = \lambda_{\varepsilon,1}(0)u_{\varepsilon,1}^0$, one has

$$d_\varepsilon(\psi) - \lambda_{\varepsilon,1}(0) \int_\Lambda |\psi|^2 ds dt = \int_\Lambda \left(\frac{|\partial_s \phi|^2}{h_\varepsilon^2} + \frac{|\partial_t \phi|^2}{\varepsilon^2} \right) |u_{\varepsilon,1}^0|^2 ds dt \geq 0.$$

Then, we obtain (17). \square

Proof of Theorem 1. Apply Propositions 2 and 3. \square

Remark 4. For the sequence $\{\lambda_{\varepsilon,1}(0)\}_\varepsilon$, we have the estimate: there exists $K > 0$ so that

$$\int_{-1}^1 |\partial_t v|^2 dt \leq \varepsilon^2 d_\varepsilon(0)(v) = \int_{-1}^1 (|\partial_t v|^2 + \varepsilon^2 Y_\varepsilon^0 |v|^2) dt \leq \int_{-1}^1 |\partial_t v|^2 dt + K \varepsilon^2 \int_{-1}^1 |v|^2 dt,$$

for all $v \in H_0^1(-1, 1)$, for all $\varepsilon > 0$ small enough. Consequently,

$$\left(\frac{\pi}{2} \right)^2 \leq \varepsilon^2 \lambda_{\varepsilon,1}(0) \leq \left(\frac{\pi}{2} \right)^2 + O(\varepsilon^2).$$

4 Discrete spectrum

Based in [19], the main idea in the proofs of Theorems 2 and 3 is to find a function $\psi \in \text{dom } c_\varepsilon$ so that $(c_\varepsilon(\psi) - \lambda_{\varepsilon,1}(0)\|\psi\|^2)/\|\psi\|^2 < 0$.

Proof of Theorem 2. Take $\delta > 0$. Define $\psi_\delta(s, t) := \phi(s)u_{\varepsilon,1}^0(t)$, where

$$\phi(s) := \begin{cases} e^{\delta(s+s_0)}, & s \leq -s_0, \\ 1, & -s_0 \leq s \leq s_0, \\ e^{-\delta(s-s_0)}, & s \geq s_0. \end{cases}$$

Note that $\psi_\delta \in \text{dom } c_\varepsilon$. Some calculations show that

$$\begin{aligned} c_\varepsilon(\psi_\delta) - \lambda_{\varepsilon,1}(0) \int_\Lambda |\psi_\delta|^2 ds dt &= \int_\Lambda \frac{|\partial_s \psi_\delta|^2}{f_\varepsilon^2} ds dt + \frac{1}{4} \int_\Lambda \frac{(\partial_s f_\varepsilon)^2}{f_\varepsilon^4} |\psi_\delta|^2 ds dt - \mathcal{R} \int_\Lambda \frac{\partial_s f_\varepsilon}{f_\varepsilon^3} \overline{\psi_\delta} \partial_s \psi_\delta ds dt \\ &\quad + \frac{1}{\varepsilon^2} \int_\Lambda |\partial_t \psi_\delta|^2 ds dt - \lambda_{\varepsilon,1}(0) \int_\Lambda |\psi_\delta|^2 ds dt + \int_\Lambda V_\varepsilon |\psi_\delta|^2 ds dt \\ &= \int_\Lambda \frac{|\partial_s \psi_\delta|^2}{f_\varepsilon^2} ds dt + \frac{1}{4} \int_\Lambda \frac{(\partial_s f_\varepsilon)^2}{f_\varepsilon^4} |\psi_\delta|^2 ds dt - \mathcal{R} \int_\Lambda \frac{\partial_s f_\varepsilon}{f_\varepsilon^3} \overline{\psi_\delta} \partial_s \psi_\delta ds dt \\ &\quad + \int_\Lambda (V_\varepsilon - Y_\varepsilon^0) |\psi_\delta|^2 ds dt. \end{aligned}$$

Now, note that $f_\varepsilon \rightarrow 1$, $\partial_s f_\varepsilon \rightarrow 0$, $(V_\varepsilon - Y_\varepsilon^0) \rightarrow (|\Theta'|^2 - \gamma^2)/2$, uniformly, as $\varepsilon \rightarrow 0$. Then,

$$c_\varepsilon(\psi_\delta) - \lambda_{\varepsilon,1}(0)\|\psi_\delta\|^2 \rightarrow \delta + \frac{1}{2} \int_{-s_0}^{s_0} (|\Theta'(s)|^2 - \gamma^2) ds,$$

as $\varepsilon \rightarrow 0$. Since $\|\psi_\delta\|^2 = 2s_0 + \delta^{-1}$, one has

$$\frac{c_\varepsilon(\psi_\delta) - \lambda_{\varepsilon,1}(0)\|\psi_\delta\|^2}{\|\psi_\delta\|^2} \rightarrow O(\delta^2) + \frac{\delta}{2} \int_{-s_0}^{s_0} (|\Theta'(s)|^2 - \gamma^2) ds, \quad (18)$$

as $\varepsilon \rightarrow 0$.

Since $\int_{-s_0}^{s_0} (|\Theta'(s)|^2 - \gamma^2) ds < 0$, we can choose δ small enough so that the limit in (18) is negative. Consequently, there exists $\varepsilon_1 > 0$ so that

$$\frac{c_\varepsilon(\psi_\delta) - \lambda_{\varepsilon,1}(0)\|\psi_\delta\|^2}{\|\psi_\delta\|^2} < 0,$$

for all $\varepsilon \in (0, \varepsilon_1)$. □

Proof of Theorem 3. Given $\delta > 0$ and $\eta > 0$, define $\psi_{\delta,\eta}(s, t) := \phi_\eta(s)u_{\varepsilon,1}^0(t)$, where

$$\phi_\eta(s) := \begin{cases} e^{\delta(s+s_0)}, & s \leq -s_0, \\ 1 + \eta(\gamma - |\Theta'(s)|), & -s_0 \leq s \leq s_0, \\ e^{-\delta(s-s_0)}, & s \geq s_0. \end{cases}$$

Note that $\psi_{\delta,\eta} \in \text{dom } c_\varepsilon$. Similarly as in the proof of Theorem 2, we can show that

$$c_\varepsilon(\psi_{\delta,\eta}) - \lambda_{\varepsilon,1}(0)\|\psi_{\delta,\eta}\|^2 \rightarrow \delta + O(\eta^2) - \eta \int_{-s_0}^{s_0} (|\Theta'(s)| - \gamma)^2 (|\Theta'(s)| + \gamma) ds,$$

as $\varepsilon \rightarrow 0$. Since $\|\psi_{\delta,\eta}\|^2 = 2s_0 + \delta^{-1} + O(\eta^2)$, one has

$$\frac{c_\varepsilon(\psi_{\delta,\eta}) - \lambda_{\varepsilon,1}(0)\|\psi_{\delta,\eta}\|^2}{\|\psi_{\delta,\eta}\|^2} \rightarrow O(\delta^2) + \delta O(\eta^2) - \delta\eta \int_{-s_0}^{s_0} (|\Theta'(s)| - \gamma)^2 (|\Theta'(s)| + \gamma) ds, \quad (19)$$

as $\varepsilon \rightarrow 0$. Taking $\eta = \sqrt{\delta}$, again we can choose δ small enough so that the limit in (19) is negative. Then, there exists $\varepsilon_2 > 0$ so that

$$\frac{c_\varepsilon(\psi_{\delta,\eta}) - \lambda_{\varepsilon,1}(0)\|\psi_{\delta,\eta}\|^2}{\|\psi_{\delta,\eta}\|^2} < 0,$$

for all $\varepsilon \in (0, \varepsilon_2)$. □

5 Thin strips

In this section we present the proof of Theorem 4 stated in the Introduction. The strategy will be to establish upper and lower bounds for the eigenvalues $\lambda_j(-\tilde{\Delta}_\varepsilon)$. Recall $-\tilde{\Delta}_\varepsilon$ is the self-adjoint operator associated with the quadratic form $\tilde{a}_\varepsilon(\varphi)$; see (12). Then, the analysis will be based on estimates for $\tilde{a}_\varepsilon(\varphi)$.

Define

$$\tilde{f}_\varepsilon(s, t) := \sqrt{1 + |\Theta'_\varepsilon(s)|^2 \varepsilon^2 t^2},$$

and consider the Hilbert space $\mathcal{H}_\varepsilon := L^2(\Lambda, \tilde{f}_\varepsilon ds dt)$; the norm in this space is denoted by $\|\cdot\|_{\mathcal{H}_\varepsilon}$. Performing a change of coordinates similar to that in Section 2, $\tilde{a}_\varepsilon(\varphi)$ becomes

$$\tilde{b}_\varepsilon(\psi) := \int_\Lambda \frac{|\partial_s \psi|^2}{\tilde{f}_\varepsilon} ds dt + \frac{1}{\varepsilon^2} \int_\Lambda |\partial_t \psi|^2 \tilde{f}_\varepsilon ds dt,$$

$\text{dom } \tilde{b}_\varepsilon = H_0^1(\Lambda) \subset \mathcal{H}_\varepsilon$.

Upper bound. Denote by $\chi_1(t) := \cos(\pi t/2)$ the first eigenfunction of the Dirichlet Laplacian $-\Delta_{(-1,1)}^D$ in $L^2(-1, 1)$; $(\pi/2)^2$ is the eigenvalue associated with χ_1 . Consider the subspace

$$\mathcal{A}_\varepsilon := \{\varphi_w := w(s)\chi_1(t)(\tilde{f}_\varepsilon(s, t))^{-1/2} : w \in H^1(\mathbb{R})\}$$

of the Hilbert space \mathcal{H}_ε . The identification $w \mapsto \varphi_w$, $w \in H^1(\mathbb{R})$, motivates the definition of the one-dimensional quadratic form

$$m_\varepsilon(w) := \tilde{b}_\varepsilon(\varphi_w) - (\pi/2\varepsilon)^2 \|\varphi_w\|_{\mathcal{H}_\varepsilon}^2,$$

$\text{dom } m_\varepsilon := H^1(\mathbb{R})$. Denote by M_ε the self-adjoint operator associated with $m_\varepsilon(w)$. In particular, for each $j \in \mathbb{N}$,

$$\lambda_j(-\tilde{\Delta}_\varepsilon) - \left(\frac{\pi}{2\varepsilon}\right)^2 \leq \lambda_j(M_\varepsilon). \quad (20)$$

We are going to get upper bounds for the values $\lambda_j(M_\varepsilon)$.

Define the function

$$W_\varepsilon(s, t) := -\frac{7|(\Theta'_\varepsilon \cdot \Theta''_\varepsilon)(s)|^2 \varepsilon^4 t^4}{4\tilde{f}_\varepsilon^6(s, t)} + \frac{(2|\Theta''_\varepsilon(s)|^2 + 2(\Theta'_\varepsilon \cdot \Theta'''_\varepsilon)(s) - 3|\Theta'_\varepsilon(s)|^4) \varepsilon^2 t^2}{4\tilde{f}_\varepsilon^4(s, t)} + \frac{|\Theta'_\varepsilon(s)|^2}{2\tilde{f}_\varepsilon^2(s, t)}.$$

Recall that we have the condition (V) in the Introduction, we get the estimates

$$\|(1/\tilde{f}_\varepsilon^2) - 1\|_{L^\infty(\Lambda)} \leq \varepsilon^{2-2a}, \quad \|W_\varepsilon - |\Theta'_\varepsilon|^2/2\|_{L^\infty(\Lambda)} \leq K(\varepsilon^{4-2(a+b)} + \varepsilon^{2-2b} + \varepsilon^{2-(a+c)} + \varepsilon^{2-4a}),$$

for some $K > 0$, for all $\varepsilon > 0$ small enough. Finally, some calculations show that

$$\begin{aligned} m_\varepsilon(w) &= \int_\Lambda \frac{|w'\chi_1|^2}{\tilde{f}_\varepsilon^2} dsdt + \int_\Lambda W_\varepsilon |w\chi_1|^2 dsdt \\ &\leq \int_{\mathbb{R}} \left(|w'|^2 + \frac{|\Theta'_\varepsilon(s)|^2}{2} |w|^2 \right) ds + O(\varepsilon^d) \int_{\mathbb{R}} |w|^2 ds, \end{aligned}$$

for all $w \in H^1(\mathbb{R})$, for all $\varepsilon > 0$ small enough, where $d = \min\{4 - 2(a+b), 2 - 2b, 2 - (a+c), 2 - 4a\}$. As a consequence, for each $j \in \mathbb{N}$,

$$\lambda_j(M_\varepsilon) \leq \lambda_j \left(-\Delta_{\mathbb{R}} + \frac{|\Theta'_\varepsilon(s)|^2}{2} \mathbf{1} \right) + O(\varepsilon^d). \quad (21)$$

By (20) and (21), for each $j \in \mathbb{N}$,

$$\lambda_j(-\tilde{\Delta}_\varepsilon) - \left(\frac{\pi}{2\varepsilon} \right)^2 \leq \lambda_j \left(-\Delta_{\mathbb{R}} + \frac{|\Theta'_\varepsilon(s)|^2}{2} \mathbf{1} \right) + O(\varepsilon^d). \quad (22)$$

Lower bound. For each $\varepsilon \geq 0$ small enough, consider the one-dimensional self-adjoint operator

$$(S_\varepsilon v)(t) := -v''(t) - \frac{\varepsilon t}{1 + \varepsilon t^2} v'(t), \quad \text{dom } S_\varepsilon = H_0^1(-1, 1),$$

acting in the Hilbert space $L^2((-1, 1), \sqrt{1 + \varepsilon t^2} dt)$. The particular case $\varepsilon = 0$ corresponds to the Dirichlet Laplacian operator $-\Delta_{(-1,1)}^D$ in $L^2(-1, 1)$.

Denote by $\Sigma(\varepsilon)$ the first eigenvalue of S_ε . By the analytic perturbation theory, we can write

$$\Sigma(\varepsilon) = \left(\frac{\pi}{2} \right)^2 + \delta(\varepsilon)\varepsilon + O(\varepsilon^2),$$

where

$$\delta(\varepsilon) := - \int_{-1}^1 \frac{t}{\sqrt{1 + \varepsilon t^2}} \chi_1'(t) \chi_1(t) dt;$$

see [20] for more details. Consequently, for each $\psi \in H_0^1(\Lambda)$, we have the estimate

$$\tilde{b}_\varepsilon(\psi) \geq \int_\Lambda \left(\frac{|\partial_s \psi|^2}{\tilde{f}_\varepsilon} + \frac{\Sigma(\varepsilon^2 |\Theta'_\varepsilon(s)|^2)}{\varepsilon^2} \tilde{f}_\varepsilon |\psi|^2 \right) dsdt.$$

More exactly,

$$\tilde{b}_\varepsilon(\psi) - \left(\frac{\pi}{2\varepsilon} \right)^2 \|\psi\|_{\mathcal{H}_\varepsilon}^2 \geq \int_\Lambda \left(\frac{|\partial_s \psi|^2}{\tilde{f}_\varepsilon} + |\Theta'_\varepsilon(s)|^2 \delta(\varepsilon^2 |\Theta'_\varepsilon(s)|^2) \tilde{f}_\varepsilon |\psi|^2 \right) dsdt + O(\varepsilon^{2-4a}) \|\psi\|_{\mathcal{H}_\varepsilon}^2. \quad (23)$$

Now, define the quadratic form

$$n_\varepsilon(\psi) := \int_\Lambda \left(\frac{|\partial_s \psi|^2}{\tilde{f}_\varepsilon} + |\Theta'_\varepsilon(s)|^2 \delta(\varepsilon^2 |\Theta'_\varepsilon(s)|^2) \tilde{f}_\varepsilon |\psi|^2 \right) dsdt,$$

dom $n_\varepsilon = H_0^1(\Lambda)$. Denote by N_ε the self-adjoint operator associated with $n_\varepsilon(\psi)$. For each $j \in \mathbb{N}$, inequality (23) implies

$$\lambda_j(N_\varepsilon) + O(\varepsilon^{2-4a}) \leq \lambda_j(-\tilde{\Delta}_\varepsilon) - \left(\frac{\pi}{2\varepsilon} \right)^2. \quad (24)$$

The next step is to find lower bounds for the values $\lambda_j(N_\varepsilon)$.

Lemma 2. *There exists a number $K > 0$ so that*

$$\left\| |\Theta'_\varepsilon(s)|^2 \delta(\varepsilon^2 |\Theta'_\varepsilon(s)|^2) \tilde{f}_\varepsilon - \frac{|\Theta'_\varepsilon(s)|^2}{2} \right\|_{L^\infty(\Lambda)} \leq K \varepsilon^{1-3a},$$

for all $\varepsilon > 0$ small enough.

Proof. At first, note that

$$\left| |\Theta'_\varepsilon(s)|^2 \delta(\varepsilon^2 |\Theta'_\varepsilon(s)|^2) \tilde{f}_\varepsilon - \frac{|\Theta'_\varepsilon(s)|^2}{2} \right| \leq |\Theta'_\varepsilon(s)|^2 \left(\left| \delta(\varepsilon^2 |\Theta'_\varepsilon(s)|^2) - \frac{1}{2} \right| |\tilde{f}_\varepsilon| + \frac{1}{2} |\tilde{f}_\varepsilon - 1| \right),$$

for all $(s, t) \in \Lambda$. Some calculations show that

$$\delta(\varepsilon^2 |\Theta'_\varepsilon(s)|^2) - \frac{1}{2} = \frac{\pi}{2} \int_{-1}^1 \left(\frac{1}{\tilde{f}_\varepsilon} - 1 \right) t \sin\left(\frac{\pi t}{2}\right) \cos\left(\frac{\pi t}{2}\right) dt.$$

Since $\|(1/\tilde{f}_\varepsilon) - 1\|_{L^\infty(\Lambda)} \leq \varepsilon^{1-a}$, we have the estimate $\|\delta(\varepsilon^2 |\Theta'_\varepsilon(s)|^2) - 1/2\|_{L^\infty(\Lambda)} \leq \varepsilon^{1-a}/2$. Thus, along with the condition $|\Theta'_\varepsilon(s)| \leq K/\varepsilon^a$ and the estimate $\|\tilde{f}_\varepsilon - 1\|_{L^\infty(\Lambda)} \leq \varepsilon^{1-a}$, we get the result. \square

Using Lemma 2 and the estimate $\|1/\tilde{f}_\varepsilon - 1\|_{L^\infty(\Lambda)} \leq \varepsilon^{1-a}$, one has

$$n_\varepsilon(\psi) \geq (1 + O(\varepsilon^{1-a})) \int_\Lambda |\partial_s \psi|^2 ds dt + \int_\Lambda \frac{|\Theta'_\varepsilon(s)|^2}{2} |\psi|^2 ds dt + O(\varepsilon^{1-3a}) \int_\Lambda |\psi|^2 ds dt.$$

Since $|\Theta'_\varepsilon(s)| \leq K/\varepsilon^a$, for all $s \in \mathbb{R}$, it follows that

$$n_\varepsilon(\psi) \geq (1 + O(\varepsilon^{1-a})) \int_\Lambda \left(|\partial_s \psi|^2 + \frac{|\Theta'_\varepsilon(s)|^2}{2} |\psi|^2 \right) ds dt,$$

for all $\psi \in H_0^1(\Lambda)$, for all $\varepsilon > 0$ small enough.

As a consequence, for each $j \in \mathbb{N}$,

$$(1 + O(\varepsilon^{1-a})) \lambda_j \left(-\Delta_{\mathbb{R}} + \frac{|\Theta'_\varepsilon(s)|^2}{2} \mathbf{1} \right) \leq \lambda_j(N_\varepsilon). \quad (25)$$

Inequalities (24) and (25) ensure that, for each $j \in \mathbb{N}$,

$$(1 + O(\varepsilon^{1-a})) \lambda_j \left(-\Delta_{\mathbb{R}} + \frac{|\Theta'_\varepsilon(s)|^2}{2} \mathbf{1} \right) + O(\varepsilon^{2-4a}) \leq \lambda_j(-\tilde{\Delta}_\varepsilon) - \left(\frac{\pi}{2\varepsilon} \right)^2. \quad (26)$$

Due to inequalities (22) and (26), we will study the spectral problem of the operator $-\Delta_{\mathbb{R}} + (|\Theta'_\varepsilon(s)|^2/2)\mathbf{1}$.

Proposition 4. *For each $j \in \mathbb{N}$,*

$$\lim_{\varepsilon \rightarrow 0} \lambda_j \left(-\Delta_{\mathbb{R}} + \frac{|\Theta'_\varepsilon(s)|^2}{2} \mathbf{1} \right) = \lambda_j \left(-\Delta_{\mathbb{R}} + \frac{|\Theta'(s)|^2}{2} \mathbf{1} \right). \quad (27)$$

Proof. Recall the conditions (I)-(VII) in the Introduction. Consider the quadratic forms

$$y(w) := \int_{\mathbb{R}} \left(|w'|^2 + \frac{|\Theta'(s)|^2}{2} |w|^2 \right) ds, \quad \text{dom } y = \{w \in H^1(\mathbb{R}) : y(w) < \infty\},$$

and, for each $\varepsilon > 0$ small enough,

$$y_\varepsilon(w) := \int_{\mathbb{R}} \left(|w'|^2 + \frac{|\Theta'_\varepsilon(s)|^2}{2} |w|^2 \right) ds, \quad \text{dom } y_\varepsilon = H^1(\mathbb{R});$$

denote by Y and Y_ε the self-adjoint operators associated with $y(w)$ and $y_\varepsilon(w)$, respectively. In particular, the condition (III) implies that $|\Theta'_\varepsilon(s)| \rightarrow |\Theta'(s)|$ pointwise, as $\varepsilon \rightarrow 0$. Thus, for each $w \in C_0^\infty(\mathbb{R})$, $Y_\varepsilon w \rightarrow Yw$, as $\varepsilon \rightarrow 0$. Since $C_0^\infty(\mathbb{R})$ is a core of Y , given a constant $c > 0$,

$$(Y_\varepsilon + c\mathbf{1})^{-1}u \rightarrow (Y + c\mathbf{1})^{-1}u, \quad \forall u \in L^2(\mathbb{R}). \quad (28)$$

Now, denote by \mathcal{C} the ideal of all compact operators in the algebra of all bounded operators in $L^2(\mathbb{R})$. Fix $\varepsilon^* > 0$ small enough. By (11) and (V), one has

$$\text{dist}((Y_{\varepsilon^*} + c\mathbf{1})^{-1}, \mathcal{C}) \leq \left(\liminf_{|s| \rightarrow \infty} \frac{|\Theta'_{\varepsilon^*}(s)|^2}{2} \right)^{-1} \leq 2(\varepsilon^*)^{2a}. \quad (29)$$

The condition (VII) implies that

$$(Y_\varepsilon + c\mathbf{1})^{-1} \leq (Y_{\varepsilon^*} + c\mathbf{1})^{-1}, \quad (30)$$

for all $\varepsilon < \varepsilon^*$.

As a consequence of (28), (29) and (30), Proposition 5.3 of [16] ensures that

$$\limsup_{\varepsilon \rightarrow 0} \|(Y_\varepsilon + c\mathbf{1})^{-1} - (Y + c\mathbf{1})^{-1}\| \leq 2(\varepsilon^*)^{2a}.$$

Taking $\varepsilon^* \rightarrow 0$,

$$\|(Y_\varepsilon + c\mathbf{1})^{-1} - (Y + c\mathbf{1})^{-1}\| \rightarrow 0, \quad \text{as } \varepsilon \rightarrow 0.$$

Then, we obtain (27). □

Proof of Theorem 4. It just to apply inequalities (22) and (26), and Proposition 4. □

6 Appendix

Stability of the essential spectrum

The results of this appendix are simple adaptations of Lemma 4.1 and Proposition 4.2 of [3], and Lemma 4.2 of [9].

Lemma 3. *A real number λ belongs to the essential spectrum of C_ε if and only if there exists a sequence $\{\psi_n\}_{n \in \mathbb{N}} \subset \text{dom } c_\varepsilon$ satisfying the following conditions:*

(i) $\|\psi_n\| = 1$, for all $n \in \mathbb{N}$;

(ii) $(C_\varepsilon - \lambda\mathbf{1})\psi_n \rightarrow 0$, as $n \rightarrow \infty$, in the norm of the dual space $(\text{dom } c_\varepsilon)^*$;

(iii) $\text{supp } \psi_n \subset \Lambda \setminus (-n, n) \times (-1, 1)$, for all $n \in \mathbb{N}$.

Proof. It is known that $\lambda \in \sigma_{\text{ess}}(C_\varepsilon)$ if and only if there exists a sequence $\{\xi_n\}_{n \in \mathbb{N}} \subset \text{dom } c_\varepsilon$ satisfying (i), (ii), and (iii'). $\xi_n \rightarrow 0$ in $L^2(\Lambda)$, as $n \rightarrow \infty$; see, e.g., Theorem 5 in [26]. Let $\{\psi_n\}_{n \in \mathbb{N}}$ be a sequence satisfying the conditions (i), (ii) and (iii). Consequently, it satisfies (i), (ii) and (iii').

Now, let $\{\xi_n\}_{n \in \mathbb{N}} \subset \text{dom } c_\varepsilon$ be a sequence satisfying (i), (ii), and (iii'). Take $\eta \in C^\infty(\mathbb{R}; \mathbb{R})$, $0 \leq \eta \leq 1$, $\eta = 0$ in $[-1, 1]$, and $\eta = 1$ in $\mathbb{R} \setminus (-2, 2)$. Define the sequence $\{\eta_k\}_{k \in \mathbb{N}} \subset C^\infty(\Lambda)$, where $\eta_k(s, t) := \eta(s/k)$.

Since $(1-\eta_k)(C_\varepsilon+1)^{-1}$ is compact in $L^2(\Lambda)$, by (iii'), we have $(1-\eta_k)(C_\varepsilon+1)^{-1}\xi_n \rightarrow 0$ in $L^2(\Lambda)$, as $n \rightarrow \infty$, for all $k \in \mathbb{N}$. Then there exists a subsequence $\{\xi_{n_k}\}_{k \in \mathbb{N}}$ of $\{\xi_n\}_{n \in \mathbb{N}}$ so that $(1-\eta_k)(C_\varepsilon+1)^{-1}\xi_{n_k} \rightarrow 0$ in $L^2(\Lambda)$, as $k \rightarrow \infty$. By writing

$$\xi_{n_k} = (C_\varepsilon + \mathbf{1})^{-1}(C_\varepsilon - \lambda \mathbf{1})\xi_{n_k} + (\lambda + 1)(C_\varepsilon + \mathbf{1})^{-1}\xi_{n_k},$$

and using (ii), it follows that $(1-\eta_k)\xi_{n_k} \rightarrow 0$ in $L^2(\Lambda)$, as $k \rightarrow \infty$. Thus, we can assume that $\|\eta_k \xi_{n_k}\| \geq 1/2$, for all $k \in \mathbb{N}$. Finally, define

$$\psi_k := \frac{\eta_k \xi_{n_k}}{\|\eta_k \xi_{n_k}\|}, \quad k \in \mathbb{N}.$$

The sequence $\{\psi_k\}_{k \in \mathbb{N}} \subset \text{dom } c_\varepsilon$ satisfies the conditions (i) and (iii). It remains to verify (ii), i.e.,

$$\sup_{\substack{\phi \in H_0^1(\Lambda) \\ \phi \neq 0}} \frac{|c_\varepsilon(\phi, \psi_k) - \lambda \langle \phi, \psi_k \rangle|}{\|\phi\|_+} \rightarrow 0, \quad (31)$$

as $k \rightarrow \infty$, where $\|\phi\|_+^2 := c_\varepsilon(\phi) + \|\phi\|^2$.

Some calculations show that

$$\begin{aligned} c_\varepsilon(\phi, \eta_k \xi_{n_k}) - \lambda \langle \phi, \eta_k \xi_{n_k} \rangle &= c_\varepsilon(\eta_k \phi, \xi_{n_k}) - \lambda \langle \eta_k \phi, \xi_{n_k} \rangle + \int_\Lambda \frac{1}{f_\varepsilon^2} \phi \partial_s^2 \eta_k \overline{\xi_{n_k}} \, ds dt \\ &\quad + 2 \int_\Lambda \frac{1}{f_\varepsilon^2} \left(\partial_s \phi - \frac{\partial_s f_\varepsilon}{f_\varepsilon} \phi \right) \partial_s \eta_k \overline{\xi_{n_k}} \, ds dt. \end{aligned}$$

Since $\{\xi_{n_k}\}_{k \in \mathbb{N}}$ satisfies (ii), we have

$$\sup_{\substack{\phi \in H_0^1(\Lambda) \\ \phi \neq 0}} \frac{|c_\varepsilon(\eta_k \phi, \xi_{n_k}) - \lambda \langle \eta_k \phi, \xi_{n_k} \rangle|}{\|\phi\|_+} \leq \sup_{\substack{\phi \in H_0^1(\Lambda) \\ \eta_k \phi \neq 0}} \frac{|c_\varepsilon(\eta_k \phi, \xi_{n_k}) - \lambda \langle \eta_k \phi, \xi_{n_k} \rangle|}{\|\eta_k \phi\|_+} \rightarrow 0,$$

as $k \rightarrow \infty$. By Hölder's Inequality and by the estimates $\|\phi\| \leq \|\phi\|_+$ and $c_\varepsilon(\phi) \leq \|\phi\|_+^2$, we get

$$\sup_{\substack{\phi \in H_0^1(\Lambda) \\ \phi \neq 0}} \left\{ \frac{1}{\|\phi\|_+} \int_\Lambda \frac{1}{f_\varepsilon^2} |\phi| |\partial_s^2 \eta_k| |\xi_{n_k}| \, ds dt \right\} \leq \|\partial_s^2 \eta_k\|_{L^\infty(\Lambda)} = k^{-2} \|\eta''\|_{L^\infty(\mathbb{R})} \rightarrow 0,$$

and

$$\sup_{\substack{\phi \in H_0^1(\Lambda) \\ \phi \neq 0}} \left\{ \frac{1}{\|\phi\|_+} \int_\Lambda \frac{1}{f_\varepsilon^2} \left| \partial_s \phi - \frac{\partial_s f_\varepsilon}{2f_\varepsilon} \phi \right| |\partial_s \eta_k| |\xi_{n_k}| \, ds dt \right\} \leq \|\partial_s \eta_k\|_{L^\infty(\Lambda)} = k^{-1} \|\eta'\|_{L^\infty(\mathbb{R})} \rightarrow 0,$$

as $k \rightarrow \infty$. Finally, since $\beta, \beta' \in L^\infty(\mathbb{R})$, we have

$$\sup_{\substack{\phi \in H_0^1(\Lambda) \\ \phi \neq 0}} \left\{ \frac{1}{\|\phi\|_+} \int_\Lambda \left| \frac{\partial_s f_\varepsilon}{f_\varepsilon^3} \right| |\phi| |\partial_s \eta_k| |\xi_{n_k}| \, ds dt \right\} \leq \|\partial_s \eta_k\|_{L^\infty(\Lambda)} = k^{-1} \|\eta'\|_{L^\infty(\mathbb{R})} \rightarrow 0,$$

as $k \rightarrow \infty$. Then, (31) holds true. \square

Remark 5. The same conclusion of Lemma 3 holds true for the operator D_ε .

Proof of Proposition 2. Let $\lambda \in \sigma_{\text{ess}}(D_\varepsilon)$, then there exists a sequence $\{\psi_n\}_{n \in \mathbb{N}} \subset \text{dom } d_\varepsilon$ so that $\|\psi_n\| = 1$, $\text{supp } \psi_n \subset \Lambda \setminus (-n, n) \times (-1, 1)$, for all $n \in \mathbb{N}$, and $(D_\varepsilon - \lambda \mathbf{1})\psi_n \rightarrow 0$, as $n \rightarrow \infty$, in the norm of

the dual space $(\text{dom } d_\varepsilon)^*$. Recall $\text{supp } \beta \subset [-s_0, s_0]$, for some $s_0 > 0$. Take $n_0 \in \mathbb{N}$ so that $n_0 > s_0$. For each $\phi \in H_0^1(\Lambda)$, one has

$$c_\varepsilon(\phi, \psi_n) - \lambda \langle \phi, \psi_n \rangle = d_\varepsilon(\phi, \psi_n) - \lambda \langle \phi, \psi_n \rangle,$$

for all $n \geq n_0$. Consequently,

$$\sup_{\substack{\phi \in H_0^1(\Lambda) \\ \phi \neq 0}} \frac{|c_\varepsilon(\phi, \psi_n) - \lambda \langle \phi, \psi_n \rangle|}{\|\phi\|_+} \longrightarrow 0,$$

as $n \rightarrow \infty$. The Lemma 3 implies that $\lambda \in \sigma_{ess}(C_\varepsilon)$. In a similar way, it is possible to show that $\sigma_{ess}(C_\varepsilon) \subset \sigma_{ess}(D_\varepsilon)$. \square

References

- [1] D. Borisov, P. Exner, R. Gadyl'shin, and D. Krejčířík, 'Bound states in weakly deformed strips and layers', *Ann. Henri Poincaré* **2**(3) (2001), 553-572.
- [2] D. Borisov, P. Exner, and R. Gadyl'shin, 'Geometric coupling thresholds in a two-dimensional strip', *J. Math. Phys.* **43**(12) (2002), 6265-6278.
- [3] P. Briet, H. Abdou-Soimadou, and D. Krejčířík, 'Spectral analysis of sheared nanoribbons', *Z. Angew. Math. Phys.* **70** (2) (2019), Article ID 48, 18 pages.
- [4] B. Chenaud, P. Duclos, P. Freitas, and D. Krejčířík, 'Geometrically induced discrete spectrum in curved tubes', *Differential Geom. Appl.* **23**(2) (2005), 95-105.
- [5] I. J. Clark and A. J. Bracken, 'Effective potentials of quantum strip waveguides and their dependence upon torsion', *J. Phys. A: Math. Gen.* **29**(2) (1996), 339-348.
- [6] M. Dauge, Y. Lafranche, and N. Raymond, 'Quantum waveguides with corners', *ESAIM Proceedings, EDP Sciences* **35** (2012), 14-45.
- [7] M. Dauge, and N. Raymond, 'Plane waveguides with corners in the small angle limit', *J. Math. Phys.* **53**(12) (2012), Article ID 123529, 34 pages.
- [8] M. Dauge, T. Ourmières-Bonafos, and N. Raymond, 'Spectral Asymptotics of the Dirichlet Laplacian in a conical layer', *Commun. Pure Appl. Anal.* **15**(3) (2015), 1239-1258.
- [9] Y. Dermenjian, M. Durand, and V. Iftimie, 'Spectral analysis of an acoustic multistratified perturbed cylinder', *Commun. Partial Differ. Equations* **23**(1-2) (1998), 141-169.
- [10] P. Duclos and P. Exner, 'Curvature-induced bound states in quantum waveguides in two and three dimensions', *Rev. Math. Phys.* **07**(1) (1995), 73-102.
- [11] P. Duclos, P. Exner, and D. Krejčířík, 'Bound states in curved quantum layers', *Commun. Math. Phys.* **223**(1) (2001), 13-28.
- [12] L. C. Evans, *Partial Differential Equations: Second Edition* (American Mathematical Society, Providence, RI, 2010).
- [13] P. Exner and H. Kovařík, 'Spectrum of the Schrödinger operator in a perturbed periodically twisted tube', *Lett. Math. Phys.* **73**(3) (2005), 183-192.
- [14] P. Exner and H. Kovařík, *Quantum waveguides* (Springer, 2015).

- [15] P. Exner and P. Šeba, ‘Bound states in curved quantum waveguides’, *J. Math. Phys.* **30**(11) (1989), 2574-2580.
- [16] L. Friedlander and M. Solomyak, ‘On the spectrum of the Dirichlet Laplacian in a narrow infinite strip’, *Amer. Math. Soc. Transl.* **225** (2008), 103-116.
- [17] L. Friedlander and M. Solomyak, ‘On the spectrum of the Dirichlet Laplacian in a narrow strip’, *Isr. J. Math.* **170** (2009), 337-354.
- [18] L. Friedlander and M. Solomyak, ‘On the spectrum of narrow periodic waveguides’, *Russ. J. Math. Phys.* **15**(2) (2008), 238–242.
- [19] J. Goldstone and R. L. Jaffe, ‘Bound states in twisting tubes’, *Phys. Rev. B* **45** (1992), 14100-14107.
- [20] T. Kato, ‘On the convergence of the perturbation method’, *Journal of the Faculty of Science, University of Tokyo Section 1 Mathematics, Astronomy, Physics, Chemistry* **6**(3) (1951), 145-226.
- [21] T. Kato, *Perturbation Theory for Linear Operators* (Springer-Verlag, Berlin, 1995).
- [22] D. Krejčířík, ‘Quantum strips on surfaces’, *J. Geom. Phys.* **45**(1-2) (2003), 203-217.
- [23] D. Krejčířík, ‘Hardy inequalities in strips on ruled surfaces’, *J. Inequal. Appl.* **2006** (2006), Article ID 46409, 10 pages.
- [24] D. Krejčířík and R. T. de Aldecoa, ‘Ruled strips with asymptotically diverging twisting’, *Ann. Henri Poincaré* **19**(7) (2018), 2069-2086.
- [25] D. Krejčířík and J. Kříž, ‘On the spectrum of curved planar waveguides’, *Publ. Res. Inst. Math. Sci.* **41**(3) (2005), 757-791.
- [26] D. Krejčířík and Z. Lu, ‘Location of the essential spectrum in curved quantum layers’, *J. Math. Phys.* **55**(8) (2014), Article ID 083520, 13 pages.
- [27] D. Krejčířík and H. Šediváková, ‘The effective Hamiltonian in curved quantum waveguides under mild regularity assumptions’, *Rev. Math. Phys.* **24**(7) (2012), Article ID 1250018, 39 pages.
- [28] D. Krejčířík and K. Zahradová, ‘Quantum strips in higher dimensions’, *Oper. Matrices* **14**(3) (2020), 635-665.
- [29] M. Reed and B. Simon, *Methods of Modern Mathematical Physics, IV: Analysis of Operators* (Academic Press, New York, 1978).
- [30] W. Renger and W. Bulla, ‘Existence of bound states in quantum waveguides under weak conditions’, *Lett. Math. Phys.* **35**(1) (1995), 1–12.
- [31] A. A. Verri, ‘Dirichlet Laplacian in a thin twisted strip’, *Int. J. Math.* **30**(2) (2019), Article ID 1950006, 17 pages.

RAFAEL T. AMORIM, Departamento de Matemática,
 Universidade Federal de São Carlos - UFSCar, São Carlos, SP, 13565-905, Brazil
 e-mail: rafaelamorim@estudante.ufscar.br

ALESSANDRA A. VERRI, Departamento de Matemática,
 Universidade Federal de São Carlos - UFSCar, São Carlos, SP, 13565-905, Brazil
 e-mail: alessandraverri@ufscar.br