

# Closed Range Integral Operators on Hardy, BMOA and Besov Spaces

Kostas Panteris

Department of Mathematics and Applied Mathematics, University of Crete, University Campus Voutes, 70013 Heraklion, Greece

## ARTICLE HISTORY

Compiled December 7, 2021

## ABSTRACT

If  $g \in H^\infty$ , the integral operator  $S_g$  on  $H^p$ , BMOA and  $B^p$  (Besov) spaces, is defined as  $S_g f(z) = \int_0^z f'(w)g(w)dw$ . In this paper, we prove three necessary and sufficient conditions for the operator  $S_g$  to have closed range on  $H^p$  ( $1 < p < \infty$ ), BMOA and  $B^p$  ( $1 < p < \infty$ ).

## 1. Introduction and Preliminaries

Let  $\mathbb{D}$  denote the open unit disk in the complex plane,  $\mathbb{T}$  the unit circle,  $A$  the normalized area Lebesgue measure in  $\mathbb{D}$  and  $m$  the normalized length Lebesgue measure in  $\mathbb{T}$ . For  $1 \leq p < \infty$  the Hardy space  $H^p$  is defined as the set of all analytic functions  $f$  in  $\mathbb{D}$  for which

$$\sup_{0 \leq r < 1} \int_{\mathbb{T}} |f(r\zeta)|^p dm(\zeta) < +\infty$$

and the corresponding norm in  $H^p$  is defined by

$$\|f\|_{H^p}^p = \sup_{0 \leq r < 1} \int_{\mathbb{T}} |f(r\zeta)|^p dm(\zeta).$$

When  $p = \infty$ , we define  $H^\infty$  to be the space of bounded analytic functions  $f$  in  $\mathbb{D}$  and  $\|f\|_\infty = \sup\{|f(z)| : z \in \mathbb{D}\}$ .

In this work we will mainly make use of the following equivalent norm (see Calderon's theorem in [3], page 213) in  $H^p$ ,  $1 \leq p < \infty$ :

$$\|f\|_{H^p}^p = |f(0)|^p + \int_{\mathbb{T}} \left( \iint_{\Gamma_\beta(\zeta)} |f'(z)|^2 dA(z) \right)^{\frac{p}{2}} dm(\zeta), \quad (1)$$

where  $\Gamma_\beta(\zeta)$  is the Stolz angle at  $\zeta \in \mathbb{T}$ , the conelike region with aperture  $\beta \in (0, 1)$ ,

which is defined as

$$\Gamma_\beta(\zeta) = \{z \in \mathbb{D} : |z| < \beta\} \cup \bigcup_{|z| < \beta} [z, \zeta).$$

The BMOA space is defined as the set of all analytic functions  $f$  in  $\mathbb{D}$  for which

$$\sup_{\beta \in \mathbb{D}} \iint_{\mathbb{D}} \frac{1 - |\beta|^2}{|1 - \bar{\beta}z|^2} |f'(z)|^2 \log \frac{1}{|z|} dA(z) < \infty$$

and we may define the corresponding norm in BMOA by

$$\|f\|_*^2 = |f(0)|^2 + \sup_{\beta \in \mathbb{D}} \iint_{\mathbb{D}} \frac{1 - |\beta|^2}{|1 - \bar{\beta}z|^2} |f'(z)|^2 \log \frac{1}{|z|} dA(z).$$

For  $1 < p < \infty$  the Besov space  $B^p$  is defined as the set of all analytic functions  $f$  in  $\mathbb{D}$  for which

$$\iint_{\mathbb{D}} |f'(z)|^p (1 - |z|^2)^{p-2} dA(z) < +\infty$$

and the corresponding norm in  $B^p$  is defined by

$$\|f\|_{B^p}^p = |f(0)|^p + \iint_{\mathbb{D}} |f'(z)|^p (1 - |z|^2)^{p-2} dA(z).$$

Let  $g : \mathbb{D} \rightarrow \mathbb{C}$  be an analytic function. If  $X$  is a space of analytic functions  $f$  in  $\mathbb{D}$  (in particular, in this paper,  $X = H^p$  or  $X = BMOA$  or  $X = B^p$ ) then, the integral operator  $S_g : X \rightarrow X$ , induced by  $g$ , is defined as

$$S_g f(z) = \int_0^z f'(w) g(w) dw, \quad z \in \mathbb{D},$$

for every  $f \in X$ .

Let  $\rho(z, w)$  denote the pseudo-hyperbolic distance between  $z, w \in \mathbb{D}$ ,

$$\rho(z, w) = \left| \frac{z - w}{1 - \bar{z}w} \right|,$$

$D_\eta(a)$  denote the pseudo-hyperbolic disk of center  $a \in \mathbb{D}$  and radius  $\eta < 1$ :

$$D_\eta(a) = \{z \in \mathbb{D} : \rho(a, z) < \eta\},$$

and  $\Delta_\eta(\alpha)$  denote the euclidean disk of center  $a \in \mathbb{D}$  and radius  $\eta(1 - |\alpha|)$ ,  $\eta < 1$ :

$$\Delta_\eta(\alpha) = \{z \in \mathbb{D} : |z - \alpha| < \eta(1 - |\alpha|)\}.$$

In the following,  $C$  denotes a positive and finite constant which may change from one occurrence to another. Moreover, by writing  $K(z) \asymp L(z)$  for the non-negative

quantities  $K(z)$  and  $L(z)$  we mean that  $K(z)$  is comparable to  $L(z)$  if  $z$  belongs to a specific set: there are positive constants  $C_1$  and  $C_2$  independent of  $z$  such that

$$C_1K(z) \leq L(z) \leq C_2K(z).$$

## 2. Closed range integral operators on Hardy spaces

Let  $g : \mathbb{D} \rightarrow \mathbb{C}$  be an analytic function and, for  $c > 0$ , let  $G_c = \{z \in \mathbb{D} : |g(z)| > c\}$ . It is well known (see [1]) that the integral operator  $S_g : H^p \rightarrow H^p$  ( $1 \leq p < \infty$ ) is bounded if and only if  $g \in H^\infty$ .

We say that  $S_g$ , on  $H^p$ , is bounded below, if there is  $C > 0$  such that  $\|S_g f\|_{H^p} > C\|f\|_{H^p}$  for every  $f \in H^p$ . Since  $S_g$  maps every constant function to the 0 function, if we want to study the property of being bounded below for  $S_g$ , we are obliged to consider spaces of analytic functions modulo the constants or, equivalently, spaces of analytic functions  $f$  such that  $f(0) = 0$ . Theorem 2.3 in [1] states that  $S_g$  is bounded below on  $H^p/\mathbb{C}$  if and only if it has closed range on  $H^p/\mathbb{C}$ . Next, we denote  $H^p/\mathbb{C}$  as  $H_0^p$ .

Corollary 3.6 in [1] states that  $S_g : H_0^2 \rightarrow H_0^2$  has closed range if and only if there exist  $c > 0$ ,  $\delta > 0$  and  $\eta \in (0, 1)$  such that

$$A(G_c \cap D_\eta(a)) \geq \delta A(D_\eta(a))$$

for all  $a \in \mathbb{D}$ .

In the end of [1] A. Anderson posed the question, if the above condition for  $H_0^2$  holds also for all  $H_0^p$ . In this paper, theorem 2.2 gives an affirmative answer to this question, for the case  $1 < p < \infty$ . Although the answer in case  $p = 2$  is an immediate consequence of D. Luecking's theorem (see [1], Proposition 3.5), the answer in case  $1 < p < \infty$  requires much more effort.

For  $\lambda \in (0, 1)$  and  $f \in H^p$  we set

$$E_\lambda(\alpha) = \{z \in \Delta_\eta(\alpha) : |f'(z)|^2 > \lambda|f'(\alpha)|^2\}$$

and

$$B_\lambda f(\alpha) = \frac{1}{A(E_\lambda(\alpha))} \iint_{E_\lambda(\alpha)} |f'(z)|^2 dA(z).$$

Lemma 2.1 is due to D. Luecking (see [2], lemma 1).

**Lemma 2.1.** *Let  $f$  analytic in  $\mathbb{D}$ ,  $a \in \mathbb{D}$  and  $\lambda \in (0, 1)$ . Then*

$$\frac{A(E_\lambda(\alpha))}{A(\Delta_\eta(\alpha))} \geq \frac{\log \frac{1}{\lambda}}{\log \frac{B_\lambda f(\alpha)}{|f'(\alpha)|^2} + \log \frac{1}{\lambda}}.$$

Moreover in [2], the following sentence is proved: If  $\alpha \in \mathbb{D}$  and  $\frac{2\eta}{1+\eta^2} \leq r < 1$  then

$$\Delta_\eta(\alpha) \subseteq D_r(\alpha). \tag{2}$$

We proceed with the main result of this section.

**Theorem 2.2.** *Let  $1 < p < \infty$  and  $g \in H^\infty$ . Then the following are equivalent:*

- (i)  $S_g : H_0^p \rightarrow H_0^p$  has closed range
- (ii) There exist  $c > 0$ ,  $\delta > 0$  and  $\eta \in (0, 1)$  such that

$$A(G_c \cap D_\eta(a)) \geq \delta A(D_\eta(a)) \quad (3)$$

for all  $a \in \mathbb{D}$ .

- (iii) There exist  $c > 0$ ,  $\delta > 0$  and  $\eta \in (0, 1)$  such that

$$A(G_c \cap \Delta_\eta(a)) \geq \delta A(\Delta_\eta(a)) \quad (4)$$

for all  $a \in \mathbb{D}$ .

We first prove two lemmas which will play an important role in the proof of theorem 2.2.

For  $\zeta \in \mathbb{T}$  and  $0 < \beta < \beta' < 1$  we consider the Stolz angles  $\Gamma_\beta(\zeta)$  and  $\Gamma_{\beta'}(\zeta)$ , where  $\beta'$  has been chosen so that  $\Delta_\eta(\alpha) \subset \Gamma_{\beta'}(\zeta)$  for every  $\alpha \in \Gamma_\beta(\zeta)$ .

**Lemma 2.3.** *Let  $\varepsilon > 0$ ,  $f$  analytic in  $\mathbb{D}$  and*

$$A = \left\{ \alpha \in \mathbb{D} : |f'(\alpha)|^2 < \frac{\varepsilon}{A(\Delta_\eta(\alpha))} \iint_{\Delta_\eta(\alpha)} |f'(z)|^2 dA(z) \right\}.$$

There is  $C > 0$  depending only on  $\eta$  such that

$$\iint_{A \cap \Gamma_\beta(\zeta)} |f'(z)|^2 dA(z) \leq \varepsilon C \iint_{\Gamma_{\beta'}(\zeta)} |f'(z)|^2 dA(z)$$

**Proof.** Integrating

$$|f'(\alpha)|^2 < \frac{\varepsilon}{A(\Delta_\eta(\alpha))} \iint_{\Delta_\eta(\alpha)} |f'(z)|^2 dA(z)$$

over  $\alpha \in A \cap \Gamma_\beta(\zeta)$  and using Fubini's theorem on the right side, we get

$$\iint_{A \cap \Gamma_\beta(\zeta)} |f'(\alpha)|^2 dA(\alpha) < \varepsilon \iint_{\Gamma_{\beta'}(\zeta)} |f'(z)|^2 \left[ \iint_{A \cap \Gamma_\beta(\zeta)} \frac{\chi_{\Delta_\eta(\alpha)}(z)}{A(\Delta_\eta(\alpha))} dA(\alpha) \right] dA(z)$$

Using (2) with  $r = \frac{2\eta}{1+\eta^2}$ , we have  $\chi_{\Delta_\eta(\alpha)}(z) \leq \chi_{D_r(\alpha)}(z) = \chi_{D_r(z)}(\alpha)$ . We have that  $A(D_r(z)) \asymp (1 - |z|)^2$  and, for  $\alpha \in D_\eta(z)$ , we have  $(1 - |z|) \asymp (1 - |\alpha|)$ , where the underlying constants in these relations depend only on  $\eta$ . In addition,  $A(\Delta_\eta(\alpha)) =$

$\eta^2(1 - |\alpha|)^2$ . So,

$$\begin{aligned} \iint_{A \cap \Gamma_\beta(\zeta)} \frac{\chi_{\Delta_\eta(\alpha)}(z)}{A(\Delta_\eta(\alpha))} dA(\alpha) &\leq \iint_{A \cap \Gamma_\beta(\zeta)} \frac{\chi_{D_r(z)}(\alpha)}{\eta^2(1 - |\alpha|)^2} dA(\alpha) \\ &\leq C \iint_{D_r(z)} \frac{1}{\eta^2(1 - |z|)^2} dA(\alpha) = C \frac{A(D_r(z))}{\eta^2(1 - |z|)^2} \leq C, \end{aligned} \quad (5)$$

where  $C > 0$  depends only on  $\eta$ . □

**Lemma 2.4.** *Let  $0 < \varepsilon < 1$ ,  $f$  analytic in  $\mathbb{D}$ ,  $0 < \lambda < \frac{1}{2}$  and*

$$B = \left\{ \alpha \in \mathbb{D} : |f'(\alpha)|^2 < \varepsilon^3 B_\lambda f(\alpha) \right\}.$$

*There is  $C > 0$  depending only on  $\eta$  such that*

$$\iint_{B \cap \Gamma_\beta(\zeta)} |f'(z)|^2 dA(z) \leq \varepsilon C \iint_{\Gamma_{\beta'}(\zeta)} |f'(z)|^2 dA(z)$$

**Proof.** We write

$$\iint_{B \cap \Gamma_\beta(\zeta)} |f'(\alpha)|^2 dA(\alpha) = \iint_{B \cap \Gamma_\beta(\zeta) \cap A} |f'(\alpha)|^2 dA(\alpha) + \iint_{(B \cap \Gamma_\beta(\zeta)) \setminus A} |f'(\alpha)|^2 dA(\alpha),$$

where  $A$  is as in lemma 2.3. The first integral is estimated by lemma 2.3, so it remains to show the desired result for the second integral. Integrating the relation

$$|f'(\alpha)|^2 < \varepsilon^3 B_\lambda f(\alpha) = \varepsilon^3 \frac{1}{A(E_\lambda(\alpha))} \iint_{E_\lambda(\alpha)} |f'(z)|^2 dA(z)$$

over the set  $(B \cap \Gamma_\beta(\zeta)) \setminus A$  and using Fubini's theorem on the right side, we get

$$\begin{aligned} \iint_{(B \cap \Gamma_\beta(\zeta)) \setminus A} |f'(\alpha)|^2 dA(\alpha) &\leq \varepsilon^3 \iint_{\Gamma_{\beta'}(\zeta)} |f'(z)|^2 \left[ \iint_{(B \cap \Gamma_\beta(\zeta)) \setminus A} \frac{1}{A(E_\lambda(\alpha))} \chi_{E_\lambda(\alpha)}(z) dA(\alpha) \right] dA(z) \\ &\leq \varepsilon^3 \iint_{\Gamma_{\beta'}(\zeta)} |f'(z)|^2 \left[ \iint_{(B \cap \Gamma_\beta(\zeta)) \setminus A} \frac{1}{A(E_\lambda(\alpha))} \chi_{\Delta_\eta(\alpha)}(z) dA(\alpha) \right] dA(z) \end{aligned} \quad (6)$$

where the last inequality is justified by  $E_\lambda(\alpha) \subseteq \Delta_\eta(\alpha)$ . Let  $\alpha \notin A$ , i.e.

$$|f'(\alpha)|^2 \geq \frac{\varepsilon}{A(\Delta_\eta(\alpha))} \iint_{\Delta_\eta(\alpha)} |f'(z)|^2 dA(z). \quad (7)$$

Set  $r = \eta(1 - |\alpha|)$  and suppose  $\lambda < \frac{1}{2}$  and  $|z - \alpha| < \frac{r}{4}$ . We have that

$$\begin{aligned} |f'(z)^2 - f'(\alpha)^2| &= \frac{1}{2\pi} \left| \int_{|w-\alpha|=\frac{r}{2}} f'(w)^2 \left( \frac{1}{w-z} - \frac{1}{w-\alpha} \right) dw \right| \\ &= \frac{1}{2\pi} \left| \int_{|w-\alpha|=\frac{r}{2}} f'(w)^2 \frac{z-\alpha}{(w-z)(w-\alpha)} dw \right|. \end{aligned} \quad (8)$$

For  $|w - \alpha| = \frac{r}{2}$ , by the subharmonicity of  $|f'|^2$  we have

$$|f'(w)|^2 < \frac{1}{\frac{r^2}{4}} \iint_{|u-w| \leq \frac{r}{2}} |f'(u)|^2 dA(u) \leq \frac{C}{A(\Delta_\eta(\alpha))} \iint_{\Delta_\eta(\alpha)} |f'(u)|^2 dA(u).$$

Since  $|w - z| > \frac{r}{4}$  when  $|w - \alpha| = \frac{r}{2}$ , from (8) we get

$$|f'(z)^2 - f'(\alpha)^2| \leq \frac{C|z-\alpha|}{r} \frac{1}{A(\Delta_\eta(\alpha))} \iint_{\Delta_\eta(\alpha)} |f'(u)|^2 dA(u).$$

Since we may assume that  $C > 2$ , taking  $|z - \alpha| < \frac{\varepsilon r}{2C}$ , then we have  $|z - \alpha| < \frac{r}{4}$  and we get

$$|f'(z)^2 - f'(\alpha)^2| \leq \frac{\varepsilon}{2A(\Delta_\eta(\alpha))} \iint_{\Delta_\eta(\alpha)} |f'(u)|^2 dA(u). \quad (9)$$

Combining (7) and (9), we get

$$|f'(z)|^2 > \frac{1}{2}|f'(\alpha)|^2 > \lambda|f'(\alpha)|^2.$$

This means that if  $\Delta' = \{z \in \mathbb{D} : |z - \alpha| < \frac{\varepsilon r}{2C}\}$  then  $\Delta' \subset E_\lambda(\alpha)$  and

$$A(E_\lambda(\alpha)) \geq A(\Delta') = \frac{\varepsilon^2}{4C^2} r^2 = \frac{\varepsilon^2}{4C^2} A(\Delta_\eta(\alpha)).$$

We finally use this last inequality in (6) and we complete the proof.  $\square$

**Proof of theorem 2.2.** (ii)  $\Leftrightarrow$  (iii) This is easy and it is proved in [2].

(iii)  $\Rightarrow$  (i) Let  $\alpha \in \mathbb{D} \setminus B$ , where  $B$  is as in lemma 2.4, where  $0 < \varepsilon < 1$ ,  $0 < \lambda < \frac{1}{2}$ .

Then  $\frac{B_\lambda f(\alpha)}{|f'(\alpha)|^2} \leq \frac{1}{\varepsilon^3}$  and, if we choose  $\lambda < \varepsilon^{\frac{6}{5}}$ , then, from lemma 2.1, we get that

$$\frac{A(E_\lambda(\alpha))}{A(\Delta_\eta(\alpha))} > \frac{\frac{2}{\delta} \log \frac{1}{\varepsilon^3}}{\log \frac{1}{\varepsilon^3} + \frac{2}{\delta} \log \frac{1}{\varepsilon^3}} > 1 - \frac{\delta}{2}. \quad (10)$$

Combining (4) and (10), we get

$$\begin{aligned}
A(G_c \cap E_\lambda(\alpha)) &= A(G_c \cap \Delta_\eta(\alpha)) - A(G_c \cap (\Delta_\eta(\alpha) \setminus E_\lambda(\alpha))) \\
&\geq \delta A(\Delta_\eta(\alpha)) - A(\Delta_\eta(\alpha) \setminus E_\lambda(\alpha)) \\
&= \delta A(\Delta_\eta(\alpha)) - A(\Delta_\eta(\alpha)) + A(E_\lambda(\alpha)) \\
&\geq \delta A(\Delta_\eta(\alpha)) - A(\Delta_\eta(\alpha)) + A(\Delta_\eta(\alpha)) - \frac{\delta}{2} A(\Delta_\eta(\alpha)) \\
&= \frac{\delta}{2} A(\Delta_\eta(\alpha))
\end{aligned}$$

Now let  $f \in H_0^p$ ,  $\zeta \in \mathbb{T}$  and  $\alpha \in \Gamma_\beta(\zeta) \setminus B$ . Then, using the last relation and  $E_\lambda(\alpha) \subset \Delta_\eta(\alpha) \subset \Gamma_{\beta'}(\zeta)$ , we get

$$\begin{aligned}
\frac{1}{A(\Delta_\eta(\alpha))} \iint_{G_c \cap \Gamma_{\beta'}(\zeta)} \chi_{\Delta_\eta(\alpha)}(z) |f'(z)|^2 dA(z) \\
\geq \frac{\delta}{2A(G_c \cap E_\lambda(\alpha))} \iint_{G_c \cap E_\lambda(\alpha)} \chi_{\Delta_\eta(\alpha)}(z) |f'(z)|^2 dA(z) \\
= \frac{\delta}{2A(G_c \cap E_\lambda(\alpha))} \iint_{G_c \cap E_\lambda(\alpha)} |f'(z)|^2 dA(z) \geq \frac{\delta\lambda}{2} |f'(\alpha)|^2.
\end{aligned}$$

Integrating the last relation over the set  $\Gamma_\beta(\zeta) \setminus B$  and using Fubini's theorem on the left side, we have

$$\iint_{G_c \cap \Gamma_{\beta'}(\zeta)} |f'(z)|^2 \left[ \iint_{\Gamma_\beta(\zeta) \setminus B} \frac{\chi_{\Delta_\eta(\alpha)}(z)}{A(\Delta_\eta(\alpha))} dA(\alpha) \right] dA(z) \geq \frac{\delta\lambda}{2} \iint_{\Gamma_\beta(\zeta) \setminus B} |f'(\alpha)|^2 dA(\alpha).$$

With similar arguments as in relation (5), we can show that the integral in the brackets is bounded above from a constant  $C > 0$  depending only on  $\eta$ . So, we have that

$$\begin{aligned}
\iint_{G_c \cap \Gamma_{\beta'}(\zeta)} |f'(z)|^2 dA(z) &\geq \frac{C\delta\lambda}{2} \iint_{\Gamma_\beta(\zeta) \setminus B} |f'(\alpha)|^2 dA(\alpha) \\
&= \frac{C\delta\lambda}{2} \iint_{\Gamma_\beta(\zeta)} |f'(\alpha)|^2 dA(\alpha) - \frac{C\delta\lambda}{2} \iint_{\Gamma_\beta(\zeta) \cap B} |f'(\alpha)|^2 dA(\alpha).
\end{aligned}$$

Because of lemma 2.4 we have that

$$\iint_{G_c \cap \Gamma_{\beta'}(\zeta)} |f'(z)|^2 dA(z) \geq \frac{C\delta\lambda}{2} \iint_{\Gamma_\beta(\zeta)} |f'(\alpha)|^2 dA(\alpha) - \varepsilon \frac{C'\delta\lambda}{2} \iint_{\Gamma_{\beta'}(\zeta)} |f'(\alpha)|^2 dA(\alpha)$$

and so

$$\iint_{G_c \cap \Gamma_{\beta'}(\zeta)} |f'(z)|^2 dA(z) + \varepsilon \frac{C'\delta\lambda}{2} \iint_{\Gamma_{\beta'}(\zeta)} |f'(\alpha)|^2 dA(\alpha) \geq \frac{C\delta\lambda}{2} \iint_{\Gamma_\beta(\zeta)} |f'(\alpha)|^2 dA(\alpha).$$

Hence,

$$\begin{aligned} & \left( \iint_{G_c \cap \Gamma_{\beta'}(\zeta)} |f'(z)|^2 dA(z) \right)^{\frac{1}{2}} + \left( \frac{C' \varepsilon \delta \lambda}{2} \right)^{\frac{1}{2}} \left( \iint_{\Gamma_{\beta'}(\zeta)} |f'(\alpha)|^2 dA(\alpha) \right)^{\frac{1}{2}} \\ & \geq \left( \frac{C \delta \lambda}{2} \right)^{\frac{1}{2}} \left( \iint_{\Gamma_{\beta}(\zeta)} |f'(\alpha)|^2 dA(\alpha) \right)^{\frac{1}{2}}. \end{aligned}$$

Applying Minkowski's inequality, we get

$$\begin{aligned} & \left[ \int_{\mathbb{T}} \left( \iint_{G_c \cap \Gamma_{\beta'}(\zeta)} |f'(z)|^2 dA(z) \right)^{\frac{p}{2}} dm(\zeta) \right]^{\frac{1}{p}} + \left( \frac{C' \varepsilon \delta \lambda}{2} \right)^{\frac{1}{2}} \left[ \int_{\mathbb{T}} \left( \iint_{\Gamma_{\beta'}(\zeta)} |f'(\alpha)|^2 dA(\alpha) \right)^{\frac{p}{2}} dm(\zeta) \right]^{\frac{1}{p}} \\ & \geq \left( \frac{C \delta \lambda}{2} \right)^{\frac{1}{2}} \left[ \int_{\mathbb{T}} \left( \iint_{\Gamma_{\beta}(\zeta)} |f'(\alpha)|^2 dA(\alpha) \right)^{\frac{p}{2}} dm(\zeta) \right]^{\frac{1}{p}} \end{aligned}$$

and so

$$\begin{aligned} & \left[ \int_{\mathbb{T}} \left( \iint_{G_c \cap \Gamma_{\beta'}(\zeta)} |f'(z)|^2 dA(z) \right)^{\frac{p}{2}} dm(\zeta) \right]^{\frac{1}{p}} \geq \left( \frac{C \delta \lambda}{2} \right)^{\frac{1}{2}} \left[ \int_{\mathbb{T}} \left( \iint_{\Gamma_{\beta}(\zeta)} |f'(\alpha)|^2 dA(\alpha) \right)^{\frac{p}{2}} dm(\zeta) \right]^{\frac{1}{p}} \\ & \quad - \left( \frac{C' \varepsilon \delta \lambda}{2} \right)^{\frac{1}{2}} \left[ \int_{\mathbb{T}} \left( \iint_{\Gamma_{\beta'}(\zeta)} |f'(\alpha)|^2 dA(\alpha) \right)^{\frac{p}{2}} dm(\zeta) \right]^{\frac{1}{p}}. \end{aligned} \tag{11}$$

According to (1), both integrals at the right side of (11), represent equivalent norms in  $H_0^p$ . Due to the relation between  $\beta$  and  $\beta'$  there is  $C'' > 0$  which depends only on  $\eta$ , such that

$$\left[ \int_{\mathbb{T}} \left( \iint_{\Gamma_{\beta'}(\zeta)} |f'(\alpha)|^2 dA(\alpha) \right)^{\frac{p}{2}} dm(\zeta) \right]^{\frac{1}{p}} \leq C'' \left[ \int_{\mathbb{T}} \left( \iint_{\Gamma_{\beta}(\zeta)} |f'(\alpha)|^2 dA(\alpha) \right)^{\frac{p}{2}} dm(\zeta) \right]^{\frac{1}{p}}. \tag{12}$$

Combining relations (11) and (12), we get

$$\begin{aligned} & \left[ \int_{\mathbb{T}} \left[ \left( \iint_{G_c \cap \Gamma_{\beta'}(\zeta)} |f'(z)|^2 dA(z) \right)^{\frac{p}{2}} dm(\zeta) \right]^{\frac{1}{p}} \geq \left( \frac{C \delta \lambda}{2} \right)^{\frac{1}{2}} \left[ \int_{\mathbb{T}} \left( \iint_{\Gamma_{\beta}(\zeta)} |f'(\alpha)|^2 dA(\alpha) \right)^{\frac{p}{2}} dm(\zeta) \right]^{\frac{1}{p}} \\ & \quad - \left( \frac{C' \varepsilon \delta \lambda}{2} \right)^{\frac{1}{2}} C'' \left[ \int_{\mathbb{T}} \left( \iint_{\Gamma_{\beta}(\zeta)} |f'(\alpha)|^2 dA(\alpha) \right)^{\frac{p}{2}} dm(\zeta) \right]^{\frac{1}{p}} \\ & \quad = \left( \frac{\delta \lambda}{2} \right)^{\frac{1}{2}} [C^{\frac{1}{2}} - \varepsilon^{\frac{1}{2}} C'^{\frac{1}{2}} C''] \|f\|_{H_0^p}. \end{aligned}$$



Choosing  $\varepsilon$  small enough so that  $C - \varepsilon^{\frac{1}{2}}C'^{\frac{1}{2}}C'' > 0$ , we have that

$$\left[ \int_{\mathbb{T}} \left( \iint_{G_c \cap \Gamma_{\beta'}(\zeta)} |f'(z)|^2 dA(z) \right)^{\frac{p}{2}} dm(\zeta) \right]^{\frac{1}{p}} \geq C \|f\|_{H_0^p},$$

and since  $G_c = \{z \in \mathbb{D} : |g(z)| > c\}$ , we have

$$\begin{aligned} \|S_g f\|_{H_0^p} &\asymp \left[ \int_{\mathbb{T}} \left( \iint_{\Gamma_{\beta'}(\zeta)} |(S_g f(z))'|^2 dA(z) \right)^{\frac{p}{2}} dm(\zeta) \right]^{\frac{1}{p}} \\ &= \left[ \int_{\mathbb{T}} \left( \iint_{\Gamma_{\beta'}(\zeta)} |f'(z)|^2 |g(z)|^2 dA(z) \right)^{\frac{p}{2}} dm(\zeta) \right]^{\frac{1}{p}} \\ &\geq c \left[ \int_{\mathbb{T}} \left( \iint_{G_c \cap \Gamma_{\beta'}(\zeta)} |f'(z)|^2 dA(z) \right)^{\frac{p}{2}} dm(\zeta) \right]^{\frac{1}{p}} \geq C \|f\|_{H_0^p}. \end{aligned}$$

So the integral operator  $S_g$  has closed range.

(i)  $\Rightarrow$  (ii) Let  $\alpha \in \mathbb{D}$ ,  $\zeta \in \mathbb{T}$ ,  $\eta \in (0, 1)$ ,  $E(z_0; r) = \{z \in \mathbb{D} : |z - z_0| < r\}$ ,  $C(z_0, r) = \{z \in \mathbb{D} : |z - z_0| = r\}$  and the arc  $I_\alpha = \{\zeta \in \mathbb{T} : \Gamma_{\frac{1}{2}}(\zeta) \cap D_\eta(\alpha) \neq \emptyset\}$ . It's easy to see that  $\zeta \in I_\alpha$  is equivalent to  $\alpha \in \Gamma_{\eta'}(\zeta)$ , where  $\eta'$  depends only on  $\eta$ . In fact, an elementary geometric argument shows that  $1 - \eta' \asymp 1 - \eta$ , where the underlying constants are absolute.

Set  $R_0 = \frac{1+\eta'}{2}$ . We continue with the proof by considering two cases for  $\alpha$ : **(a)**  $R_0 \leq |\alpha| < 1$  and **(b)**  $0 \leq |\alpha| \leq R_0$ .

**Case (a)**  $R_0 \leq |\alpha| \leq 1$ : Then another simple geometric argument gives  $m(I_\alpha) \asymp \frac{1-|\alpha|}{(1-\eta')^{\frac{1}{2}}}$  and hence:

$$m(I_\alpha) \asymp \frac{1-|\alpha|}{(1-\eta)^{\frac{1}{2}}}. \quad (13)$$

If  $S_g$  has closed range on  $H_0^p$  then there exists  $C > 0$  such that for every  $f \in H_0^p$  we have

$$C \|S_g f\|_{H_0^p}^p \geq \|f\|_{H_0^p}^p. \quad (14)$$

Let

$$\psi_\alpha(z) = \frac{\alpha - z}{1 - \bar{\alpha}z}.$$

Then, after some calculations, we get that  $\|\psi_\alpha - \alpha\|_{H^p}^p \asymp (1 - |\alpha|)$ .

Setting  $f = \psi_\alpha - \alpha$  in (14) and using  $(x + y)^p \leq 2^{p-1}(x^p + y^p)$ , we get

$$\begin{aligned}
1 - |\alpha| &\leq C \|S_g(\psi_\alpha - \alpha)\|_{H_0^p}^p = C \int_{\mathbb{T}} \left( \iint_{\Gamma_{\frac{1}{2}}(\zeta)} |\psi'_\alpha(z)|^2 |g(z)|^2 dA(z) \right)^{\frac{p}{2}} dm(\zeta) \\
&\leq C \int_{I_\alpha} \left( \iint_{\Gamma_{\frac{1}{2}}(\zeta) \cap G_c \cap D_\eta(\alpha)} |\psi'_\alpha(z)|^2 |g(z)|^2 dA(z) \right)^{\frac{p}{2}} dm(\zeta) \\
&\quad + C \int_{I_\alpha} \left( \iint_{\Gamma_{\frac{1}{2}}(\zeta) \cap (D_\eta(\alpha) \setminus G_c)} |\psi'_\alpha(z)|^2 |g(z)|^2 dA(z) \right)^{\frac{p}{2}} dm(\zeta) \\
&\quad + C \int_{I_\alpha} \left( \iint_{\Gamma_{\frac{1}{2}}(\zeta) \setminus D_\eta(\alpha)} |\psi'_\alpha(z)|^2 |g(z)|^2 dA(z) \right)^{\frac{p}{2}} dm(\zeta) \\
&\quad + C \int_{\mathbb{T} \setminus I_\alpha} \left( \iint_{\Gamma_{\frac{1}{2}}(\zeta)} |\psi'_\alpha(z)|^2 |g(z)|^2 dA(z) \right)^{\frac{p}{2}} dm(\zeta) \\
&= C(I_1 + I_2 + I_3 + I_4). \tag{15}
\end{aligned}$$

Using  $A(D_\eta(\alpha)) = \frac{(1-|\alpha|^2)^2}{(1-\eta^2|\alpha|^2)^2} \eta^2 \leq \frac{(1-|\alpha|^2)^2}{(1-\eta^2)^2}$ , we get

$$\begin{aligned}
I_1 &\leq \|g\|_\infty^p \int_{I_\alpha} \left( \iint_{G_c \cap D_\eta(\alpha)} \frac{(1-|\alpha|^2)^2}{|1-\bar{\alpha}z|^4} dA(z) \right)^{\frac{p}{2}} dm(\zeta) \\
&\leq \|g\|_\infty^p m(I_\alpha) \left( \frac{A(G_c \cap D_\eta(\alpha))}{(1-|\alpha|^2)^2} \right)^{\frac{p}{2}} \\
&\leq \|g\|_\infty^p m(I_\alpha) \frac{1}{(1-\eta^2)^p} \left( \frac{A(G_c \cap D_\eta(\alpha))}{A(D_\eta(\alpha))} \right)^{\frac{p}{2}}.
\end{aligned}$$

Using  $|g(z)| \leq c$  in  $\mathbb{D} \setminus G_c$  and making the change of variables  $w = \psi_\alpha(z)$ , we get

$$I_2 \leq c^p \int_{I_\alpha} \left( \iint_{\mathbb{D}} |\psi'_\alpha(z)|^2 dA(z) \right)^{\frac{p}{2}} dm(\zeta) = c^p \int_{I_\alpha} \left( \iint_{\mathbb{D}} dA(w) \right)^{\frac{p}{2}} dm(\zeta) = c^p m(I_\alpha).$$

We increase  $I_3$  by extending it over  $\mathbb{D} \setminus D_\eta(\alpha)$  and then we make the change of variables  $w = \psi_\alpha(z)$  to get

$$I_3 \leq \|g\|_\infty^p \int_{I_\alpha} \left( \iint_{\mathbb{D} \setminus D_\eta(0)} dA(w) \right)^{\frac{p}{2}} dm(\zeta) = \|g\|_\infty^p m(I_\alpha) (1-\eta^2)^{\frac{p}{2}}.$$

In order to estimate  $I_4$  we have first to estimate  $\iint_{\Gamma_{\frac{1}{2}}(\zeta)} |\psi'_\alpha(z)|^2 dA(z)$ , when  $\zeta \in \mathbb{T} \setminus I_\alpha$ .

Without loss of generality we may assume that  $\alpha \in [R_0, 1)$ . For  $j \in \mathbb{N}, j \geq 2$ , we define  $r_j = 1 - \frac{1}{2^j}$  and consider the sets  $\Omega_1 = E(0; \frac{1}{2})$  and  $\Omega_j = (E(0; r_j) \setminus E(0; r_{j-1})) \cap \Gamma_{\frac{1}{2}}(\zeta)$ .

Then we have that  $\Gamma_{\frac{1}{2}}(\zeta) = \bigcup_{j=1}^{+\infty} \Omega_j$  and  $A(\Omega_j) \asymp \frac{1}{4^j}$ , when  $j \geq 1$ . We fix  $z_j \in \Omega_j$  such

that  $Arg(z_j) = Arg(\zeta)$ . Then, if  $z \in \Omega_j$ , we have  $|1 - \alpha z| \asymp |\frac{1}{\alpha} - z| \asymp |\frac{1}{\alpha} - z_j|$ . Also we have that  $|\frac{1}{\alpha} - z_j| \asymp |\frac{1}{2^j} + 1 - |\alpha| + |Arg(\zeta)|$ . In all these relations, the underlying constants are absolute. If  $\zeta \in \mathbb{T} \setminus I_\alpha$ , then  $a \notin \Gamma_{\frac{1}{2}}(\zeta)$  which means that  $1 - |\alpha| < |Arg(\zeta)|$ , so we have that  $|\frac{1}{\alpha} - z_j| \asymp |\frac{1}{2^j} + |Arg(\zeta)||$ . There is some  $j_0$  so that  $\frac{1}{2^{j_0}} \leq |Arg(\zeta)| \leq \frac{1}{2^{j_0-1}}$ . For  $j < j_0$  we have  $|Arg(\zeta)| < \frac{1}{2^j}$  which implies that  $|\frac{1}{\alpha} - z_j| \asymp \frac{1}{2^j}$  and for  $j > j_0$  we have  $|Arg(\zeta)| > \frac{1}{2^j}$  which implies that  $|\frac{1}{\alpha} - z_j| \asymp |Arg(\zeta)|$ . Therefore

$$\begin{aligned}
\iint_{\Gamma_{\frac{1}{2}}(\zeta)} |\psi'_\alpha(z)|^2 dA(z) &= \iint_{\Omega_1} \frac{(1 - |\alpha|^2)^2}{|1 - \alpha z|^4} dA(z) + \sum_{j=2}^{+\infty} \iint_{\Omega_j} \frac{(1 - |\alpha|^2)^2}{|1 - \alpha z|^4} dA(z) \\
&\asymp (1 - |\alpha|^2)^2 + \sum_{j=2}^{j_0} \iint_{\Omega_j} \frac{(1 - |\alpha|^2)^2}{|\frac{1}{\alpha} - z_j|^4} dA(z) + \sum_{j=j_0}^{+\infty} \iint_{\Omega_j} \frac{(1 - |\alpha|^2)^2}{|\frac{1}{\alpha} - z_j|^4} dA(z) \\
&\asymp (1 - |\alpha|^2)^2 + \sum_{j=2}^{j_0} A(\Omega_j) (1 - |\alpha|^2)^2 (2^j)^4 + \sum_{j=j_0}^{+\infty} A(\Omega_j) \frac{(1 - |\alpha|^2)^2}{|Arg(\zeta)|^4} \\
&\asymp (1 - |\alpha|^2)^2 + (1 - |\alpha|^2)^2 \sum_{j=2}^{j_0} \frac{1}{4^j} 16^j + \frac{(1 - |\alpha|^2)^2}{|Arg(\zeta)|^4} \sum_{j=j_0}^{+\infty} \frac{1}{4^j}.
\end{aligned}$$

But  $\sum_{j=2}^{j_0} 4^j \asymp 4^{j_0} \asymp \frac{1}{|Arg(\zeta)|^2}$  and  $\sum_{j=j_0}^{+\infty} \frac{1}{4^j} \asymp \frac{1}{4^{j_0}} \asymp |Arg(\zeta)|^2$ . Therefore

$$\iint_{\Gamma_{\frac{1}{2}}(\zeta)} |\psi'_\alpha(z)|^2 dA(z) \asymp (1 - |\alpha|^2)^2 + \frac{(1 - |\alpha|^2)^2}{|Arg(\zeta)|^2}.$$

Since  $\alpha$  is positive, there is  $\phi_0$  such that  $\mathbb{T} \setminus I_\alpha = [\phi_0, 2\pi - \phi_0]$  and  $\phi_0 \asymp m(I_\alpha)$ . Therefore

$$\begin{aligned}
I_4 &\asymp \int_{\phi_0}^{\pi} (1 - |\alpha|^2)^p d\phi + \int_{\phi_0}^{\pi} \frac{(1 - |\alpha|^2)^p}{\phi^p} d\phi \\
&\asymp (1 - |\alpha|^2)^p + \frac{(1 - |\alpha|^2)^p}{\phi_0^{p-1}} \\
&\asymp (1 - |\alpha|^2)^p + \frac{(1 - |\alpha|^2)^p}{m(I_\alpha)^{p-1}}.
\end{aligned}$$

Substituting the estimates for  $I_1, I_2, I_3, I_4$  in (15), we get

$$\begin{aligned}
1 - |\alpha| &\leq C \left[ \|g\|_\infty^p m(I_\alpha) \frac{1}{(1 - \eta^2)^p} \left( \frac{A(G_c \cap D_\eta(\alpha))}{A(D_\eta(\alpha))} \right)^{\frac{p}{2}} + c^p m(I_\alpha) \right. \\
&\quad \left. + \|g\|_\infty^p m(I_\alpha) (1 - \eta^2)^{\frac{p}{2}} + (1 - |\alpha|^2)^p + \frac{(1 - |\alpha|^2)^p}{m(I_\alpha)^{p-1}} \right].
\end{aligned}$$

Using (13) we get

$$\begin{aligned}
1 - |\alpha| &\leq C \left[ \|g\|_\infty^p \frac{1 - |\alpha|}{(1 - \eta)^{\frac{1}{2}}} \frac{1}{(1 - \eta^2)^p} \left( \frac{A(G_c \cap D_\eta(\alpha))}{A(D_\eta(\alpha))} \right)^{\frac{p}{2}} \right. \\
&\quad + c^p \frac{1 - |\alpha|}{(1 - \eta)^{\frac{1}{2}}} + \|g\|_\infty^p \frac{1 - |\alpha|}{(1 - \eta)^{\frac{1}{2}}} (1 - \eta^2)^{\frac{p}{2}} \\
&\quad \left. + (1 - |\alpha|^2)(1 - \eta^2)^{p-1} + (1 - |\alpha|^2)(1 - \eta)^{\frac{p-1}{2}} \right].
\end{aligned}$$

Thus

$$\begin{aligned}
C &\leq \|g\|_\infty^p \frac{1}{(1 - \eta)^{\frac{2p+1}{2}}} \left( \frac{A(G_c \cap D_\eta(\alpha))}{A(D_\eta(\alpha))} \right)^{\frac{p}{2}} + \frac{c^p}{(1 - \eta)^{\frac{1}{2}}} \\
&\quad + \|g\|_\infty^p (1 - \eta)^{\frac{p-1}{2}} + (1 - \eta)^{p-1} + (1 - \eta)^{\frac{p-1}{2}}.
\end{aligned}$$

Choose  $\eta$  close enough to 1 so that  $\|g\|_\infty^p (1 - \eta)^{\frac{p-1}{2}} + (1 - \eta)^{p-1} + (1 - \eta)^{\frac{p-1}{2}} < \frac{C}{4}$  and then set  $C_\eta = \frac{1}{(1 - \eta)^{\frac{1}{2}}}$ . We have that

$$\frac{3C}{4} \leq \frac{\|g\|_\infty^p}{C_\eta^{2p+1}} \left( \frac{A(G_c \cap D_\eta(\alpha))}{A(D_\eta(\alpha))} \right)^{\frac{p}{2}} + \frac{c^p}{C_\eta^{\frac{1}{2}}}.$$

Choose  $c$  small enough so that  $\frac{c^p}{C_\eta^{\frac{1}{2}}} < \frac{C}{4}$ . Then

$$\frac{C}{2} \leq \frac{\|g\|_\infty^p}{C_\eta^{2p+1}} \left( \frac{A(G_c \cap D_\eta(\alpha))}{A(D_\eta(\alpha))} \right)^{\frac{p}{2}}$$

and finally

$$\left( \frac{CC_\eta^{2p+1}}{2\|g\|_\infty^p} \right)^{\frac{2}{p}} \leq \frac{A(G_c \cap D_\eta(\alpha))}{A(D_\eta(\alpha))}$$

or

$$A(G_c \cap D_\eta(\alpha)) \geq \delta A(D_\eta(\alpha)),$$

for every  $\alpha$  with  $R_0 \leq |\alpha| < 1$ .

**Case (b)**  $0 \leq |\alpha| \leq R_0$ : There exists  $\eta_1$ , depending only on  $\eta$ , such that  $D_\eta(R_0) \subseteq D_{\eta_1}(0)$ . Take  $\alpha'$  so that  $|\alpha'| = R_0$  and  $\text{Arg}(\alpha') = \text{Arg}(\alpha)$ . Then  $D_\eta(\alpha') \subseteq D_{\eta_1}(\alpha)$ . Set  $\eta_2 = \max\{\eta, \eta_1\}$ . Then from case (a) for  $\alpha'$  we have

$$\begin{aligned}
A(G_c \cap D_{\eta_2}(\alpha)) &\geq A(G_c \cap D_{\eta_1}(\alpha)) \geq A(G_c \cap D_\eta(\alpha')) \\
&\geq \delta A(D_\eta(\alpha')) \geq C\delta A(D_{\eta_1}(\alpha)) \geq C\delta A(D_{\eta_2}(\alpha)),
\end{aligned}$$

where the constants  $C > 0$  depend only on  $\eta$ .

Moreover, when  $R_0 \leq |\alpha| < 1$ , we have

$$A(G_c \cap D_{\eta_2}(\alpha)) \geq A(G_c \cap D_\eta(\alpha)) \geq \delta A(D_\eta(\alpha)) \geq C\delta A(D_{\eta_2}(\alpha)),$$

where the constant  $C > 0$  depends only on  $\eta$ . So, we have proved that there are  $\eta_2 \in (0, 1)$ ,  $c > 0$  and  $C > 0$  such that

$$A(G_c \cap D_{\eta_2}(\alpha)) \geq CA(D_{\eta_2}(\alpha)),$$

for every  $\alpha \in \mathbb{D}$ , which is what we had to prove.  $\square$

**Remark 1.** As we can observe, the proof of the implication (ii)  $\Rightarrow$  (i) in theorem 2.2 is valid even in the case  $p = 1$ .

### 3. Closed range integral operators on BMOA space

Let denote as  $BMOA_0$  the space  $BMOA/\mathbb{C}$ . In [1], A. Anderson posed the question of finding a necessary and sufficient condition for the operator  $S_g$  to have closed range on  $BMOA_0$ . Next, we answer this question, proving that conditions (ii) and (iii) of theorem 2.2, for  $H_0^p$ , are also necessary and sufficient for the integral operator  $S_g$  to have closed range on  $BMOA_0$ .

Let  $z_0 \in \mathbb{D}$ . The point evaluation functional of the derivative, on  $BMOA$ , induced by  $z_0$ , is defined as  $\Lambda_{z_0}f = f'(z_0)$ ,  $f \in BMOA$ . It is easy to check that  $\Lambda_{z_0}$  is bounded on  $BMOA$ . Therefore, using Theorem 2.2 and Corollary 2.3 in [1], we conclude that the operator  $S_g : BMOA_0 \rightarrow BMOA_0$  is bounded if and only if  $g \in H^\infty$ . So, we consider  $g \in H^\infty$  and set again  $G_c = \{z \in \mathbb{D} : |g(z)| > c\}$ .

The following theorem is the main result of this section.

**Theorem 3.1.** *Let  $g \in H^\infty$ . Then the following are equivalent:*

- (i) *The operator  $S_g : BMOA_0 \rightarrow BMOA_0$  has closed range*
- (ii) *There exist  $c > 0$ ,  $\delta > 0$  and  $\eta \in (0, 1)$  such that*

$$A(G_c \cap D_\eta(a)) \geq \delta A(D_\eta(a)) \tag{16}$$

for all  $a \in \mathbb{D}$ .

Recall that the weighted Bergman space  $\mathbb{A}_\gamma^p$ ,  $\gamma > -1$ , is defined as the set of all analytic functions  $f$  in  $\mathbb{D}$  such that

$$\iint_{\mathbb{D}} |f(z)|^p (1 - |z|^2)^\gamma dA(z) < \infty.$$

We will make use of the following theorem of D. Luecking (see [2]).

**Theorem 3.2.** *Let  $p \geq 1$ ,  $\gamma > -1$  and measurable  $G \subseteq \mathbb{D}$ . The following assertions are equivalent.*

- (i) *There exists  $C > 0$  such that*

$$\iint_G |f(z)|^p (1 - |z|^2)^\gamma dA(z) \geq C \iint_{\mathbb{D}} |f(z)|^p (1 - |z|^2)^\gamma dA(z) \tag{17}$$

for every  $f \in \mathbb{A}_\gamma^p$ .

(ii) There exist  $c > 0$ ,  $\delta > 0$  and  $\eta \in (0, 1)$  such that

$$A(G \cap D_\eta(a)) \geq \delta A(D_\eta(a))$$

for all  $a \in \mathbb{D}$ .

In the proof of theorem 3.1, we will use the fact that  $\log \frac{1}{|z|} \asymp 1 - |z|^2$ , when  $0 < \delta \leq |z| < 1$ , where  $\delta$  is fixed but arbitrary.

**Proof of theorem 3.1.** (ii)  $\Rightarrow$  (i) If (16) holds then, because of theorem 3.2, (17) also holds for  $G = G_c$ . For  $\beta \in \mathbb{D}$ ,  $z \in \mathbb{D}$  and  $f \in BMOA_0$ , we consider the function  $h_\beta(z) = \frac{(1-|\beta|^2)^{\frac{1}{2}}}{1-\bar{\beta}z} f'(z)$ . It's easy to see that if  $f \in BMOA_0$  then  $h_\beta \in \mathbb{A}_1^2$ . Indeed

$$\begin{aligned} \|h_\beta\|_{\mathbb{A}_1^2}^2 &= \iint_{\mathbb{D}} \frac{1-|\beta|^2}{|1-\bar{\beta}z|^2} |f'(z)|^2 (1-|z|^2) dA(z) \\ &\leq C \iint_{\mathbb{D}} \frac{1-|\beta|^2}{|1-\bar{\beta}z|^2} |f'(z)|^2 \log \frac{1}{|z|} dA(z) \leq \|f\|_{BMOA_0}^2 < \infty. \end{aligned}$$

Let  $\beta \in \mathbb{D}$ . We have that

$$\begin{aligned} \|S_g f\|_{BMOA_0}^2 &= \sup_{z_0 \in \mathbb{D}} \iint_{\mathbb{D}} \frac{1-|z_0|^2}{|1-\bar{z}_0 z|^2} |(S_g f(z))'|^2 \log \frac{1}{|z|} dA(z) \\ &= \sup_{z_0 \in \mathbb{D}} \iint_{\mathbb{D}} \frac{1-|z_0|^2}{|1-\bar{z}_0 z|^2} |f'(z)|^2 |g(z)|^2 \log \frac{1}{|z|} dA(z) \\ &\geq \iint_{\mathbb{D}} \frac{1-|\beta|^2}{|1-\bar{\beta}z|^2} |f'(z)|^2 |g(z)|^2 \log \frac{1}{|z|} dA(z) \\ &\geq c^2 \iint_{G_c} \frac{1-|\beta|^2}{|1-\bar{\beta}z|^2} |f'(z)|^2 \log \frac{1}{|z|} dA(z) \\ &= c^2 \iint_{G_c} |h_\beta(z)|^2 \log \frac{1}{|z|} dA(z) \\ &\geq C \iint_{G_c} |h_\beta(z)|^2 (1-|z|^2) dA(z) \\ &\geq C \iint_{\mathbb{D}} |h_\beta(z)|^2 (1-|z|^2) dA(z), \end{aligned}$$

where the last inequality is justified by theorem 3.2. So

$$\|S_g f\|_{BMOA_0}^2 \geq C \iint_{\mathbb{D}} \frac{1-|\beta|^2}{|1-\bar{\beta}z|^2} |f'(z)|^2 \log \frac{1}{|z|} dA(z).$$

Taking the supremum over  $\beta \in \mathbb{D}$  in the last relation we get

$$\|S_g f\|_{BMOA_0}^2 \geq C \|f\|_{BMOA_0}^2.$$

(i)  $\Rightarrow$  (ii) If  $S_g$  has closed range then there exist  $C_1 > 0$  such that for every  $f \in BMOA_0$  we have

$$\|S_g f\|_{BMOA_0}^2 \geq C_1 \|f\|_{BMOA_0}^2.$$

For  $\alpha \in \mathbb{D}$ , if we set  $f = \psi_\alpha - \alpha$  in the last inequality, just as in the case of Hardy spaces and observe that  $\|\psi_\alpha - \alpha\|_{BMOA} \asymp 1$  and  $\frac{(1-|\beta|^2)(1-|z|^2)}{|1-\bar{\beta}z|^2} < 1$ , for every  $z, \beta \in \mathbb{D}$ , then we have

$$\begin{aligned} C_1 &\leq \|S_g(\psi_\alpha - \alpha)\|_{BMOA_0}^2 \\ &= \sup_{\beta \in \mathbb{D}} \iint_{\mathbb{D}} \frac{1-|\beta|^2}{|1-\bar{\beta}z|^2} |(S_g(\psi_\alpha - \alpha)(z))'|^2 \log \frac{1}{|z|} dA(z) \\ &\leq C \sup_{\beta \in \mathbb{D}} \iint_{\mathbb{D}} \frac{1-|\beta|^2}{|1-\bar{\beta}z|^2} |\psi'_\alpha(z)|^2 |g(z)|^2 (1-|z|^2) dA(z) \\ &\leq C \iint_{\mathbb{D}} |\psi'_\alpha(z)|^2 |g(z)|^2 dA(z) \\ &\leq C \left[ \|g\|_\infty^2 \iint_{G_c \cap D_\eta(\alpha)} \frac{(1-|\alpha|^2)^2}{|1-\bar{\alpha}z|^4} dA(z) + c^2 \iint_{D_\eta(\alpha) \setminus G_c} |\psi'_\alpha(z)|^2 dA(z) \right. \\ &\quad \left. + \|g\|_\infty^2 \iint_{\mathbb{D} \setminus D_\eta(\alpha)} |\psi'_\alpha(z)|^2 dA(z) \right] \\ &\leq C \left[ \|g\|_\infty^2 \iint_{G_c \cap D_\eta(\alpha)} \frac{1}{(1-|\alpha|^2)^2} dA(z) + c^2 \iint_{\mathbb{D}} |\psi'_\alpha(z)|^2 dA(z) \right. \\ &\quad \left. + \|g\|_\infty^2 \iint_{\mathbb{D} \setminus D_\eta(\alpha)} |\psi'_\alpha(z)|^2 dA(z) \right] \\ &= C \left[ \|g\|_\infty^2 \frac{A(G_c \cap D_\eta(\alpha))}{(1-|\alpha|^2)^2} + c^2 \iint_{\mathbb{D}} dA(w) + \|g\|_\infty^2 \iint_{\mathbb{D} \setminus D_\eta(0)} dA(w) \right] \\ &\leq C \left[ C' \|g\|_\infty^2 \frac{A(G_c \cap D_\eta(\alpha))}{A(D_\eta(\alpha))} + c^2 + \|g\|_\infty^2 (1-\eta^2) \right], \end{aligned}$$

where  $C'$  depends only on  $\eta$  and  $C$  is absolute. Therefore

$$C_1 \leq C' \|g\|_\infty^2 \frac{A(G_c \cap D_\eta(\alpha))}{A(D_\eta(\alpha))} + c^2 + \|g\|_\infty^2 (1-\eta^2).$$

First, we choose  $\eta$  close enough to 1 so that  $\|g\|_\infty^2 (1-\eta^2) < \frac{C_1}{4}$  and  $c$  small enough

so that  $c < \frac{C_1}{4}$ . So

$$A(G_c \cap D_\eta(\alpha)) \geq \frac{C}{2\|g\|_\infty^2} A(D_\eta(\alpha)) = \delta A(D_\eta(\alpha)),$$

where  $C$  depends only on  $\eta$ . □

**Remark 2.** The  $Q_p$  space,  $0 < p < \infty$ , is defined as the set of all analytic functions  $f$  in  $\mathbb{D}$  for which

$$\sup_{\beta \in \mathbb{D}} \iint_{\mathbb{D}} \frac{(1 - |\beta|^2)^p}{|1 - \beta z|^{2p}} |f'(z)|^2 (1 - |z|^2)^p dA(z) < \infty.$$

Let denote as  $Q_{p,0}$  the space  $Q_p/\mathbb{C}$ . For  $\beta, z \in \mathbb{D}$  and  $f \in Q_{p,0}$ , we consider the functions  $h_\beta(z) = \frac{(1 - |\beta|^2)^{\frac{p}{2}}}{(1 - \beta z)^p} f'(z)$ . It's easy to see that if  $f \in Q_{p,0}$  then  $h_\beta \in \mathbb{A}_p^2$  and using similar arguments as in the proof of theorem 3.1, we can prove that (16) is also necessary and sufficient for  $S_g$  to have closed range on  $Q_{p,0}$  ( $0 < p < \infty$ ).

#### 4. Closed range integral operators on Besov spaces

Let denote as  $B_0^p$  the space  $B^p/\mathbb{C}$ . With similar arguments as in the case of *BMOA* space we can see that the operator  $S_g : B_0^p \rightarrow B_0^p$  ( $1 < p < \infty$ ) is bounded if and only if  $g \in H^\infty$ . So, we consider  $g \in H^\infty$  and  $G_c = \{z \in \mathbb{D} : |g(z)| > c\}$ . We will prove that condition (16) is also necessary and sufficient for the operator  $S_g$  to have closed range on  $B_0^p$ . For the sufficiency, we observe that, if  $f \in B^p$  then  $f' \in \mathbb{A}_{p-2}^p$ , the weighted Bergman space defined in the previous section, so we can use theorem 3.2. We have

$$\begin{aligned} \|S_g f\|_{B_0^p}^p &= \iint_{\mathbb{D}} |(S_g f(z))'|^p (1 - |z|^2)^{p-2} dA(z) \\ &\geq \iint_{G_c} |f'(z)|^p |g(z)|^p (1 - |z|^2)^{p-2} dA(z) \\ &\geq c^p \iint_{G_c} |f'(z)|^p (1 - |z|^2)^{p-2} dA(z) \\ &\geq C \iint_{\mathbb{D}} |f'(z)|^p (1 - |z|^2)^{p-2} dA(z) \\ &= C \|f\|_{B_0^p}^p, \end{aligned}$$

where the last inequality is justified by theorem 3.2. So  $S_g$  has closed range on  $B_0^p$ .

If  $S_g$  has closed range on  $B_0^p$  then there exist  $C_1 > 0$  such that for every  $f \in B_0^p$  we have

$$\|S_g f\|_{B_0^p}^p \geq C_1 \|f\|_{B_0^p}^p.$$



For  $\alpha \in \mathbb{D}$ , if we set  $f = f_\alpha = \frac{(1-|\alpha|^2)^{\frac{2}{p}}}{\frac{2\bar{\alpha}}{p}(1-\bar{\alpha}z)^{\frac{2}{p}}} - \frac{(1-|\alpha|^2)^{\frac{2}{p}}}{\frac{2\alpha}{p}}$  in the last inequality, just as in the case of *BMOA*, and observe that  $\|f_\alpha\|_{B_0^p} \asymp 1$  and  $|f'_\alpha(z)| = \frac{(1-|\alpha|^2)^{\frac{2}{p}}}{|1-\bar{\alpha}z|^{\frac{2}{p}+1}}$ , then we have

$$\begin{aligned}
C_1 &\leq \|S_g f_\alpha\|_{B_0^p}^p = \iint_{\mathbb{D}} |f'_\alpha(z)|^p |g(z)|^p (1-|z|)^{p-2} dA(z) \\
&\leq \|g\|_\infty^p \iint_{G_c \cap D_\eta(\alpha)} \frac{(1-|\alpha|^2)^2}{|1-\bar{\alpha}z|^{2+p}} (1-|z|)^{p-2} dA(z) + c^p \iint_{D_\eta(\alpha) \setminus G_c} |f'_\alpha(z)|^p (1-|z|)^{p-2} dA(z) \\
&\quad + \|g\|_\infty^p \iint_{\mathbb{D} \setminus D_\eta(\alpha)} \frac{(1-|\alpha|^2)^2}{|1-\bar{\alpha}z|^{2+p}} (1-|z|)^{p-2} dA(z) \\
&\leq \|g\|_\infty^p \iint_{G_c \cap D_\eta(\alpha)} \frac{1}{(1-|\alpha|^2)^2} dA(z) + c^p \iint_{\mathbb{D}} |f'_\alpha(z)|^p (1-|z|)^{p-2} dA(z) \\
&\quad + \|g\|_\infty^p \iint_{\mathbb{D} \setminus D_\eta(0)} \frac{(1-|\alpha|^2)^2}{|1-\bar{\alpha}\psi_\alpha(w)|^{2+p}} (1-|\psi_\alpha(w)|)^{p-2} |\psi'_\alpha(w)|^2 dA(w) \\
&= \|g\|_\infty^p \iint_{G_c \cap D_\eta(\alpha)} \frac{1}{(1-|\alpha|^2)^2} dA(z) + c^p \|f_\alpha\|_{B_0^p}^p + \|g\|_\infty^p \iint_{\mathbb{D} \setminus D_\eta(0)} \frac{(1-|w|^2)^{p-2}}{|1-\bar{\alpha}w|^{p-2}} dA(w) \\
&\leq \|g\|_\infty^p \frac{A(G_c \cap D_\eta(\alpha))}{(1-|\alpha|^2)^2} + c^p \|f_\alpha\|_{B_0^p}^p + \|g\|_\infty^p \iint_{\mathbb{D} \setminus D_\eta(0)} dA(w) \\
&\leq C' \|g\|_\infty^p \frac{A(G_c \cap D_\eta(\alpha))}{A(D_\eta(\alpha))} + Cc^p + \|g\|_\infty^p (1-\eta^2),
\end{aligned}$$

where  $C'$  depends only on  $\eta$  and  $C$  is absolute. So we have

$$C_1 \leq C' \|g\|_\infty^p \frac{A(G_c \cap D_\eta(\alpha))}{A(D_\eta(\alpha))} + Cc^p + \|g\|_\infty^p (1-\eta^2).$$

Choosing  $\eta$  close enough to 1 so that  $\|g\|_\infty^p (1-\eta^2) < \frac{C_1}{4}$ , and  $c$  small enough so that  $Cc^p < \frac{C_1}{4}$ , we get

$$A(G_c \cap D_\eta(\alpha)) \geq \frac{C_1}{2C' \|g\|_\infty^p} A(D_\eta(\alpha)) = \delta A(D_\eta(\alpha)).$$

## Acknowledgements

Many thanks to Prof. Michael Papadimitrakis for discussions about the mathematical content of this paper. His contribution was essential in order for it to take its final form.

## References

- [1] ANDERSON, A. *Some Closed Range Integral Operators on Spaces of Analytic Functions*, Integr. Equ. Oper. Theory 69, 87–99 (2011)
- [2] LUECKING D.H., *Inequalities on Bergman spaces*, Illinois J. Math. 25, 1–11 (1981)
- [3] PAVLOVIC M., *Function Classes on the Unit Disc*, Walter de Gruyter GmbH, Berlin/Boston, (2014)