

ORTHOGONALITY IN BANACH SPACES VIA PROJECTIVE TENSOR PRODUCT

KOUSIK DHARA, NARAYAN RAKSHIT, JAYDEB SARKAR, AND ARYAMAN SENSARMA

ABSTRACT. Let X be a complex Banach space and $x, y \in X$. By definition, we say that x is Birkhoff-James orthogonal to y if $\|x + \lambda y\|_X \geq \|x\|_X$ for all $\lambda \in \mathbb{C}$. We prove that x is Birkhoff-James orthogonal to y if and only if there exists a semi-inner product φ on X such that $\|\varphi\| = 1$, $\varphi(x, x) = \|x\|^2$ and $\varphi(x, y) = 0$. A similar result holds for C^* -algebras. A key point in our approach to orthogonality is the representations of bounded bilinear maps via projective tensor product spaces.

1. INTRODUCTION

This paper deals with the notion of Birkhoff-James orthogonality [2, 6, 7] for vectors in Banach spaces via projective tensor product of Banach spaces. Let X be a normed linear space over the scalar field \mathbb{K} , where \mathbb{K} is either \mathbb{R} or \mathbb{C} . Suppose $x, y \in X$. We say that x is *Birkhoff-James orthogonal* (or simply *orthogonal*) to y if

$$\|x + \lambda y\|_X \geq \|x\|_X,$$

for all $\lambda \in \mathbb{K}$. We denote this by $x \perp_B y$. If X is a Hilbert space, then this notion coincides with the usual orthogonality, that is, $x \perp_B y$ if and only if

$$\langle x, y \rangle_X = 0.$$

The objective of this note is to present an abstract characterization of orthogonality of vectors in Banach spaces over \mathbb{K} . We are motivated by Bhatia and Šemrl's investigation [3, Theorem 1.1 and Remark 3.1] of orthogonality in the setting of bounded linear operators on Hilbert spaces:

Theorem 1.1. (*Bhatia and Šemrl*) *Let S and T be bounded operators on a Hilbert space \mathcal{H} .*

(i) *Then $S \perp_B T$ if and only if there exists a sequence $\{h_n\}$ of unit vectors such that $\|Sh_n\| \rightarrow \|S\|$, and $\langle Sh_n, Th_n \rangle \rightarrow 0$ as $n \rightarrow \infty$.*

(ii) *If we additionally assume that \mathcal{H} is a finite dimensional Hilbert space, then $S \perp_B T$ if and only if there exists a unit vector $h \in \mathcal{H}$ such that*

$$\|Sh\| = \|S\| \quad \text{and} \quad \langle Sh, Th \rangle = 0.$$

2010 *Mathematics Subject Classification.* 46B28, 47A30, 46B20, 46B22, 47B01, 47L05.

Key words and phrases. Orthogonality, projective tensor product, semi-inner product.

We treat the above Bhatia-Šemrl result (more specifically, part (ii) of Theorem 1.1) as a paradigm and examine orthogonality of vectors in Banach spaces in terms of semi-inner products. Our main result, in the setting of Banach spaces, is the following:

Theorem 1.2. *Let X be a Banach space, $x, y \in X$, and $x \neq 0$. Then $x \perp_B y$ if and only if there exists a semi-inner product $\varphi : X \oplus_\infty X \rightarrow \mathbb{K}$ such that $\|\varphi\| = 1$ and*

$$\varphi(x, x) = \|x\|^2 \quad \text{and} \quad \varphi(x, y) = 0.$$

As an immediate consequence of the above, we also prove an orthogonality result in the setting of C^* -algebras (see Corollary 2.2). Here, for a pair of Banach spaces X and Y , we define $X \oplus_\infty Y$ to be the Banach space

$$X \times Y = \{(x, y) : x \in X, y \in Y\},$$

with the norm

$$\|(x, y)\|_\infty = \max\{\|x\|_X, \|y\|_Y\},$$

for all $x \in X$ and $y \in Y$. Also recall that a semi-inner product on a vector space V is scalar-valued function $\varphi : V \times V \rightarrow \mathbb{K}$ such that for all $x, y, z \in V$ and $\alpha, \beta \in \mathbb{K}$, we have

- (a) $\varphi(\alpha x + \beta y, z) = \alpha\varphi(x, z) + \beta\varphi(y, z)$,
- (b) $\varphi(x, y) = \overline{\varphi(y, x)}$,
- (c) $\varphi(x, x) \geq 0$.

It is easy to check that, if φ is a semi-inner product, then φ is an inner product if and only if $\varphi(x, x) = 0$ implies $x = 0$.

The main ingredients of our approach to the orthogonality problem are: (1) Bhatia and Šemrl's orthogonality of bounded linear operators on infinite-dimensional Hilbert spaces (part (i) of Theorem 1.1 above), (2) projective tensor product techniques (see Theorem 1.3 below), and (3) some standard Banach space techniques (like Banach-Alaoglu theorem).

Before we proceed with the main content of the paper, let us shortly review the existing literature on orthogonality. The notion of Birkhoff-James orthogonality is an active research area. In fact, Bhatia and Šemrl's paper [3] on orthogonality of operators on Hilbert spaces has stimulated extensive research for the past two decades. For instance, orthogonality of a pair of compact operators acting on a reflexive Banach space to a normed linear space has been studied by Sain, Paul and Mal [13]. Their investigation involves finer geometric Banach space techniques. Also see [11] on orthogonality of linear operators on finite dimensional Banach spaces, [12] on norm attainment and orthogonality, and [9] on approximate orthogonality. More recent advances can be found, for instance, in the quickly growing literature [1, 4, 5, 8, 14, 15] (also see the references therein).

Given Banach spaces X and Y , we denote the Banach space of all bounded linear operators from X to Y by $\mathcal{B}(X, Y)$, and we let X_1 denote the closed unit ball in X . If $Y = X$, then we write $\mathcal{B}(X)$. A bilinear (sesquilinear) map $B : X \times Y \rightarrow Z$ is said to be bounded if there exists $M > 0$ such that

$$\|B(x, y)\| \leq M \quad (x \in X_1, y \in Y_1).$$

We denote the Banach space of all bounded bilinear (sesquilinear) maps from $X \times Y$ to Z by $\text{Bil}(X \times Y, Z)$ ($\text{Ses}(X \times Y, Z)$). Here

$$\|B\| = \sup\{\|B(x, y)\| : x \in X_1, y \in Y_1\},$$

for all $B \in \text{Bil}(X \times Y, Z)$ ($B \in \text{Ses}(X \times Y, Z)$). As a tool for the proof of the main result, we use the notion of projective tensor product. The projective tensor product $X \hat{\otimes}_\pi Y$ of Banach spaces X and Y is the completion of the algebraic tensor product $X \otimes Y$ under the projective norm

$$\|u\|_\pi = \inf \left\{ \sum_{i=1}^n \|x_i\| \|y_i\| : u = \sum_{i=1}^n x_i \otimes y_i \right\}.$$

Our key point is the following result [10, Theorem 2.9] concerning representations of bounded bilinear maps via projective tensor product spaces.

Theorem 1.3. *Let X, Y and Z be Banach spaces, and let $B \in \text{Bil}(X \times Y, Z)$. Then there exists a unique $\tilde{B} \in \mathcal{B}(X \hat{\otimes}_\pi Y, Z)$ such that*

$$\tilde{B}(x \otimes y) = B(x, y) \quad (x \in X, y \in Y).$$

Moreover, the correspondence $B \longleftrightarrow \tilde{B}$ is an isometric isomorphism between $\text{Bil}(X \times Y, Z)$ and $\mathcal{B}(X \hat{\otimes}_\pi Y, Z)$.

We fix some more notation that we will use from now on. Given a vector space X over \mathbb{K} , we denote by \overline{X} the complex conjugate vector space of X . That is, $\overline{X} = X$ with the same additive group structures, but with the scalar multiplication \star defined by

$$(1.1) \quad \alpha \star x = \overline{\alpha}x,$$

for all $\alpha \in \mathbb{K}$ and $x \in \overline{X}$. Clearly, if $(X, \|\cdot\|)$ is normed linear space over \mathbb{K} , then there is an anti-linear isometric isomorphism between $(X, \|\cdot\|)$ and $(\overline{X}, \|\cdot\|)$. If \mathcal{H} is a Hilbert space, then $\overline{\mathcal{H}}$ is identified with the dual (the space of continuous linear functionals) of \mathcal{H} by Riesz representation theorem.

Note that the space $\text{Bil}(X \oplus_\infty \overline{X}, \mathbb{K})$ is isometrically isomorphic to the space $\text{Ses}(X \oplus_\infty X, \mathbb{K})$. The correspondence is given by

$$(1.2) \quad \psi(x, y) = \varphi(x, y),$$

where $\psi \in \text{Bil}(X \oplus_\infty X, \mathbb{K})$, $\varphi \in \text{Ses}(X \oplus_\infty X, \mathbb{K})$, $x \in X$ and $y \in Y$.

2. MAIN RESULTS

We begin with the proof of Theorem 1.2. But before we do so, let us recall the classical Banach–Alaoglu theorem: Let X be a Banach space over \mathbb{K} . Then the closed unit ball $(X^*)_1$ of the dual X^* is compact with respect to the weak* topology on X^* .

Proof of Theorem 1.2: Suppose $x \perp_B y$. Note that the Banach space X is isometrically isomorphic to a closed subspace of $C((X^*)_1)$, where $(X^*)_1$ endowed with the weak* topology is a compact set. Here the correspondence is given by the formula $X \ni u \mapsto \hat{u}$, where

$$\hat{u}(f) = f(u) \quad (f \in (X^*)_1).$$

Next we note that the commutative Banach algebra $C((X^*)_1)$ is isometrically isomorphic to a closed subspace of $\mathcal{B}(L^2(\mu))$ for some σ -finite measure μ . Here the correspondence is given by the formula $C((X^*)_1) \ni g \mapsto M_g$, where $M_g : L^2(\mu) \rightarrow L^2(\mu)$ is the multiplication operator defined by

$$M_g(h) = gh \quad (h \in L^2(\mu)).$$

Using the above identifications, we have $M_{\hat{x}}, M_{\hat{y}} \in \mathcal{B}(L^2(\mu))$ and

$$M_{\hat{x}} \perp_B M_{\hat{y}}.$$

By the infinite dimensional part of Theorem 1.1, there exists a sequence $\{h_n\}$ of unit vectors in $L^2(\mu)$ such that

$$\|M_{\hat{x}}(h_n)\|_{L^2(\mu)} \rightarrow \|M_{\hat{x}}\|_{\mathcal{B}(L^2(\mu))},$$

and

$$\langle M_{\hat{x}}(h_n), M_{\hat{y}}(h_n) \rangle_{L^2(\mu)} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

For each $n \geq 1$, define $\psi_n : X \oplus_\infty \overline{X} \rightarrow \mathbb{K}$ by

$$\psi_n(z, w) = \langle M_{\hat{z}}(h_n), M_{\hat{w}}(h_n) \rangle_{L^2(\mu)} \quad (z \in X, w \in \overline{X}).$$

Now we prove that ψ_n is bilinear. Clearly, ψ_n is linear in its first variable. Suppose $\alpha_1, \alpha_2 \in \mathbb{K}$ and $z, w_1, w_2 \in X$. Then

$$\begin{aligned} \psi_n(z, (\alpha_1 \star w_1 + \alpha_2 \star w_2)) &= \langle M_{\hat{z}}(h_n), M_{\widehat{\alpha_1 \star w_1 + \alpha_2 \star w_2}}(h_n) \rangle \\ &= \langle M_{\hat{z}}(h_n), M_{\widehat{\alpha_1 w_1 + \alpha_2 w_2}}(h_n) \rangle \\ &= \langle \hat{z}h_n, (\alpha_1 \hat{w}_1 + \alpha_2 \hat{w}_2)h_n \rangle \\ &= \alpha_1 \langle \hat{z}h_n, \hat{w}_1 h_n \rangle + \alpha_2 \langle \hat{z}h_n, \hat{w}_2 h_n \rangle, \end{aligned}$$

that is

$$\psi_n(z, (\alpha_1 \star w_1 + \alpha_2 \star w_2)) = \alpha_1 \psi_n(z, w_1) + \alpha_2 \psi_n(z, w_2),$$

and hence ψ_n is bilinear for all $n \geq 1$. To prove that ψ_n is bounded, suppose $(z, w) \in X \oplus_\infty \overline{X}$ and $\|(z, w)\|_\infty \leq 1$. Then

$$\begin{aligned} |\psi_n(z, w)| &\leq \|M_{\hat{z}}\|_{\mathcal{B}(L^2(\mu))} \|M_{\hat{w}}\|_{\mathcal{B}(L^2(\mu))} \|h_n\|_{L^2(\mu)}^2 \\ &= \|\hat{z}\|_{C((X^*)_1)} \|\hat{w}\|_{C((X^*)_1)} \\ &= \|z\|_X \|w\|_X, \end{aligned}$$

which implies that $\|\psi_n\| \leq 1$ for all n . Consequently

$$\{\psi_n\}_{n \geq 1} \subseteq \text{Bil}(X \oplus_\infty \overline{X}, \mathbb{K}).$$

Note, by Theorem 1.3, that

$$\text{Bil}(X \oplus_\infty \overline{X}, \mathbb{K}) \simeq \mathcal{B}(X \hat{\otimes}_\pi \overline{X}, \mathbb{K}).$$

Here the correspondence is given by $\text{Bil}(X \oplus_\infty \overline{X}, \mathbb{K}) \ni \psi \mapsto \tilde{\psi}$, where

$$\tilde{\psi}(z \otimes w) = \psi(z, w) \quad (z, w \in X).$$

Since $\|\psi_n\| \leq 1$, we have $\|\tilde{\psi}_n\| \leq 1$ for each n . By applying the Banach-Alaoglu theorem, one can find a subsequence $\{\tilde{\psi}_{n_k}\}$ of $\{\tilde{\psi}_n\}$ and a map $\tilde{\psi} \in \mathcal{B}(X \hat{\otimes}_\pi \overline{X}, \mathbb{K})$ with $\|\tilde{\psi}\| \leq 1$ such that

$$\tilde{\psi}_{n_k} \xrightarrow{w^*} \tilde{\psi}.$$

This yields $\psi_{n_k} \rightarrow \psi$ in the pointwise topology, that is

$$\psi_{n_k}(z, w) \rightarrow \psi(z, w) \quad (z, w \in X).$$

By isometry, $\|\psi\| \leq 1$. Observe that

$$\psi_{n_k}(x, x) = \langle M_{\hat{x}}(h_{n_k}), M_{\hat{x}}(h_{n_k}) \rangle_{L^2(\mu)} = \|M_{\hat{x}}(h_{n_k})\|_{L^2(\mu)}^2.$$

But since

$$\|M_{\hat{x}}(h_{n_k})\|_{L^2(\mu)}^2 \rightarrow \|M_{\hat{x}}\|_{\mathcal{B}(L^2(\mu))}^2 = \|\hat{x}\|^2,$$

it follows that

$$\psi_{n_k}(x, x) \rightarrow \|x\|^2,$$

for all $x \in X$. On the other hand, $\psi_{n_k}(x, x) \rightarrow \psi(x, x)$. Hence

$$\psi(x, x) = \|x\|^2 \quad (x \in X).$$

Since

$$\psi_{n_k}(x, y) = \langle M_{\hat{x}}(h_{n_k}), M_{\hat{y}}(h_{n_k}) \rangle_{L^2(\mu)} \rightarrow 0,$$

we have $\psi(x, y) = 0$. Note that if $\|\psi\| < 1$, then $\psi(x, x) < \|x\|^2$. Hence $\|\psi\| = 1$. Now observe that the identification in (1.2) provides a sesquilinear map

$$\varphi \in \text{Ses}(X \oplus_\infty X, \mathbb{K}),$$

with

$$\varphi(x, y) = \psi(x, y) \quad (x, y \in X).$$

This gives $\varphi(z, z) \geq 0$ for all $z \in X$, and $\varphi(z, w) = \overline{\varphi(w, z)}$ for all $z, w \in X$. Hence φ is a semi-inner product on X such that $\|\varphi\| = 1$, $\varphi(x, x) = \|x\|^2$ and $\varphi(x, y) = 0$.

Conversely, suppose that such a φ exists. Then for all $\lambda \in \mathbb{K}$,

$$\|x\|^2 = |\varphi(x, x + \lambda y)| \leq \|\varphi\| \|x\| \|x + \lambda y\|,$$

which implies that $\|x\| \leq \|x + \lambda y\|$, that is, $x \perp_B y$. This completes the proof of the theorem.

Remark 2.1. It is worth pointing out that besides Theorem 1.1 and Theorem 1.3, the inclusion

$$X \hookrightarrow C((X^*)_1) \hookrightarrow \mathcal{B}(L^2(\mu)),$$

also plays an important role in our proof.

Now we turn to orthogonality of elements in C^* -algebras. The following result follows from Theorem 1.2.

Corollary 2.2. *Let \mathcal{A} be a C^* -algebra and $a, b \in \mathcal{A}$ and $a \neq 0$. Then $a \perp_B b$ if and only if there exists a bilinear map $\psi : \mathcal{A} \oplus_\infty \mathcal{A} \rightarrow \mathbb{C}$ such that $\|\psi\| = 1$, $\psi(a, a^*) = \|a\|^2$ and $\psi(a, b^*) = 0$.*

Proof. Suppose $a \perp_B b$. Then by Theorem 1.2, there exists a semi-inner product $\varphi : \mathcal{A} \oplus_\infty \mathcal{A} \rightarrow \mathbb{C}$ such that $\|\varphi\| = 1$, $\varphi(a, a) = \|a\|^2$ and $\varphi(a, b) = 0$. Define $\psi : \mathcal{A} \oplus_\infty \mathcal{A} \rightarrow \mathbb{C}$ by

$$\psi(a, b) = \varphi(a, b^*).$$

Then ψ is the required map. To prove the converse, suppose such a ψ exists. Then

$$\begin{aligned} \|a\|^2 &= |\psi(a, (a + \lambda b)^*)| \\ &\leq \|\psi\| \|a\| \|(a + \lambda b)^*\| \\ &= \|a\| \|a + \lambda b\|, \end{aligned}$$

that is, $\|(a + \lambda b)\| \geq \|a\|$ for all $\lambda \in \mathbb{C}$

□

It is worth pointing out that Theorem 1.2 is also applicable to finite dimensional Banach spaces. Clearly, Theorem 1.2 and Corollary 2.2 are analogous to Bhatia and Šemrl classifications (see part (ii) of Theorem 1.1) of orthogonality of matrices on finite dimensional Hilbert spaces. On the other hand, our results uses the Bhatia and Šemrl classifications of orthogonality in the setting of infinite dimensional Hilbert spaces (see part (i) of Theorem 1.1). In addition, it is not completely clear if our results recovers the Bhatia and Šemrl classifications of orthogonality in the setting of finite matrices.

All in all, on one hand our results are valid for general Banach spaces and rather abstract, and on the other hand our approach is intimately related to the delicate

structure of projective tensor product of Banach spaces (see Chapter 2 in [10]). We also believe that our approach to orthogonality via projective tensor product is of independent interest and may have other applications.

Acknowledgement: The research of the third named author is supported in part by NBHM grant NBHM/R.P.64/2014, and the Mathematical Research Impact Centric Support (MATRICS) grant, File No: MTR/2017/000522 and Core Research Grant, File No: CRG/2019/000908, by the Science and Engineering Research Board (SERB), Department of Science & Technology (DST), Government of India. The research of the fourth author is supported by the NBHM postdoctoral fellowship, File No: 0204/3/2020/R&D-II/2445.

REFERENCES

- [1] L. Arambašić and R. Rajić, *The Birkhoff-James orthogonality in Hilbert C^* -modules*, Linear Algebra Appl. 437 (2012) 1913–1929.
- [2] G. Birkhoff, *Orthogonality in linear metric spaces*, Duke Math. J. 1 (1935), 169–172.
- [3] R. Bhatia and P. Šemrl, *Orthogonality of matrices and some distance problems*, Linear Algebra Appl. 287 (1999), 77–85.
- [4] T. Bhattacharyya and P. Grover, *Characterization of Birkhoff-James orthogonality*, J. Math. Anal. Appl. 407 (2013), 350–358.
- [5] J. Chmieliński, T. Stypuła and P. Wójcik, *Approximate orthogonality in normed spaces and its applications*, Linear Algebra Appl. 531 (2017), 305–317
- [6] R. C. James, *Orthogonality in normed linear spaces*, Duke Math. J. 12 (1945), 291–302.
- [7] R. C. James, *Orthogonality and linear functionals in normed linear spaces*, Trans. Amer. Math. Soc. 61 (1947), 265–292.
- [8] N. Komuro, K. Saito and R. Tanaka, *On symmetry of Birkhoff orthogonality in the positive cones of C^* -algebras with applications*, J. Math. Anal. Appl. 474 (2019), 1488–1497.
- [9] A. Mal, K. Paul, T.S.S.R.K. Rao and D. Sain, *Approximate Birkhoff-James orthogonality and smoothness in the space of bounded linear operators*, Monatsh. Math. 190 (2019), 549–558.
- [10] R. Ryan, *Introduction to tensor products of Banach spaces*, Springer Monographs in Mathematics (2002).
- [11] D. Sain, *Birkhoff-James orthogonality of linear operators on finite dimensional Banach spaces*, J. Math. Anal. Appl. 447 (2017), 860–866.
- [12] D. Sain and K. Paul, *Operator norm attainment and inner product spaces*, Linear Algebra Appl. 439 (2013), 2448–2452.
- [13] D. Sain, K. Paul and A. Mal, *A complete characterization of Birkhoff-James orthogonality in infinite dimensional normed space*, J. Operator Theory. 80 (2018), 399–413.
- [14] A. Turnšek, *A remark on orthogonality and symmetry of operators in $B(H)$* , Linear Algebra Appl. 535 (2017), 141–150.
- [15] P. Wójcik, *The Birkhoff orthogonality in pre-Hilbert C^* -modules*, Oper. Matrices 10 (3) (2016) 713–729.

KOUSIK DHARA, INDIAN STATISTICAL INSTITUTE, STATISTICS AND MATHEMATICS UNIT, 8TH MILE, MYSORE ROAD, BANGALORE, 560 059, INDIA.

Email address: kousik.dhara1@gmail.com

NARAYAN RAKSHIT, INDIAN STATISTICAL INSTITUTE, STATISTICS AND MATHEMATICS UNIT, 8TH MILE, MYSORE ROAD, BANGALORE, 560 059, INDIA.

Email address: narayan753@gmail.com

JAYDEB SARKAR, INDIAN STATISTICAL INSTITUTE, STATISTICS AND MATHEMATICS UNIT, 8TH MILE, MYSORE ROAD, BANGALORE, 560 059, INDIA.

Email address: jaydeb@gmail.com, jay@isibang.ac.in

ARYAMAN SENSARMA, INDIAN STATISTICAL INSTITUTE, STATISTICS AND MATHEMATICS UNIT, 8TH MILE, MYSORE ROAD, BANGALORE, 560 059, INDIA.

Email address: aryamansensarma@gmail.com