

Structure and cohomology of 3-Lie-Rinehart superalgebras

Abdelkader Ben Hassine^{1,2}, Taoufik Chtioui²,
Sami Mabrouk³, Sergei Silvestrov^{4*}

¹Department of Mathematics, Faculty of Science and Arts at Belqarn,
P. O. Box 60, Sabt Al-Alaya 61985, University of Bisha, Kingdom of Saudi Arabia

²Faculty of Sciences, University of Sfax, BP 1171, 3000 Sfax, Tunisia

³Faculty of Sciences, University of Gafsa, BP 2100, Gafsa, Tunisia

⁴Division of Applied Mathematics, School of Education, Culture and Communication,
Mälardalen University, Box 883, 72123 Västerås, Sweden.

Abstract

We introduce the concept of 3-Lie-Rinehart superalgebra and systematically describe a cohomology complex by considering coefficient modules. Furthermore, we study the relationships between a Lie-Rinehart superalgebra and its induced 3-Lie-Rinehart superalgebra. The deformations of 3-Lie-Rinehart superalgebra are considered via the cohomology theory.

Introduction

The notion of Lie-Rinehart algebras was introduced by J. Herz in [31] and mainly developed in [48, 49]. A Lie-Rinehart algebra can be thought as a Lie \mathbb{K} -algebra, which is simultaneously an A -module, where A is an associative and commutative \mathbb{K} -algebra, in such a way that both structures are related by a compatibility condition. A first approach to this class of algebras can be found in [33, 35].

G. Rinehart developed in [49] a formalism of differential forms for general commutative algebras which relies on the notion of (\mathbb{K}, A) -Lie algebra where \mathbb{K} is a commutative ring with unit and A a commutative \mathbb{K} -algebra. The notion of Lie-Rinehart algebra includes an abstract algebraic characterization of the algebraic structure which underlies a Lie algebroid [32, 33, 43, 44]. Thus a Lie-Rinehart algebra is an algebraic generalization of the notion of a Lie algebroid: the space of sections of a vector bundle is replaced by a module over a ring, a vector field by a derivation of the ring. For further details and a history of the notion of Lie-Rinehart algebra,

2020 *Mathematics subject classification*: 17A40; 17B05; 17B60; 17B66; 17B61; 53D17

Keywords: Lie-Rinehart superalgebra, 3-Lie-Rinehart superalgebra, Cohomology, Deformation.

*Corresponding author: sergei.silvestrov@mdh.se

we refer to [36]. Lie-Rinehart algebras have been investigated further in many papers [18, 21, 25, 33–35, 37–39, 43].

Some generalizations of Lie-Rinehart algebras, such as Lie-Rinehart superalgebras [24] or restricted Lie-Rinehart algebras [26], have been recently studied. The cohomology $H_{Rin}^*(L; M)$ of a Lie-Rinehart algebra L with coefficients in a Lie-Rinehart module M was first defined by Rinehart [49] and further developed in [33].

In [21–23], the authors introduced the notion of cross modules of Lie-Rinehart algebras. Hom-Lie-Rinehart algebras and their extensions in the small dimension cohomology space was introduced and studied in [45]. Hom-Rinehart algebras have close relations with Hom-Gerstenhaber Algebras and Hom-Lie Algebroids [46, 47].

The study of 3-Lie algebras [28] gets a lot of attention since it has close relationships with Lie algebras, Hom-Lie algebras, commutative associative algebras, and cubic matrices [14–16, 19]. For example, it is applied to the study of Nambu mechanics and the study of supersymmetry and gauge symmetry transformations of the world-volume theory of multiple coincident M2-branes [10, 45, 50, 51].

In 1996, the concept of n -Lie superalgebras was firstly introduced by Daletskii and Kushnirievich in [27]. Moreover, Cantarini and Kac gave a more general concept of n -Lie superalgebras again in 2010 (see [20]). n -Lie superalgebras are more general structures including n -Lie algebras, n -ary Nambu-Lie superalgebras, and Lie superalgebras. The construction of $(n + 1)$ -Lie algebras induced by n -Lie algebras using combination of bracket multiplication with a trace, motivated by the works on the quantization of the Nambu brackets [11], was generalized using the brackets of general Hom-Lie algebra or n -Hom-Lie algebra and trace-like linear forms satisfying some conditions depending on the linear maps defining the Hom-Lie or n -Hom-Lie algebras in [8, 9]. The structure of 3-Lie algebras induced by Lie algebras, classification of 3-Lie algebras and application to constructions of B.R.S. algebras have been considered in [1, 3, 4]. Interesting constructions of ternary Lie superalgebras in connection to superspace extension of Nambu-Hamilton equation is considered in [5]. In [17], a method was demonstrated of how to construct n -ary multiplications from the binary multiplication of a Hom-Lie algebra and a $(n - 2)$ -linear function satisfying certain compatibility conditions. Solvability and Nilpotency for n -Hom-Lie Algebras and $(n + 1)$ -Hom-Lie Algebras induced by n -Hom-Lie Algebras have been considered in [42]. In [1, 2, 5], 3-Lie superalgebras were constructed starting with a Lie superalgebras and various properties of such algebras considered. Related constructions for n -ary hom-Lie algebras and for n -ary hom-Lie superalgebras can be found in [6–9, 17, 29, 41]. The ternary case of (Hom-)Lie Rinehart algebras was developed in [12, 13, 30].

This paper is organized as follows. In Section 1, we recall the notion of Lie-Rinehart superalgebra and define their cohomology group. We generalize the Lie-Rinehart superalgebra to the ternary case. Section 2 is devoted to some construction results. We begin by constructing 3-Lie-Rinehart superalgebras starting with a Lie-Rinehart superalgebras. We construct some new 3-Lie-Rinehart superalgebras from a given 3-Lie-Rinehart superalgebra. The notion of a module for a 3-Lie-Rinehart superalgebras appeared in Section 3, and subsequently we introduce a cochain complex and cohomology of a 3-Lie-Rinehart superalgebras with coefficients in a module. Then we study relations between 1 and 2 cocycles of a Lie-Rinehart superalgebra and the induced 3-Lie-Rinehart superalgebra. At the end of this section, we study the deformation of 3-Lie-Rinehart superalgebras.

1 Definitions and Notations

In this section, we review basic definitions of Lie superalgebras, 3-Lie superalgebras, Lie-Rinehart superalgebras and generalize the notion of 3-Lie-Rinehart algebras to the super case.

Let $V = V_{\bar{0}} \oplus V_{\bar{1}}$ be a \mathbb{Z}_2 -graded vector space. If $v \in V$ is a homogenous element, then its degree will be denoted by \bar{v} , where $\bar{v} \in \mathbb{Z}_2$ and $\mathbb{Z}_2 = \{\bar{0}, \bar{1}\}$. Let $End(V)$ be the \mathbb{Z}_2 -graded vector space of endomorphisms of a \mathbb{Z}_2 -graded vector space $V = V_{\bar{0}} \oplus V_{\bar{1}}$. Denoted by $\mathcal{H}(V)$ the set of homogenous elements of V . The composition of two endomorphisms $a \circ b$ determines the structure of Lie superalgebra in $End(V)$ and the graded binary commutator $[a, b] = a \circ b - (-1)^{\bar{a}\bar{b}} b \circ a$ induces the structure of Lie superalgebra in $End(V)$.

1.1 Lie-Rinehart superalgebras

Definition 1.1 ([40]). A Lie superalgebra is a pair $(\mathfrak{g}, [\cdot, \cdot])$ consisting of a \mathbb{Z}_2 -graded vector space $\mathfrak{g} = \mathfrak{g}_{\bar{0}} \oplus \mathfrak{g}_{\bar{1}}$, an even bilinear map $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ satisfying the following identities:

$$[x, y] = -(-1)^{\bar{x}\bar{y}}[y, x], \text{ super skew-symmetric} \quad (1.1)$$

$$\cup_{x,y,z} (-1)^{\bar{x}\bar{z}}[x, [y, z]] = 0, \text{ super-Jacobi} \quad (1.2)$$

where x, y and z are homogeneous elements in \mathfrak{g} .

Definition 1.2. A representation of a Lie superalgebra $(\mathfrak{g}, [\cdot, \cdot])$ on a \mathbb{Z}_2 -graded vector space $V = V_{\bar{0}} \oplus V_{\bar{1}}$ is an even linear map $\mu : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$, such that for all homogeneous $x, y \in \mathfrak{g}$, the following identity is satisfied:

$$\mu([x, y]) = \mu(x)\mu(y) - (-1)^{\bar{x}\bar{y}}\mu(y)\mu(x). \quad (1.3)$$

Now, we recall the definition of Lie-Rinehart superalgebra [24].

Definition 1.3. A Lie-Rinehart superalgebra L over (an associative supercommutative superalgebra) A is a Lie superalgebra over \mathbb{K} with an A -module structure and an even linear map $\mu : L \rightarrow Der(A)$, such that that following conditions hold:

- (i) μ is a representation of $(L, [\cdot, \cdot])$ on A .
- (ii) $\mu(ax) = a\mu(x)$ for all $a \in A, x \in L$.
- (iii) The compatibility condition:

$$\forall a \in \mathcal{H}(A), x, y \in \mathcal{H}(L) : [x, ay] = \mu(x)ay + (-1)^{\bar{a}\bar{x}}a[x, y]. \quad (1.4)$$

Example 1.4. Let us observe that Lie-Rinehart superalgebras over A with trivial map $\mu : L \rightarrow Der(A)$ are exactly Lie superalgebras. If $A = \mathbb{K}$, then $Der(A) = 0$ and there is no difference between Lie and Lie-Rinehart superalgebras. Therefore the concept of Lie-Rinehart superalgebras generalizes the concept of Lie superalgebras.

Example 1.5. Let A an associative supercommutative superalgebra. Then, the A -module $Der(A) \oplus A$ is a Lie-Rinehart superalgebra over A with the bracket

$$[(D, a), (D', a')] = ([D, D'], D(a') - (-1)^{\bar{D}'\bar{a}}D'(a)),$$

and an even map $p_1 : Der(A) \oplus A \rightarrow Der(A)$, the projection onto the first factor.

Now, we define the cohomology of a Lie-Rinehart superalgebra. First we introduce the notion of left module over a Lie-Rinehart superalgebra.

Definition 1.6. Let M be an A -module. Then M is a left module over a Lie-Rinehart superalgebra L if there exists an even map $\theta : L \otimes M \rightarrow M$ such that:

- (i) θ is a representation of the Lie superalgebra $(L, [\cdot, \cdot])$ on M .
- (ii) $\theta(ax, m) = a\theta(x, m)$ for all $a \in A, x \in L, m \in M$.
- (iii) $\theta(x, am) = (-1)^{\bar{x}\bar{a}}a\theta(x, m) + \rho(x)(a)(m)$ for all $x \in \mathcal{H}(L), a \in \mathcal{H}(A), m \in M$.

Let $(L, A, [\cdot, \cdot], \mu)$ be a Lie-Rinehart superalgebra, and let M be a left module over L . For Lie-Rinehart superalgebra L with coefficients in M , consider the \mathbb{Z}_+ -graded space of \mathbb{K} -modules

$$C^*(L; M) := \bigoplus_{n \geq 0} C^n(L; M)$$

where $C^n(L; M) \subset \text{Hom}(\wedge^n L, M)$ consisting of elements satisfying:

$$f(x_1, \dots, ax_i, \dots, x_n) = (-1)^{\bar{a}(\bar{x}_1 + \bar{x}_2 + \dots + \bar{x}_{i-1} + \bar{f})} af(x_1, \dots, x_i, \dots, x_n),$$

for all $x_i \in \mathcal{H}(L), i = 1, \dots, n$ and $a \in \mathcal{H}(A)$.

Define the even linear map $\delta_{LR} : C^n(L; M) \rightarrow C^{n+1}(L; M)$ given by

$$\begin{aligned} \delta_{LR}f(x_1, \dots, x_{n+1}) &:= \sum_{i=1}^{n+1} (-1)^{i+1+\bar{x}_i(\bar{f}+\bar{x}_1+\dots+\bar{x}_{i-1})} \theta(x_i, f(x_1, \dots, \widehat{x}_i, \dots, x_{n+1})) \\ &+ \sum_{1 \leq i < j \leq n+1} (-1)^{(\bar{x}_i+\bar{x}_j)(\bar{x}_1+\dots+\bar{x}_{i-1}+\bar{x}_{i+1}+\dots+\bar{x}_{j-1})} f([x_i, x_j], x_1, \dots, \widehat{x}_i, \dots, \widehat{x}_j, \dots, x_{n+1}) \end{aligned}$$

for all $f \in C^n(L; M), x_i \in \mathcal{H}(L)$, where $1 \leq i \leq n+1$.

With the above notation, the map δ_{LR} gives rise to a coboundary map.

Proposition 1.7. If $f \in C^n(L; M)$, then $\delta_{LR}^2 f \in C^{n+1}(L; M)$ and $\delta_{LR}^2 = 0$.

By the above proposition, $(C^*(L, M), \delta_{LR})$ is a cochain complex. The resulting cohomology of the cochain complex we define to be the cohomology space of Lie-Rinehart superalgebra $(L, A, [\cdot, \cdot], \mu)$ with coefficients in M , and we denote this cohomology as $H^*(L, M)$.

1.2 3-Lie-Rinehart superalgebras

Definition 1.8 ([20]). A \mathbb{Z}_2 -graded vector space $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ is said to be a 3-Lie superalgebra, if it is endowed with an even trilinear map (bracket) $[\cdot, \cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$, satisfying the following conditions:

$$[x, y, z] = -(-1)^{\bar{x}\bar{y}}[y, x, z], \quad [x, y, z] = -(-1)^{\bar{y}\bar{z}}[x, z, y], \quad (1.5)$$

$$\begin{aligned} [x, y, [z, u, v]] &= [[x, y, z], u, v] + (-1)^{\bar{z}(\bar{x}+\bar{y})}[z, [x, y, u], v] \\ &+ (-1)^{(\bar{z}+\bar{u})(\bar{x}+\bar{y})}[z, u, [x, y, v]], \end{aligned} \quad (1.6)$$

where $x, y, z, u, v \in \mathfrak{g}$ are homogeneous elements.

Proposition 1.9. Let \mathfrak{g} be a \mathbb{Z}_2 -graded vector space together with a super skew-symmetric even linear map $[\cdot, \cdot, \cdot] : \otimes^3 \mathfrak{g} \rightarrow \mathfrak{g}$. Then $(\mathfrak{g}, [\cdot, \cdot, \cdot])$ is a 3-Lie superalgebra if and only if the following identities hold:

$$[[x_1, x_2, x_3], x_4, x_5] - (-1)^{\bar{x}_3 \bar{x}_4} [[x_1, x_2, x_4], x_3, x_5] + (-1)^{\bar{x}_2(\bar{x}_3 + \bar{x}_4)} [[x_1, x_3, x_4], x_2, x_5] - (-1)^{\bar{x}_1(\bar{x}_2 + \bar{x}_3 + \bar{x}_4)} [[x_2, x_3, x_4], x_1, x_5] = 0, \quad (1.7)$$

$$\begin{aligned} & (-1)^{(\bar{x}_4 + \bar{x}_5)(\bar{x}_1 + \bar{x}_2 + \bar{x}_3)} [[x_4, x_5, x_1], x_2, x_3] + (-1)^{(\bar{x}_4 + \bar{x}_5)(\bar{x}_2 + \bar{x}_3)} [x_1, [x_4, x_5, x_2], x_3] \\ & + (-1)^{(\bar{x}_1 + \bar{x}_2)(\bar{x}_3 + \bar{x}_4)} [[x_3, x_5, x_1], x_2, x_4] - (-1)^{(\bar{x}_3 + \bar{x}_5)\bar{x}_2 + \bar{x}_4 \bar{x}_5} [x_1, [x_5, x_3, x_2], x_4] \\ & + (-1)^{\bar{x}_3(\bar{x}_4 + \bar{x}_5)} [x_1, x_2, [x_4, x_5, x_3]] = 0, \end{aligned} \quad (1.8)$$

for all $x_i \in \mathcal{H}(\mathfrak{g})$, $1 \leq i \leq 5$.

Proof. If $(\mathfrak{g}, [\cdot, \cdot, \cdot])$ is a 3-Lie superalgebra, then applying (1.6) to the last term in (1.6) for $x_i \in \mathcal{H}(\mathfrak{g})$, $1 \leq i \leq 5$ and using (1.5),

$$\begin{aligned} [x_1, x_2, [x_3, x_4, x_5]] &= [[x_1, x_2, x_3], x_4, x_5] + (-1)^{\bar{x}_3(\bar{x}_1 + \bar{x}_2)} [x_3, [x_1, x_2, x_4], x_5] \\ &\quad + (-1)^{(\bar{x}_3 + \bar{x}_4)(\bar{x}_1 + \bar{x}_2)} [x_3, x_4, [x_1, x_2, x_5]] \\ &= [[x_1, x_2, x_3], x_4, x_5] + (-1)^{\bar{x}_3(\bar{x}_1 + \bar{x}_2)} [x_3, [x_1, x_2, x_4], x_5] \\ &\quad + (-1)^{(\bar{x}_3 + \bar{x}_4)(\bar{x}_1 + \bar{x}_2)} [[x_3, x_4, x_1], x_2, x_5] \\ &\quad + (-1)^{\bar{x}_2(\bar{x}_3 + \bar{x}_4)} [x_1, [x_3, x_4, x_2], x_5] \\ &\quad + [x_1, x_2, [x_3, x_4, x_5]], \end{aligned}$$

yields (1.7).

Similarly, applying (1.6) to the first and second terms on the right hand side of (1.6) for $x_i \in \mathcal{H}(\mathfrak{g})$, $1 \leq i \leq 5$ and using (1.5),

$$\begin{aligned} [x_1, x_2, [x_3, x_4, x_5]] &= (-1)^{(\bar{x}_4 + \bar{x}_5)(\bar{x}_1 + \bar{x}_2 + \bar{x}_3)} [x_4, x_5, [x_1, x_2, x_3] \\ &\quad - (-1)^{(\bar{x}_3 + \bar{x}_5)(\bar{x}_1 + \bar{x}_2) + \bar{x}_4 \bar{x}_5} [x_3, x_5, [x_1, x_2, x_4]] \\ &\quad + (-1)^{(\bar{x}_3 + \bar{x}_4)(\bar{x}_1 + \bar{x}_2)} [x_3, x_4, [x_1, x_2, x_5]] \\ &= (-1)^{(\bar{x}_4 + \bar{x}_5)(\bar{x}_1 + \bar{x}_2 + \bar{x}_3)} [[x_4, x_5, x_1], x_2, x_3] \\ &\quad + (-1)^{(\bar{x}_4 + \bar{x}_5)(\bar{x}_2 + \bar{x}_3)} [x_1, [x_4, x_5, x_2], x_3] \\ &\quad + (-1)^{\bar{x}_3(\bar{x}_4 + \bar{x}_5)} [x_1, x_2, [x_4, x_5, x_3]] \\ &\quad - (-1)^{(\bar{x}_3 + \bar{x}_5)(\bar{x}_1 + \bar{x}_2) + \bar{x}_4 \bar{x}_5} [[x_3, x_5, x_1], x_2, x_4] \\ &\quad - (-1)^{(\bar{x}_3 + \bar{x}_5)\bar{x}_2 + \bar{x}_4 \bar{x}_5} [x_1, [x_3, x_5, x_2], x_4] \\ &\quad - (-1)^{\bar{x}_4 \bar{x}_5} [x_1, x_2, [x_3, x_5, x_4]] \\ &\quad + (-1)^{(\bar{x}_1 + \bar{x}_2)(\bar{x}_3 + \bar{x}_4)} [x_3, x_4, [x_1, x_2, x_5]], \end{aligned}$$

yields (1.8).

Conversely, suppose that (1.7) and (1.8) hold. First (1.7) gives

$$\begin{aligned} [[x_1, x_2, x_3], x_4, x_5] &= (-1)^{\bar{x}_3 \bar{x}_4} [[x_1, x_2, x_4], x_3, x_5] \\ &\quad - (-1)^{\bar{x}_2(\bar{x}_3 + \bar{x}_4)} [[x_1, x_3, x_4], x_2, x_5] + (-1)^{\bar{x}_1(\bar{x}_2 + \bar{x}_3 + \bar{x}_4)} [[x_2, x_3, x_4], x_1, x_5] \\ &= (-1)^{(\bar{x}_3 + \bar{x}_4 + \bar{x}_5)(\bar{x}_1 + \bar{x}_2)} [[x_3, x_4, x_5], x_1, x_2] \\ &\quad + (-1)^{\bar{x}_1(\bar{x}_2 + \bar{x}_4 + \bar{x}_5) + \bar{x}_3(\bar{x}_4 + \bar{x}_5)} [[x_2, x_4, x_5], x_1, x_3] + (-1)^{(\bar{x}_2 + \bar{x}_3)(\bar{x}_4 + \bar{x}_5)} [[x_1, x_4, x_5], x_2, x_3] \end{aligned}$$

by (1.8). Thus, \mathfrak{g} is a 3-Lie superalgebra. \square

In the following we recall that given a Lie superalgebra analogues of supertrace one can construct a 3-Lie superalgebra [1]. Let $(\mathfrak{g}, [\cdot, \cdot])$ be a Lie superalgebra and $\tau : \mathfrak{g} \rightarrow \mathbb{K}$ an even linear form. We say that τ is a supertrace of \mathfrak{g} if $\tau([\cdot, \cdot]) = 0$. For any $x_1, x_2, x_3 \in \mathcal{H}(\mathfrak{g})$, we define the 3-ary bracket by

$$[x_1, x_2, x_3]_\tau = \tau(x_1)[x_2, x_3] - (-1)^{\bar{x}_1\bar{x}_2}\tau(x_2)[x_1, x_3] + (-1)^{\bar{x}_3(\bar{x}_1+\bar{x}_2)}\tau(x_3)[x_1, x_2]. \quad (1.9)$$

Theorem 1.10. *For any Lie superalgebra $(\mathfrak{g}, [\cdot, \cdot])$ and supertrace τ , the pair $(\mathfrak{g}, [\cdot, \cdot, \cdot]_\tau)$ is a 3-Lie superalgebra.*

Definition 1.11. *Let \mathfrak{g} be a 3-Lie superalgebra, V be a graded vector space and $\rho : \mathfrak{g} \wedge \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ be an even linear mapping. If ρ satisfies*

$$\begin{aligned} \rho(x_1, x_2)\rho(x_3, x_4) - (-1)^{(\bar{x}_1+\bar{x}_2)(\bar{x}_3+\bar{x}_4)}\rho(x_3, x_4)\rho(x_1, x_2) \\ = \rho([x_1, x_2, x_3], x_4) - (-1)^{\bar{x}_3\bar{x}_4}\rho([x_1, x_2, x_4], x_3), \end{aligned} \quad (1.10)$$

$$\begin{aligned} \rho([x_1, x_2, x_3], x_4) = \rho(x_1, x_2)\rho(x_3, x_4) + (-1)^{\bar{x}_1(\bar{x}_2+\bar{x}_3)}\rho(x_2, x_3)\rho(x_1, x_4) \\ + (-1)^{\bar{x}_3(\bar{x}_1+\bar{x}_2)}\rho(x_3, x_1)\rho(x_2, x_4), \end{aligned} \quad (1.11)$$

for all $x_i \in \mathcal{H}(\mathfrak{g}), 1 \leq i \leq 4$, then (V, ρ) is called a representation of \mathfrak{g} , or (V, ρ) is an \mathfrak{g} -module.

Define

$$\text{ad} : \mathfrak{g} \wedge \mathfrak{g} \rightarrow \mathfrak{gl}(\mathfrak{g}), \quad \text{ad}_{x,y}(z) = [x, y, z]. \quad (1.12)$$

Using (1.6), we can see that $(\mathfrak{g}, \text{ad})$ is a representation of the 3-Lie superalgebra \mathfrak{g} and it is called the adjoint representation of \mathfrak{g} .

Proposition 1.12. *Let (V, μ) be a representation of a Lie superalgebra $(\mathfrak{g}, [\cdot, \cdot])$ and τ be a supertrace of \mathfrak{g} . Then (V, ρ_τ) is a representation of the 3-Lie superalgebra $(\mathfrak{g}, [\cdot, \cdot, \cdot]_\tau)$ where $\rho_\tau : \mathfrak{g} \wedge \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ is defined by*

$$\rho_\tau(x, y) = \tau(x)\mu(y) - (-1)^{\bar{x}\bar{y}}\tau(y)\mu(x), \quad \forall x, y \in \mathcal{H}(\mathfrak{g}). \quad (1.13)$$

Proof. For all $x_1, x_2, x_3, x_4 \in \mathcal{H}(\mathfrak{g})$, the left hand of (1.10) becomes

$$\begin{aligned} \rho_\tau(x_1, x_2)\rho_\tau(x_3, x_4) - (-1)^{(\bar{x}_1+\bar{x}_2)(\bar{x}_3+\bar{x}_4)}\rho_\tau(x_3, x_4)\rho_\tau(x_1, x_2) \\ = \left(\tau(x_1)\mu(x_2) - (-1)^{\bar{x}_1\bar{x}_2}\tau(x_2)\mu(x_1)\right)\left(\tau(x_3)\mu(x_4) - (-1)^{\bar{x}_3\bar{x}_4}\tau(x_4)\mu(x_3)\right) \\ - (-1)^{(\bar{x}_1+\bar{x}_2)(\bar{x}_3+\bar{x}_4)}\left(\tau(x_3)\mu(x_4) - (-1)^{\bar{x}_3\bar{x}_4}\tau(x_4)\mu(x_3)\right)\left(\tau(x_1)\mu(x_2) - (-1)^{\bar{x}_1\bar{x}_2}\tau(x_2)\mu(x_1)\right) \\ = \tau(x_1)\mu(x_2)\tau(x_3)\mu(x_4) - (-1)^{\bar{x}_3\bar{x}_4}\tau(x_1)\mu(x_2)\tau(x_4)\mu(x_3) \\ - (-1)^{\bar{x}_1\bar{x}_2}\tau(x_2)\mu(x_1)\tau(x_3)\mu(x_4) - (-1)^{\bar{x}_1\bar{x}_2+\bar{x}_3\bar{x}_4}\tau(x_2)\mu(x_1)\tau(x_4)\mu(x_3) \\ - (-1)^{(\bar{x}_1+\bar{x}_2)(\bar{x}_3+\bar{x}_4)}\tau(x_3)\mu(x_4)\tau(x_1)\mu(x_2) + (-1)^{(\bar{x}_1+\bar{x}_2)(\bar{x}_3+\bar{x}_4)+\bar{x}_1\bar{x}_2}\tau(x_3)\mu(x_4)\tau(x_2)\mu(x_1) \\ + (-1)^{(\bar{x}_1+\bar{x}_2)(\bar{x}_3+\bar{x}_4)+\bar{x}_3\bar{x}_4}\tau(x_4)\mu(x_3)\tau(x_1)\mu(x_2) \\ - (-1)^{(\bar{x}_1+\bar{x}_2)(\bar{x}_3+\bar{x}_4)+\bar{x}_1\bar{x}_2+\bar{x}_3\bar{x}_4}\tau(x_4)\mu(x_3)\tau(x_2)\mu(x_1). \end{aligned}$$

Using (1.9) and (1.13), the right hand of (1.10) can be written as follows:

$$\begin{aligned} \rho_\tau([x_1, x_2, x_3], x_4) - (-1)^{\bar{x}_3\bar{x}_4}\rho_\tau([x_1, x_2, x_4], x_3) \\ = -(-1)^{\bar{x}_4(\bar{x}_2+\bar{x}_3)}\tau(x_1)\tau(x_4)\mu([x_2, x_3]) + (-1)^{\bar{x}_4(\bar{x}_2+\bar{x}_3)+\bar{x}_1\bar{x}_2}\tau(x_2)\tau(x_4)\mu([x_1, x_3]) \end{aligned}$$

$$\begin{aligned}
& - (-1)^{(\bar{x}_1+\bar{x}_2)(\bar{x}_3+\bar{x}_4)} \tau(x_3) \tau(x_4) \mu([x_1, x_2]) + (-1)^{\bar{x}_2 \bar{x}_3} \tau(x_1) \tau(x_3) \mu([x_2, x_4]) \\
& + (-1)^{\bar{x}_1 \bar{x}_2 + \bar{x}_3(\bar{x}_1+\bar{x}_4)} \tau(x_2) \tau(x_3) \mu([x_1, x_4]) - (-1)^{(\bar{x}_1+\bar{x}_2)(\bar{x}_3+\bar{x}_4)} \tau(x_4) \tau(x_3) \mu([x_1, x_2]).
\end{aligned}$$

By direct identification, we proof that

$$\begin{aligned}
\rho_\tau(x_1, x_2) \rho_\tau(x_3, x_4) - (-1)^{(\bar{x}_1+\bar{x}_2)(\bar{x}_3+\bar{x}_4)} \rho_\tau(x_3, x_4) \rho_\tau(x_1, x_2) \\
= \rho_\tau([x_1, x_2, x_3], x_4) - (-1)^{\bar{x}_3 \bar{x}_4} \rho_\tau([x_1, x_2, x_4], x_3).
\end{aligned}$$

Similarly, we can proof (1.11). \square

We generalize the Lie-Rinehart superalgebra to the ternary case.

Definition 1.13. A 3-Lie-Rinehart superalgebra over A is a tuple $(L, A, [\cdot, \cdot, \cdot], \rho)$, where A is an associative supercommutative superalgebra, L is an A -module, $[\cdot, \cdot, \cdot] : L \times L \times L \rightarrow L$ is an even super skew-symmetric trilinear map, and the \mathbb{K} -map $\rho : L \times L \rightarrow \text{Der}(A)$ such that the following conditions hold:

- (i) $(L, [\cdot, \cdot, \cdot])$ is a 3-Lie superalgebra.
- (ii) ρ is a representation of $(L, [\cdot, \cdot, \cdot])$ on A .
- (iii) For all $x, y \in \mathcal{H}(L)$, $a \in \mathcal{H}(A)$,

$$\rho(ax, y) = (-1)^{\bar{a}\bar{x}} \rho(x, ay) = a\rho(x, y). \quad (1.14)$$

- (iv) The compatibility condition:

$$[x, y, az] = (-1)^{\bar{a}(\bar{x}+\bar{y})} a[x, y, z] + \rho(x, y)az, \quad \forall x, y, z \in \mathcal{H}(L), a \in \mathcal{H}(A). \quad (1.15)$$

Remark 1.14. If $\rho = 0$, then $(L, A, [\cdot, \cdot, \cdot])$ is called a 3-Lie A -superalgebra.

Remark 1.15. If the condition (1.14) is not satisfied, then we call $(L, A, [\cdot, \cdot, \cdot], \rho)$ a weak 3-Lie-Rinehart superalgebra over A .

Definition 1.16. Let $(L, A, [\cdot, \cdot, \cdot]_L, \rho)$ and $(L', A', [\cdot, \cdot, \cdot]_{L'}, \rho')$ be two 3-Lie-Rinehart superalgebras, then a 3-Lie-Rinehart superalgebra homomorphism is defined as a pair of maps (g, f) , where $g : A \rightarrow A'$ and $f : L \rightarrow L'$ are two \mathbb{K} -algebra homomorphisms such that

- 1) $f(ax) = g(a)f(x)$ for all $x \in L, a \in A$,
- 2) $g(\rho(x, y)(a)) = \rho'(f(x), f(y))(g(a))$ for all $x \in L, a \in A$.

2 Some constructions of 3-Lie-Rinehart superalgebras

In this section, we give some construction results. We begin by constructing 3-Lie-Rinehart superalgebras starting with a Lie-Rinehart superalgebras. Then, we establish the converse sense. At the end of this section, we construct some new 3-Lie-Rinehart superalgebras from a given 3-Lie-Rinehart superalgebra.

The following Theorem generalizes results in [2] to Lie-Rinehart case.

Theorem 2.1. Let $(L, A, [\cdot, \cdot], \mu)$ be a Lie-Rinehart superalgebra and τ is a supertrace. If the condition

$$\tau(ax)y = \tau(x)ay \quad (2.1)$$

is satisfied for any $x, y \in \mathcal{H}(L)$, $a \in \mathcal{H}(A)$, then $(L, A, [\cdot, \cdot, \cdot]_\tau, \rho_\tau)$ is a 3-Lie-Rinehart superalgebra, where $[\cdot, \cdot, \cdot]_\tau$ and ρ_τ are defined in (1.9) and (1.12) respectively. We say that $(L, A, [\cdot, \cdot, \cdot]_\tau, \rho_\tau)$ is induced by $(L, A, [\cdot, \cdot], \mu)$ and is denoted by L_τ .

Proof. Proposition 1.12 gives that (A, ρ_τ) is a representation of the 3-Lie superalgebra $(L, [\cdot, \cdot, \cdot]_\tau)$ and $\rho_\tau(x, y) \in \text{Der}(A)$. For all $x, y, z \in \mathcal{H}(L)$ and $a \in \mathcal{H}(A)$,

$$\begin{aligned} [x, y, az]_\tau &= \tau(x)[y, az] - (-1)^{\bar{x}\bar{y}}\tau(y)[x, az] + (-1)^{(\bar{a}+\bar{z})(\bar{x}+\bar{y})}\tau(az)[x, y] \\ &= (-1)^{\bar{a}(\bar{x}+\bar{y})}a\tau(x)[y, z] + \tau(x)\mu(y)az - (-1)^{\bar{a}(\bar{x}+\bar{y})+\bar{x}\bar{y}}a\tau(y)[x, z] \\ &\quad - (-1)^{\bar{x}\bar{y}}\tau(y)\mu(x)az + (-1)^{(\bar{a}+\bar{z})(\bar{x}+\bar{y})}a\tau(z)[x, y] \\ &= (-1)^{\bar{a}(\bar{x}+\bar{y})}a[x, y, z]_\tau + \rho_\tau(x, y)az. \end{aligned}$$

So, we obtain (1.15). Since (2.1) is satisfied and

$$\begin{aligned} \rho_\tau(ax, y) &= \tau(ax)\mu(y) - (-1)^{(\bar{a}+\bar{x})\bar{y}}\tau(y)\mu(ax), \\ (-1)^{\bar{a}\bar{x}}\rho_\tau(x, ay) &= (-1)^{\bar{a}\bar{x}}\tau(x)\mu(ay) - (-1)^{(\bar{a}+\bar{y})\bar{x}+\bar{a}\bar{x}}\tau(ay)\mu(x), \\ a\rho_\tau(x, y) &= a\tau(x)\mu(y) - (-1)^{\bar{x}\bar{y}}a\tau(y)\mu(x), \end{aligned}$$

the condition (1.14) holds. \square

Conversely, we can construct a Lie-Rinehart superalgebra structure from a given 3-Lie-Rinehart superalgebra.

Proposition 2.2. *Let $(L, A, [\cdot, \cdot, \cdot], \rho)$ be a 3-Lie-Rinehart superalgebra. Let $x_0 \in L_{\bar{0}}$. Define the bracket $[\cdot, \cdot]_{x_0} = [x_0, \cdot, \cdot]$ and $\rho_{x_0}(x)(v) = \rho(x_0, x)v$. Then $(L, A, [\cdot, \cdot]_{x_0}, \rho_{x_0})$ is a Lie-Rinehart superalgebra.*

Proof. It is easy to check that the bracket is super skew-symmetric. For any $x, y, z \in \mathcal{H}(L)$, we have

$$\begin{aligned} [x, [y, z]_{x_0}]_{x_0} &= [x_0, x, [x_0, y, z]] \\ &= [[x_0, x, x_0], y, z] + [x_0, [x_0, x, y], z] + (-1)^{\bar{x}\bar{y}}[x_0, y, [x_0, x, z]] \\ &= [x_0, [x_0, x, y], z] + (-1)^{\bar{x}\bar{y}}[x_0, y, [x_0, x, z]] \\ &= [[x, y]_{x_0}, z]_{x_0} + (-1)^{\bar{x}\bar{y}}[y, [x, z]_{x_0}]_{x_0}. \end{aligned}$$

It is obvious to see that ρ_{x_0} is a representation of L on A and $\rho_{x_0}(x) \in \text{Der}(A)$. It remains to prove (1.4). Using (1.15), we obtain

$$\begin{aligned} [x, ay]_{x_0} = [x_0, x, ay] &= (-1)^{\bar{a}\bar{x}}a[x_0, x, y] + \rho(x_0, x)ay \\ &= (-1)^{\bar{a}\bar{x}}a[x, y]_{x_0} + \rho_{x_0}(x)ay. \end{aligned}$$

which completes the proof. \square

Definition 2.3. *Let $(L, A, [\cdot, \cdot, \cdot], \rho)$ be a 3-Lie-Rinehart superalgebra.*

- 1) *If S is a subalgebra of the 3-Lie superalgebra $(L, [\cdot, \cdot, \cdot])$ satisfying $AS \subset S$, then $(S, A, [\cdot, \cdot, \cdot], \rho|_{S \wedge S})$ is a 3-Lie-Rinehart superalgebra, which is called a subalgebra of the 3-Lie-Rinehart superalgebra $(L, A, [\cdot, \cdot, \cdot], \rho)$.*
- 2) *If I is an ideal of the 3-Lie superalgebra $(L, [\cdot, \cdot, \cdot])$ and satisfies $AI \subset I$ and $\rho(I, L)(A)L \subset I$, then $(I, A, [\cdot, \cdot, \cdot], \rho|_{I \wedge I})$ is a 3-Lie-Rinehart superalgebra, which is called an ideal of the 3-Lie-Rinehart superalgebra $(L, A, [\cdot, \cdot, \cdot], \rho)$.*
- 3) *If a 3-Lie-Rinehart superalgebra $(L, A, [\cdot, \cdot, \cdot], \rho)$ cannot be decomposed into the direct sum of two nonzero ideals, then L is called an indecomposable 3-Lie-Rinehart superalgebra.*

Proposition 2.4. *If $(L, A, [\cdot, \cdot, \cdot], \rho)$ is a 3-Lie-Rinehart superalgebra. Then*

$$\text{Ker}\rho = \{x \in \mathcal{H}(L) \mid \rho(x, L) = 0\}$$

is an ideal, which is called the kernel of the representation ρ .

Proof. By (1.10) and (1.11), for all homogeneous elements $x \in \text{Ker}\rho$, $y, z, w \in \mathcal{H}(L)$,

$$\rho([x, y, z], w) = \rho(x, y)\rho(z, w) + (-1)^{\bar{x}(\bar{y}+\bar{z})}\rho(y, z)\rho(x, w) + (-1)^{\bar{z}(\bar{x}+\bar{y})}\rho(z, x)\rho(y, w) = 0.$$

Therefore, $[x, L, L] \subseteq \text{Ker}\rho$. By (1.14), for all $a \in \mathcal{H}(A)$,

$$\rho(ax, y) = (-1)^{\bar{x}\bar{a}}\rho(x, ay) = a\rho(x, y) = 0, \quad \rho(\rho(x, y)az, w) = \rho(x, y)(a)\rho(z, w) = 0.$$

We get $ax \in \text{Ker}\rho$, $\rho(x, y)az \in \text{Ker}\rho$, that is, $A\text{Ker}\rho \subset \text{Ker}\rho$, $\rho(\text{Ker}\rho, L)(A)L \subset \text{Ker}\rho$.

Therefore, $\text{Ker}\rho$ is an ideal of the 3-Lie-Rinehart superalgebra $(L, A, [\cdot, \cdot, \cdot], \rho)$. \square

Theorem 2.5. *Let $(L, A, [\cdot, \cdot, \cdot], \rho)$ be a 3-Lie-Rinehart superalgebra. Then the following identities hold, for all $a, b, c \in \mathcal{H}(A)$, $x_i \in \mathcal{H}(L)$, $1 \leq i \leq 5$,*

$$\begin{aligned} & \rho(x_2, x_3)a[x_1, x_4, x_5] + (-1)^{(\bar{x}_1+\bar{x}_4)(\bar{x}_2+\bar{x}_3)+\bar{a}(\bar{x}_1+\bar{x}_4+\bar{x}_2+\bar{x}_3)}\rho(x_1, x_4)a[x_2, x_3, x_5] \\ & + (-1)^{\bar{x}_2(\bar{x}_3+\bar{x}_1)+\bar{a}(\bar{x}_1+\bar{x}_2)}\rho(x_3, x_1)a[x_2, x_4, x_5] \\ & + (-1)^{\bar{x}_4(\bar{x}_3+\bar{x}_1)+\bar{a}(\bar{x}_3+\bar{x}_4)}\rho(x_2, x_4)a[x_3, x_1, x_5] \\ & + (-1)^{\bar{x}_1(\bar{x}_2+\bar{x}_3)+\bar{a}(\bar{x}_1+\bar{x}_3)}\rho(x_1, x_2)a[x_3, x_4, x_5] \\ & + (-1)^{\bar{x}_2(\bar{x}_1+\bar{x}_3+\bar{x}_4)+\bar{x}_4\bar{x}_1+\bar{a}(\bar{x}_2+\bar{x}_4)}\rho(x_3, x_4)a[x_1, x_2, x_5] = 0. \end{aligned} \quad (2.2)$$

$$\begin{aligned} & \rho(x_2, x_3)(a_4a_5)[x_1, x_4, x_5] + (-1)^{\bar{x}_2(\bar{a}_1+\bar{a}_4+\bar{a}_5+\bar{x}_3)}\rho(x_3, x_1)(a_4a_5)[x_2, x_4, x_5] \\ & + (-1)^{\bar{x}_1(\bar{a}_4+\bar{a}_5+\bar{x}_2+\bar{x}_3)}\rho(x_1, x_2)(a_4a_5)[x_3, x_4, x_5] \\ & + (-1)^{(\bar{a}_4+\bar{x}_5)(\bar{x}_2+\bar{x}_3)+(\bar{x}_1+\bar{x}_4)(\bar{x}_2+\bar{x}_3+\bar{a}_5)}a_4\rho(x_1, x_4)a_5[x_5, x_2, x_3] \\ & + (-1)^{\bar{a}_4(\bar{x}_2+\bar{x}_3)+(\bar{x}_4+\bar{x}_5)(\bar{x}_1+\bar{x}_3)+\bar{x}_4\bar{a}_5}a_4\rho(x_2, x_4)a_5[x_5, x_3, x_1] \\ & + (-1)^{\bar{a}_4(\bar{x}_2+\bar{x}_3)+(\bar{x}_4+\bar{x}_5)(\bar{x}_1+\bar{x}_2)+\bar{x}_4\bar{a}_5}a_4\rho(x_3, x_4)a_5[x_5, x_1, x_2] \\ & - (-1)^{(\bar{x}_1+\bar{a}_5+\bar{x}_5)(\bar{a}_4+\bar{x}_2+\bar{x}_3)+\bar{x}_4(\bar{x}_2+\bar{x}_3)}a_5\rho(x_1, x_5)a_4[x_4, x_2, x_3] \\ & - (-1)^{(\bar{a}_5+\bar{x}_5)(\bar{x}_2+\bar{x}_3)+(\bar{a}_4+\bar{x}_5)(\bar{x}_3+\bar{x}_5)}a_5\rho(x_2, x_5)a_4[x_4, x_3, x_1] \\ & - (-1)^{\bar{a}_5(\bar{x}_2+\bar{x}_3+\bar{a}_4)+(\bar{x}_2+\bar{x}_5)(\bar{x}_1+\bar{x}_4+\bar{a}_4)+\bar{x}_2(\bar{x}_3+\bar{x}_5)+\bar{x}_1\bar{x}_4}a_5\rho(x_3, x_5)a_4[x_4, x_1, x_2] = 0. \end{aligned} \quad (2.3)$$

$$\begin{aligned} & \rho(x_1, x_2)\rho(x_3, x_4) + (-1)^{(\bar{x}_1+\bar{x}_2)(\bar{x}_3+\bar{x}_4)}\rho(x_3, x_4)\rho(x_1, x_2) \\ & + (-1)^{x_1(\bar{x}_2+\bar{x}_3)}\rho(x_2, x_3)\rho(x_1, x_4) + (-1)^{\bar{x}_4(\bar{x}_2+\bar{x}_3)}\rho(x_1, x_4)\rho(x_2, x_3) \\ & + (-1)^{\bar{x}_3(\bar{x}_1+\bar{x}_2)}\rho(x_3, x_1)\rho(x_2, x_4) + (-1)^{\bar{x}_1(\bar{x}_2+\bar{x}_3+\bar{x}_4)+\bar{x}_3\bar{x}_4}\rho(x_2, x_4)\rho(x_3, x_1) = 0. \end{aligned} \quad (2.4)$$

$$\begin{aligned} & (\rho(x_1, x_2)b)\rho(x_3, x_4) + (-1)^{\bar{x}_3(\bar{x}_1+\bar{x}_2+\bar{b})+\bar{b}\bar{x}_2}(\rho(x_3, x_1)b)\rho(x_2, x_4) \\ & + (-1)^{\bar{x}_1(\bar{x}_2+\bar{x}_3+\bar{b})+\bar{x}_1\bar{b}}(\rho(x_2, x_3)b)\rho(x_1, x_4) = 0. \end{aligned} \quad (2.5)$$

Proof. Using (1.15) we have

$$\begin{aligned} & \rho(x_2, x_3)a[x_1, x_4, x_5] + (-1)^{(\bar{x}_1+\bar{x}_4)(\bar{x}_2+\bar{x}_3)+\bar{a}(\bar{x}_1+\bar{x}_4+\bar{x}_2+\bar{x}_3)}\rho(x_1, x_4)a[x_2, x_3, x_5] \\ & + (-1)^{\bar{x}_2(\bar{x}_3+\bar{x}_1)+\bar{a}(\bar{x}_1+\bar{x}_2)}\rho(x_3, x_1)a[x_2, x_4, x_5] \end{aligned}$$

$$\begin{aligned}
& +(-1)^{\bar{x}_4(\bar{x}_3+\bar{x}_1)+\bar{a}(\bar{x}_2+\bar{x}_4)}\rho(x_2, x_4)a[x_3, x_1, x_5] \\
& +(-1)^{\bar{x}_1(\bar{x}_2+\bar{x}_3)+\bar{a}(\bar{x}_1+\bar{x}_3)}\rho(x_1, x_2)a[x_3, x_4, x_5] \\
& +(-1)^{\bar{x}_2(\bar{x}_1+\bar{x}_3+\bar{x}_4)+\bar{x}_4\bar{x}_1+\bar{a}(\bar{x}_2+\bar{x}_4)}\rho(x_3, x_4)a[x_1, x_2, x_5] \\
& = [x_2, x_3, a[x_1, x_4, x_5]] - \underbrace{(-1)^{\bar{a}(\bar{x}_2+\bar{x}_3)}a[x_2, x_3, [x_1, x_4, x_5]]}_{I_1} \\
& +(-1)^{(\bar{x}_1+\bar{x}_4)(\bar{x}_2+\bar{x}_3)+\bar{a}(\bar{x}_1+\bar{x}_4+\bar{x}_2+\bar{x}_3)}\left([x_1, x_4, a[x_2, x_3, x_5]] - \underbrace{(-1)^{\bar{a}(\bar{x}_1+\bar{x}_4)}a[x_1, x_4, [x_2, x_3, x_5]]}_{I_2}\right) \\
& +(-1)^{\bar{x}_2(\bar{x}_3+\bar{x}_1)+\bar{a}(\bar{x}_1+\bar{x}_2)}\left([x_3, x_1, a[x_2, x_4, x_5]] - \underbrace{(-1)^{\bar{a}(\bar{x}_3+\bar{x}_1)}a[x_4, x_1, [x_2, x_4, x_5]]}_{I_3}\right) \\
& +(-1)^{\bar{x}_4(\bar{x}_3+\bar{x}_1)+\bar{a}(\bar{x}_3+\bar{x}_4)}\left([x_2, x_4, a[x_3, x_1, x_5]] - \underbrace{(-1)^{\bar{a}(\bar{x}_2+\bar{x}_4)}a[x_2, x_4, [x_3, x_1, x_5]]}_{I_4}\right) \\
& +(-1)^{\bar{x}_1(\bar{x}_2+\bar{x}_3)+\bar{a}(\bar{x}_1+\bar{x}_3)}\left([x_1, x_2, a[x_3, x_4, x_5]] - \underbrace{(-1)^{\bar{a}(\bar{x}_1+\bar{x}_2)}a[x_1, x_2, [x_3, x_4, x_5]]}_{I_5}\right) \\
& +(-1)^{\bar{x}_2(\bar{x}_1+\bar{x}_3+\bar{x}_4)+\bar{x}_4\bar{x}_1+\bar{a}(\bar{x}_2+\bar{x}_4)}\left([x_3, x_4, a[x_1, x_2, x_5]] - \underbrace{(-1)^{\bar{a}(\bar{x}_3+\bar{x}_4)}a[x_3, x_4, [x_1, x_2, x_5]]}_{I_6}\right)
\end{aligned}$$

Thanks to Proposition 1.9, we have $\sum_{i=1}^6 I_i = 0$. Using (1.15), (1.10) and (1.11), we obtain (2.2). Similarly, (2.3)- (2.5) can be verified by a direct computation according to (2.2) and Definition 1.13. \square

Now we construct some new 3-Lie-Rinehart superalgebra from a given 3-Lie-Rinehart superalgebra $(L, A, [\cdot, \cdot, \cdot], \rho)$.

Theorem 2.6. *Let $(L, A, [\cdot, \cdot, \cdot], \rho)$ be a 3-Lie-Rinehart superalgebra and $B = A \otimes L = \{ax \mid a \in \mathcal{H}(A), x \in \mathcal{H}(L)\}$. Then $(B, A, [\cdot, \cdot, \cdot]', \rho')$ is a 3-Lie-Rinehart superalgebra, where the multiplication is defined by, for all $x_1, x_2, x_3 \in \mathcal{H}(L)$, $a_1, a_2, a_3 \in \mathcal{H}(A)$,*

$$\begin{aligned}
[a_1x_1, a_2x_2, a_3x_3]' & = (-1)^{\bar{a}_2\bar{x}_1+\bar{a}_3(\bar{x}_1+\bar{x}_2)}a_1a_2a_3[x_1, x_2, x_3] + (-1)^{\bar{a}_2\bar{x}_1}a_1a_2\rho(x_1, x_2)(a_3)x_3 \\
& + (-1)^{\bar{a}_3\bar{x}_2+(\bar{a}_1+\bar{x}_1)(\bar{a}_2+\bar{a}_3+\bar{x}_2+\bar{x}_3)}a_2a_3\rho(x_2, x_3)(a_1)x_1 \\
& + (-1)^{\bar{a}_3\bar{x}_1+(\bar{a}_2+x_2)(\bar{a}_3+\bar{x}_3)}a_1a_3\rho(x_1, x_3)(a_2)x_2,
\end{aligned} \tag{2.6}$$

and $\rho' : (A \otimes L) \wedge (A \otimes L) \rightarrow \text{Der}(A)$ is defined by $\rho'(a_1x_1, a_2x_2) = (-1)^{\bar{a}_2\bar{x}_1}a_1a_2\rho(x_1, x_2)$.

Note that $B = B_{\bar{0}} \oplus B_{\bar{1}}$ where $B_{\bar{0}} = A_{\bar{0}} \otimes L_{\bar{0}} \oplus A_{\bar{1}} \otimes L_{\bar{1}}$ and $B_{\bar{1}} = A_{\bar{0}} \otimes L_{\bar{1}} \oplus A_{\bar{1}} \otimes L_{\bar{0}}$.

Proof. For the super skew-symmetry of the bracket we have

$$\begin{aligned}
[a_1x_1, a_2x_2, a_3x_3]' & = (-1)^{\bar{a}_2\bar{x}_1+\bar{a}_3(\bar{x}_1+\bar{x}_2)}a_1a_2a_3[x_1, x_2, x_3] + (-1)^{\bar{a}_2\bar{x}_1}a_1a_2\rho(x_1, x_2)(a_3)x_3 \\
& + (-1)^{\bar{a}_3\bar{x}_2+(\bar{a}_1+\bar{x}_1)(\bar{a}_2+\bar{a}_3+\bar{x}_2+\bar{x}_3)}a_2a_3\rho(x_2, x_3)(a_1)x_1 \\
& + (-1)^{\bar{a}_3\bar{x}_1+(\bar{a}_2+x_2)(\bar{a}_3+\bar{x}_3)}a_1a_3\rho(x_1, x_3)(a_2)x_2 \\
& = -(-1)^{(\bar{a}_1+\bar{x}_1)(\bar{a}_2+\bar{x}_2)}\left((-1)^{\bar{a}_1\bar{x}_2+\bar{a}_3(\bar{x}_1+\bar{x}_2)}a_2a_1a_3[x_2, x_1, x_3]\right) \\
& + (-1)^{\bar{a}_1\bar{x}_2}a_2a_1\rho(x_2, x_1)(a_3)x_3
\end{aligned}$$

$$\begin{aligned}
& +(-1)^{\bar{a}_3\bar{x}_1+(\bar{a}_2+\bar{x}_2)(\bar{a}_1+\bar{a}_3+\bar{x}_2+\bar{x}_3)} a_1 a_3 \rho(x_1, x_3)(a_2)x_2 \\
& +(-1)^{\bar{a}_3\bar{x}_2+(\bar{a}_1+x_1)(\bar{a}_3+\bar{x}_3)} a_2 a_3 \rho(x_2, x_3)(a_1)x_1 \\
= & -(-1)^{(\bar{a}_1+\bar{x}_1)(\bar{a}_2+\bar{x}_2)} [a_2 x_2, a_1 x_1, a_3 x_3]'.
\end{aligned}$$

Similarly, we obtain $[a_1 x_1, a_2 x_2, a_3 x_3]' = -(-1)^{(\bar{a}_2+\bar{x}_2)(\bar{a}_3+\bar{x}_3)} [a_1 x_1, a_3 x_3, a_2 x_2]'$.

Using (2.2) and (2.3), we can deduce that $(B, [\cdot, \cdot, \cdot]')$ is a 3-Lie superalgebra and B is an A -module. Thanks to (1.10) and (1.11), for all $x_1, x_2, x_3 \in \mathcal{H}(L)$ and $b, a_1, a_2, a_3 \in \mathcal{H}(A)$, it is easy to show that (A, ρ') is a representation of B . Indeed,

$$\begin{aligned}
& \rho'(a_1 x_1, a_2 x_2) \rho'(a_3 x_3, a_4 x_4) - (-1)^{(\bar{a}_1+\bar{x}_1+\bar{a}_2+\bar{x}_2)(\bar{x}_3+\bar{a}_3+\bar{x}_4+\bar{x}_4)} \rho'(a_3 x_3, a_4 x_4) \rho'(a_1 x_1, a_2 x_2) \\
& = \rho'([a_1 x_1, a_2 x_2, a_3 x_3]', a_4 x_4) + (-1)^{(\bar{a}_1+\bar{x}_1+\bar{a}_2+\bar{x}_2)(\bar{a}_3+\bar{x}_3)} \rho'(a_3 x_3, [a_1 x_1, a_2 x_2, a_4 x_4]'), \\
& \rho'(a_1 x_1, a_2 x_2) \rho'(a_3 x_3, a_4 x_4) + (-1)^{(\bar{a}_1+\bar{x}_1)(\bar{a}_2+\bar{x}_2+\bar{x}_3+\bar{x}_3)} \rho'(a_2 x_2, a_3 x_3) \rho'(a_1 x_1, a_4 x_4) \\
& + (-1)^{(\bar{a}_1+\bar{x}_1+\bar{a}_2+\bar{x}_2)(\bar{a}_3+\bar{x}_3)} \rho'(a_3 x_3, a_1 x_1) \rho'(a_2 x_2, a_4 x_4) = \rho'([a_1 x_1, a_2 x_2, a_3 x_3]', a_4 x_4).
\end{aligned}$$

To prove the compatibility condition (1.15), we compute as follows

$$\begin{aligned}
& [a_1 x_1, a_2 x_2, b(a_3 x_3)]' = (-1)^{\bar{a}_2\bar{x}_1+(\bar{a}_3+\bar{b})(\bar{x}_1+\bar{x}_3)} a_1 a_2 b a_3 [x_1, x_2, x_3] \\
& + (-1)^{\bar{a}_2\bar{x}_1} a_1 a_2 \rho(x_1, x_2)(b a_3) x_3 \\
& + (-1)^{(\bar{b}+\bar{a}_3)\bar{x}_2+(\bar{a}_1+\bar{x}_1)(\bar{a}_2+\bar{x}_2+\bar{a}_3+\bar{x}_3)} a_2 b a_3 \rho(x_2, x_3) a_1 x_1 \\
& + (-1)^{(\bar{b}+\bar{a}_3)\bar{x}_1+(\bar{a}_2+\bar{x}_2)(\bar{a}_3+\bar{b}+\bar{x}_3)} a_1 b a_3 \rho(x_1, x_3) a_2 x_2 \\
= & (-1)^{\bar{b}(\bar{a}_1+\bar{x}_1+\bar{a}_2+\bar{x}_2)} b (a_1 a_2 a_3 [x_1, x_2, x_3] + (-1)^{\bar{a}_2\bar{x}_1} a_1 a_2 \rho(x_1, x_2) a_3 x_3 \\
& + (-1)^{\bar{a}_3\bar{x}_2+(\bar{a}_1+\bar{x}_1)(\bar{a}_2+\bar{x}_2+\bar{a}_3+\bar{x}_3)} a_2 a_3 \rho(x_2, x_3) a_1 x_1 \\
& + (-1)^{\bar{a}_3\bar{x}_1+(\bar{a}_2+\bar{x}_2)(\bar{a}_3+\bar{x}_3)} a_1 a_3 \rho(x_1, x_3) a_2 x_2) + (-1)^{\bar{a}_2\bar{x}_1} a_1 a_2 \rho(x_1, x_2) b(a_3 x_3) \\
= & (-1)^{\bar{b}(\bar{a}_1+\bar{x}_1+\bar{a}_2+\bar{x}_2)} b [a_1 x_1, a_2 x_2, a_3 x_3]' + \rho'(a_1 x_1, a_2 x_2) b(a_3 x_3).
\end{aligned}$$

It is obvious to show that

$$\rho'(b(a_1 x_1), a_2 x_2) = b \rho'(a_1 x_1, a_2 x_2) = (-1)^{\bar{b}(\bar{a}_1+\bar{x}_1)} \rho'(a_1 x_1, b(a_2 x_2)).$$

Therefore, $(B, A, [\cdot, \cdot, \cdot]', \rho')$ is a 3-Lie-Rinehart superalgebra. \square

Theorem 2.7. *Let $(L, A, [\cdot, \cdot, \cdot], \rho)$ be a 3-Lie-Rinehart superalgebra and*

$$E = L \oplus A = \{(x, a) \mid x \in L, a \in A\}.$$

Then $(E, A, [\cdot, \cdot, \cdot]_1, \rho_1)$ is a 3-Lie-Rinehart superalgebra, where for any $a, b, c \in \mathcal{H}(A)$, $x, y, z \in \mathcal{H}(L)$ and $k \in \mathbb{K}$,

$$k(x, a) = (kx, ka), (x, a) + (y, b) = (x + y, a + b), a(y, b) = (ay, ab), \quad (2.7)$$

$$[(x, a), (y, b), (z, c)]_1 = ([x, y, z], \rho(x, y)c - (-1)^{\bar{y}\bar{z}} \rho(x, z)b + (-1)^{\bar{x}(\bar{y}+\bar{z})} \rho(y, z)a), \quad (2.8)$$

$$\rho_1 : E \wedge E \rightarrow \text{Der}(A), \rho_1((x, a), (y, b)) = \rho(x, y). \quad (2.9)$$

Note that $E = E_{\bar{0}} \oplus E_{\bar{1}}$, where $E_{\bar{0}} = L_{\bar{0}} \oplus A_{\bar{0}}$ and $E_{\bar{1}} = L_{\bar{1}} \oplus A_{\bar{1}}$.

Proof. Let $x_i \in \mathcal{H}(L), a_i \in \mathcal{H}(A), i = 1, \dots, 5$. Thanks to (2.7), E is an A -module, and the 3-ary linear multiplication defined by (2.8) is super skew-symmetric. We have

$$\begin{aligned}
& [(x_1, a_1), (x_2, a_2), [(x_3, a_3), (x_4, a_4), (x_5, a_5)]_1]_1 \\
&= [(x_1, a_1), (x_2, a_2), ([x_3, x_4, x_5], \rho(x_3, x_4)a_5 - (-1)^{\bar{x}_4\bar{x}_5}\rho(x_3, x_5)a_4 \\
&\quad + (-1)^{\bar{x}_3(\bar{x}_4+\bar{x}_5)}\rho(x_4, x_5)a_3)]_1 \\
&= ([x_1, x_2, [x_3, x_4, x_5]], \rho(x_1, x_2)\rho(x_3, x_4)a_5 - (-1)^{\bar{x}_4\bar{x}_5}\rho(x_1, x_2)\rho(x_3, x_5)a_4 \\
&\quad + (-1)^{\bar{x}_3(\bar{x}_4+\bar{x}_5)}\rho(x_1, x_2)\rho(x_4, x_5)a_3 - (-1)^{\bar{x}_2(\bar{x}_3+\bar{x}_4+\bar{x}_5)}\rho(x_1, [x_3, x_4, x_5])a_2 \\
&\quad - (-1)^{\bar{x}_1(\bar{x}_2+\bar{x}_3+\bar{x}_4+\bar{x}_5)}\rho(x_2, [x_3, x_4, x_5])a_1).
\end{aligned}$$

Similarly we obtain

$$\begin{aligned}
& [(x_1, a_1), (x_2, a_2), (x_3, a_3)]_1, (x_4, a_4), (x_5, a_5)]_1 \\
&+ (-1)^{\bar{x}_3(\bar{x}_1+\bar{x}_2)}[(x_3, a_3), [(x_1, a_1), (x_2, a_2), (x_4, a_4)]_1, (x_5, a_5)]_1 \\
&+ (-1)^{(\bar{x}_3+\bar{x}_4)(\bar{x}_1+\bar{x}_2)}[(x_3, a_3), (x_4, a_4), [(x_1, a_1), (x_2, a_2), (x_5, a_5)]_1]_1 \\
&= [([x_1, x_2, x_3], \rho(x_1, x_2)a_3 - (-1)^{\bar{x}_2\bar{x}_3}\rho(x_1, x_3)a_2 \\
&\quad + (-1)^{\bar{x}_1(\bar{x}_2+\bar{x}_3)}\rho(x_2, x_3)a_2), (x_4, a_4), (x_5, a_5)]_1 \\
&\quad + (-1)^{\bar{x}_3(\bar{x}_1+\bar{x}_2)}([x_3, a_3], ([x_1, x_2, x_4], \rho(x_1, x_2)a_4 - (-1)^{\bar{x}_2\bar{x}_4}\rho(x_1, x_4)a_2 \\
&\quad + (-1)^{\bar{x}_1(\bar{x}_2+\bar{x}_4)}\rho(x_2, x_4)a_2), (x_5, a_5)]_1) \\
&\quad + (-1)^{(\bar{x}_3+\bar{x}_4)(\bar{x}_1+\bar{x}_2)}([x_3, a_3), (x_4, a_5), ([x_1, x_2, x_5], \rho(x_1, x_2)a_5 - (-1)^{\bar{x}_2\bar{x}_5}\rho(x_1, x_5)a_2 \\
&\quad + (-1)^{\bar{x}_1(\bar{x}_2+\bar{x}_5)}\rho(x_2, x_5)a_2)]_1) \\
&= ([x_1, x_2, [x_3, x_4, x_5]], \rho(x_1, x_2)\rho(x_3, x_4)a_5 - (-1)^{\bar{x}_4\bar{x}_5}\rho(x_1, x_2)\rho(x_3, x_5)a_4 \\
&\quad + (-1)^{\bar{x}_4(\bar{x}_4+\bar{x}_5)}\rho(x_1, x_2)\rho(x_4, x_5)a_3 - (-1)^{\bar{x}_2(\bar{x}_3+\bar{x}_4+\bar{x}_5)}\rho(x_1, [x_3, x_4, x_5])a_2 \\
&\quad - (-1)^{\bar{x}_1(\bar{x}_2+\bar{x}_3+\bar{x}_4+\bar{x}_5)}\rho(x_2, [x_3, x_4, x_5])a_1) \\
&\quad + (-1)^{\bar{x}_3(\bar{x}_1+\bar{x}_2)}([x_3, [x_1, x_2, x_4], x_5], \rho(x_3, [x_1, x_2, x_4])a_5 \\
&\quad - (-1)^{\bar{x}_5(\bar{x}_1+\bar{x}_2+\bar{x}_4)}(\rho(x_3, x_5)\rho(x_1, x_2)a_4 - (-1)^{\bar{x}_2\bar{x}_4}\rho(x_3, x_5)\rho(x_1, x_4)a_2 \\
&\quad + (-1)^{\bar{x}_1(\bar{x}_2+\bar{x}_4)}\rho(x_3, x_5)\rho(x_2, x_4)a_1) + (-1)^{\bar{x}_3(\bar{x}_1+\bar{x}_2+\bar{x}_4+\bar{x}_5)}\rho([x_1, x_2, x_4], x_5)a_3) \\
&\quad + (-1)^{(\bar{x}_3+\bar{x}_4)(\bar{x}_1+\bar{x}_2)}([x_3, x_4, [x_1, x_2, x_5]], \rho(x_3, x_4)\rho(x_1, x_2)a_5 - (-1)^{\bar{x}_2\bar{x}_5}\rho(x_3, x_4)\rho(x_1, x_5)a_2 \\
&\quad + (-1)^{\bar{x}_1(\bar{x}_2+\bar{x}_5)}\rho(x_3, x_4)\rho(x_2, x_5)a_1 - (-1)^{\bar{x}_4(\bar{x}_1+\bar{x}_2+\bar{x}_5)}\rho(x_3, [x_1, x_2, x_5])a_4 \\
&\quad + (-1)^{\bar{x}_3(\bar{x}_1+\bar{x}_2+\bar{x}_4+\bar{x}_5)}\rho(x_4, [x_1, x_2, x_5])a_3).
\end{aligned}$$

Then (1.6) holds thanks to (1.10). Therefore, $(E, [\cdot, \cdot, \cdot]_1)$ is a 3-Lie superalgebra.

Now we prove that ρ_1 is a representation of E over A . By (2.9), we have

$$\begin{aligned}
& \rho_1([(x_1, a_1), (x_2, a_2), (x_3, a_3)], (x_4, a_4)) - (-1)^{\bar{x}_3\bar{x}_4}\rho_1((x_3, a_3), [(x_1, a_1), (x_2, a_2), (x_4, a_4)]) \\
&= \rho_1([([x_1, x_2, x_3], \rho(x_1, x_2)a_3 - (-1)^{\bar{x}_2\bar{x}_3}\rho(x_1, x_2)a_3 + (-1)^{\bar{x}_1(\bar{x}_2+\bar{x}_3)}\rho(x_2, x_3)a_1), (x_4, a_4)] \\
&\quad - (-1)^{\bar{x}_3\bar{x}_4}(\rho_1((x_3, a_3), ([x_1, x_2, x_4], \rho(x_1, x_2)a_4 - (-1)^{\bar{x}_2\bar{x}_4}\rho(x_1, x_2)a_4 \\
&\quad + (-1)^{\bar{x}_1(\bar{x}_2+\bar{x}_4)}\rho(x_2, x_4)a_1)))
\end{aligned}$$

$$\begin{aligned}
&= \rho([x_1, x_2, x_3], x_4) - (-1)^{\bar{x}_3\bar{x}_4} \rho(x_3, [x_1, x_2, x_4]) \\
&= \rho(x_1, x_2) \rho(x_3, x_4) - (-1)^{(\bar{x}_1+\bar{x}_2)(\bar{x}_3+\bar{x}_4)} \rho(x_3, x_4) \rho(x_1, x_2) \\
&= \rho_1((x_1, a_1), (x_2, a_2)) \rho_1((x_3, a_3), (x_4, a_4)) \\
&\quad - (-1)^{(\bar{x}_1+\bar{x}_2)(\bar{x}_3+\bar{x}_4)} \rho((x_3, a_3), (x_4, a_4)) \rho((x_1, a_1), (x_2, a_2)).
\end{aligned}$$

Therefore, (1.10) holds. Similarly, we can prove (1.11). Then ρ_1 is a representation of E over A . For all $b \in \mathcal{H}(A)$, we have

$$\begin{aligned}
&[(x_1, a_1), (x_2, a_2), b(x_3, a_3)]_1 = [(x_1, a_1), (x_2, a_2), (bx_3, ba_3)]_1 \\
&= ([x_1, x_2, bx_3], \rho(x_1, x_2)(ba_3) - (-1)^{\bar{x}_2\bar{x}_3} \rho(x_1, bx_3)a_2 + (-1)^{\bar{x}_1(\bar{x}_2+\bar{x}_3)} \rho(x_2, bx_3)a_1) \\
&= ((-1)^{\bar{b}(\bar{x}_1+\bar{x}_2)} b[x_1, x_2, x_3] + \rho(x_1, x_2)bx_3, \rho(x_1, x_2)(ba_3) - (-1)^{\bar{x}_2\bar{x}_3} \rho(x_1, bx_3)a_2 \\
&\quad + (-1)^{\bar{x}_1(\bar{x}_2+\bar{x}_3)} \rho(x_2, bx_3)a_1) \\
&= (-1)^{\bar{b}(\bar{x}_1+\bar{x}_2)} b[(x_1, a_1), (x_2, a_2), (x_3, a_3)] + \rho_1(x_1, x_2)b(x_3, a_3).
\end{aligned}$$

Moreover, we have

$$\begin{aligned}
\rho_1(c(x, a), (y, b)) &= \rho_1((cx, ca), (y, b)) = \rho(cx, y) \\
&= (-1)^{\bar{c}\bar{x}} \rho(x, cy) = (-1)^{\bar{c}\bar{x}} \rho_1((x, a), (cy, cb)) = (-1)^{\bar{c}\bar{x}} \rho_1((x, a), c(y, b)).
\end{aligned}$$

Similarly, we obtain $\rho_1(c(x, a), (y, b)) = c\rho_1((x, a), (y, b))$. Then $(E, A, [\cdot, \cdot, \cdot]_1, \rho_1)$ is a 3-Lie-Rinehart superalgebra. \square

3 Cohomology and deformations of 3-Lie-Rinehart superalgebras

In this section, we study the notion of a module for a 3-Lie-Rinehart superalgebras and subsequently we introduce a cochain complex and cohomology of a 3-Lie-Rinehart superalgebras with coefficients in a module, then we study relations between 1 and 2 cocycles of a Lie-Rinehart superalgebra and the induced 3-Lie-Rinehart superalgebra. At the end of this section, we study the deformation of 3-Lie-Rinehart superalgebras

3.1 Representations and cohomology of 3-Lie-Rinehart superalgebras

Definition 3.1. Let M be an A -module. and $\psi : L \otimes L \rightarrow \text{End}(M)$ be an even linear map. The pair (M, ψ) is called a left module of the 3-Lie-Rinehart superalgebra $(L, A, [\cdot, \cdot, \cdot], \rho)$ if the following conditions hold:

- (i) ψ is a representation of $(L, [\cdot, \cdot, \cdot])$ on M ,
- (ii) $\psi(a \cdot x, y) = (-1)^{\bar{a}\bar{x}} \psi(x, a \cdot y) = a \cdot \psi(x, y)$, for all $a \in \mathcal{H}(A)$ and $x, y \in \mathcal{H}(L)$,
- (iii) $\psi(x, y)(a \cdot m) = (-1)^{\bar{a}(\bar{x}+\bar{y})} a \cdot \psi(x, y)(m) + \rho(x, y)(a)m$, for all $x, y \in \mathcal{H}(L)$, $a \in \mathcal{H}(A)$ and $m \in M$.

Example 3.2. A is a left module over L since ρ is a representation of $(L, [\cdot, \cdot, \cdot])$ over A and the other conditions are satisfied automatically by definition of the map ρ .

Example 3.3. The pair (L, ad) is a left module over L , which is called the adjoint representation of $(L, A, [\cdot, \cdot, \cdot], \rho)$.

Proposition 3.4. Let $(L, A, [\cdot, \cdot, \cdot], \rho)$ be a 3-Lie-Rinehart superalgebra. Then (M, ψ) is a left module over $(L, A, [\cdot, \cdot, \cdot], \rho)$ if and only if $(L \oplus M, A, [\cdot, \cdot, \cdot]_{L \oplus M}, \rho_{L \oplus M})$ is a 3-Lie-Rinehart superalgebra with the multiplication:

$$\begin{aligned} [x_1 + m_1, x_2 + m_2, x_3 + m_3]_{L \oplus M} &:= [x_1, x_2, x_3] + \psi(x_1, x_2)m_3 + (-1)^{\bar{x}_1(\bar{x}_2 + \bar{x}_3)}\psi(x_2, x_3)m_1 \\ &\quad + (-1)^{\bar{x}_2\bar{x}_3}\psi(x_3, x_1)m_2, \\ \rho_{L \oplus M} : (L \oplus M) \otimes (L \oplus M) &\rightarrow Der(A), \quad \rho_{L \oplus M}(x_1 + m_1, x_2 + m_2) := \rho(x_1, x_2), \end{aligned}$$

for any $x_1, x_2, x_3 \in \mathcal{H}(L)$ and $m_1, m_2, m_3 \in M$.

Note that $(L \oplus M)_{\bar{0}} = L_{\bar{0}} \oplus M_{\bar{0}}$, implying that if $x + m \in \mathcal{H}(L \oplus M)$, then $\overline{x + m} = \bar{x} = \bar{m}$.

Proof. Since L and M are A -modules, then $L \oplus M$ is an A -module via

$$a(x + m) = ax + am, \forall a \in A, x \in L, m \in M.$$

If (M, ψ) is a left module over $(L, A, [\cdot, \cdot, \cdot], \rho)$. Then $(L \oplus M, [\cdot, \cdot, \cdot]_{L \oplus M})$ is a 3-Lie superalgebra. It is obvious that $\rho_{L \oplus M}$ is a representation of the 3-Lie superalgebra $(L \oplus M, [\cdot, \cdot, \cdot]_{L \oplus M})$ over A .

For any $x_1, x_2, x_3 \in \mathcal{H}(L)$, $m_1, m_2, m_3 \in \mathcal{H}(M)$ and $a \in \mathcal{H}(A)$,

$$\begin{aligned} [x_1 + m_1, x_2 + m_2, a(x_3 + m_3)]_{L \oplus M} &= [x_1 + m_1, x_2 + m_2, ax_3 + am_3]_{L \oplus M} \\ &= [x_1, x_2, ax_3] + \psi(x_1, x_2)am_3 + (-1)^{\bar{x}_1(\bar{x}_2 + \bar{x}_3)}\psi(x_2, ax_3)m_1 + (-1)^{\bar{x}_2\bar{x}_3}\psi(ax_3, x_1)m_2 \\ &= (-1)^{\bar{a}(\bar{x}_1 + \bar{x}_2)}a[x_1, x_2, x_3] + \psi(x_1, x_2)am_3 + (-1)^{\bar{x}_1(\bar{x}_2 + \bar{x}_3)}\psi(x_2, ax_3)m_1 \\ &\quad + (-1)^{\bar{x}_2\bar{x}_3}\psi(ax_3, x_1)m_2 + \rho(x_1, x_2)(am_3) \\ &= (-1)^{\bar{a}(\bar{x}_1 + \bar{x}_2)}a([x_1, x_2, x_3] + \psi(x_1, x_2)m_3 + (-1)^{\bar{x}_1(\bar{x}_2 + \bar{x}_3)}\psi(x_2, x_3)m_1 + (-1)^{\bar{x}_2\bar{x}_3}\psi(x_3, x_1)m_2) \\ &\quad + \rho(x_1, x_2)(am_3) \\ &= (-1)^{\bar{a}(\bar{x}_1 + \bar{x}_2)}a[x_1 + m_1, x_2 + m_2, x_3 + m_3]_{L \oplus M} + \rho_{L \oplus M}(x_1 + m_1, x_2 + m_2)a(x_3 + m_3). \end{aligned}$$

Moreover,

$$\begin{aligned} \rho_{L \oplus M}(a(x_1 + m_1), x_2 + m_2) &= \rho_{L \oplus M}(ax_1 + am_1, x_2 + m_2) = \rho(ax_1, x_2) = (-1)^{\bar{a}\bar{x}_1}\rho(x_1, ax_2) \\ &= (-1)^{\bar{a}\overline{\bar{x}_1 + \bar{m}_1}}\rho_{L \oplus M}(x_1 + m_1, ax_2 + am_2) = (-1)^{\bar{a}\overline{\bar{x}_1 + \bar{m}_1}}\rho_{L \oplus M}(x_1 + m_1, a(x_2 + m_2)). \end{aligned}$$

Similarly, we obtain $\rho_{L \oplus M}(a(x_1 + m_1), x_2 + m_2) = a\rho_{L \oplus M}(x_1 + m_1, x_2 + m_2)$. Therefore, $(L \oplus M, A, [\cdot, \cdot, \cdot]_{L \oplus M}, \rho_{L \oplus M})$ is a 3-Lie-Rinehart superalgebra. The sufficient condition can be done in the same way. \square

Let (M, ψ) be a left module of the 3-Lie-Rinehart superalgebra $(L, A, [\cdot, \cdot, \cdot], \rho)$ and $C^n(L, M)$ the space of all linear maps $f : \wedge^2 L \otimes \cdots \otimes \wedge^2 L \wedge L \rightarrow M$ satisfying the conditions below:

- (i) $f(x_1, \dots, x_i, x_{i+1}, \dots, x_{2n}, x_{2n+1}) = -(-1)^{\bar{x}_i\bar{x}_{i+1}}f(x_1, \dots, x_{i+1}, x_i, \dots, x_{2n+1})$, for all $x_i \in \mathcal{H}(L)$, $1 \leq i \leq 2n + 1$,
- (ii) $f(x_1, \dots, a \cdot x_i, \dots, x_{2n+1}) = (-1)^{\bar{a}(\bar{x}_1 + \cdots + \bar{x}_{i-1} + \bar{f})}af(x_1, \dots, x_i, \dots, x_{2n+1})$, for all $x_i \in \mathcal{H}(L)$, $1 \leq i \leq 2n + 1$ and $a \in \mathcal{H}(A)$.

Next we consider the \mathbb{Z}_+ -graded space of \mathbb{K} -modules

$$C^*(L, M) := \bigoplus_{n \geq 0} C^n(L, M).$$

Define the \mathbb{K} -linear maps $\delta_{3LR} : C^{n-1}(L, M) \rightarrow C^n(L, M)$ given by

$$\begin{aligned} \delta_{3LR} f(x_1, \dots, x_{2n+1}) = & (-1)^{n+(\bar{f}+\bar{x}_1+\bar{x}_2+\dots+\bar{x}_{2n-2})(\bar{x}_{2n-1}+\bar{x}_{2n+1})+\bar{x}_{2n+1}} \bar{x}_{2n} \psi(x_{2n-1}, x_{2n+1}) f(x_1, \dots, x_{2n-2}, x_{2n}) \\ & + (-1)^{n+(\bar{f}+\bar{x}_1+\bar{x}_2+\dots+\bar{x}_{2n-1})(\bar{x}_{2n}+\bar{x}_{2n+1})} \psi(x_{2n}, x_{2n+1}) f(x_1, \dots, x_{2n-1}) \\ & + \sum_{k=1}^n (-1)^{k+(\bar{f}+\bar{x}_1+\bar{x}_2+\dots+\bar{x}_{2k-2})(\bar{x}_{2k-1}+\bar{x}_{2k})} \psi(x_{2k-1}, x_{2k}) f(x_1, \dots, \widehat{x}_{2k-1}, \widehat{x}_{2k}, \dots, x_{2n+1}) \\ & + \sum_{k=1}^n \sum_{j=2k+1}^{2n+1} (-1)^{k+(\bar{x}_{2k+1}+\dots+\bar{x}_{j-1})(\bar{x}_{2k-1}+\bar{x}_{2k})} f(x_1, \dots, \widehat{x}_{2k-1}, \widehat{x}_{2k}, \dots, [x_{2k-1}, x_{2k}, x_j], \dots, x_{2n+1}). \end{aligned}$$

Proposition 3.5. *If $f \in C^{n-1}(L, M)$, then $\delta_{3LR} f \in C^n(L, M)$ and $\delta_{3LR}^2 = 0$.*

Proof. Let f be a homogenous element in $C^{n-1}(L, M)$, it is obvious that $\delta_{3LR} f$ is skew-symmetric. For all $x_1, x_2, \dots, x_{2n+1} \in \mathcal{H}(L)$, $a \in \mathcal{H}(A)$ and for $i < 2n-1$,

$$\begin{aligned} \delta_{3LR} f(x_1, \dots, ax_i, \dots, x_{2n+1}) = & (-1)^{n+(\bar{f}+\bar{x}_1+\bar{x}_2+\dots+\bar{x}_{2n-2})(\bar{x}_{2n-1}+\bar{x}_{2n+1})+\bar{x}_{2n+1}} \bar{x}_{2n} \psi(x_{2n-1}, x_{2n+1}) f(x_1, \dots, ax_i, \dots, x_{2n-2}, x_{2n}) \\ & + (-1)^{n+(\bar{f}+\bar{x}_1+\bar{x}_2+\dots+\bar{x}_{2n-1})(\bar{x}_{2n}+\bar{x}_{2n+1})} \psi(x_{2n}, x_{2n+1}) f(x_1, \dots, ax_i, \dots, x_{2n-1}) \\ & + \sum_{\substack{k=1 \\ i < 2k-1}}^n (-1)^{k+(\bar{f}+\bar{a}+\bar{x}_1+\bar{x}_2+\dots+\bar{x}_{2k-2})(\bar{x}_{2k-1}+\bar{x}_{2k})} \psi(x_{2k-1}, x_{2k}) f(x_1, \dots, ax_i, \dots, \widehat{x}_{2k-1}, \widehat{x}_{2k}, \dots, x_{2n+1}) \\ & + \sum_{\substack{k=1 \\ i > 2k}}^n (-1)^{k+(\bar{f}+\bar{x}_1+\bar{x}_2+\dots+\bar{x}_{2k-2})(\bar{x}_{2k-1}+\bar{x}_{2k})} \psi(x_{2k-1}, x_{2k}) f(x_1, \dots, \widehat{x}_{2k-1}, \widehat{x}_{2k}, \dots, ax_i, \dots, x_{2n+1}) \\ & + \sum_{k=1}^n (-1)^{k+(\bar{f}+\bar{x}_1+\bar{x}_2+\dots+\bar{x}_{2k-2})(\bar{x}_{2k-1}+\bar{a}+\bar{x}_{2k})} \psi(ax_{2k-1}, x_{2k}) f(x_1, \dots, \widehat{x}_{2k-1}, \widehat{x}_{2k}, \dots, x_{2n+1}) \\ & + \sum_{k=1}^n (-1)^{k+(\bar{f}+\bar{x}_1+\bar{x}_2+\dots+\bar{x}_{2k-2})(\bar{x}_{2k-1}+\bar{a}+\bar{x}_{2k})} \psi(x_{2k-1}, ax_{2k}) f(x_1, \dots, \widehat{x}_{2k-1}, \widehat{x}_{2k}, \dots, x_{2n+1}) \\ & + \sum_{\substack{k=1 \\ i < 2k}}^n \sum_{j=2k+1}^{2n+1} (-1)^{k+(\bar{x}_{2k+1}+\dots+\bar{x}_{j-1})(\bar{x}_{2k-1}+\bar{x}_{2k})} \\ & \quad f(x_1, \dots, ax_i, \dots, \widehat{x}_{2k-1}, \widehat{x}_{2k}, \dots, [x_{2k-1}, x_{2k}, x_j], \dots, x_{2n+1}) \\ & + \sum_{k=1}^n \sum_{\substack{j=2k+1 \\ 2k-1 < i < j}}^{2n+1} (-1)^{k+(\bar{a}+\bar{x}_{2k+1}+\dots+\bar{x}_{j-1})(\bar{x}_{2k-1}+\bar{x}_{2k})} \\ & \quad f(x_1, \dots, \widehat{x}_{2k-1}, \widehat{x}_{2k}, \dots, ax_i, \dots, [x_{2k-1}, x_{2k}, x_j], \dots, x_{2n+1}) \\ & + \sum_{k=1}^n \sum_{\substack{j=2k+1 \\ j < i}}^{2n+1} (-1)^{k+(\bar{x}_{2k+1}+\dots+\bar{x}_{j-1})(\bar{x}_{2k-1}+\bar{x}_{2k})} \end{aligned}$$

$$\begin{aligned}
& f(x_1, \dots, \widehat{x}_{2k-1}, \widehat{x}_{2k}, \dots, [x_{2k-1}, x_{2k}, x_j], \dots, ax_i, \dots, x_{2n+1}) \\
& + \sum_{k=1}^n \sum_{j=2k+1}^{2n+1} (-1)^{k+(\bar{x}_{2k+1}+\dots+\bar{x}_{j-1})(\bar{x}_{2k-1}+\bar{a}+\bar{x}_{2k})} f(x_1, \dots, \widehat{x}_{2k-1}, \widehat{x}_{2k}, \dots, [ax_{2k-1}, x_{2k}, x_j], \dots, x_{2n+1}) \\
& + \sum_{k=1}^n \sum_{j=2k+1}^{2n+1} (-1)^{k+(\bar{x}_{2k+1}+\dots+\bar{x}_{j-1})(\bar{x}_{2k-1}+\bar{a}+\bar{x}_{2k})} f(x_1, \dots, \widehat{x}_{2k-1}, \widehat{x}_{2k}, \dots, [x_{2k-1}, ax_{2k}, x_j], \dots, x_{2n+1}) \\
& + \sum_{k=1}^n \sum_{j=2k+1}^{2n+1} (-1)^{k+(\bar{x}_{2k+1}+\dots+\bar{x}_{j-1})(\bar{x}_{2k-1}+\bar{x}_{2k})} f(x_1, \dots, \widehat{x}_{2k-1}, \widehat{x}_{2k}, \dots, [x_{2k-1}, x_{2k}, ax_j], \dots, x_{2n+1}).
\end{aligned}$$

Using Definition 3.1 and (1.15), we obtain

$$\delta_{3LR}f(x_1, \dots, ax_i, \dots, x_{2n+1}) = (-1)^{\bar{a}(\bar{x}_1+\dots+\bar{x}_{i-1}+\bar{f})} a\delta_{3LR}f(x_1, \dots, x_i, \dots, x_{2n+1}),$$

Similarly, we can proof the same result if $i = 2n - 1, 2n, 2n + 1$. Then δ_{3LR} is well-defined. Further, $\delta_{3LR}^2 = 0$ follows from the direct but a long calculation. \square

By the above proposition, $(C^*(L, M), \delta_{3LR})$ is a cochain complex. The resulting cohomology of the cochain complex can be defined as the cohomology space of 3-Lie-Rinehart superalgebra $(L, A, [\cdot, \cdot, \cdot], \rho)$ with coefficients in (M, ψ) , and we denote this cohomology as $H^*(L, M)$.

Definition 3.6. Let $(L, A, [\cdot, \cdot, \cdot], \rho)$ be a 3-Lie-Rinehart superalgebra and (M, ψ) be a left module over L . If $v \in H_{3LR}^1(L, M)$ satisfies

$$\begin{aligned}
& (-1)^{\bar{v}(\bar{x}_1+\bar{x}_2)}\psi(x_1, x_2)v(x_3) + (-1)^{\bar{x}_2\bar{x}_3+\bar{v}(\bar{x}_2+\bar{x}_3)}\psi(x_1, x_3)v(x_2) \\
& + (-1)^{(\bar{x}_1+\bar{v})(\bar{x}_2+\bar{x}_3)}\psi(x_2, x_3)v(x_1) + v([x_1, x_2, x_3]) = 0,
\end{aligned}$$

for any $x_1, x_2, x_3 \in \mathcal{H}(L)$, then v is called a 1-cocycle associated with ψ .

Definition 3.7. Let $(L, A, [\cdot, \cdot, \cdot], \rho)$ be a 3-Lie-Rinehart superalgebra and (M, ψ) be a left module over L . If $\omega \in H_{3LR}^2(L, M)$ satisfies

$$\begin{aligned}
& (-1)^{(\bar{\omega}+\bar{x}_1+\bar{x}_2)(\bar{x}_3+\bar{x}_5)+\bar{x}_4\bar{x}_5}\psi(x_3, x_5)\omega(x_1, x_2, x_4) + (-1)^{(\bar{\omega}+\bar{x}_1+\bar{x}_2+\bar{x}_3)(\bar{x}_4+\bar{x}_5)}\psi(x_4, x_5)\omega(x_1, x_2, x_3) \\
& - (-1)^{\bar{\omega}(\bar{x}_1+\bar{x}_2)}\psi(x_1, x_2)\omega(x_3, x_4, x_5) + (-1)^{(\bar{\omega}+\bar{x}_1+\bar{x}_2)(\bar{x}_3+\bar{x}_4)}\psi(x_3, x_4)\omega(x_1, x_2, x_5) \\
& - \omega([x_1, x_2, x_3], x_4, x_5) - (-1)^{\bar{x}_3(\bar{x}_1+\bar{x}_2)}\omega(x_3, [x_1, x_2, x_4], x_5) \\
& - (-1)^{(\bar{x}_3+\bar{x}_4)(\bar{x}_1+\bar{x}_2)}\omega(x_3, x_4, [x_1, x_2, x_5]) + \omega(x_1, x_2, [x_3, x_4, x_5]) = 0,
\end{aligned}$$

for any $x_1, x_2, x_3, x_4, x_5 \in \mathcal{H}(L)$, then ω is called a 2-cocycle associated with ψ .

Theorem 3.8. Let $(L, A, [\cdot, \cdot, \cdot], \mu)$ be a Lie-Rinehart superalgebra, τ be a supertrace and $\varphi \in Z_{LR}^2(L, L)$ such that

$$\begin{aligned}
& \forall x, y, z \in \mathcal{H}(A) : \\
& \tau(x)\tau(\varphi(y, z)) - (-1)^{\bar{x}\bar{y}}\tau(y)\tau(\varphi(x, z)) + (-1)^{\bar{z}(\bar{x}+\bar{y})}\tau(z)\tau(\varphi(x, y)) = 0.
\end{aligned}$$

Define the linear map $\phi : \otimes^3 L \rightarrow L$ by

$$\phi(x, y, z) = \tau(x)\varphi(y, z) - (-1)^{\bar{x}\bar{y}}\tau(y)\varphi(x, z) + (-1)^{\bar{z}(\bar{x}+\bar{y})}\tau(z)\varphi(x, y).$$

Then ϕ is a 2-cocycle of the induced 3-Lie-Rinehart superalgebra $(L, A, [\cdot, \cdot, \cdot]_{\tau}, \rho_{\tau})$.

Proof. Let $\varphi \in Z_{LR}^2(L, L)$, it is obvious that ϕ is skew-symmetric and $\bar{\phi} = \bar{\varphi}$. Set $x, y, z \in \mathcal{H}(L)$ then we have

$$\begin{aligned}\phi(ax, y, z) &= \tau(ax) \varphi(y, z) - (-1)^{\bar{x}\bar{y}} \tau(y) \varphi(ax, z) + (-1)^{\bar{z}(\bar{x}+\bar{y})} \tau(z) \varphi(ax, y) \\ &= a\tau(x) \varphi(y, z) - (-1)^{\bar{x}\bar{y}+\bar{a}\bar{\varphi}} \tau(y) a\varphi(x, z) + (-1)^{\bar{z}(\bar{x}+\bar{y})+\bar{a}\bar{\varphi}} \tau(z) a\varphi(x, y) \\ &= (-1)^{\bar{a}\bar{\varphi}} a \left(\tau(x) \varphi(y, z) - (-1)^{\bar{x}\bar{y}} \tau(y) \varphi(x, z) + (-1)^{\bar{z}(\bar{x}+\bar{y})} \tau(z) \varphi(x, y) \right) \\ &= (-1)^{\bar{a}\bar{\varphi}} a\phi(x, y, z).\end{aligned}$$

Similarly, $\phi(x, ay, z) = (-1)^{\bar{a}(\bar{x}+\bar{\phi})} a\phi(x, y, z)$ and $\phi(x, y, az) = (-1)^{\bar{a}(\bar{x}+\bar{y}+\bar{\phi})} a\phi(x, y, z)$. On the other hand, let $x_1, x_2, y_1, y_2, z \in \mathcal{H}(L)$. Then

$$\begin{aligned}\delta_{3LR}\phi(x_1, x_2, y_1, y_2, z) &= \phi(x_1, x_2, [y_1, y_2, z]_\tau) - \phi([x_1, x_2, y_1]_\tau, y_2, z) \\ &- (-1)^{\bar{y}_1(\bar{x}_1+\bar{x}_2)} \phi(y_1, [x_1, x_2, y_2]_\tau, z) - (-1)^{(\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2)} \phi(y_1, y_2, [x_1, x_2, z]_\tau) \\ &+ (-1)^{(\bar{x}_1+\bar{x}_2)\bar{\phi}} \left[x_1, x_2, \phi(y_1, y_2, z) \right]_\tau - \left[\phi(x_1, x_2, y_1), y_2, z \right]_\tau \\ &- (-1)^{\bar{y}_1(\bar{x}_1+\bar{x}_2+\bar{\phi})} \left[y_1, \phi(x_1, x_2, y_2), z \right]_\tau - (-1)^{(\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2+\bar{\phi})} \left[y_1, y_2, \phi(x_1, x_2, z) \right]_\tau \\ &= \tau(y_1) \phi(x_1, x_2, [y_2, z]) - (-1)^{\bar{y}_1\bar{y}_2} \tau(y_2) \phi(x_1, x_2, [z, y_1]) \\ &+ (-1)^{\bar{z}(\bar{y}_1+\bar{y}_2)} \tau(z) \phi(x_1, x_2, [y_1, y_2]) - \tau(x_1) \phi([x_2, y_1], y_2, z) \\ &+ (-1)^{\bar{x}_1\bar{x}_2} \tau(x_2) \phi([x_1, y_1], y_2, z) - (-1)^{\bar{y}_1(\bar{x}_1+\bar{x}_2)} \tau(y_1) \phi([x_1, x_2], y_2, z) \\ &- (-1)^{\bar{y}_1(\bar{x}_1+\bar{x}_2)} \tau(x_1) \phi(y_1, [x_2, y_2], z) + (-1)^{\bar{y}_1(\bar{x}_1+\bar{x}_2)+\bar{x}_1\bar{x}_2} \tau(x_2) \phi(y_1, [x_1, y_2], z) \\ &- (-1)^{(\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2)} \tau(y_2) \phi(y_1, [x_1, x_2], z) - (-1)^{(\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2)} \tau(x_1) \phi(y_1, y_2, [x_2, z]) \\ &+ (-1)^{(\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2)+\bar{x}_1\bar{x}_2} \tau(x_2) \phi(y_1, y_2, [x_1, z]) - (-1)^{(\bar{z}+\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2)} \tau(z) \phi(y_1, y_2, [x_1, x_2]) \\ &+ (-1)^{\bar{\phi}(\bar{x}_1+\bar{x}_2)} \tau(x_1) \left[x_2, \phi(y_1, y_2, z) \right] - (-1)^{\bar{\phi}(\bar{x}_1+\bar{x}_2)+\bar{x}_1\bar{x}_2} \tau(x_2) \left[x_1, \phi(y_1, y_2, z) \right] \\ &+ (-1)^{(\bar{x}_1+\bar{x}_2)(\bar{y}_1+\bar{y}_2+\bar{z})} \tau(\phi(y_1, y_2, z)) [x_1, x_2] - \tau(\phi(x_1, x_2, y_1)) [y_2, z] \\ &+ (-1)^{\bar{y}_2(\bar{x}_1+\bar{x}_2+\bar{y}_1+\bar{\phi})} \tau(y_2) \left[\phi(x_1, x_2, y_1), z \right] - (-1)^{\bar{z}(\bar{x}_1+\bar{x}_2+\bar{y}_1+\bar{y}_2+\bar{\phi})} \tau(z) \left[\phi(x_1, x_2, y_1), y_2 \right] \\ &- (-1)^{\bar{y}_1(\bar{x}_1+\bar{x}_2+\bar{\phi})} \tau(y_1) \left[\phi(x_1, x_2, y_2), z \right] + (-1)^{\bar{y}_1\bar{y}_2} \tau(\phi(x_1, x_2, y_2)) [y_1, z] \\ &- (-1)^{(\bar{y}_1+\bar{z})(\bar{x}_1+\bar{x}_2+\bar{\phi})+\bar{z}(\bar{y}_1+\bar{y}_2)} \tau(z) \left[y_1, \phi(x_1, x_2, y_2) \right] \\ &- (-1)^{(\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2+\bar{\phi})} \tau(y_1) \left[y_2, \phi(x_1, x_2, z) \right] \\ &+ (-1)^{(\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2+\bar{\phi})+\bar{y}_1\bar{y}_2} \tau(y_2) \left[y_1, \phi(x_1, x_2, z) \right] \\ &- (-1)^{\bar{z}(\bar{y}_1+\bar{y}_2)} \tau(\phi(x_1, x_2, z)) [y_1, y_2] \\ &= \tau(y_1) \left(\tau(x_1) \varphi(x_2, [y_2, z]) - (-1)^{\bar{x}_1\bar{x}_2} \tau(x_2) \varphi(x_1, [y_2, z]) \right) \\ &+ (-1)^{(\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2)} \tau(y_2) \varphi([x_1, x_2], z) - (-1)^{(\bar{y}_1+\bar{z})(\bar{x}_1+\bar{x}_2)+\bar{z}\bar{y}_2} \tau(z) \varphi([x_1, x_2], y_2) \\ &- (-1)^{\bar{y}_1(\bar{x}_1+\bar{x}_2+\bar{\phi})} \tau(x_1) \left[\varphi(x_2, y_2), z \right] + (-1)^{\bar{y}_1(\bar{x}_1+\bar{x}_2+\bar{\phi})+\bar{x}_1\bar{x}_2} \tau(x_2) \left[\varphi(x_1, y_2), z \right] \\ &- (-1)^{\bar{y}_1(\bar{x}_1+\bar{x}_2+\bar{\phi})+\bar{y}_2(\bar{x}_1+\bar{x}_2)} \tau(y_2) \left[\varphi(x_1, x_2), z \right] - (-1)^{(\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2+\bar{\phi})} \tau(x_1) \left[y_2, \varphi(x_2, z) \right] \\ &+ (-1)^{(\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2+\bar{\phi})} \tau(x_1) \left[y_2, \varphi(x_2, z) \right] - (-1)^{(\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2+\bar{\phi})+\bar{z}(\bar{x}_1+\bar{x}_2)} \tau(z) \left[y_2, \varphi(x_1, x_2) \right] \\ &- (-1)^{\bar{y}_1\bar{y}_2} \tau(y_2) \left(\tau(x_1) \varphi(x_2, [y_1, z]) - (-1)^{\bar{x}_1\bar{x}_2} \tau(x_2) \varphi(x_1, [y_1, z]) \right)\end{aligned}$$

$$\begin{aligned}
& +(-1)^{(\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2)+\bar{y}_1\bar{y}_2}\tau(y_1)\varphi([x_1,x_2],z)+(-1)^{(\bar{y}_1+\bar{y}_2+\bar{z})(\bar{x}_1+\bar{x}_2)+\bar{y}_2(\bar{y}_1+\bar{z})}\tau(z)\varphi(y_1,[x_1,x_2]) \\
& -(-1)^{\bar{y}_2(\bar{x}_1+\bar{x}_2+\bar{\phi})}\tau(x_1)[\varphi(x_2,y_1),z]+(-1)^{\bar{y}_2(\bar{x}_1+\bar{x}_2+\bar{\phi})+\bar{x}_1\bar{x}_2}\tau(x_2)[\varphi(x_1,y_1),z] \\
& -(-1)^{(\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2)+\bar{y}_2\bar{\phi}}\tau(y_1)[\varphi(x_1,x_2),z]-(-1)^{(\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2+\bar{\phi})}\tau(x_1)[y_1,\varphi(x_2,z)] \\
& +(-1)^{(\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2+\bar{\phi})+\bar{x}_1\bar{x}_2}\tau(x_2)[y_1,\varphi(x_1,z)] \\
& -(-1)^{(\bar{y}_1+\bar{y}_2+\bar{z})(\bar{x}_1+\bar{x}_2)+(\bar{y}_1+\bar{y}_2)\bar{\phi}}\tau(z)[y_1,\varphi(x_1,x_2)] \\
& +(-1)^{\bar{z}(\bar{y}_1+\bar{y}_2)}\tau(z)\left(\tau(x_1)\varphi(x_2,[y_1,y_2]) - (-1)^{\bar{x}_1\bar{x}_2}\tau(x_2)\varphi(x_1,[y_1,y_2])\right) \\
& -(-1)^{(\bar{z}+\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2+\bar{z})}\tau(y_1)\varphi(y_2,[x_1,x_2])+(-1)^{(\bar{z}+\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2+\bar{z})+\bar{y}_1\bar{y}_2}\tau(y_2)\varphi(y_1,[x_1,x_2]) \\
& -(-1)^{\bar{z}(\bar{x}_1+\bar{x}_2+\bar{y}_1+\bar{y}_2+\bar{\phi})}\tau(x_1)[\varphi(x_2,y_1),y_2]+(-1)^{\bar{z}(\bar{x}_1+\bar{x}_2+\bar{y}_1+\bar{y}_2+\bar{\phi})+\bar{x}_1\bar{x}_2}\tau(x_2)[\varphi(x_1,y_1),y_2] \\
& -(-1)^{\bar{z}(\bar{x}_1+\bar{x}_2+\bar{y}_1+\bar{y}_2+\bar{\phi})+\bar{y}_1(\bar{x}_1+\bar{x}_2)}\tau(y_1)[\varphi(x_1,x_2),y_2] \\
& +(-1)^{(\bar{y}_1+\bar{z})(\bar{x}_1+\bar{x}_2+\bar{\phi})+\bar{z}(\bar{y}_1+\bar{y}_2)}\tau(x_1)[y_1,\varphi(x_2,y_2)] \\
& -(-1)^{(\bar{y}_1+\bar{z})(\bar{x}_1+\bar{x}_2+\bar{\phi})+\bar{z}(\bar{y}_1+\bar{y}_2)+\bar{x}_1+\bar{x}_2}\tau(x_2)[y_1,\varphi(x_1,y_2)] \\
& +(-1)^{(\bar{y}_1+\bar{y}_2+\bar{z})(\bar{x}_1+\bar{x}_2+\bar{\phi})+\bar{z}(\bar{y}_1+\bar{y}_2)+\bar{y}_2\bar{\phi}}\tau(y_2)[y_1,\varphi(x_1,x_2)] \\
& +\tau(x_1)\left((-1)^{\bar{y}_2(\bar{x}_2+\bar{y}_1)}\tau(y_2)\varphi([x_2,y_1],z)-(-1)^{\bar{z}(\bar{y}_1+\bar{y}_2+\bar{x}_2)}\tau(z)\varphi([x_2,y_1],y_2)\right) \\
& -(-1)^{\bar{y}_1(\bar{x}_1+\bar{x}_2)}\tau(y_1)\varphi([x_2,y_2],z)-(-1)^{\bar{y}_1(\bar{x}_1+\bar{x}_2)+\bar{z}(\bar{y}_1+\bar{y}_2+\bar{x}_2)}\tau(z)\varphi(y_1,[x_2,y_2]) \\
& -(-1)^{(\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2)}\tau(y_1)\varphi(y_2,[x_2,z])+(-1)^{(\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2)+\bar{y}_1\bar{y}_2}\tau(x_2)\varphi(y_1,[x_2,z]) \\
& +(-1)^{(\bar{x}_1+\bar{x}_2)\bar{\phi}}\tau(y_1)[x_2,\varphi(y_2,z)]-(-1)^{(\bar{x}_1+\bar{x}_2)\bar{\phi}+\bar{y}_1\bar{y}_2}\tau(y_2)[x_2,\varphi(y_1,z)] \\
& +(-1)^{(\bar{x}_1+\bar{x}_2)\bar{\phi}+\bar{z}(\bar{y}_2+\bar{y}_2)}\tau(z)[x_2,\varphi(y_1,y_2)] \\
& +(-1)^{\bar{x}_1\bar{x}_2}\tau(x_2)\left(-(-1)^{\bar{y}_2(\bar{x}_1+\bar{x}_2)}\tau(y_2)\varphi([x_1,y_1],z)+(-1)^{\bar{z}(\bar{x}_1+\bar{y}_1+\bar{y}_2)}\tau(z)\varphi([x_1,y_1],y_2)\right) \\
& +(-1)^{\bar{y}_1(\bar{x}_1+\bar{x}_2)}\tau(y_1)\varphi([x_1,y_1],z)+(-1)^{\bar{y}_1(\bar{x}_1+\bar{x}_2)+\bar{z}(\bar{x}_1+\bar{y}_1+\bar{y}_2)}\tau(z)\varphi(y_1,[x_1,y_2]) \\
& +(-1)^{(\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2)}\tau(y_1)\varphi(y_2,[x_1,z])-(-1)^{(\bar{y}_1+\bar{y}_2)(\bar{x}_1+\bar{x}_2)+\bar{y}_1\bar{y}_2}\tau(y_2)\varphi(y_1,[x_1,z]) \\
& -(-1)^{(\bar{x}_1+\bar{x}_2)\bar{\phi}}\tau(y_1)[x_1,\varphi(y_1,z)]+(-1)^{(\bar{x}_1+\bar{x}_2)\bar{\phi}+\bar{y}_1\bar{y}_2}\tau(y_2)[x_1,\varphi(y_1,z)] \\
& -(-1)^{(\bar{x}_1+\bar{x}_2)\bar{\phi}+\bar{z}(\bar{y}_1+\bar{y}_2)}\tau(z)[x_1,\varphi(y_1,y_2)] \\
& +(-1)^{(\bar{x}_1+\bar{x}_2)(\bar{y}_1+\bar{y}_2+\bar{z})}\left(\tau(y_1)\tau(\varphi(y_2,z))-(-1)^{\bar{y}_1\bar{y}_2}\tau(y_2)\tau(\varphi(y_1,z))\right) \\
& +(-1)^{\bar{z}(\bar{y}_1+\bar{y}_2)}\tau(z)\tau(\varphi(y_1,y_2))\left)[x_1,x_2\right] \\
& -\left(\tau(x_1)\tau(\varphi(x_2,y_1))-(-1)^{\bar{x}_1\bar{x}_2}\tau(x_2)\tau(\varphi(x_1,y_1))\right) \\
& +(-1)^{\bar{y}_1(\bar{x}_1+\bar{x}_2)}\tau(y_1)\tau(\varphi(x_1,x_2))\left)[y_2,z\right] \\
& +(-1)^{\bar{y}_1\bar{y}_2}\left(\tau(x_1)\tau(\varphi(x_2,y_2))-(-1)^{\bar{x}_1\bar{x}_2}\tau(x_2)\tau(\varphi(x_1,y_2))\right) \\
& +(-1)^{\bar{y}_2(\bar{x}_1+\bar{x}_2)}\tau(y_2)\tau(\varphi(x_1,x_2))\left)[y_1,z\right] \\
& -(-1)^{\bar{z}(\bar{y}_1+\bar{y}_2)}\left(\tau(x_1)\tau(\varphi(x_2,z))-(-1)^{\bar{x}_1\bar{x}_2}\tau(x_2)\tau(\varphi(x_1,z))\right) \\
& +(-1)^{\bar{z}(\bar{x}_1+\bar{x}_2)}\tau(z)\tau(\varphi(x_1,x_2))\left)[y_1,y_2\right] \\
& = -\tau(y_1)\tau(x_1)\delta_{LR}\varphi(z,y_2,x_2)-\tau(y_1)\tau(x_2)\delta_{LR}\varphi(y_2,z,x_1)
\end{aligned}$$

$$\begin{aligned}
& -\tau(y_2)\tau(x_1)\delta_{LR}\varphi(y_1, z, x_2) - \tau(y_2)\tau(x_2)\delta_{LR}\varphi(z, y_1, x_1) \\
& -\tau(z)\tau(x_1)\delta_{LR}\varphi(y_2, y_1, x_2) - \tau(z)\tau(x_2)\delta_{LR}\varphi(y_1, y_2, x_1) \\
& +(-1)^{(\bar{x}_1+\bar{x}_2)(\bar{y}_1+\bar{y}_2+\bar{z})}\left(\tau(y_1)\tau(\varphi(y_2, z)) - (-1)^{\bar{y}_1\bar{y}_2}\tau(y_2)\tau(\varphi(y_1, z))\right) \\
& +(-1)^{\bar{z}(\bar{y}_1+\bar{y}_2)}\tau(z)\tau(\varphi(y_1, y_2))\left)[x_1, x_2\right] \\
& -\left(\tau(x_1)\tau(\varphi(x_2, y_1)) - (-1)^{\bar{x}_1\bar{x}_2}\tau(x_2)\tau(\varphi(x_1, y_1))\right) \\
& +(-1)^{\bar{y}_1(\bar{x}_1+\bar{x}_2)}\tau(y_1)\tau(\varphi(x_1, x_2))\left)[y_2, z\right] \\
& +(-1)^{\bar{y}_1\bar{y}_2}\left(\tau(x_1)\tau(\varphi(x_2, y_2)) - (-1)^{\bar{x}_1\bar{x}_2}\tau(x_2)\tau(\varphi(x_1, y_2))\right) \\
& +(-1)^{\bar{y}_2(\bar{x}_1+\bar{x}_2)}\tau(y_2)\tau(\varphi(x_1, x_2))\left)[y_1, z\right] \\
& -(-1)^{\bar{z}(\bar{y}_1+\bar{y}_2)}\left(\tau(x_1)\tau(\varphi(x_2, z)) - (-1)^{\bar{x}_1\bar{x}_2}\tau(x_2)\tau(\varphi(x_1, z))\right) \\
& +(-1)^{\bar{z}(\bar{x}_1+\bar{x}_2)}\tau(z)\tau(\varphi(x_1, x_2))\left)[y_1, y_2\right].
\end{aligned}$$

Since for all $x, y, z \in \mathcal{H}(L)$,

$$\tau(x)\tau(\varphi(y, z)) - (-1)^{\bar{x}\bar{y}}\tau(y)\tau(\varphi(x, z)) + (-1)^{\bar{z}(\bar{x}+\bar{y})}\tau(z)\tau(\varphi(x, y)) = 0,$$

we get $\delta_{3LR}\phi = 0$. \square

Corollary 3.9. Let $\varphi \in Z_{LR}^2(L, \mathbb{K})$. Then

$$\psi(x, y, z) = \tau(x)\varphi(y, z) - (-1)^{\bar{x}\bar{y}}\tau(y)\varphi(x, z) + (-1)^{\bar{z}(\bar{x}+\bar{y})}\tau(z)\varphi(x, y)$$

is a 2-cocycle of the induced 3-Lie-Rinehart superalgebra.

Theorem 3.10. Every 1-cocycle for the scalar cohomology of the Lie-Rinehart superalgebra $(L, A, [\cdot, \cdot], \mu)$ is a 1-cocycle for the scalar cohomology of the induced 3-Lie-Rinehart superalgebra $(L, A, [\cdot, \cdot]_\tau, \rho_\tau)$.

Proof. Let $\omega \in Z_{LR}^1(L, \mathbb{K})$. Then,

$$\forall x, y \in L : \delta_{LR}\omega(x, y) = \omega([x, y]) = 0,$$

which is equivalent to $[L, L] \subset \ker \omega$. It is obvious to prove that $[L, L, L]_\tau \subset [L, L]$ and then $[L, L, L]_\tau \subset \ker \omega$, that is

$$\forall x, y, z \in L : \omega([x, y, z]_\tau) = \delta_{3LR}\omega(x, y, z) = 0,$$

which means that ω is a 1-cocycle for the scalar cohomology of $(L, A, [\cdot, \cdot]_\tau, \rho_\tau)$. \square

Lemma 3.11. Let $\varphi \in C^1(L, \mathbb{K})$. Then, for all $x, y, z \in L$,

$$\delta_{3LR}\varphi(x, y, z) = \tau(x)\delta_{LR}\varphi(y, z) - (-1)^{\bar{x}\bar{y}}\tau(y)\delta_{LR}\varphi(x, z) + (-1)^{\bar{z}(\bar{x}+\bar{y})}\tau(z)\delta_{LR}\varphi(x, y).$$

Proof. Let $\varphi \in C^1(L, \mathbb{K})$, $x, y, z \in L$. Then

$$\begin{aligned}
\delta_{3LR}\varphi(x, y, z) &= \varphi([x, y, z]_\tau) \\
&= \tau(x)\varphi([y, z]) - (-1)^{\bar{x}\bar{y}}\tau(y)\varphi([x, z]) + (-1)^{\bar{z}(\bar{x}+\bar{y})}\tau(z)\varphi([x, y]) \\
&= \tau(x)\delta_{LR}\varphi(y, z) - (-1)^{\bar{x}\bar{y}}\tau(y)\delta_{LR}\varphi(x, z) + (-1)^{\bar{z}(\bar{x}+\bar{y})}\tau(z)\delta_{LR}\varphi(x, y).
\end{aligned}$$

completing the proof. \square

Proposition 3.12. Let $\varphi_1, \varphi_2 \in Z_{LR}^2(L, \mathbb{K})$. If φ_1, φ_2 are in the same cohomology class then ψ_1, ψ_2 defined by:

$$\psi_i(x, y, z) = \tau(x) \varphi_i(y, z) - (-1)^{\bar{x}\bar{y}} \tau(y) \varphi_i(x, z) + (-1)^{\bar{z}(\bar{x}+\bar{y})} \tau(z) \varphi_i(x, y), i = 1, 2$$

are in the same cohomology class.

Proof. If $\varphi_1, \varphi_2 \in Z_{LR}^2(L, \mathbb{K})$ are two cocycles in the same cohomology class, that is $\varphi_2 - \varphi_1 = \delta_{LR} v, v \in C^1(L, \mathbb{K})$, then

$$\begin{aligned} \psi_2(x, y, z) - \psi_1(x, y, z) &= \tau(x) \varphi_2(y, z) - (-1)^{\bar{x}\bar{y}} \tau(y) \varphi_2(x, z) + (-1)^{\bar{z}(\bar{x}+\bar{y})} \tau(z) \varphi_2(x, y) \\ &\quad - \tau(x) \varphi_1(y, z) - (-1)^{\bar{x}\bar{y}} \tau(y) \varphi_1(x, z) + (-1)^{\bar{z}(\bar{x}+\bar{y})} \tau(z) \varphi_1(x, y) \\ &= \tau(x) (\varphi_2 - \varphi_1)(y, z) - (-1)^{\bar{x}\bar{y}} \tau(y) (\varphi_2 - \varphi_1)(x, z) + (-1)^{\bar{z}(\bar{x}+\bar{y})} \tau(z) (\varphi_2 - \varphi_1)(x, y) \\ &= \tau(x) \delta_{LR} v(y, z) - (-1)^{\bar{x}\bar{y}} \tau(y) \delta_{LR} v(x, z) + (-1)^{\bar{z}(\bar{x}+\bar{y})} \tau(z) \delta_{LR} v(x, y) = \delta_{3LR} v(x, y, z), \end{aligned}$$

which means that ψ_1 and ψ_2 are in the same cohomology class. \square

3.2 Deformations of 3-Lie-Rinehart superalgebras

Let $(L, A, [\cdot, \cdot, \cdot], \rho)$ be a 3-Lie-Rinehart superalgebra. Denote by $\mathbb{K}[[t]]$ the space of formal power series ring with parameter t .

Definition 3.13. A deformation of a 3-Lie-Rinehart superalgebra $(L, A, [\cdot, \cdot, \cdot], \rho)$ is a $\mathbb{K}[[t]]$ -trilinear map

$$[\cdot, \cdot, \cdot]_t : L[[t]] \times L[[t]] \times L[[t]] \rightarrow L[[t]], \quad [x, y, z]_t = \sum_{i \geq 0} t^i m_i(x, y, z),$$

with $m_0 = [\cdot, \cdot, \cdot]$ and m_i are even trilinear maps for $i \geq 0$, satisfying (1.6), (1.14) and (1.15).

Let $[\cdot, \cdot, \cdot]_t$ be a deformation of $[\cdot, \cdot, \cdot]$. Then

$$\begin{aligned} m_t(x, y, m_t(z, u, v)) - m_t(m_t(x, y, u), v, w) - (-1)^{\bar{u}(\bar{x}+\bar{y})} m_t(u, m_t(x, y, v), w) \\ - (-1)^{(\bar{u}+\bar{v})(\bar{x}+\bar{y})} m_t(u, v, m_t(x, y, w)) = 0. \end{aligned}$$

Comparing the coefficients of $t^n, n \geq 0$, we get the following equation:

$$\begin{aligned} \sum_{i,j=0}^n (m_i(x, y, m_j(z, u, v)) - m_i(m_j(x, y, u), v, w) - (-1)^{\bar{u}(\bar{x}+\bar{y})} m_i(u, m_j(x, y, v), w) \\ - (-1)^{(\bar{u}+\bar{v})(\bar{x}+\bar{y})} m_i(u, v, m_j(x, y, w))) = 0. \end{aligned}$$

Definition 3.14. The 3-cochain m_1 is called the infinitesimal of the deformation $[\cdot, \cdot, \cdot]_t$. More generally, if $m_i = 0$ for $1 \leq i \leq (n-1)$ and m_n is non zero cochain, then m_n is called the n -infinitesimal of the deformation $[\cdot, \cdot, \cdot]_t$.

Definition 3.15. Two deformations $[\cdot, \cdot, \cdot]_t$ and $[\cdot, \cdot, \cdot]'_t$ are said to be equivalent if there exists a formal automorphism

$$\Phi_t : L[[t]] \rightarrow L[[t]], \quad \Phi_t = id_L + \sum_{i \geq 1} t^i \phi_i,$$

where $\phi_i : L \rightarrow L$ is a even \mathbb{K} -linear maps such that

$$[x, y, z]'_t = \Phi_t^{-1}[\Phi_t(x), \Phi_t(y), \Phi_t(z)]_t.$$

Definition 3.16. A deformation is called trivial if it is equivalent to the deformation $m_0 = [\cdot, \cdot, \cdot]$.

Definition 3.17. A 3-Lie-Rinehart superalgebra is said to be rigid if every deformation of it is trivial.

Theorem 3.18. The cohomology class of the infinitesimal of a deformation $[\cdot, \cdot, \cdot]_t$ is determined by the equivalence class of $[\cdot, \cdot, \cdot]_t$.

Proof. Let Φ_t represents an equivalence of deformation given by m_t and \tilde{m}_t . Then

$$\tilde{m}_t(x, y, z) = \Phi_t^{-1} m_t(\Phi_t(x), \Phi_t(y), \Phi_t(z)).$$

Comparing the coefficients of t from both sides of the above equation we have

$$\begin{aligned} \tilde{m}_1(x, y, z) + \Phi_1([x, y, z]) &= m_1(x, y, z) + [\Phi_1(x), y, z] + (-1)^{\bar{x}\bar{\Phi}} [x, \Phi_1(y), z] \\ &\quad + (-1)^{(\bar{x}+\bar{y})\bar{\Phi}} [x, y, \Phi_1(z)], \end{aligned}$$

or equivalently, $m_1 - \tilde{m}_1 = \delta_{3LR}(\phi_1)$. So, cohomology class of infinitesimal of the deformation is determined by the equivalence class of deformation of m_t . \square

Theorem 3.19. A non-trivial deformation of a 3-Lie-Rinehart superalgebra is equivalent to a deformation whose n -infinitesimal cochain is not a coboundary for some $n \geq 1$.

Proof. Let m_t be a deformation of 3-Lie-Rinehart superalgebra with n -infinitesimal m_n for some $n \geq 1$. Assume that there exists a 3-cochain $\phi \in C^2(L, L)$ with $\delta(\phi) = m_n$. Set $\Phi_t = id_L + \phi t^n$ and $\tilde{m} = \Phi_t^{-1} \circ m_t \circ \Phi_t$. Comparing the coefficients of t^n , we get the following equation:

$$\tilde{m}_n - m_n = -\delta_{3LR}(\phi).$$

So $\tilde{m}_n = 0$. Then, one has the deformation whose n -infinitesimal is not a coboundary for some $n \geq 1$. \square

References

- [1] Abramov, V., Super 3-Lie algebras induced by super Lie algebras, *Adv. Appl. Clifford Algebras*, 27 (1), 9-16 (2017)
- [2] Abramov, V., 3-Lie superalgebras induced by Lie superalgebras, *Axioms*, 8 (1), 21 (2019)
- [3] Abramov, V., Weil algebra, 3-Lie algebra and B.R.S. algebra, In: Silvestrov, S., Malyarenko, A., Rancic, M. (Eds.), *Algebraic Structures and Applications*, Springer Proceedings in Mathematics and Statistics, Vol 317, Springer, Ch 1, 1-12 (2020).
- [4] Abramov, V., Lätt, P., Classification of low dimensional 3-Lie superalgebras, In: Silvestrov S., Rancic M. (eds), *Engineering Mathematics II*, Springer Proceedings in Mathematics and Statistics, Vol 179, Springer, 1-12 (2016)
- [5] Abramov, V., Lätt, P., Ternary Lie superalgebras and Nambu-Hamilton equation in superspace, In: Silvestrov S., Malyarenko A., Rancic, M. (eds), *Algebraic Structures and Applications*. Springer Proceedings in Mathematics and Statistics, vol 317, Springer, Ch 3, 47-80 (2020)

- [6] Abramov, V., Silvestrov, S., 3-Hom-Lie algebras based on σ -derivation and involution, *Adv. Appl. Clifford Algebras*, 30, 45 (2020)
- [7] Arnlind, J., Kitouni, A., Makhlouf, A., Silvestrov, S., Structure and cohomology of 3-Lie algebras induced by Lie algebras, In: Makhlouf, A., Paal, E., Silvestrov, S. D., Stolin, A., *Algebra, Geometry and Mathematical Physics*, Springer Proceedings in Mathematics and Statistics, vol 85, Springer, 123-144 (2014)
- [8] Arnlind, J., Makhlouf, A., Silvestrov, S., Ternary Hom-Nambu-Lie algebras induced by Hom-Lie algebras, *J. Math. Phys.* 51, 043515, 11 pp. (2010)
- [9] Arnlind, J., Makhlouf, A., Silvestrov, S., Construction of n -Lie algebras and n -ary Hom-Nambu-Lie algebras, *J. Math. Phys.* 52, 123502, 13 pp. (2011)
- [10] Albuquerque, H., Barreiro, E., Calderón, A. J., Sánchez, J. M., Split Lie-Rinehart algebras, *J. Algebra Appl.* DOI:10.1142/S0219498821501644 (2020), arXiv:1706.07084 (2017)
- [11] Awata, H., Li, M., Minic, D., Yoneya, T., On the quantization of Nambu brackets, *J. High Energy Phys.* 2, Paper 13, 17 pp. (2001)
- [12] Bai, R., Li, X., Wu, Y., 3-Lie-Rinehart algebras, arXiv:1903.12283v1 (2019)
- [13] Bai, R., Li, X., Wu, Y., Hom 3-Lie-Rinehart algebras, arXiv:2001.07570 (2020)
- [14] Bai, R., Bai, C., Wang, J., Realizations of 3-Lie algebras, *J. Math. Phys.* 51, 063505 (2010)
- [15] Bai, R., Wu, Y., Constructions of 3-Lie algebras, *Lin. Multilin. Alg.* 63 (11), 2171-2186 (2015)
- [16] Bai, R., Li, Z., Wang, W., Infinite-dimensional 3-Lie algebras and their connections to Harish-Chandra module, *Front. Math. China*, 12 (3), 515-530 (2017)
- [17] Ben Hassine, A., Mabrouk, S., Ncib, O., Some constructions of multiplicative n -ary Hom-Nambu algebras, *Adv. Appl. Clifford Algebras*, 29 (5), 88 (2019)
- [18] Bkouche, R., Structures (K, A) -linéaires, *C. R. Acad. Sci. Paris Sér. A-B* 262, A373-A376 (1966) (in French)
- [19] Calderon Martin, A. J., Forero Piulestan, M., Split 3-Lie algebras, *J. Math. Phys.* 52 (12) 123503 (2011)
- [20] Cantarini, N., Kac, V. G., Classification of simple linearly compact n -Lie superalgebras, *Comm. Math. Phys.* 298(3), 833-853 (2010)
- [21] Casas, J. M., Obstructions to Lie-Rinehart algebra extensions, *Algebra Colloq.* 18 (1), 83-104 (2011)
- [22] Casas, J. M., Ladra, M., Pirashvili, T., Crossed modules for Lie-Rinehart algebras, *J. Algebra*, 274(1), 192-201 (2004)
- [23] Casas, J. M., Ladra, M., Pirashvili, T., Triple cohomology of Lie-Rinehart algebras and the canonical class of associative algebras, *J. Algebra*, 291 (1), 144-163 (2005)
- [24] Chemla, S., Operations for modules on Lie-Rinehart superalgebras, *Manuscripta Math.* 87 (2), 199-224 (1995)
- [25] Chen, Z., Liu, Z., Zhong, D., Lie-Rinehart bialgebras for crossed products. *J. Pure Appl. Algebra* 215 (6), 1270-1283 (2011)

- [26] Dokas, I., Cohomology of restricted Lie-Rinehart algebras. *Adv. Math.* 231 (5), 2573-2592 (2012)
- [27] Daletskii, Y. L., Kushnirenich, V. A., Inclusion of the Nambu-Takhtajan algebra in the structure of formal differential geometry, *Dopov. Akad. Nauk Ukr.* 1996, 12-17 (1996)
- [28] Filippov, V., n -Lie algebras, *Sib. Mat. Zh.* 26, 126-140 (1985)
- [29] Guan, B., Chen L., Sun, B., 3-ary Hom-Lie superalgebras induced by Hom-Lie superalgebras, *Adv. Appl. Clifford Algebras*, 27, 3063-3082 (2017)
- [30] Guo, S., Zhang, X., Wang, S., On 3-Hom-Lie-Rinehart algebras, arXiv:1911.10992 (2019)
- [31] Herz, J., Pseudo-algebras de Lie. I, II, *C. R. Acad. Sci. Paris* 236, 1935-1937, 2289-2291 (1953)
- [32] Higgins, P. J., Mackenzie, K., Algebraic constructions in the category of Lie algebroids, *J. Algebra*, 129 (1), 194-230 (1990)
- [33] Huebschmann, J., Poisson cohomology and quantization. *J. Reine Angew. Math.* 408, 57-113 (1990)
- [34] Huebschmann, J., Lie-Rinehart algebras, Gerstenhaber algebras and Batalin-Vilkovisky algebras, *Ann. Inst. Fourier (Grenoble)*, 48, 425-440 (1998)
- [35] Huebschmann, J., Duality for Lie-Rinehart algebras and the modular class. *J. Reine Angew. Math.* 510, 103-159 (1999)
- [36] Huebschmann, J., Lie-Rinehart algebras, descent, and quantization, In: *Galois Theory, Hopf Algebras, and Semiabelian Categories*, Fields Inst. Commun., vol. 43, Amer. Math. Soc., Providence, RI, 295-316 (2004)
- [37] Huebschmann, J., Extensions of Lie-Rinehart algebras and cotangent bundle reduction, *Proc. Lond. Math. Soc.* 107 (5), 1135-1172 (2013)
- [38] Huebschmann, J., Multi derivation Maurer-Cartan algebras and sh Lie-Rinehart algebras, *J. Algebra*, 472, 437-479 (2017)
- [39] Kraemer, U., Rovi, A., A Lie-Rinehart algebra with no antipode, *Comm. Algebra*, 43 (10), 4049-4053 (2015)
- [40] Kac, V. G., A sketch of Lie superalgebra theory, *Commun. Math. Phys.* 53 (1), 31-64 (1977)
- [41] Kitouni, A., Makhlouf, A., Silvestrov, S., On $(n + 1)$ -Hom-Lie algebras induced by n -Hom-Lie algebras, *Georgian Math. J.* 23 (1), 75-95 (2016)
- [42] Kitouni, A., Makhlouf, A., Silvestrov, S., On solvability and nilpotency for n -Hom-Lie algebras and $(n + 1)$ -Hom-Lie algebras induced by n -Hom-Lie algebras, In: *Silvestrov, S., Malyarenko, A., Rancic, M. (Eds.), Algebraic Structures and Applications*, Springer Proceedings in Mathematics and Statistics, Vol 317, Springer, Ch 6, 127-157 (2020)
- [43] Mackenzie, K., Lie groupoids and Lie algebroids in differential geometry, *London Math. Soc. Lecture Note Series* 124, Cambridge University Press, Cambridge (1987)
- [44] Mackenzie, K. C. H., Lie algebroids and Lie pseudoalgebras, *Bull. London Math. Soc.* 27 (2) 97-147 (1995)

- [45] Mandal, A., Mishra, S. K., Hom-Lie-Rinehart algebras, *Comm. Algebra*, 46 (9), 3722-3744 (2016)
- [46] Mandal, A., Mishra, S. K., On hom-Gerstenhaber algebras, and hom-Lie algebroids, *J. Geom. Phys.* 133, 287-302 (2018)
- [47] Mishra, S. K., Silvestrov, S., A review on Hom-Gerstenhaber algebras and Hom-Lie algebroids, In: Silvestrov S., Malyarenko A., Rančić, M. (eds), *Algebraic Structures and Applications*, Springer Proceedings in Mathematics and Statistics, vol 317. Springer, Ch 11, 285-315 (2020)
- [48] Palais, R., The cohomology of Lie rings, *Proc. Symp. Pure Math.*, Amer. Math. Soc., Providence, R. I., 130-137 (1961)
- [49] Rinehart, G., Differential forms on general commutative algebras, *Trans. Amer. Math. Soc.* 108, 195-222 (1963)
- [50] Rota, G., Baxter algebras and combinatorial identities I, II, *Bull. Amer. Math. Soc.* 75, 325-329, 330-334 (1969)
- [51] Takhtajan, L., On foundation of the generalized Nambu mechanics, *Comm. Math. Phys.* 160, 295-315 (1994)