

Statistical complexity is maximized close to criticality in cortical dynamics

F. A. B. Coutinho¹, P. Carelli¹, M. A. A. S. Pereira¹, M. A. A. S. Pereira², M. A. A. S. Pereira³, M. A. A. S. Pereira^{3,*}, M. A. A. S. Pereira^{1,†}

¹Departamento de Física, Universidade Federal de Pernambuco, Recife PE 50670-901, Brazil.

²Departamento de Ciências Fundamentais e Sociais,
Universidade Federal da Paraíba, Areia PB 58397-000 Brazil.

³Instituto de Física, Universidade Federal de Alagoas, Maceió, Alagoas 57072-970 Brazil.

Complex systems are typically characterized as an intermediate situation between a complete regular structure and a random system. Brain signals can be studied as a striking example of such systems: cortical states can range from highly synchronous and ordered neuronal activity (with higher spiking variability) to desynchronized and disordered regimes (with lower spiking variability). It has been recently shown, by testing independent signatures of criticality, that a phase transition occurs in a cortical state of intermediate spiking variability. Here, we use a symbolic information approach to show that, despite the monotonical increase of the Shannon entropy between ordered and disordered regimes, we can determine an intermediate state of maximum complexity based on the Jensen disequilibrium measure. More specifically, we show that statistical complexity is maximized close to criticality for cortical spiking data of urethane-anesthetized rats, as well as for a network model of excitable elements that presents a critical point of a non-equilibrium phase transition.

PACS numbers:

I. INTRODUCTION

Statistical complexity is a measure of the amount of information contained in a system's state space. It is defined as the Shannon entropy of the system's state space, normalized by the system's state space size. In this paper, we show that statistical complexity is maximized close to criticality in cortical dynamics. We use a symbolic information approach to show that, despite the monotonical increase of the Shannon entropy between ordered and disordered regimes, we can determine an intermediate state of maximum complexity based on the Jensen disequilibrium measure. More specifically, we show that statistical complexity is maximized close to criticality for cortical spiking data of urethane-anesthetized rats, as well as for a network model of excitable elements that presents a critical point of a non-equilibrium phase transition.

Statistical complexity is a measure of the amount of information contained in a system's state space. It is defined as the Shannon entropy of the system's state space, normalized by the system's state space size. In this paper, we show that statistical complexity is maximized close to criticality in cortical dynamics. We use a symbolic information approach to show that, despite the monotonical increase of the Shannon entropy between ordered and disordered regimes, we can determine an intermediate state of maximum complexity based on the Jensen disequilibrium measure. More specifically, we show that statistical complexity is maximized close to criticality for cortical spiking data of urethane-anesthetized rats, as well as for a network model of excitable elements that presents a critical point of a non-equilibrium phase transition.

Statistical complexity is a measure of the amount of information contained in a system's state space. It is defined as the Shannon entropy of the system's state space, normalized by the system's state space size. In this paper, we show that statistical complexity is maximized close to criticality in cortical dynamics. We use a symbolic information approach to show that, despite the monotonical increase of the Shannon entropy between ordered and disordered regimes, we can determine an intermediate state of maximum complexity based on the Jensen disequilibrium measure. More specifically, we show that statistical complexity is maximized close to criticality for cortical spiking data of urethane-anesthetized rats, as well as for a network model of excitable elements that presents a critical point of a non-equilibrium phase transition.

Statistical complexity is a measure of the amount of information contained in a system's state space. It is defined as the Shannon entropy of the system's state space, normalized by the system's state space size. In this paper, we show that statistical complexity is maximized close to criticality in cortical dynamics. We use a symbolic information approach to show that, despite the monotonical increase of the Shannon entropy between ordered and disordered regimes, we can determine an intermediate state of maximum complexity based on the Jensen disequilibrium measure. More specifically, we show that statistical complexity is maximized close to criticality for cortical spiking data of urethane-anesthetized rats, as well as for a network model of excitable elements that presents a critical point of a non-equilibrium phase transition.

Statistical complexity is a measure of the amount of information contained in a system's state space. It is defined as the Shannon entropy of the system's state space, normalized by the system's state space size. In this paper, we show that statistical complexity is maximized close to criticality in cortical dynamics. We use a symbolic information approach to show that, despite the monotonical increase of the Shannon entropy between ordered and disordered regimes, we can determine an intermediate state of maximum complexity based on the Jensen disequilibrium measure. More specifically, we show that statistical complexity is maximized close to criticality for cortical spiking data of urethane-anesthetized rats, as well as for a network model of excitable elements that presents a critical point of a non-equilibrium phase transition.

Statistical complexity is a measure of the amount of information contained in a system's state space. It is defined as the Shannon entropy of the system's state space, normalized by the system's state space size. In this paper, we show that statistical complexity is maximized close to criticality in cortical dynamics. We use a symbolic information approach to show that, despite the monotonical increase of the Shannon entropy between ordered and disordered regimes, we can determine an intermediate state of maximum complexity based on the Jensen disequilibrium measure. More specifically, we show that statistical complexity is maximized close to criticality for cortical spiking data of urethane-anesthetized rats, as well as for a network model of excitable elements that presents a critical point of a non-equilibrium phase transition.

*fernanda@fis.ufal.br
†pedro.carelli@ufpe.br

FIG. 1

II. METHODS

A. Information-theoretical quantifiers

A time series $X(t)$ is a sequence of values $X(t) \equiv \{x_t; t = 1, 2, \dots, M\}$, where M is the length of the series. The probability distribution $P \equiv \{p_j; j = 1, 2, \dots, N\}$ is defined by $\sum_{j=1}^N p_j = 1$ and $p_j \geq 0$.

$$S[P] = - \sum_{j=1}^N p_j \ln p_j. \quad (1)$$

The entropy $H[P]$ is defined as $H[P] = S[P]/S[P_e]$, where $P_e = \{p_j = 1/N; j = 1, 2, \dots, N\}$ is the uniform distribution. The quantity $Q_J[P, P_e]$ is defined as $Q_J[P, P_e] = J[P, P_e]/J[P_e, P_e]$, where $J[P, P_e] = S[P, P_e]/S[P_e]$ is the joint entropy. The quantity C_+ is defined as $C_+ = 1 - Q_J[P, P_e]$.


$$C[P] = Q_J[P, P_e] \cdot H[P]. \quad (2)$$

$$Q_J[P, P_e] = Q_0 J[P, P_e], \quad (3)$$


$$J[P, P_e] = S \left[\frac{P + P_e}{2} \right] - \frac{S[P]}{2} - \frac{S[P_e]}{2}, \quad (4)$$

where Q_0 is a constant (usually $Q_0 = 1$) and $0 \leq Q_J \leq 1$.

(a)



(b)



(c)




FIG. 1: Characterizing the symbolic representation of time series. (a) The six possible symbols associated with permutations π_j for ordinal patterns of length $D = 3$. (b) Example of a very simple time series $X(t)$ and (c) its own non-normalized probability density function (PDF).

The joint entropy $J[P, P_e]$ is defined as $J[P, P_e] = S[P, P_e]/S[P_e]$, where $S[P, P_e]$ is the joint entropy of P and P_e . The quantity C_+ is defined as $C_+ = 1 - Q_J[P, P_e]$.

B. Symbolic representation of a time series

A time series $X(t)$ of length M is converted into a symbolic representation $s(t)$ of length D , where $s(t) \in \{0, 1, 2\}^D$ is a sequence of symbols of length D . The symbolic representation is defined as $s(t) = \{x_{t-(D-1)}, x_{t-(D-2)}, \dots, x_{t-1}, x_t\}$.




FIG. 2: Characterizing cortical states and neuronal variability. (a) Firing rates F calculated using 10 ms interval. (b) Raster plots for all 274 recorded neurons. (c) Coefficient of variation (CV) of the spiking activity calculated for 10-s-long non-overlapping windows of a single animal and the CV histogram. Symbols (circle, triangle, and square) respectively indicate the level of spiking variability of three representative examples: high ($CV = 2.60$, highly synchronous and ordered neuronal activity) intermediate ($CV = 1.30$), and low ($CV = 0.484$, desynchronized and disordered regime). (d) The entropy and (e) the complexity of each window and their histograms. The three illustrative examples exhibit: $H = 0.241$ and $C = 0.190$ (green circle), $H = 0.596$ and $C = 0.304$ (red triangle), $H = 0.917$ and $C = 0.206$ (blue square).

$CV = 2.60$,
 $CV = 1.30$,
 $CV = 0.484$,
 CV by

$CV =$
 $H =$
 $CV = 2.60$,
 $H = 0.241$,
 $H = 0.596$,
 $H = 0.917$,
 $CV = 1.30$,
 $CV = 0.484$,
 CV, H

B. Statistical complexity is maximized between the synchronized and desynchronized cortical states

$C = 0.190$,
 $C = 0.206$,
 $C = 0.304$,
 C versus CV ,
 $CV = 0.15$,
 $H = 0.05$,
 $CV_{C_{max}} = 1.31 \pm 0.15$,
 $C_{max} = 0.330 \pm 0.001$

$$H_{C_{max}} = 0.66 \pm 0.05.$$

Fig. 4. Dependence of the symbolic Shannon entropy H and the statistical complexity C on the coefficient of variation CV of the spiking activity. The data points are shown for different levels of spiking variability. The error bars represent the standard deviation. The values of C_{max} and $H_{C_{max}}$ are indicated in the text.

The dependence of the symbolic Shannon entropy H and the statistical complexity C on the coefficient of variation CV of the spiking activity is shown in Fig. 4. The data points are shown for different levels of spiking variability. The error bars represent the standard deviation. The values of C_{max} and $H_{C_{max}}$ are indicated in the text.

C. Statistical complexity is maximized close to the phase-transition in a network model

The network model is defined by the following equations:

$$s_i(t+1) = (s_i(t) + \sum_{j=1}^N p_{ij} s_j(t) - r \delta t) \bmod 2$$

where $s_i(t) \in \{0, 1\}$ is the state of the i -th node at time t , p_{ij} is the transition probability, r is the reset rate, and δt is the time step. The network size is $N=10^5$.




FIG. 3: Information-theory quantifiers for different levels of spiking variability of the cortical states of a single animal. (a) symbolic Shannon entropy and (b) statistical complexity versus coefficient of variation (CV) of the spiking activity. (c) Complexity–entropy plane. The peak in complexity occurs for $C_{max} = 0.330 \pm 0.001$, $\langle CV \rangle_{C_{max}} = 1.31 \pm 0.15$, $H_{C_{max}} = 0.66 \pm 0.05$.

The network model is defined by the following equations:

$$s_i(t+1) = (s_i(t) + \sum_{j=1}^N p_{ij} s_j(t) - r \delta t) \bmod 2$$

where $s_i(t) \in \{0, 1\}$ is the state of the i -th node at time t , p_{ij} is the transition probability, r is the reset rate, and δt is the time step. The network size is $N=10^5$.




FIG. 4: Information-theory quantifiers as in Fig. 3 for all animals together. (a) Shannon entropy and (b) statistical complexity plotted along with the respective coefficient of variation (CV) of the spiking activity for each window of time. (c) Complexity–entropy plane. The dashed line represents the mean values of C and H . The peak in complexity occurs for $C_{max} = 0.333 \pm 0.001$, $\langle CV \rangle_{C_{max}} = 1.35 \pm 0.15$, $H_{C_{max}} = 0.70 \pm 0.05$




FIG. 5: Comparison between shuffled (dashed line) and real (continuous line) data for all rats. (a) Average (lines) and standard deviation (shading) of complexity C plotted against the CV associated with each window before the shuffling. (b) Average (lines) and standard deviation (shading) of complexity and entropy in the complexity-entropy plane.

$\langle CV \rangle_{C_{max}} = 1.58 \pm 0.15$, $H_{C_{max}} = 0.60 \pm 0.05$, $C_{max} = 0.347 \pm 0.001$.

IV. CONCLUDING REMARKS

- 012408 (2020).
- [15] M. E. Hasselmo, *Behavioural brain research* **67**, 1 (1995).
- [16] N. M. Timme, N. J. Marshall, N. Bennett, M. Ripp, E. Lautzenhiser, and J. M. Beggs, *Frontiers in physiology* **7**, 425 (2016).
- [17] C. Shannon and W. Weaver, *The mathematical theory of communication* (Champaign, IL: University of Illinois Press, 1949).
- [18] P. W. Lamberti, M. T. Martín, A. Plastino, and O. A. Rosso, *Physica A: Statistical Mechanics and its Applications* **334**, 119 (2004).
- [19] R. Lopez-Ruiz, H. L. Mancini, and X. Calbet, *Physics letters A* **209**, 321 (1995).
- [20] I. Grosse, P. Bernaola-Galván, P. Carpena, R. Román-Roldán, J. Oliver, and H. E. Stanley, *Physical Review E* **65**, 041905 (2002).
- [21] S. N. Kadir, D. F. Goodman, and K. D. Harris, *Neural Comput* **26**, 2379 (2014).
- [22] C. Rossant, S. N. Kadir, D. F. Goodman, J. Schulman, M. L. Hunter, A. B. Saleem, A. Grosmark, M. Belluscio, G. H. Denfield, A. S. Ecker, et al., *Nat. Neurosci* **19**, 634 (2016).
- [23] W. T. Grandy and P. W. Milonni (eds.), *Physics and probability: essays in honor of Edwin T. Jaynes* (Cambridge University Press, 1993).
- [24] O. Kinouchi and M. Copelli, *Nat. Phys.* **2**, 348 (2006).
- [25] T. T. A. Carvalho, A. J. Fontenele, M. Girardi-Schappo, T. Feliciano, L. A. A. Aguiar, T. P. L. Silva, N. A. P. de Vasconcelos, P. V. Carelli, and M. Copelli, *Subsampled directed-percolation models explain scaling relations experimentally observed in the brain* (2020), 2007.13813.
- [26] R. V. Williams-García, M. Moore, J. M. Beggs, and G. Ortiz, *Phys. Rev. E* **90**, 062714 (2014), URL <https://link.aps.org/doi/10.1103/PhysRevE.90.062714>.