

CHÂTELET SURFACES AND NON-INVARIANCE OF THE BRAUER-MANIN OBSTRUCTION FOR 3-FOLDS

HAN WU

ABSTRACT. In this paper, we construct three kinds of Châtelet surfaces, which have some given arithmetic properties with respect to field extensions of number fields. We then use these constructions to study the properties of weak approximation with Brauer-Manin obstruction and the Hasse principle with Brauer-Manin obstruction for 3-folds, which are pencils of Châtelet surfaces parameterized by a curve, with respect to field extensions of number fields. We give general constructions (conditional on a conjecture of M. Stoll) to negatively answer some questions, and illustrate these constructions and some exceptions with some explicit unconditional examples.

1. INTRODUCTION

1.1. Background. Let X be a proper algebraic variety defined over a number field K . Let Ω_K be the set of all nontrivial places of K . Let $\infty_K \subset \Omega_K$ be the subset of all archimedean places, and let $S \subset \Omega_K$ be a finite subset. Let K_v be the completion of K at $v \in \Omega_K$. Let \mathbb{A}_K be the ring of adèles of K . If the set of K -rational points $X(K) \neq \emptyset$, then the set of adelic points $X(\mathbb{A}_K) \neq \emptyset$. The converse is known as the Hasse principle. We say that X is a counterexample to the Hasse principle if $X(\mathbb{A}_K) \neq \emptyset$ whereas $X(K) = \emptyset$. A well known counterexample to the Hasse principle is Selmer's cubic curve defined over \mathbb{Q} by $3w_0^3 + 4w_1^3 + 5w_2^3 = 0$ with homogeneous coordinates $(w_0 : w_1 : w_2) \in \mathbb{P}^2$. Let $pr^S: \mathbb{A}_K \rightarrow \mathbb{A}_K^S$ be the natural projection of the ring of adèles and adèles without S components, which induces a natural projection $pr^S: X(\mathbb{A}_K) \rightarrow X(\mathbb{A}_K^S)$ if $X(\mathbb{A}_K) \neq \emptyset$. For X is proper, the set of adelic points $X(\mathbb{A}_K^S)$ is equal to $\prod_{v \in \Omega_K \setminus S} X(K_v)$, and the adelic topology of $X(\mathbb{A}_K^S)$ is indeed the product topology of v -adic topologies. Viewing $X(K)$ as a subset of $X(\mathbb{A}_K)$ (respectively of $X(\mathbb{A}_K^S)$) by the diagonal embedding, we say that X satisfies weak approximation (respectively weak approximation off S) if $X(K)$ is dense in $X(\mathbb{A}_K)$ (respectively in $X(\mathbb{A}_K^S)$), cf. [Sko01, Chapter 5.1]. Cohomological obstructions have been used to explain failures of the Hasse principle and nondensity of $X(K)$ in $X(\mathbb{A}_K^S)$. Let $\text{Br}(X) = H_{\text{ét}}^2(X, \mathbb{G}_m)$ be the Brauer group of X . The Brauer-Manin pairing

$$X(\mathbb{A}_K) \times \text{Br}(X) \rightarrow \mathbb{Q}/\mathbb{Z},$$

suggested by Brauer-Manin [Man71], between $X(\mathbb{A}_K)$ and $\text{Br}(X)$, is provided by local class field theory. The left kernel of this pairing is denoted by $X(\mathbb{A}_K)^{\text{Br}}$, which is a closed subset of $X(\mathbb{A}_K)$. By the global reciprocity in class field theory, there is an exact sequence:

$$0 \rightarrow \text{Br}(K) \rightarrow \bigoplus_{v \in \Omega_K} \text{Br}(K_v) \rightarrow \mathbb{Q}/\mathbb{Z} \rightarrow 0.$$

It induces an inclusion: $X(K) \subset pr^S(X(\mathbb{A}_K)^{\text{Br}})$. We say that the failure of the Hasse principle of X is explained by the Brauer-Manin obstruction if $X(\mathbb{A}_K) \neq \emptyset$ and $X(\mathbb{A}_K)^{\text{Br}} = \emptyset$. For the failure of the Hasse principle of a smooth, projective, and geometrically connected curve over K , if the Tate-Shafarevich group and the rational points set of its Jacobian are finite, then this failure can be explained by the Brauer-Manin obstruction, cf. [Sch99]. A counterexample to the Hasse principle such that its failure of the Hasse principle is

2020 *Mathematics Subject Classification.* 11G35, 14G12, 14G25, 14G05.

Key words and phrases. rational points, Hasse principle, weak approximation, Brauer-Manin obstruction, Châtelet surfaces, Châtelet surface bundles over curves.

not explained by the Brauer-Manin obstruction, was constructed firstly by Skorobogatov [Sko99], subsequently by Poonen [Poo10], Harpaz and Skorobogatov [HS14], Colliot-Thélène, Pál and Skorobogatov [CTPS16] and so on. We say that X satisfies weak approximation with Brauer-Manin obstruction (respectively with Brauer-Manin obstruction off S) if $X(K)$ is dense in $X(\mathbb{A}_K)^{\text{Br}}$ (respectively in $pr^S(X(\mathbb{A}_K)^{\text{Br}})$). For an elliptic curve defined over \mathbb{Q} , if its analytic rank equals one, then this elliptic curve satisfies weak approximation with Brauer-Manin obstruction, cf. [Wan96]. For an abelian variety defined over K , if its Tate-Shafarevich group is finite, then this abelian variety satisfies weak approximation with Brauer-Manin obstruction off ∞_K , cf. [Sko01, Proposition 6.2.4]. For any smooth, proper and rationally connected variety X defined over an number field, it is conjectured by J.-L. Colliot-Thélène that X satisfies weak approximation with Brauer-Manin obstruction. The Colliot-Thélène's conjecture holds for Châtelet surfaces, cf. [CTSSD87a, CTSSD87b]. For any smooth, projective, and geometrically connected curve C defined over an number field K , it is conjectured by Stoll [Sto07] that C satisfies weak approximation with Brauer-Manin obstruction off ∞_K : see Conjecture 4.0.1 for more details. Before the Stoll's conjecture, if this curve C is a counterexample to the Hasse principle, Skorobogatov [Sko01, Chapter 6.3] asked a weaker open question: is the failure of the Hasse principle of C explained by the Brauer-Manin obstruction?

1.2. Questions. Given a nontrivial extension of number fields L/K , and a finite subset $S \subset \Omega_K$, let $S_L \subset \Omega_L$ be the subset of all places above S . Let X be a smooth, projective, and geometrically connected variety defined over K . Let $X_L = X \times_{\text{Spec } K} \text{Spec } L$ be the base change of X by L . In this paper, we consider the following questions.

Question 1.2.1. If the variety X has a K -rational point, and satisfies weak approximation with Brauer-Manin obstruction off S , must X_L also satisfy weak approximation with Brauer-Manin obstruction off S_L ?

Question 1.2.2. Assume that the varieties X and X_L are counterexamples to the Hasse principle. If the failure of the Hasse principle of X is explained by the Brauer-Manin obstruction, must the failure of the Hasse principle of X_L also be explained by the Brauer-Manin obstruction?

1.3. Main results for Châtelet surfaces. A Châtelet surface defined over \mathbb{Q} , which is a counterexample to the Hasse principle, was constructed by Iskovskikh [Isk71]. Poonen [Poo09] generalized it to any given number field. For any number field K , Liang [Lia18] constructed a Châtelet surface defined over K , which has a K -rational point and does not satisfy weak approximation off ∞_K . By using weak approximation and strong approximation off all 2-adic places for \mathbb{A}^1 (cf. Lemma 2.0.1) to choose elements in K , and using Čebotarev's density theorem (cf. Theorem 2.0.2) to add some splitting conditions, we construct three kinds of Châtelet surfaces to generalize their arguments.

Proposition 1.3.1. For any extension of number fields L/K , and any finite subset $S \subset \Omega_K \setminus \{\text{all complex and 2-adic places}\}$ splitting completely in L , there exist Châtelet surfaces V_1, V_2, V_3 defined over K , which have the following properties.

- The subset $V_1(\mathbb{A}_K^S) \subset V_1(\mathbb{A}_L^{S_L})$ is nonempty, but $V_1(K_v) = V_1(L_{v'}) = \emptyset$ for all $v \in S$ and all $v' \in S_L$, cf. Proposition 3.1.1.
- The Brauer group $\text{Br}(V_2)/\text{Br}(K) \cong \text{Br}(V_{2L})/\text{Br}(L) \cong \mathbb{Z}/2\mathbb{Z}$, is generated by an element $A \in \text{Br}(V_2)$. The subset $V_2(K) \subset V_2(L)$ is nonempty. For any $v \in S$, there exist P_v and Q_v in $V_2(K_v)$ such that the local invariants $\text{inv}_v(A(P_v)) = 0$ and $\text{inv}_v(A(Q_v)) = 1/2$. For any other $v \notin S$, and any $P_v \in V_2(K_v)$, the local invariant $\text{inv}_v(A(P_v)) = 0$. For any $v' \in S_L$, there exist $P_{v'}$ and $Q_{v'}$ in $V_2(L_{v'})$ such that the local invariants $\text{inv}_{v'}(A(P_{v'})) = 0$ and $\text{inv}_{v'}(A(Q_{v'})) = 1/2$. For any other $v' \notin S_L$, and any $P_{v'} \in V_2(L_{v'})$, the local invariant $\text{inv}_{v'}(A(P_{v'})) = 0$, cf. Proposition 3.2.1.
- The Brauer group $\text{Br}(V_3)/\text{Br}(K) \cong \text{Br}(V_{3L})/\text{Br}(L) \cong \mathbb{Z}/2\mathbb{Z}$, is generated by an element $A \in \text{Br}(V_3)$. The subset $V_3(\mathbb{A}_K) \subset V_3(\mathbb{A}_L)$ is nonempty. For any $v \in \Omega_K$, and any $P_v \in V_3(K_v)$, the local invariant $\text{inv}_v(A(P_v)) = 0$ if

$v \notin S$; the local invariant $\text{inv}_v(A(P_v)) = 1/2$ if $v \in S$.
 For any $v' \in \Omega_L$, and any $P_{v'} \in V_3(L_{v'})$, the local invariant $\text{inv}_{v'}(A(P_{v'})) = 0$ if $v' \notin S_L$; the local invariant $\text{inv}_{v'}(A(P_{v'})) = 1/2$ if $v' \in S_L$, cf. Proposition 3.3.1.

Combining our construction method with the global reciprocity law, we have the following results for Châtelet surfaces.

Corollary 1.3.2 (Corollary 3.2.4). For any extension of number fields L/K , and any finite nonempty subset $S \subset \Omega_K \setminus \{\text{all complex and 2-adic places}\}$ splitting completely in L , there exists a Châtelet surface V defined over K such that $V(K) \neq \emptyset$. For any subfield $L' \subset L$ over K , the Brauer group $\text{Br}(V)/\text{Br}(K) \cong \text{Br}(V_{L'})/\text{Br}(L') \cong \mathbb{Z}/2\mathbb{Z}$. And the surface $V_{L'}$ has the following properties.

- For any finite subset $T' \subset \Omega_{L'}$ such that $T' \cap S_{L'} \neq \emptyset$, the surface $V_{L'}$ satisfies weak approximation off T' .
- For any finite subset $T' \subset \Omega_{L'}$ such that $T' \cap S_{L'} = \emptyset$, the surface $V_{L'}$ does not satisfy weak approximation off T' . In particular, the surface $V_{L'}$ does not satisfy weak approximation.

Corollary 1.3.3 (Corollary 3.3.3). For any extension of number fields L/K , there exists a Châtelet surface V defined over K such that $V(\mathbb{A}_K) \neq \emptyset$. For any subfield $L' \subset L$ over K , the Brauer group $\text{Br}(V)/\text{Br}(K) \cong \text{Br}(V_{L'})/\text{Br}(L') \cong \mathbb{Z}/2\mathbb{Z}$. And the surface $V_{L'}$ has the following properties.

- If the degree $[L' : K]$ is odd, then the surface $V_{L'}$ is a counterexample to the Hasse principle. In particular, the surface V is a counterexample to the Hasse principle.
- If the degree $[L' : K]$ is even, then the surface $V_{L'}$ satisfies weak approximation. In particular, in this case, the set $V(L') \neq \emptyset$.

1.4. Main results for Châtelet surface bundles over curves. We will apply our results for Châtelet surfaces to construct Châtelet surface bundles over curves to give negative answers to Questions 1.2.

1.4.1. *A negative answer to Question 1.2.1.* For any quadratic extension of number fields L/K , and assuming the Stoll's conjecture, a Châtelet surface bundle over a curve was constructed by Liang[Lia18] to give a negative answer to Question 1.2.1. Also an unconditional example with explicit equations was given for $L = \mathbb{Q}(\sqrt{5})$ and $K = \mathbb{Q}$ in loc. cit. His method only works for quadratic extensions. In this paper, we generalize it to any nontrivial extension of number fields.

For any nontrivial extension of number fields L/K , assuming the Stoll's conjecture, we have the following theorem to give a negative answer to Question 1.2.1.

Theorem 1.4.1.1 (Theorem 6.2.1). For any nontrivial extension of number fields L/K , and any finite subset $T \subset \Omega_L$, assuming the Stoll's conjecture, there exists a Châtelet surface bundle over a curve: $X \rightarrow C$ defined over K such that

- X has a K -rational point, and satisfies weak approximation with Brauer-Manin obstruction off ∞_K ,
- X_L does not satisfy weak approximation with Brauer-Manin obstruction off T .

For $K = \mathbb{Q}$ and $L = \mathbb{Q}(\sqrt{3})$, based on the method given in Theorem 6.2.1, we give an explicit unconditional example in Subsection 7.1.

1.4.2. *Negative answers to Question 1.2.2.* To the best knowledge of the author, Question 1.2.2 has not yet been seriously discussed in the literature.

For any number field K , and any nontrivial field extension L of odd degree over K , assuming the Stoll's conjecture, we have the following theorem to give a negative answer to Question 1.2.2.

Theorem 1.4.2.1 (Theorem 6.3.1). For any number field K , and any nontrivial field extension L of odd degree over K , assuming the Stoll's conjecture, there exists a Châtelet surface bundle over a curve: $X \rightarrow C$ defined over K such that

- X is a counterexample to the Hasse principle, and its failure of the Hasse principle is explained by the Brauer-Manin obstruction,
- X_L is a counterexample to the Hasse principle, but its failure of the Hasse principle cannot be explained by the Brauer-Manin obstruction.

Let ζ_7 be a primitive 7-th root of unity. For $K = \mathbb{Q}$ and $L = \mathbb{Q}(\zeta_7 + \zeta_7^{-1})$, based on the method given in Theorem 6.3.1, we give an explicit unconditional example in Subsection 7.2. The 3-fold X is a smooth compactification of the following 3-dimensional affine subvariety given by equations

$$\begin{cases} y^2 - 377z^2 = 14(x^4 - 89726)y'^2 + (x^2 - 878755181)(5x^2 - 4393775906) \\ y'^2 = x'^3 - 343x' - 2401 \end{cases}$$

in $(x, y, z, x', y') \in \mathbb{A}^5$.

For any number field K having a real place, and any nontrivial field extension L/K having a real place, assuming the Stoll's conjecture, we have the following theorem to give a negative answer to Question 1.2.2.

Theorem 1.4.2.2 (Theorem 6.3.2). For any number field K having a real place, and any nontrivial field extension L/K having a real place, assuming the Stoll's conjecture, there exists a Châtelet surface bundle over a curve: $X \rightarrow C$ defined over K such that

- X is a counterexample to the Hasse principle, and its failure of the Hasse principle is explained by the Brauer-Manin obstruction,
- X_L is a counterexample to the Hasse principle, but its failure of the Hasse principle cannot be explained by the Brauer-Manin obstruction.

For $K = \mathbb{Q}$ and $L = \mathbb{Q}(\sqrt{3})$, based on the method given in Theorem 6.3.2, we give an explicit unconditional example in Subsection 7.3. The 3-fold X is a smooth compactification of the following 3-dimensional affine subvariety given by equations

$$\begin{cases} y^2 + 23z^2 = 5(x^4 + 805)(x' - 4)^2 - 5(x^4 + 115) \\ y'^2 = x'^3 - 16 \end{cases}$$

in $(x, y, z, x', y') \in \mathbb{A}^5$.

Exceptions 1.4.2.3 (Subsection 7.4). For Question 1.2.2, besides cases of Theorem 6.3.1 and Theorem 6.3.2, there are some exceptions. When the degree $[L : K]$ is even and L has no real place, we can give some unconditional examples, case by case, to give negative answers to Question 1.2.2, although we do not have a uniform way to construct them. In Subsection 7.4, we give an example to explain how it works for the case that $K = \mathbb{Q}$ and $L = \mathbb{Q}(i)$. The 3-fold X , is a smooth compactification of the following 3-dimensional affine subvariety given by equations

$$\begin{cases} y^2 + 15z^2 = (x^4 - 10x^2 + 15)(y'^2 + 32)/8 - (5x^4 - 39x^2 + 75)y'/2 \\ y'^2 = x'^3 - 16 \end{cases}$$

in $(x, y, z, x', y') \in \mathbb{A}^5$.

1.4.3. Main ideas behind our constructions in the proof of theorems. Given a nontrivial extension of number fields L/K , we start with a curve C such that $C(K)$ and $C(L)$ are finite, nonempty and $C(K) \neq C(L)$. Using our results for Châtelet surfaces and a construction method from Poonen [Poo10], we construct a Châtelet surface bundle over this curve: $\beta: X \rightarrow C$ such that the fiber of each $C(K)$ point is isomorphic to V_∞ , and that the fiber of each $C(L) \setminus C(K)$ point is isomorphic to V_0 . Assuming the Stoll's conjecture, by a main proposition from [Poo10, Proposition 5.4] and the functoriality of Brauer-Manin pairing, the Brauer-Manin sets of X , roughly speaking, is the union of adelic points sets

of rational fibers. Using some fibration arguments, the arithmetic properties of V_∞ and V_0 will determine the arithmetic properties of X . For Theorem 6.2.1 and 6.3.1, these ideas will be enough. For Theorems 6.3.2, we choose the curve C with an additional connected condition at one real place, and make full use of this real place information.

2. NOTATION AND PRELIMINARIES

Given a number field K , let \mathcal{O}_K be the ring of its integers, and let Ω_K be the set of all its nontrivial places. Let $\infty_K \subset \Omega_K$ be the subset of all archimedean places, and let $2_K \subset \Omega_K$ be the subset of all 2-adic places. Let $\infty_K^r \subset \infty_K$ be the subset of all real places, and let $\infty_K^c \subset \infty_K$ be the subset of all complex places. Let $\Omega_K^f = \Omega_K \setminus \infty_K$ be the set of all finite places of K . Let K_v be the completion of K at $v \in \Omega_K$. For $v \in \infty_K$, let $\tau_v: K \hookrightarrow K_v$ be the embedding of K into its completion. For $v \in \Omega_K^f$, let \mathcal{O}_{K_v} be its valuation ring, and let \mathbb{F}_v be its residue field. Let $S \subset \Omega_K$ be a finite subset, and let $\mathcal{O}_S = \bigcap_{v \in \Omega_K^f \setminus S} (K \cap \mathcal{O}_{K_v})$ be the ring of S -integers. Let $\mathbb{A}_K, \mathbb{A}_K^S$ be the ring of adèles and adèles without S components of K . A strong approximation theorem [CF67, Chapter II §15] states that K is dense in \mathbb{A}_K^S for any nonempty S . In this paper, we only use the following special case:

Lemma 2.0.1. Let K a number field. The set K is dense in \mathbb{A}_K^{2K} .

In this paper, we always assume that a field L is a finite extension of K . Let $S_L \subset \Omega_L$ be the subset of all places above S .

It is not difficult to generalize [Neu99, Theorem 13.4] to the following version of Čebotarev's density theorem.

Theorem 2.0.2 (Čebotarev). Let L/K be an extension of number fields. Then the set of places of K splitting completely in L , has positive density.

2.1. Hilbert symbol. We use Hilbert symbol $(a, b)_v \in \{\pm 1\}$, for $a, b \in K_v^\times$ and $v \in \Omega_K$. By definition, $(a, b)_v = 1$ if and only if $x_0^2 - ax_1^2 - bx_2^2 = 0$ has a K_v -solution in \mathbb{P}^2 with homogeneous coordinates $(x_0 : x_1 : x_2)$, which equivalently means that the curve defined over K_v by the equation $x_0^2 - ax_1^2 - bx_2^2 = 0$ in \mathbb{P}^2 , is isomorphic to \mathbb{P}^1 . The Hilbert symbol gives a symmetric bilinear form on $K_v^\times / K_v^{\times 2}$ with value in $\mathbb{Z}/2\mathbb{Z}$, cf. [Ser79, Chapter XIV, Proposition 7]. And this bilinear form is nondegenerate, cf. [Ser79, Chapter XIV, Corollary 7].

2.2. Preparation lemmas. We state the following lemmas for later use.

Lemma 2.2.1. Let K be a number field, and let v be an odd place of K . Let $a, b \in K_v^\times$ such that $v(a), v(b)$ are even. Then $(a, b)_v = 1$.

Proof. Choose a prime element $\pi_v \in K_v$. Let $a_1 = a\pi_v^{-v(a)}$ and $b_1 = b\pi_v^{-v(b)}$. Since the valuations $v(a)$ and $v(b)$ are even, the elements $\pi_v^{-v(a)}$ and $\pi_v^{-v(b)}$ are in $K_v^{\times 2}$. So $(a, b)_v = (a_1, b_1)_v$ and $a_1, b_1 \in \mathcal{O}_{K_v}^\times$. By Chevalley-Waring theorem (cf. [Ser73, Chapter I §2, Corollary 2]), the equation $x_0^2 - \bar{a}_1 x_1^2 - \bar{b}_1 x_2^2 = 0$ has a nontrivial solution in \mathbb{F}_v . For v is odd, by Hensel's lemma, this solution can be lifted to a nontrivial solution in \mathcal{O}_{K_v} . Hence $(a, b)_v = (a_1, b_1)_v = 1$. \square

Lemma 2.2.2. Let K be a number field, and let v be an odd place of K . Let $a, b, c \in K_v^\times$ such that $v(b) < v(c)$. Then $(a, b+c)_v = (a, b)_v$.

Proof. For $v(b) < v(c)$, we have $v(b^{-1}c) > 0$. By Hensel's lemma, we have $1 + b^{-1}c \in K_v^{\times 2}$. So $(a, b+c)_v = (a, b(1 + b^{-1}c))_v = (a, b)_v$. \square

Lemma 2.2.3. Let K be a number field, and let $v \in \Omega_K$. Then $K_v^{\times 2}$ is an open subgroup of K_v^\times . If $v \in \Omega_K^f$, then $\mathcal{O}_{K_v}^\times$ is also an open subgroup of K_v^\times . So, they are nonempty open subset of K_v .

Proof. It is obvious for $v \in \infty_K$. Consider $v \in \Omega_K^f$. Let p be the prime number such that $v|p$ in K . Then by Hensel's lemma, the group $K_v^{\times 2} \cap \mathcal{O}_{K_v}^\times$ contains the set $1 + p^3 \mathcal{O}_{K_v}$, which is an open subgroup of K_v^\times . Hence $K_v^{\times 2}$ and $\mathcal{O}_{K_v}^\times$ are open subgroups of K_v^\times . \square

Lemma 2.2.4. Let K be a number field, and let $v \in \Omega_K^f$. For any $n \in \mathbb{Z}$, the set $\{x \in K_v | v(x) = n\}$ is a nonempty open subset of K_v .

Proof. By Lemma 2.2.3, the set $\mathcal{O}_{K_v}^\times$ is an open subgroup of K_v^\times . Choose a prime element $\pi_v \in K_v$. Then the set $\{x \in K_v^\times | v(x) = n\} = \pi_v^n \mathcal{O}_{K_v}^\times$, so it is a nonempty open subset of K_v . \square

Lemma 2.2.5. Let K be a number field, and let $v \in \Omega_K^f$. For any $a \in K_v^\times$, the sets $\{x \in K_v^\times | (a, x)_v = 1\}$, $\{x \in K_v^\times | (a, x)_v = 1\} \cap \mathcal{O}_{K_v}$ and $\{x \in \mathcal{O}_{K_v}^\times | (a, x)_v = 1\}$ are nonempty open subsets of K_v .

Proof. For the unit 1 belongs to these sets, they are nonempty. By Lemma 2.2.3, the sets $K_v^{\times 2}$ and $\mathcal{O}_{K_v}^\times$ are nonempty open subsets of K_v . The set $\{x \in K_v^\times | (a, x)_v = 1\}$ is a union of cosets of $K_v^{\times 2}$ in the group K_v^\times . So the sets are open in K_v . \square

Lemma 2.2.6. Let K be a number field, and let $v \in \Omega_K^f$. For any $a \in K_v^\times$, the sets $\{x \in K_v^\times | (a, x)_v = -1\}$ and $\{x \in K_v^\times | (a, x)_v = -1\} \cap \mathcal{O}_{K_v}$ are open subsets of K_v . Furthermore, if $a \notin K_v^{\times 2}$, then they are nonempty.

Proof. If the set $\{x \in K_v^\times | (a, x)_v = -1\} \neq \emptyset$, then it is a union of cosets of $K_v^{\times 2}$ in the group K_v^\times . By Lemma 2.2.3, it is an open subset of K_v . For \mathcal{O}_{K_v} is open in K_v , the sets are open subsets of K_v^\times . Nonemptiness is from the nondegeneracy of the bilinear form given by the Hilbert symbol, and from multiplying a square element in \mathcal{O}_{K_v} to denominate an element in K_v^\times . \square

Lemma 2.2.7. Let K be a number field, and let $v \in \Omega_K^f$. For any $a \in K_v^\times$ with $v(a)$ odd, the set $\{x \in \mathcal{O}_{K_v}^\times | (a, x)_v = -1\}$ is a nonempty open subset of K_v .

Proof. By Lemmas 2.2.3 and 2.2.6, the set is open in K_v . We need to show that it is nonempty. For $a \notin K_v^2$, by the nondegeneracy of the bilinear form given by the Hilbert symbol, there exists an element $b \in K_v^\times$ such that $(a, b)_v = -1$. If $v(b)$ is odd, let $b' = -ab$. Then $(a, b')_v = (a, -ab)_v = (a, -a)_v (a, b)_v = -1$. Replacing b by b' if necessary, we can assume that $v(b)$ is even. Choose a prime element $\pi_v \in K_v$. Then $\pi_v^{-v(b)} \in K_v^{\times 2}$, so the element $b\pi_v^{-v(b)}$ is in this set. \square

Lemma 2.2.8. Let L/K be an extension of number fields, and let $v \in \Omega_K \setminus \infty_K^c$. We assume that $v' \in \Omega_L$ splits over v , i.e. $L_{v'} = K_v$. Given an element $a \in K$, if v is a finite place, we assume that $v(a)$ is odd; if v is an archimedean place, we assume $\tau_v(a) < 0$. Then $a \notin L^{\times 2}$.

Proof. The condition that $v(a)$ is odd for the finite place v , or that $\tau_v(a) < 0$ for the archimedean place v , implies that $a \notin K_v^{\times 2}$. For $L_{v'} = K_v$, we have $a \notin L_{v'}^{\times 2}$, so $a \notin L^{\times 2}$. \square

3. MAIN RESULTS FOR CHÂTELET SURFACES

In this section, we will construct three kinds of Châtelet surfaces. Each kind in each following subsection has the arithmetic properties mentioned in Subsection 1.3.

Let K be a number field. Châtelet surfaces are smooth projective models of conic bundle surfaces defined by the equation

$$(1) \quad y^2 - az^2 = P(x)$$

in $K[x, y, z]$ such that $a \in K^\times$, and that $P(x)$ is a separable degree-4 polynomial in $K[x]$. Given an equation (1), let V^0 be the affine surface in \mathbb{A}_K^3 defined by this equation. The natural smooth compactification V of V^0 given in [Sko01, Section 7.1] is called the Châtelet surface given by this equation, cf. [Poo09, Section 5].

Remark 3.0.1. For any local field K_v , if $a \in K_v^{\times 2}$, then V is birational equivalent to \mathbb{P}^2 over K_v . By the implicit function theorem, there exists a K_v -point on V .

Remark 3.0.2. For any local field K_v , by smoothness of V , the implicit function theorem implies that the nonemptiness of $V^0(K_v)$ is equivalent to the nonemptiness of $V(K_v)$, and that $V^0(K_v)$ is open dense in $V(K_v)$. Given an element $A \in \text{Br}(V)$, the evaluation of A on $V(K_v)$ is locally constant. By the properness of V , the space $V(K_v)$ is compact. So the set of all possible values of the evaluation of A on $V(K_v)$ is finite. Indeed, by [Sko01, Proposition 7.1.2], there only exist two possible values. It is determined by the evaluation of A on $V^0(K_v)$. In particular, if the evaluation of A on $V^0(K_v)$ is constant, then it is constant on $V(K_v)$.

Remark 3.0.3. In [CTPS16, Proposition 6.1], it is shown that whether the Brauer-Manin set of a smooth, projective, and geometrically connected variety is empty, is determined by its birational equivalence class. Here birational equivalence means birational equivalence between smooth, projective, and geometrically connected varieties. It is proved in the papers [CTSSD87a, Theorem B; CTSSD87b] (also explained in the book [Sko01, Theorem 7.2.1]), that the Brauer-Manin obstruction to the Hasse principle and weak approximation is the only one for Châtelet surfaces. Hence, all smooth projective models of a given equation (1) are the same as to the discussion of the Hasse principle, weak approximation, the failure of the Hasse principle explained by the Brauer-Manin obstruction, and weak approximation with Brauer-Manin obstruction.

Remark 3.0.4. If the polynomial $P(x)$ has a factor $x^2 - a$, i.e. there exists a degree-2 polynomial $f(x)$ such that $P(x) = f(x)(x^2 - a)$, then $Y = \frac{xy+az}{x^2-a}$ and $Z = \frac{y+xz}{x^2-a}$ give a birational equivalence between V and a quadratic surface given by $Y^2 - aZ^2 = f(x)$ with affine coordinates (x, Y, Z) . By the Hasse-Minkowski theorem and Remark 3.0.3, the surface V satisfies weak approximation.

In this section, we always use the following way to choose an element for the parameter a in the equation (1).

3.0.1. *Choosing an element for the parameter a in the equation (1).* Given an extension of number fields L/K , and a finite subset $S \subset \Omega_K \setminus (\infty_K^c \cup 2_K)$, we will choose an element $a \in \mathcal{O}_K \setminus K^2$ with respect to these L/K and S in the following way.

If $S = \emptyset$, by Theorem 2.0.2, we can take a place $v_0 \in \Omega_K^f \setminus 2_K$ splitting completely in L . Then replace S by $\{v_0\}$ to continue the following step.

Now, suppose that $S \neq \emptyset$. For $v \in \Omega_K$, by Lemma 2.2.3, the set $K_v^{\times 2}$ is a nonempty open subset of K_v . For $v \in \Omega_K^f$, by Lemma 2.2.4, the set $\{a \in K_v | v(a) \text{ is odd}\}$ is a nonempty open subset of K_v . Using weak approximation for the affine line \mathbb{A}^1 , we can choose an element $a \in K^\times$ satisfying the following conditions:

- $\tau_v(a) < 0$ for all $v \in S \cap \infty_K$,
- $a \in K_v^{\times 2}$ for all $v \in 2_K$,
- $v(a)$ is odd for all $v \in S \setminus \infty_K$.

These conditions do not change by multiplying an element in $K^{\times 2}$, so we can assume $a \in \mathcal{O}_K$. The conditions that $v(a)$ is odd for all $v \in S \setminus \infty_K$, and that $\tau_v(a) < 0$ for all $v \in S \cap \infty_K$, imply $a \in \mathcal{O}_K \setminus K_v^2$ for all $v \in S$. So $a \in \mathcal{O}_K \setminus K^2$.

Remark 3.0.5. Let $S' = \{v \in \infty_K^r | \tau_v(a) < 0\} \cup \{v \in \Omega_K^f \setminus 2_K | v(a) \text{ is odd}\}$, then S' is a finite set. By the conditions that $\tau_v(a) < 0$ for all $v \in S \cap \infty_K$, and that $v(a)$ is odd for all $v \in S \setminus \infty_K$, we have $S' \supset S$. Then $S' \neq \emptyset$.

Remark 3.0.6. If there exists one place in S splitting completely in L or $S = \emptyset$, then by the choice of a above and Lemma 2.2.8, the element $a \in \mathcal{O}_K \setminus L^2$.

Remark 3.0.7. For the choice of a , we can enlarge S in $\Omega_K \setminus (\infty_K^c \cup 2_K)$ if necessary.

3.1. Châtelet surfaces without K_v point for any $v \in S$. In this subsection, we will construct a Châtelet surface of the first kind mentioned in Subsection 1.3.

3.1.1. *Choice of parameters for the equation (1).* Given an extension of number fields L/K , and a finite subset $S \subset \Omega_K \setminus (\infty_K^c \cup 2_K)$, we choose an element $a \in \mathcal{O}_K \setminus K^2$ as in Subsubsection 3.0.1.

If $S = \emptyset$, then let $P(x) = 1 - x^4$. Then the Châtelet surface V_1 given by $y^2 - az^2 = 1 - x^4$, has a rational point $(x, y, z) = (0, 1, 0)$.

Now, suppose that $S \neq \emptyset$. We will choose an element $b \in K^\times$ with respect to the chosen a in the following way.

Let $S' = \{v \in \infty_K^r \mid \tau_v(a) < 0\} \cup \{v \in \Omega_K^f \setminus 2_K \mid v(a) \text{ is odd}\}$ be as in Remark 3.0.5, then $S' \supset S$ is a finite set. If $v \in S' \setminus \infty_K$, then $v(a)$ is odd, which implies $a \notin K_v^{\times 2}$. Then by Lemma 2.2.6, the set $\{b \in K_v^\times \mid (a, b)_v = -1\}$ is a nonempty open subset of K_v . If $v \in S' \setminus (S \cup \infty_K)$, then by Lemma 2.2.5, the set $\{b \in K_v^\times \mid (a, b)_v = 1\}$ is a nonempty open subset of K_v . Using weak approximation for affine line \mathbb{A}^1 , we can choose an element $b \in K^\times$ satisfying the following conditions:

- $\tau_v(b) < 0$ for all $v \in S \cap \infty_K$,
- $\tau_v(b) > 0$ for all $v \in (S' \setminus S) \cap \infty_K$,
- $(a, b)_v = -1$ for all $v \in S' \setminus \infty_K$,
- $(a, b)_v = 1$ for all $v \in S' \setminus (S \cup \infty_K)$.

We will choose an element $c \in K^\times$ with respect to the chosen a, b in the following way.

Let $S'' = \{v \in \Omega_K^f \setminus 2_K \mid v(b) \text{ is odd}\}$, then S'' is a finite set. The same argument as in the previous paragraph, we can choose an element $c \in K^\times$ satisfying the following conditions:

- $c \in K_v^{\times 2}$ for all $v \in S$,
- $v(c)$ is odd for all $v \in S'' \setminus S'$.

These conditions do not change by multiplying an element in $K^{\times 2}$, so we can assume $b, c \in \mathcal{O}_K$.

Let $P(x) = b(x^4 - ac)$, and let V_1 be the Châtelet surface given by $y^2 - az^2 = b(x^4 - ac)$.

Proposition 3.1.1. For any extension of number fields L/K , and any finite subset $S \subset \Omega_K \setminus (\infty_K^c \cup 2_K)$ splitting completely in L , there exists a Châtelet surface V_1 defined over K such that $V_1(\mathbb{A}_K^S) \subset V_1(\mathbb{A}_L^{S_L})$ is nonempty, but that $V_1(K_v) = V_1(L_{v'}) = \emptyset$ for all $v \in S$ and all $v' \in S_L$.

Proof. For the extension L/K , and the finite set S , we will check that the Châtelet surface V_1 chosen as in Subsubsection 3.1.1, has the properties.

For $S = \emptyset$, it is clear.

Now, suppose that $S \neq \emptyset$. We will check these properties by local computation.

Suppose that $v \in (\infty_K \setminus S') \cup 2_K$. By the choice of a , we have $a \in K_v^{\times 2}$. By Remark 3.0.1, the surface V_1 admits a K_v -point.

Suppose that $v \in (S' \setminus S) \cap \infty_K$. Take $x_0 \in K$ such that $\tau_v(x_0^4 - ac) > 0$. By the choice of b , we have $\tau_v(b(x_0^4 - ac)) > 0$. So $(a, b(x_0^4 - ac))_v = 1$, which implies that V_1^0 admits a K_v -point with $x = x_0$.

Suppose that $v \in S' \setminus (S \cup \infty_K)$, then by the choice of b , we have $(a, b)_v = 1$. Take $x_0 \in K_v$ such that the valuation $v(x_0) < 0$, then by Lemma 2.2.2, we have $(a, x_0^4 - ac)_v = (a, x_0^4)_v = 1$. So $(a, b(x_0^4 - ac))_v = (a, b)_v = 1$, which implies that V_1^0 admits a K_v -point with $x = x_0$. Suppose that $v \in S'' \setminus S'$. By the choice of a, b, c , the valuations $v(a)$ and $v(-abc)$ are even. By Lemma 2.2.1, we have $(a, -abc)_v = 1$, which implies that V_1^0 admits a K_v -point with $x = 0$.

Suppose that $v \in \Omega_K^f \setminus (S' \cup S'' \cup 2_K)$. Take $x_0 \in K_v$ such that the valuation $v(x_0) < 0$, then by Lemma 2.2.2, we have $(a, x_0^4 - ac)_v = (a, x_0^4)_v = 1$. For $v(a)$ and $v(b)$ are even, by Lemma 2.2.1, we have $(a, b)_v = 1$. So $(a, b(x_0^4 - ac))_v = 1$, which implies that V_1^0 admits a K_v -point with $x = x_0$.

So, the subset $V_1(\mathbb{A}_K^S) \subset V_1(\mathbb{A}_L^{S_L})$ is nonempty.

Suppose that $v \in S \cap \infty_K$. Then by the choice of a, b, c , we have $\tau_v(a), \tau_v(b)$ and $\tau_v(ac)$ are negative. So $(a, b(x^4 - ac))_v = -1$ for all $x \in K_v$, which implies that V_1^0 has no K_v -point. By Remark 3.0.2, we have $V_1(K_v) = \emptyset$.

Suppose that $v \in S \setminus \infty_K$. Then by the choice of b , we have $(a, b)_v = -1$. Let $x \in K_v$. If $4v(x) < v(ac)$, then by Lemma 2.2.2, we have $(a, x^4 - ac)_v = (a, x^4)_v = 1$. If $4v(x) > v(ac)$, then by Lemma 2.2.2, we have $(a, x^4 - ac)_v = (a, -ac)_v = (a, c)_v = 1$ (by the choice of c , the last equality holds). For $c \in K_v^{\times 2}$, the valuation $v(ac)$ is odd. So, the equality $4v(x) = v(ac)$ cannot happen. In each case, we have $(a, b(x^4 - ac))_v = (a, b)_v(a, x^4 - ac)_v = -1$, which implies that V_1^0 has no K_v -point. By Remark 3.0.2, we have $V_1(K_v) = \emptyset$.

So, the set $V_1(K_v) = \emptyset$ for all $v \in S$.

Take a place $v'_0 \in S_L$. Let $v_0 \in S$ be the restriction of v'_0 on K . By the assumption that v_0 splits completely in L , we have $K_{v_0} = L_{v'_0}$. Hence $V_1(K_{v_0}) = V_1(L_{v'_0})$.

So, the set $V_1(L_{v'}) = \emptyset$ for all $v' \in S_L$. \square

Remark 3.1.2. By the choice of elements a, c in Subsubsection 3.1.1, if $v \in S \setminus \infty_K$, then by comparing the valuation, the polynomial $P(x)$ is irreducible over K_v . In this case, since the choice of c is not unique, we choose another one to get a new polynomial $P'(X)$, so the polynomials $P(x)$ and $P'(x)$ are coprime in K . For $v \in S \cap \infty_K$, the polynomial $P(x)$ is a product of two irreducible factors over K_v .

Using the construction method in Subsubsection 3.1.1, we have the following examples, which are special cases of Proposition 3.1.1. They will be used for further discussion.

Example 3.1.3. Let $K = \mathbb{Q}$, and let ζ_7 be a primitive 7-th root of unity. Let $\alpha = \zeta_7 + \zeta_7^{-1}$ with the minimal polynomial $x^3 + x^2 - 2x - 1$. Let $L = \mathbb{Q}(\alpha)$. Then the degree $[L : K] = 3$. Let $S = \{29\}$. For $29 \equiv 1 \pmod{7}$, the place 29 splits completely in L , indeed in $\mathbb{Q}(\zeta_7)$. Using the construction method in Subsubsection 3.1.1, we choose data: $S = \{29\}$, $S' = \{13, 29\}$, $S'' = \{7\}$, $a = 377$, $b = 14$, $c = 238$ and $P(x) = 14(x^4 - 89726)$. Then the Châtelet surface given by $y^2 - 377z^2 = P(x)$, has the properties of Proposition 3.1.1.

Example 3.1.4. Let $K = \mathbb{Q}$, and let $L = \mathbb{Q}(\sqrt{3})$. Using the construction method in Subsubsection 3.1.1, we choose data: $S = \{\infty_K\}$, $S' = \infty_K \cup \{23\}$, $S'' = \{5\}$, $a = -23$, $b = -5$, $c = 5$ and $P(x) = -5(x^4 + 115)$. Then the Châtelet surface given by $y^2 + 23z^2 = P(x)$, has the properties of Proposition 3.1.1.

Example 3.1.5. Let $K = \mathbb{Q}$, and let $L = \mathbb{Q}(\sqrt{3})$. Then the place 23 splits completely in L . Using the construction method in Subsubsection 3.1.1, we choose data: $S = \{23\}$, $S' = \infty_K \cup \{23\}$, $S'' = \{5\}$, $a = -23$, $b = 5$, $c = 35$ and $P(x) = 5(x^4 + 805)$. Then the Châtelet surface given by $y^2 + 23z^2 = P(x)$, has the properties of Proposition 3.1.1.

3.2. Châtelet surfaces with rational points and not satisfying weak approximation. Given an number field K , Liang [Lia18, Proposition 3.4] constructed a Châtelet surface over K , which has a K -rational point and does not satisfy weak approximation off ∞_K . Using the same method as in [Poo09, Section 5] to choose the parameters for the equation (1), he constructed a Châtelet surface, and there exists an element in the Brauer group of this surface, which has two different local invariants on a given finite place, i.e. this element gives an obstruction to weak approximation for this surface. In this subsection, we generalize it.

Next, we will construct a Châtelet surface of the second kind mentioned in Subsection 1.3.

3.2.1. Choice of parameters for the equation (1). Given an extension of number fields L/K , and a finite subset $S \subset \Omega_K \setminus (\infty_K^c \cup 2_K)$, we choose an element $a \in \mathcal{O}_K \setminus K^2$ as in Subsubsection 3.0.1.

We will choose an element $b \in K^\times$ with respect to the chosen a in the following way.

Let $S' = \{v \in \infty_K^r \mid \tau_v(a) < 0\} \cup \{v \in \Omega_K^f \setminus 2_K \mid v(a) \text{ is odd}\}$ be as in Remark 3.0.5, then $S' \supset S$ is a finite set. By Lemma 2.2.4, for $v \in S \setminus \infty_K$, the set $\{b \in K_v \mid v(b) = -v(a)\}$ is a nonempty open subset of K_v ; for $v \in S' \setminus (S \cup \infty_K)$, the set $\{b \in K_v \mid v(b) = v(a)\}$

is a nonempty open subset of \mathcal{O}_{K_v} . By Lemma 2.0.1, we can choose a nonzero element $b \in \mathcal{O}_S[1/2]$ satisfying the following conditions:

- $v(b) = -v(a)$ for all $v \in S \setminus \infty_K$,
- $v(b) = v(a)$ for all $v \in S' \setminus (S \cup \infty_K)$.

We will choose an element $c \in K^\times$ with respect to the chosen a, b in the following way.

Let $S'' = \{v \in \Omega_K^f \setminus 2_K \mid v(b) \neq 0\}$, then S'' is a finite set and $S' \setminus \infty_K \subset S''$. By Theorem 2.0.2, we can take two different finite places $v_1, v_2 \in \Omega_K^f \setminus S''$ splitting completely in L . If $v \in S \setminus \infty_K$, then $v(a)$ is odd. In this case, by Lemma 2.2.7, the set $\{c \in \mathcal{O}_{K_v}^\times \mid (a, c)_v = -1\}$ is a nonempty open subset of \mathcal{O}_{K_v} . If $v \in \{v_1, v_2\}$, then $b \in \mathcal{O}_{K_v}^\times$. In this case, by Lemma 2.2.4, the sets $\{c \in K_v \mid v(c) = 1\}$ and $\{c \in K_v \mid v(1 + cb^2) = 1\}$ are nonempty open subsets of \mathcal{O}_{K_v} . Also by Lemma 2.0.1, we can choose a nonzero element $c \in \mathcal{O}_K[1/2]$ satisfying the following conditions:

- $\tau_v(1 + cb^2) < 0$ for all $v \in S \cap \infty_K$,
- $\tau_v(c) > 0$ for all $v \in (S' \setminus S) \cap \infty_K$,
- $(a, c)_v = -1$ and $v(c) = 0$ for all $v \in S \setminus \infty_K$,
- $v_1(c) = 1$ and $v_2(1 + cb^2) = 1$ for the chosen v_1, v_2 above.

Let $P(x) = (cx^2 + 1)((1 + cb^2)x^2 + b^2)$, and let V_2 be the Châtelet surface given by $y^2 - az^2 = (cx^2 + 1)((1 + cb^2)x^2 + b^2)$.

Proposition 3.2.1. For any extension of number fields L/K , and any finite subset $S \subset \Omega_K \setminus (\infty_K \cup 2_K)$ splitting completely in L , there exists a Châtelet surface V_2 defined over K , which has the following properties.

- The Brauer group $\text{Br}(V_2)/\text{Br}(K) \cong \text{Br}(V_{2L})/\text{Br}(L) \cong \mathbb{Z}/2\mathbb{Z}$, is generated by an element $A \in \text{Br}(V_2)$. The subset $V_2(K) \subset V_2(L)$ is nonempty.
- For any $v \in S$, there exist P_v and Q_v in $V_2(K_v)$ such that the local invariants $\text{inv}_v(A(P_v)) = 0$ and $\text{inv}_v(A(Q_v)) = 1/2$. For any other $v \notin S$, and any $P_v \in V_2(K_v)$, the local invariant $\text{inv}_v(A(P_v)) = 0$.
- For any $v' \in S_L$, there exist $P_{v'}$ and $Q_{v'}$ in $V_2(L_{v'})$ such that the local invariants $\text{inv}_{v'}(A(P_{v'})) = 0$ and $\text{inv}_{v'}(A(Q_{v'})) = 1/2$. For any other $v' \notin S_L$, and any $P_{v'} \in V_2(L_{v'})$, the local invariant $\text{inv}_{v'}(A(P_{v'})) = 0$.

Proof. For the extension L/K , and the finite set S , we will check that the Châtelet surface V_2 chosen as in Subsubsection 3.2.1, has the properties.

We need to prove the statement about the Brauer group, and find the element A in this proposition.

By the choice of the places v_1 , the polynomial $x^2 + c$ is an Eisenstein polynomial, so it is irreducible over K_{v_1} . Since $v_1(a)$ is even, we have $K(\sqrt{a})K_{v_1} \not\cong K_{v_1}[x]/(cx^2 + 1)$. So $K(\sqrt{a}) \not\cong K[x]/(cx^2 + 1)$. The same argument holds for the place v_2 and polynomial $(1 + cb^2)x^2 + b^2$. For all places of S split completely in L , then by Remark 3.0.6, we have $a \in \mathcal{O}_K \setminus L^2$. By the splitting condition of v_1, v_2 , we have $L(\sqrt{a}) \not\cong L[x]/(cx^2 + 1)$ and $L(\sqrt{a}) \not\cong L[x]/((1 + cb^2)x^2 + b^2)$. So $P(x) = (cx^2 + 1)((1 + cb^2)x^2 + b^2)$ is separable and a product of two degree-2 irreducible factors over K and L . According to [Sko01, Proposition 7.1.1], the Brauer group $\text{Br}(V_2)/\text{Br}(K) \cong \text{Br}(V_{2L})/\text{Br}(L) \cong \mathbb{Z}/2\mathbb{Z}$. Furthermore, by Proposition 7.1.2 in loc. cit, we take the quaternion algebra $A = (a, cx^2 + 1) \in \text{Br}(V_2)$ as a generator element of this group. Then we have the equality $A = (a, cx^2 + 1) = (a, (1 + cb^2)x^2 + b^2)$ in $\text{Br}(V_2)$.

For $(x, y, z) = (0, b, 0)$ is a rational point on V_2^0 , the set $V_2(K)$ is nonempty. We denote this rational point by Q_0 .

We need to compute the evaluation of A on $V_2(K_v)$ for all $v \in \Omega_K$.

For any $v \in \Omega_K$, the local invariant $\text{inv}_v(A(Q_0)) = 0$. By Remark 3.0.2, it suffices to compute the local invariant $\text{inv}_v(A(P_v))$ for all $P_v \in V_2^0(K_v)$.

Suppose that $v \in (\infty_K \setminus S') \cup 2_K$. Then $a \in K_v^{\times 2}$, so $\text{inv}_v(A(P_v)) = 0$ for all $P_v \in V_2(K_v)$.

Suppose that $v \in (S' \setminus S) \cap \infty_K$. For any $x \in K$, by the choice of c , we have $\tau_v(cx^2 + 1) > 0$. Then $(a, cx^2 + 1)_v = 1$, so $\text{inv}_v(A(P_v)) = 0$ for all $P_v \in V_2(K_v)$.

Suppose that $v \in S' \setminus (S \cup \infty_K)$. Take an arbitrary $P_v \in V_2^0(K_v)$. If $\text{inv}_v(A(P_v)) = 1/2$, then $(a, cx^2 + 1)_v = -1 = (a, (1 + cb^2)x^2 + b^2)_v$ at P_v . By Lemma 2.2.2, the first equality implies $v(x) \leq 0$. For $v(a) = v(b) > 0$ and $v(c) \geq 0$, by Lemma 2.2.2, we have $(a, (1 + cb^2)x^2 + b^2)_v = (a, x^2)_v = 1$, which is a contradiction. So $\text{inv}_v(A(P_v)) = 0$.

Suppose that $v \in \Omega_K^f \setminus (S' \cup 2K)$. Take an arbitrary $P_v \in V_2^0(K_v)$. If $\text{inv}_v(A(P_v)) = 1/2$, then $(a, cx^2 + 1)_v = -1 = (a, (1 + cb^2)x^2 + b^2)_v$ at P_v . For $v(a)$ is even, by Lemma 2.2.1, the first equality implies that $v(cx^2 + 1)$ is odd. For $c \in \mathcal{O}_K[1/2]$, we have $v(x) \leq 0$. So $v(c + x^{-2})$ is odd and positive. For $v(b) \geq 0$, by Hensel's lemma, we have $1 + b^2(c + x^{-2}) \in K_v^{\times 2}$. So $(a, (1 + cb^2)x^2 + b^2)_v = (a, x^2)_v(a, 1 + b^2(c + x^{-2}))_v = 1$, which is a contradiction. So $\text{inv}_v(A(P_v)) = 0$.

Suppose that $v \in S \cap \infty_K$. Take $P_v = Q_0$, then $\text{inv}_v(A(P_v)) = 0$. By the choice of b, c , we have $\tau_v(\frac{b^2}{-cb^2-1}) > \tau_v(\frac{1}{-c}) > 0$. Take $x_0 \in K$ such that $\tau_v(x_0) > \sqrt{\tau_v(\frac{b^2}{-cb^2-1})}$, then $\tau_v((cx_0^2 + 1)((1 + cb^2)x_0^2 + b^2)) > 0$ and $\tau_v(cx_0^2 + 1) < 0$. So there exists a $Q_v \in V_2^0(K_v)$ with $x = x_0$. Then $\text{inv}_v(A(Q_v)) = 1/2$.

Suppose that $v \in S \setminus \infty_K$. Take $P_v = Q_0$, then $\text{inv}_v(A(P_v)) = 0$. Take $x_0 \in K_v$ such that $v(x_0) < 0$. For $v(b) = -v(a) < 0$ and $v(c) = 0$, by Lemma 2.2.2, we have $(a, cx_0^2 + 1)_v = (a, cx_0^2)_v = (a, c)_v$ and $(a, (1 + cb^2)x_0^2 + b^2)_v = (a, cb^2x_0^2)_v = (a, c)_v$. So $(a, (cx_0^2 + 1)((1 + cb^2)x_0^2 + b^2))_v = (a, c)_v(a, c)_v = 1$. Hence, there exists a $Q_v \in V_2^0(K_v)$ with $x = x_0$. For $(a, c)_v = -1$, we have $\text{inv}_v(A(Q_v)) = 1/2$.

Finally, we need to compute the evaluation of A on $V_2(L_{v'})$ for all $v' \in \Omega_L$.

For any $v' \in \Omega_L$, the local invariant $\text{inv}_{v'}(A(Q_0)) = 0$.

Suppose that $v' \in S_L$. Let $v \in \Omega_K$ be the restriction of v' on K . By the assumption that v is split completely in L , we have $K_v = L_{v'}$. So $V_2(K_v) = V_2(L_{v'})$. By the argument already shown, there exist $P_v, Q_v \in V_2(K_v)$ such that $\text{inv}_v(A(P_v)) = 0$ and $\text{inv}_v(A(Q_v)) = 1/2$. View P_v, Q_v as elements in $V_2(L_{v'})$, and let $P_{v'} = P_v$ and $Q_{v'} = Q_v$. Then $\text{inv}_{v'}(A(P_{v'})) = \text{inv}_v(A(P_v)) = 0$ and $\text{inv}_{v'}(A(Q_{v'})) = \text{inv}_v(A(Q_v)) = 1/2$.

Suppose that $v' \in \Omega_L \setminus S_L$. This local computation is the same as the case $v \in \Omega_K \setminus S$. \square

Remark 3.2.2. For any $v \in S$, and any $P_v \in V_2(K_v)$, the local invariant of the evaluation of A on P_v is 0 or 1/2. Let $U_1 = \{P_v \in V_2(K_v) \mid \text{inv}_v(A(P_v)) = 0\}$ and $U_2 = \{P_v \in V_2(K_v) \mid \text{inv}_v(A(P_v)) = 1/2\}$. Then U_1 and U_2 are nonempty disjoint open subsets of $V_2(K_v)$, and $V_2(K_v) = U_1 \sqcup U_2$.

The following proposition states that the surface V_2 in Proposition 3.2.1, has the following weak approximation properties.

Proposition 3.2.3. Given an extension of number fields L/K , and a finite subset $S \subset \Omega_K \setminus (\infty_K^c \cup 2K)$ splitting completely in L , let V_2 be a Châtelet surface satisfying those properties of Proposition 3.2.1. If $S = \emptyset$, then V_2 and V_{2L} satisfy weak approximation. If $S \neq \emptyset$, then V_2 satisfies weak approximation off S' for any finite subset $S' \subset \Omega_K$ such that $S' \cap S \neq \emptyset$, while it fails for any finite subset $S' \subset \Omega_K$ such that $S' \cap S = \emptyset$. And in the case $S \neq \emptyset$, the surface V_{2L} satisfies weak approximation off T for any finite subset $T \subset \Omega_L$ such that $T \cap S_L \neq \emptyset$, while it fails for any finite subset $T \subset \Omega_L$ such that $T \cap S_L = \emptyset$.

Proof. According to [CTSSD87a, Theorem B; CTSSD87b], the Brauer-Manin obstruction to the Hasse principle and weak approximation is the only one for Châtelet surfaces, so $V_2(K)$ is dense in $V_2(\mathbb{A}_K)^{\text{Br}}$.

Suppose that $S = \emptyset$, then for any $(P_v)_{v \in \Omega_K} \in V_2(\mathbb{A}_K)$, by Proposition 3.2.1, the sum $\sum_{v \in \Omega_K} \text{inv}_v(A(P_v)) = 0$. For $\text{Br}(V_2)/\text{Br}(K)$ is generated by the element A , we have $V_2(\mathbb{A}_K)^{\text{Br}} = V_2(\mathbb{A}_K)$. So $V_2(K)$ is dense in $V_2(\mathbb{A}_K)^{\text{Br}} = V_2(\mathbb{A}_K)$, i.e. the surface V_2 satisfies weak approximation.

Suppose that $S' \cap S \neq \emptyset$. Take $v_0 \in S' \cap S$. For any finite subset $R \subset \Omega_K \setminus \{v_0\}$, take a nonempty open subset $M = V_2(K_{v_0}) \times \prod_{v \in R} U_v \times \prod_{v \notin R \cup \{v_0\}} V_2(K_v) \subset V_2(\mathbb{A}_K)$. Take an element $(P_v)_{v \in \Omega_K} \in M$. By Proposition 3.2.1 and $v_0 \in S$, we can take an

element $P'_{v_0} \in V_2(K_{v_0})$ such that $\text{inv}_{v_0} A(P'_{v_0}) = 1/2$. By Proposition 3.2.1, the sum $\sum_{v \in \Omega_K \setminus \{v_0\}} \text{inv}_v(A(P_v))$ is 0 or $1/2$ in \mathbb{Q}/\mathbb{Z} . If it is $1/2$, then we replace P_{v_0} by P'_{v_0} . In this way, we get a new element $(P_v)_{v \in \Omega_K} \in M$. And the sum $\sum_{v \in \Omega_K} \text{inv}_v(A(P_v)) = 0$ in \mathbb{Q}/\mathbb{Z} . So $(P_v)_{v \in \Omega_K} \in V_2(\mathbb{A}_K)^{\text{Br}} \cap M$. For $V_2(K)$ is dense in $V_2(\mathbb{A}_K)^{\text{Br}}$, the set $V_2(K) \cap M \neq \emptyset$, which implies that V_2 satisfies weak approximation off $\{v_0\}$. So V_2 satisfies weak approximation off S' .

Suppose that $S \neq \emptyset$ and $S' \cap S = \emptyset$. Take $v_0 \in S$, and let $U_{v_0} = \{P_{v_0} \in V_2(K_{v_0}) \mid \text{inv}_{v_0}(A(P_{v_0})) = 1/2\}$. For $v \in S \setminus \{v_0\}$, let $U_v = \{P_v \in V_2(K_v) \mid \text{inv}_v(A(P_v)) = 0\}$. For any $v \in S$, by Remark 3.2.2, the set U_v is a nonempty open subset of $V_2(K_v)$. Let $M = \prod_{v \in S} U_v \times \prod_{v \notin S} V_2(K_v)$. It is a nonempty open subset of $V_2(\mathbb{A}_K)$. For any $(P_v)_{v \in \Omega_K} \in M$, by Proposition 3.2.1 and the choice of U_v , the sum $\sum_{v \in \Omega_K} \text{inv}_v(A(P_v)) = 1/2$ is nonzero in \mathbb{Q}/\mathbb{Z} . So $V_2(\mathbb{A}_K)^{\text{Br}} \cap M = \emptyset$, which implies $V_2(K) \cap M = \emptyset$. Hence V_2 does not satisfy weak approximation off S' .

The same argument applies to V_{2L} . □

Applying the construction method in Subsubsection 3.2.1, we have the following weak approximation properties for Châtelet surfaces.

Corollary 3.2.4. For any extension of number fields L/K , and any finite nonempty subset $S \subset \Omega_K \setminus (\infty_K^c \cup 2_K)$ splitting completely in L , there exists a Châtelet surface V defined over K such that $V(K) \neq \emptyset$. For any subfield $L' \subset L$ over K , the Brauer group $\text{Br}(V)/\text{Br}(K) \cong \text{Br}(V_{L'})/\text{Br}(L') \cong \mathbb{Z}/2\mathbb{Z}$. And the surface $V_{L'}$ has the following properties.

- For any finite subset $T' \subset \Omega_{L'}$ such that $T' \cap S_{L'} \neq \emptyset$, the surface $V_{L'}$ satisfies weak approximation off T' .
- For any finite subset $T' \subset \Omega_{L'}$ such that $T' \cap S_{L'} = \emptyset$, the surface $V_{L'}$ does not satisfy weak approximation off T' . In particular, the surface $V_{L'}$ does not satisfy weak approximation.

Proof. For the extension L/K , and S , let V be the Châtelet surface chosen as in Subsubsection 3.2.1. Applying the same argument about the field L to its subfield L' , the properties that we list, are just what we have explained in Proposition 3.2.1 and Proposition 3.2.3. □

Using the construction method in Subsubsection 3.2.1, we have the following example, which is a special case of Proposition 3.2.1. It will be used for further discussion.

Example 3.2.5. Let $K = \mathbb{Q}$, $L = \mathbb{Q}(\sqrt{3})$, and let $S = \{73\}$. The prime numbers 11, 23, 73 split completely in L . Using the construction method in Subsubsection 3.2.1, we choose data: $S = S' = S'' = 73$, $v_1 = 11$, $v_2 = 23$, $a = 73$, $b = 1/73$, $c = 99$ and $P(x) = (99x^2 + 1)(5428x^2/5329 + 1/5329)$. Then the Châtelet surface given by $y^2 - 73z^2 = P(x)$, has the properties of Proposition 3.2.1, Proposition 3.2.3.

3.3. Châtelet surfaces related to the Hasse principle. Iskovskikh [Isk71] gave an example of the intersection of two quadratic hypersurfaces in $\mathbb{P}_{\mathbb{Q}}^4$, which is a Châtelet surface over \mathbb{Q} given by $y^2 + z^2 = (x^2 - 2)(-x^2 + 3)$. He showed that this Châtelet surface is a counterexample to the Hasse principle. Similarly, Skorobogatov [Sko01, Pages 145-146] gave a family of Châtelet surfaces with a parameter over \mathbb{Q} . He discussed the property of the Hasse principle for this family. Poonen [Poo09, Proposition 5.1] generalized their arguments to any number field. Given an number field K , he constructed a Châtelet surface defined over K , which is a counterexample to the Hasse principle. He used the Čebotarev's density theorem for some ray class fields to choose the parameters for the equation (1). The Châtelet surface that he constructed, has the property of [Poo09, Lemma 5.5] (a special situation of the following Proposition 3.3.1: the case when $S = \{v_0\}$ for some place v_0 associated to some large prime element in \mathcal{O}_K), which is the main ingredient in the proof of [Poo09, Proposition 5.1]. In this subsection, we generalize them.

Next, we will construct a Châtelet surface of the third kind mentioned in Subsection 1.3.

3.3.1. *Choice of parameters for the equation (1).* Given an extension of number fields L/K , and a finite subset $S \subset \Omega_K \setminus (\infty_K^c \cup 2_K)$, we choose an element $a \in \mathcal{O}_K \setminus K^2$ as in Subsubsection 3.0.1.

We will choose an element $b \in K^\times$ with respect to the chosen a in the following way.

Let $S' = \{v \in \infty_K^r \mid \tau_v(a) < 0\} \cup \{v \in \Omega_K^f \setminus 2_K \mid v(a) \text{ is odd}\}$ be as in Remark 3.0.5, then $S' \supset S$ is a finite set. If $v \in S' \setminus \infty_K$, then $v(a)$ is odd. Then by Lemma 2.2.7, the set $\{b \in \mathcal{O}_{K_v}^\times \mid (a, b)_v = -1\}$ is a nonempty open subset of \mathcal{O}_{K_v} . If $v \in S' \setminus (S \cup \infty_K)$, then by Lemma 2.2.5, the set $\{b \in \mathcal{O}_{K_v}^\times \mid (a, b)_v = 1\}$ is a nonempty open subset of \mathcal{O}_{K_v} . By Lemma 2.0.1, we can choose a nonzero element $b \in \mathcal{O}_K[1/2]$ satisfying the following conditions:

- $\tau_v(b) < 0$ for all $v \in S \cap \infty_K$,
- $\tau_v(b) > 0$ for all $v \in (S' \setminus S) \cap \infty_K$,
- $(a, b)_v = -1$ and $v(b) = 0$ for all $v \in S' \setminus \infty_K$,
- $(a, b)_v = 1$ and $v(b) = 0$ for all $v \in S' \setminus (S \cup \infty_K)$.

We will choose an element $c \in K^\times$ with respect to the chosen a, b in the following way.

Let $S'' = \{v \in \Omega_K^f \setminus 2_K \mid v(b) \neq 0\}$, then S'' is a finite set and $S' \cap S'' = \emptyset$. By Theorem 2.0.2, we can take two different finite places $v_1, v_2 \in \Omega_K^f \setminus (S' \cup S'' \cup 2_K)$ splitting completely in L . If $v \in (S' \setminus \infty_K) \cup \{v_1, v_2\}$, then $b \in \mathcal{O}_{K_v}^\times$. In this case, by Lemma 2.2.4, the sets $\{c \in K_v \mid v(bc+1) = v(a)+2\}$, $\{c \in K_v \mid v(c) = 1\}$ and $\{c \in K_v \mid v(bc+1) = 1\}$ are nonempty open subsets of \mathcal{O}_{K_v} . If $v \in S''$, by Lemma 2.2.5, the set $\{c \in \mathcal{O}_{K_v}^\times \mid (a, c)_v = 1\}$ is a nonempty open subset of \mathcal{O}_{K_v} . Also by Lemma 2.0.1, we can choose a nonzero element $c \in \mathcal{O}_K[1/2]$ satisfying the following conditions:

- $0 < \tau_v(c) < -1/\tau_v(b)$ for all $v \in S \cap \infty_K$,
- $\tau_v(bc+1) < 0$ for all $v \in (S' \setminus S) \cap \infty_K$,
- $v(bc+1) = v(a)+2$ for all $v \in S' \setminus \infty_K$,
- $(a, c)_v = 1$ for all $v \in S''$,
- $v_1(c) = 1$ and $v_2(bc+1) = 1$ for the chosen v_1, v_2 above.

Let $P(x) = (x^2 - c)(bx^2 - bc - 1)$, and let V_3 be the Châtelet surface given by $y^2 - az^2 = (x^2 - c)(bx^2 - bc - 1)$.

Proposition 3.3.1. For any extension of number fields L/K , and any finite subset $S \subset \Omega_K \setminus (\infty_K^c \cup 2_K)$ splitting completely in L , there exists a Châtelet surface V_3 defined over K , which has the following properties.

- The Brauer group $\text{Br}(V_3)/\text{Br}(K) \cong \text{Br}(V_{3L})/\text{Br}(L) \cong \mathbb{Z}/2\mathbb{Z}$, is generated by an element $A \in \text{Br}(V_3)$. The subset $V_3(\mathbb{A}_K) \subset V_3(\mathbb{A}_L)$ is nonempty.
- For any $v \in \Omega_K$, and any $P_v \in V_3(K_v)$,

$$\text{inv}_v(A(P_v)) = \begin{cases} 0 & \text{if } v \notin S, \\ 1/2 & \text{if } v \in S. \end{cases}$$

- For any $v' \in \Omega_L$, and any $P_{v'} \in V_3(L_{v'})$,

$$\text{inv}_{v'}(A(P_{v'})) = \begin{cases} 0 & \text{if } v' \notin S_L, \\ 1/2 & \text{if } v' \in S_L. \end{cases}$$

Proof. For the extension L/K , and the finite set S , we will check that the Châtelet surface V_3 chosen as in Subsubsection 3.3.1, has the properties.

Firstly, we need to check that V_3 has an \mathbb{A}_K -adelic point.

Suppose that $v \in (\infty_K \setminus S') \cup 2_K$. Then $a \in K_v^{\times 2}$. By Remark 3.0.1, the surface V_3 admits a K_v -point.

Suppose that $v \in (S' \setminus S) \cap \infty_K$. Let $x_0 = 0$. For $\tau_v(b) > 0$ and $\tau_v(bc+1) < 0$, we have $\tau_v(c) < 0$ and $\tau_v((x_0^2 - c)(bx_0^2 - bc - 1)) = \tau_v(c(bc+1)) > 0$, which implies that V_3^0 admits a K_v -point with $x = 0$.

Suppose that $v \in S' \setminus (S \cup \infty_K)$. Take $x_0 \in K_v$ such that the valuation $v(x_0) < 0$. For $b \in \mathcal{O}_{K_v}^\times$ and $c \in \mathcal{O}_K[1/2]$, by Lemma 2.2.2, we have $(a, x_0^2 - c)_v = (a, x_0^2)_v = 1$ and

$(a, bx_0^2 - bc - 1)_v = (a, bx_0^2)_v = (a, b)_v$. By the choice of b , we have $(a, b)_v = 1$. Hence $(a, (x_0^2 - c)(bx_0^2 - bc - 1))_v = (a, b)_v = 1$, which implies that V_3^0 admits a K_v -point with $x = x_0$.

Suppose that $v \in S''$. By the choice of a, b, c , we have $(a, c)_v = 1$, $v(a)$ even, and $bc + 1 \in \mathcal{O}_{K_v}^\times$. By Lemma 2.2.1, we have $(a, bc+1)_v = 1$. Let $x_0 = 0$. Then $(a, (x_0^2 - c)(bx_0^2 - bc - 1))_v = (a, c(bc+1))_v = (a, c)_v(a, bc+1)_v = 1$, which implies that V_3^0 admits a K_v -point with $x = 0$.

Suppose that $v \in \Omega_K^f \setminus (S' \cup S'' \cup 2K)$. Then $v(b) = 0$. Take $x_0 \in K_v$ such that the valuation $v(x_0) < 0$. For $b \in \mathcal{O}_{K_v}^\times$ and $c \in \mathcal{O}_K[1/2]$, by Lemma 2.2.2, we have $(a, x_0^2 - c)_v = (a, x_0^2)_v = 1$ and $(a, bx_0^2 - bc - 1)_v = (a, bx_0^2)_v = (a, b)_v$. For $v(a)$ and $v(b)$ are even, by Lemma 2.2.1, we have $(a, b)_v = 1$. So $(a, (x_0^2 - c)(bx_0^2 - bc - 1))_v = (a, b)_v = 1$, which implies that V_3^0 admits a K_v -point with $x = x_0$.

Suppose that $v \in S \cap \infty_K$. Let $x_0 = 0$. Then by the choice of a, b, c , we have $\tau_v(a) < 0$, $\tau_v(c) > 0$ and $\tau_v(bc + 1) > 0$. So $(a, (x_0^2 - c)(bx_0^2 - bc - 1))_v = (a, c(bc + 1))_v = 1$, which implies that V_3^0 admits a K_v -point with $x = 0$.

Suppose that $v \in S \setminus \infty_K$. Choose a prime element π_v and take $x_0 = \pi_v$. By the choice of a, b, c , we have $b, c \in \mathcal{O}_{K_v}^\times$, $v(bx_0^2) = 2$, and $v(bc + 1) = v(a) + 2 \geq 3$. By Lemma 2.2.2, we have $(a, x_0^2 - c)_v = (a, -c)_v$ and $(a, bx_0^2 - bc - 1)_v = (a, bx_0^2)_v$. By Hensel's lemma, we have $-bc = 1 - (bc + 1) \in K_v^{\times 2}$. So $(a, (x_0^2 - c)(bx_0^2 - bc - 1))_v = (a, -bcx_0^2)_v = 1$, which implies that V_3^0 admits a K_v -point with $x = \pi_v$.

Secondly, we need to prove the statement about the Brauer group, and find the element A in this proposition.

By the choice of the places v_1 , the polynomial $x^2 - c$ is an Eisenstein polynomial, so it is irreducible over K_{v_1} . Since $v_1(a)$ is even, we have $K(\sqrt{a})K_{v_1} \not\cong K_{v_1}[x]/(x^2 - c)$. So $K(\sqrt{a}) \not\cong K[x]/(x^2 - c)$. The same argument holds for the place v_2 and polynomial $bx^2 - bc - 1$. For all places of S split completely in L , then by Remark 3.0.6, we have $a \in \mathcal{O}_K \setminus L^2$. By the splitting condition of v_1, v_2 , we have $L(\sqrt{a}) \not\cong L[x]/(x^2 - c)$ and $L(\sqrt{a}) \not\cong L[x]/(bx^2 - bc - 1)$. So $P(x) = (x^2 - c)(bx^2 - bc - 1)$ is separable and a product of two degree-2 irreducible factors over K and L . According to [Sko01, Proposition 7.1.1], the Brauer group $\text{Br}(V_3)/\text{Br}(K) \cong \text{Br}(V_{3L})/\text{Br}(L) \cong \mathbb{Z}/2\mathbb{Z}$. Furthermore, by Proposition 7.1.2 in loc. cit, we take the quaternion algebra $A = (a, x^2 - c) \in \text{Br}(V_3)$ as a generator element of this group. Then we have the equality $A = (a, x^2 - c) = (a, bx^2 - bc - 1)$ in $\text{Br}(V_3)$.

Thirdly, We need to compute the evaluation of A on $V_3(K_v)$ for all $v \in \Omega_K$.

By Remark 3.0.2, it suffices to compute the local invariant $\text{inv}_v(A(P_v))$ for all $P_v \in V_3^0(K_v)$ and all $v \in \Omega_K$.

Suppose that $v \in (\infty_K \setminus S') \cup 2K$. Then $a \in K_v^{\times 2}$, so $\text{inv}_v(A(P_v)) = 0$ for all $P_v \in V_3(K_v)$. Suppose that $v \in (S' \setminus S) \cap \infty_K$. By the choice of b, c , we have $\tau_v(b) > 0$ and $\tau_v(bc + 1) < 0$. So, for any $x \in K$, we have $(a, bx^2 - bc - 1)_v = 1$. Hence $\text{inv}_v(A(P_v)) = 0$ for all $P_v \in V_3^0(K_v)$. Suppose that $v \in S' \setminus (S \cup \infty_K)$. By the choice of b , we have $(a, b)_v = 1$. Take an arbitrary $P_v \in V_3^0(K_v)$. If $v(x) < 0$ at P_v , by Lemma 2.2.2, we have $(a, x^2 - c)_v = (a, x^2)_v = 1$. If $v(x) > 0$ at P_v , since $b, c \in \mathcal{O}_{K_v}^\times$ and $v(bc + 1) = v(a) + 2 \geq 3$, by Lemma 2.2.2, we have $(a, x^2 - c)_v = (a, -c)_v$. By Hensel's lemma, we have $-bc = 1 - (bc + 1) \in K_v^{\times 2}$. So $(a, x^2 - c)_v = (a, -c)_v = (a, -bc)_v = 1$. If $v(x) = 0$ at P_v , since $b \in \mathcal{O}_{K_v}^\times$ and $v(bc + 1) = v(a) + 2 \geq 3$, by Lemma 2.2.2, we have $(a, bx^2 - bc - 1)_v = (a, bx^2)_v = 1$. So $\text{inv}_v(A(P_v)) = 0$. Suppose that $v \in \Omega_K^f \setminus (S' \cup 2K)$. Take an arbitrary $P_v \in V_3^0(K_v)$. If $\text{inv}_v(A(P_v)) = 1/2$, then $(a, bx^2 - bc - 1)_v = (a, x^2 - c)_v = -1$ at P_v . For $v(a)$ is even, by Lemma 2.2.1, the last equality implies that $v(x^2 - c)$ is odd, so it is positive. So $v(bx^2 - bc - 1) = 0$. By Lemma 2.2.1, we have $(a, bx^2 - bc - 1)_v = 1$, which is a contradiction. So $\text{inv}_v(A(P_v)) = 0$.

Suppose that $v \in S \cap \infty_K$. Take an arbitrary $P_v \in V_3^0(K_v)$. If $A(P_v) = 0$, then $(a, bx^2 - bc - 1)_v = (a, x^2 - c)_v = 1$ at P_v . The last equality implies that $\tau_v(x^2 - c) > 0$. By the choice of b , we have $\tau_v(b) < 0$, so $\tau_v(bx^2 - bc - 1) < 0$, which contradicts $(a, bx^2 - bc - 1)_v = 1$. So $\text{inv}_v(A(P_v)) = 1/2$.

Suppose that $v \in S \setminus \infty_K$. By the choice of b , we have $(a, b)_v = -1$. Take an arbitrary $P_v \in V_3^0(K_v)$. If $v(x) \leq 0$ at P_v , for $b \in \mathcal{O}_{K_v}^\times$ and $v(bc + 1) = v(a) + 2 \geq 3$, by Lemma

2.2.2, we have $(a, bx^2 - bc - 1)_v = (a, bx^2)_v = -1$. If $v(x) > 0$ at P_v , for $b, c \in \mathcal{O}_{K_v}^\times$ and $v(bc + 1) = v(a) + 2 \geq 3$, by Lemma 2.2.2, we have $(a, x^2 - c)_v = (a, -c)_v$. By Hensel's lemma, we have $-bc = 1 - (bc + 1) \in K_v^{\times 2}$. So $(a, x^2 - c)_v = (a, -c)_v = -(a, -bc)_v = -1$. So $\text{inv}_v(A(P_v)) = 1/2$.

Finally, we need to compute the evaluation of A on $V_3(L_{v'})$ for all $v' \in \Omega_L$.

Suppose that $v' \in S_L$. Let $v \in \Omega_K$ be the restriction of v' on K . By the assumption that v is split completely in L , we have $K_v = L_{v'}$. So $V_3(K_v) = V_3(L_{v'})$. Then for any $P_{v'} \in V_3(L_{v'})$, denote $P_{v'}$ in $V_3(K_v)$ by P_v . Then by the argument already shown, the local invariant $\text{inv}_{v'}(A(P_{v'})) = \text{inv}_v(A(P_v)) = 1/2$.

Suppose that $v' \in \Omega_L \setminus S_L$. This local computation is the same as the case $v \in \Omega_K \setminus S$. \square

Remark 3.3.2. If the surface V_3 has a K -rational point Q , then by the global reciprocity law, the sum $\sum_{v \in \Omega_K} \text{inv}_v(A(Q)) = 0$ in \mathbb{Q}/\mathbb{Z} . If the number $\sharp S$ is odd, then from Proposition 3.3.1 that we get, this sum is $\sharp S/2$, which is nonzero in \mathbb{Q}/\mathbb{Z} . So, in this case, the surface V_3 has no K -rational point, which implies that the surface V_3 is a counterexample to the Hasse principle. If the number $\sharp S$ is even, then for any $(P_v)_{v \in \Omega_K} \in V_3(\mathbb{A}_K)$, by Proposition 3.3.1, the sum $\sum_{v \in \Omega_K} \text{inv}_v(A(P_v)) = \sharp S/2 = 0$ in \mathbb{Q}/\mathbb{Z} . For $\text{Br}(V_3)/\text{Br}(K)$ is generated by the element A , we have $V_3(\mathbb{A}_K)^{\text{Br}} = V_3(\mathbb{A}_K) \neq \emptyset$. According to [CTSSD87a, Theorem B; CTSSD87b], the Brauer-Manin obstruction to the Hasse principle and weak approximation is the only one for Châtelet surfaces. So, in this case, the set $V_3(K) \neq \emptyset$, and it is dense in $V_3(\mathbb{A}_K)^{\text{Br}} = V_3(\mathbb{A}_K)$, i.e. the surface V_3 has a K -rational point and satisfies weak approximation. In particular, if the number $\sharp S = 0$, i.e. $S = \emptyset$, though the Brauer group $\text{Br}(V_3)/\text{Br}(K)$ is nontrivial, it gives no obstruction to weak approximation for V_3 .

Applying the construction method in Subsubsection 3.3.1, we can relate the properties in Proposition 3.2.1 to the Hasse principle and weak approximation.

Corollary 3.3.3. For any extension of number fields L/K , there exists a Châtelet surface V defined over K such that $V(\mathbb{A}_K) \neq \emptyset$. For any subfield $L' \subset L$ over K , the Brauer group $\text{Br}(V)/\text{Br}(K) \cong \text{Br}(V_{L'})/\text{Br}(L') \cong \mathbb{Z}/2\mathbb{Z}$. And the surface $V_{L'}$ has the following properties.

- If the degree $[L' : K]$ is odd, then the surface $V_{L'}$ is a counterexample to the Hasse principle. In particular, the surface V is a counterexample to the Hasse principle.
- If the degree $[L' : K]$ is even, then the surface $V_{L'}$ satisfies weak approximation. In particular, in this case, the set $V(L') \neq \emptyset$.

Proof. By Theorem 2.0.2, we can take a place $v_0 \in \Omega_K \setminus (\infty_K^c \cup 2_K)$ splitting completely in L . Let $S = \{v_0\}$. Using the construction method in Subsubsection 3.3.1, there exists a Châtelet surface V defined over K having the properties of Proposition 3.3.1. By the same argument as in the proof of Proposition 3.3.1, we have $\text{Br}(V)/\text{Br}(K) \cong \text{Br}(V_{L'})/\text{Br}(L') \cong \mathbb{Z}/2\mathbb{Z}$. For v_0 splits completely in L , it also does in L' . For $\sharp S$ is odd, if $[L' : K]$ is odd, then $\sharp S_{L'}$ is odd; if $[L' : K]$ is even, then $\sharp S_{L'}$ is even. Applying the same argument about the field L to its subfield L' , the properties that we list, are just what we have explained in Remark 3.3.2. \square

Remark 3.3.4. Though the Brauer group $\text{Br}(V)/\text{Br}(K) \cong \text{Br}(V_{L'})/\text{Br}(L') \cong \mathbb{Z}/2\mathbb{Z}$ in Corollary 3.3.3, is nontrivial, it gives an obstruction to the Hasse principle for V , also $V_{L'}$ if $[L' : K]$ is odd; but no longer gives an obstruction to weak approximation for $V_{L'}$ if $[L' : K]$ is even.

Using the construction method in Subsubsection 3.3.1, we have the following example, which is a special case of Proposition 3.3.1. It will be used for further discussion.

Example 3.3.5. Let $K = \mathbb{Q}$ and $L = \mathbb{Q}(\zeta_7 + \zeta_7^{-1})$ be as in Example 3.1.3, and let $S = \{13\}$. For $13^2 \equiv 1 \pmod{7}$, $41^2 \equiv 1 \pmod{7}$ and $43 \equiv 1 \pmod{7}$, the places 13, 41, 43 split completely in L . Using the construction method in Subsubsection 3.3.1, we choose data: $S = \{13\}$, $S' = \{13, 29\}$, $S'' = \{5\}$, $v_1 = 43$, $v_2 = 41$, $a = 377$, $b = 5$, $c = 878755181$ and $P(x) = (x^2 - 878755181)(5x^2 - 4393775906)$. Then the Châtelet surface given by $y^2 - 377z^2 = P(x)$, has the properties of Proposition 3.3.1.

4. STOLL'S CONJECTURE FOR CURVES

Whether all failures of the Hasse principle of smooth, projective, and geometrically connected curves defined over a number field, are explained by the Brauer-Manin obstruction, was considered by Skorobogatov [Sko01, Chapter 6.3] and Scharaschkin [Sch99] independently. Furthermore, Stoll [Sto07, Conjecture 9.1] made the following conjecture.

Given a curve C defined over a number field K , let $C(\mathbb{A}_K)_\bullet = \prod_{v \in \infty_K} \{\text{connected components of } C(K_v)\} \times C(\mathbb{A}_K^f)$. The product topology of $\prod_{v \in \infty_K} \{\text{connected components of } C(K_v)\}$ with discrete topology and $C(\mathbb{A}_K^f)$ with adelic topology, gives a topology for $C(\mathbb{A}_K)_\bullet$. For any $A \in \text{Br}(C)$, and any $v \in \infty_K$, the evaluation of A on each connected component of $C(K_v)$ is constant. So, the notation $C(\mathbb{A}_K)_\bullet^{\text{Br}}$ makes sense.

Conjecture 4.0.1. [Sto07, Conjecture 9.1] For any smooth, projective, and geometrically connected curve C defined over a number field K , the set $C(K)$ is dense in $C(\mathbb{A}_K)_\bullet^{\text{Br}}$. In particular, the curve C satisfies weak approximation with Brauer-Manin obstruction off ∞_K .

Remark 4.0.2. If Conjecture 4.0.1 holds for a given curve, which is a counterexample to the Hasse principle, then its failure of the Hasse principle is explained by the Brauer-Manin obstruction. For an elliptic curve defined over K , if its Tate-Shafarevich group is finite, then by the dual sequence of Cassels-Tate, Conjecture 4.0.1 holds for this elliptic curve. With the effort of Kolyvagin [Kol90, Kol91], Gross and Zagier [GZ86], and many others, for an elliptic curve defined over \mathbb{Q} , if its analytic rank equals zero or one, then its Mordell-Weil rank equals its analytic rank, and its Tate-Shafarevich group is finite. So, Conjecture 4.0.1 holds for this elliptic curve.

Definition 4.0.3. Given a nontrivial extension of number fields L/K , let C be a smooth, projective, and geometrically connected curve defined over K . We say that a triple (C, K, L) is of type *I* if $C(K)$ and $C(L)$ are finite nonempty sets, $C(K) \neq C(L)$ and Stoll's Conjecture 4.0.1 holds for the curve C . We say that a triple (C, K, L) is of type *II* if (C, K, L) is of type *I*, and there exists a real place $v' \in \infty_L$ such that $C(L_{v'})$ is connected.

Lemma 4.0.4. Given a nontrivial extension of number fields L/K , if Conjecture 4.0.1 holds for all smooth, projective, and geometrically connected curves defined over K , then there exists a curve C defined over K such that the triple (C, K, L) is of type *I*. Furthermore, if L has a real place, then this triple (C, K, L) is of type *II*.

Proof. Since L is a finite separable extension over K , there exists a $\theta \in L$ such that $L = K(\theta)$. Let $f(x)$ be the minimal polynomial of θ . Let $n = \deg(f)$, then $n = [L : K] \geq 2$. Let $\tilde{f}(w_0, w_1)$ be the homogenization of f . If n is odd, we consider a curve C defined over K by a homogeneous equation: $w_2^{n+2} = \tilde{f}(w_0, w_1)(w_1^2 - w_0^2)$ with homogeneous coordinates $(w_0 : w_1 : w_2) \in \mathbb{P}^2$. For the polynomials $f(x)$ and $x^2 - 1$ are separable and coprime in $K[x]$, the curve C is smooth, projective, and geometrically connected. By genus formula for a plane curve, the genus of C equals $g(C) = n(n+1)/2 > 1$. By Faltings's theorem, the sets $C(K)$ and $C(L)$ are finite. It is easy to check that $(w_0 : w_1 : w_2) = (1 : 1 : 0) \in C(K)$ and $(\theta : 1 : 0) \in C(L) \setminus C(K)$. By the assumption that Conjecture 4.0.1 holds for all smooth, projective, and geometrically connected curves over K , we have that the triple (C, K, L) is of type *I*. Since $n+2$ is odd, the space $C(K_{v'})$ is connected for all $v' \in \infty_L$. So, if L has a real place, then this triple (C, K, L) is of type *II*. If n is even, we replace the homogeneous equation $w_2^{n+2} = \tilde{f}(w_0, w_1)(w_1^2 - w_0^2)$ by $w_2^{n+3} = \tilde{f}(w_0, w_1)w_1(w_1^2 - w_0^2)$. The same argument applies to this new curve and new triple. \square

Remark 4.0.5. For some nonsquare integer d , let $K = \mathbb{Q}$ and $L = \mathbb{Q}(\sqrt{d})$. Consider an elliptic curve E_d defined by a Weierstraß equation: $y^2 = x^3 + d$. Let $E_d^{(d)}$ be the quadratic twist of E_d by d . The curve E_d is connected over \mathbb{R} . It is easy to check that the point $(x, y) = (0, \sqrt{d}) \in C(L) \setminus C(K)$. If both $E_d(\mathbb{Q})$ and $E_d^{(d)}(\mathbb{Q})$ are finite, then the set $E_d(L)$ is finite, cf. [Sil09, Exercise 10.16]. If additionally, the Tate-Shafarevich group $\text{III}(E_d, \mathbb{Q})$ is finite, then the triple (E_d, K, L) is of type *II*.

5. POONEN'S PROPOSITION

For our result is base on Poonen's proposition [Poo10, Proposition 5.4]. We recall that paper and his general result first. There exist some remarks on it in [Lia18, Section 4.1]. Colliot-Thélène [CT10, Proposition 2.1] gave another proof of that proposition.

Recall 5.0.1. Let B be a smooth, projective, and geometrically connected variety over a number field K . Let \mathcal{L} be a line bundle on B , assuming the set of global sections $\Gamma(B, \mathcal{L}^{\otimes 2}) \neq 0$. Let $\mathcal{E} = \mathcal{O}_B \oplus \mathcal{O}_B \oplus \mathcal{L}$. Let a be a constant in K^\times , and let s be a nonzero global section in $\Gamma(B, \mathcal{L}^{\otimes 2})$. The zero locus of $(1, -a, -s) \in \Gamma(B, \mathcal{O}_B \oplus \mathcal{O}_B \oplus \mathcal{L}^{\otimes 2}) \subset \Gamma(B, \text{Sym}^2 \mathcal{E})$ in the projective space bundle $\text{Proj}(\mathcal{E})$ is a projective and geometrically integral variety, denoted by X with the natural projection $X \rightarrow B$. Let \overline{K} be an algebraic closure field of K . Denote $B \times_{\text{Spec } K} \text{Spec } \overline{K}$ by \overline{B} .

Proposition 5.0.2. [Poo10, Proposition 5.3] Given a number field K , all notations are the same as in Recall 5.0.1. Let $\alpha: X \rightarrow B$ be the natural projection. Assume that

- the closed subscheme defined by $s = 0$ in B is smooth, projective, and geometrically connected,
- $\text{Br } \overline{B} = 0$ and $X(\mathbb{A}_K) \neq \emptyset$.

Then X is smooth, projective, and geometrically connected. And $\alpha^*: \text{Br}(B) \rightarrow \text{Br}(X)$ is an isomorphism.

6. MAIN RESULTS FOR CHÂTELET SURFACE BUNDLES OVER CURVES

6.1. Preparation Lemmas. We state the following lemmas, which will be used for the proof of our theorems.

Fibration methods are used to do research on weak approximation, weak approximation with Brauer-Manin obstruction between two varieties. We modify those fibration methods to fit into our context.

Lemma 6.1.1. Given a number field K , and a finite subset $S \subset \Omega_K$, let $f: X \rightarrow Y$ be a K -morphism of proper K -varieties X and Y . We assume that

- (1) the set $Y(K)$ is finite,
- (2) the variety Y satisfies weak approximation with Brauer-Manin obstruction off S ,
- (3) for any $P \in Y(K)$, the fiber X_P of f over P satisfies weak approximation off S .

Then X satisfies weak approximation with Brauer-Manin obstruction off S .

Proof. For any finite subset $S' \subset \Omega_K \setminus S$, take an open subset $N = \prod_{v \in S'} U_v \times \prod_{v \notin S'} X(K_v) \subset X(\mathbb{A}_K)$ such that $N \cap X(\mathbb{A}_K)^{\text{Br}} \neq \emptyset$. Let $M = \prod_{v \in S'} f(U_v) \times \prod_{v \notin S'} f(X(K_v))$, then by the functoriality of Brauer-Manin pairing, $M \cap Y(\mathbb{A}_K)^{\text{Br}} \neq \emptyset$. By Assumptions (1) and (2), we have $Y(K) = \text{pr}^S(Y(\mathbb{A}_K)^{\text{Br}})$. So there exists $P_0 \in \text{pr}^S(M) \cap Y(K)$. Consider the fiber X_{P_0} . Let $L = \prod_{v \in S'} [X_{P_0}(K_v) \cap U_v] \times \prod_{v \notin S' \cup S} X_{P_0}(K_v)$, then it is a nonempty open subset of $X_{P_0}(\mathbb{A}_K^S)$. By Assumption (3), there exists $Q_0 \in L \cap X_{P_0}(K)$. So $Q_0 \in X(K) \cap N$, which implies that X satisfies weak approximation with Brauer-Manin obstruction off S . \square

Lemma 6.1.2. Given a number field K , and a finite subset $S \subset \Omega_K$, let $f: X \rightarrow Y$ be a K -morphism of proper K -varieties X and Y . We assume that

- (1) the set $Y(K)$ is finite,
- (2) the morphism $f^*: \text{Br}(Y) \rightarrow \text{Br}(X)$ is surjective,
- (3) there exists some $P \in Y(K)$ such that the fiber X_P of f over P does not satisfy weak approximation off S , and that $\prod_{v \in S} X_P(K_v) \neq \emptyset$.

Then X does not satisfy weak approximation with Brauer-Manin obstruction off S .

Proof. By Assumption (3), take a $P_0 \in Y(K)$ such that the fiber X_{P_0} does not satisfy weak approximation off S , and that $\prod_{v \in S} X_{P_0}(K_v) \neq \emptyset$. Then there exist a finite nonempty subset $S' \subset \Omega_K \setminus S$ and a nonempty open subset $L = \prod_{v \in S'} U_v \times \prod_{v \notin S'} X_{P_0}(K_v) \subset X_{P_0}(\mathbb{A}_K)$

such that $L \cap X_{P_0}(K) = \emptyset$. By Assumption (1), the set $Y(K)$ is finite, so we can take a Zariski open subset $V_{P_0} \subset Y$ such that $V_{P_0}(K) = \{P_0\}$. For any $v \in S'$, since U_v is open in $X_{P_0}(K_v) \subset f^{-1}(V_{P_0})(K_v)$, we can take an open subset W_v of $f^{-1}(V_{P_0})(K_v)$ such that $W_v \cap X_{P_0}(K_v) = U_v$. Consider the open subset $N = \prod_{v \in S'} W_v \times \prod_{v \notin S'} X(K_v) \subset X(\mathbb{A}_K)$, then $L \subset N$. By the functoriality of Brauer-Manin pairing and Assumption (2), we have $L \subset N \cap X(\mathbb{A}_K)^{\text{Br}}$. So $N \cap X(\mathbb{A}_K)^{\text{Br}} \neq \emptyset$. But $N \cap X(K) = N \cap X_{P_0}(K) = L \cap X_{P_0}(K) = \emptyset$, which implies that X does not satisfy weak approximation with Brauer-Manin obstruction off S . \square

We use the following lemma to choose a dominant morphism from a given curve to \mathbb{P}^1 .

Lemma 6.1.3. Given a nontrivial extension of number fields L/K , let C be a smooth, projective, and geometrically connected curve defined over K . Assume that the triple (C, K, L) is of type I (Definition 4.0.3). For any finite K -subscheme $R \subset \mathbb{P}^1 \setminus \{0, \infty\}$, there exists a dominant K -morphism $\gamma: C \rightarrow \mathbb{P}^1$ such that $\gamma(C(L) \setminus C(K)) = \{0\} \subset \mathbb{P}^1(K)$, $\gamma(C(K)) = \{\infty\} \subset \mathbb{P}^1(K)$, and that γ is étale over R .

Proof. Let $K(C)$ be the function field of C . For $C(K)$ and $C(L)$ are finite nonempty sets and $C(L) \setminus C(K) \neq \emptyset$, by Riemann-Roch theorem, we can choose a rational function $\phi \in K(C)^\times \setminus K^\times$ such that the set of its poles contains $C(K)$, and that the set of its zeros contains $C(L) \setminus C(K)$. This rational function ϕ gives a dominant K -morphism $\gamma_0: C \rightarrow \mathbb{P}^1$ such that $\gamma_0(C(L) \setminus C(K)) = \{0\} \subset \mathbb{P}^1(K)$ and $\gamma_0(C(K)) = \{\infty\} \subset \mathbb{P}^1(K)$. We can choose an automorphism $\varphi_{\lambda_0}: \mathbb{P}^1 \rightarrow \mathbb{P}^1$, $(u : v) \mapsto (\lambda_0 u : v)$ with $\lambda_0 \in K^\times$ such that the branch locus of γ_0 has no intersection with $\varphi_{\lambda_0}(R)$. Let $\lambda = (\varphi_{\lambda_0})^{-1} \circ \gamma_0$. Then the morphism λ is étale over R and satisfies other conditions. \square

The following lemma is well known.

Lemma 6.1.4. Let C be a curve over a field, and let $B = C \times \mathbb{P}^1$. Then $\text{Br} \overline{B} = 0$.

Proof. By [Gro68, III, Corollary 1.2], the Brauer group for a given curve over an algebraic closed field is zero. So $\text{Br}(\overline{C} \times \overline{\mathbb{P}^1}) \cong \text{Br}(\overline{C}) = 0$. \square

Definition 6.1.5. Let C be a smooth, projective, and geometrically connected curve defined over a number field. We say that a morphism $\beta: X \rightarrow C$ is a Châtelet surface bundle over the curve C , if

- X is a smooth, projective, and geometrically connected variety,
- the morphism β is faithfully flat, and proper,
- the generic fiber of β is a Châtelet surface over the function field of C .

Next, we construct Châtelet surface bundles over curves to give negative answers to Questions 1.2.

6.2. Non-invariance of weak approximation with Brauer-Manin obstruction. For any quadratic extension of number fields L/K , and assuming Conjecture 4.0.1, Liang [Lia18, Theorem 4.5] constructed a Châtelet surface bundle over a curve to give a negative answer to Question 1.2.1. Assuming that the extension L/K is quadratic, he constructed a Châtelet surface defined over K such that the property of weak approximation is not invariant under the extension of L/K . Then choosing a higher genus curve, he combined this Châtelet surface with the construction method of Poonen [Poo10] to get the result. His method only works for quadratic extensions. In this subsection, we generalize his result to any extension L/K .

Theorem 6.2.1. For any nontrivial extension of number fields L/K , and any finite subset $T \subset \Omega_L$, assuming that Conjecture 4.0.1 holds over K , there exists a Châtelet surface bundle over a curve: $X \rightarrow C$ defined over K such that

- X has a K -rational point, and satisfies weak approximation with Brauer-Manin obstruction off ∞_K ,
- X_L does not satisfy weak approximation with Brauer-Manin obstruction off T .

Proof. Firstly, we will construct two Châtelet surfaces. Let $S \subset \Omega_K$ be the set of all restrictions of T on K . By Theorem 2.0.2, we can take a finite place $v_0 \in \Omega_K^f \setminus (S \cup 2_K)$ splitting completely in L . For the extension L/K , and $S = \{v_0\}$, let V_0 be the Châtelet surface chosen as in Subsubsection 3.2.1. Then V_0 defined by $y^2 - az^2 = P_0(x)$ over K having the properties of Proposition 3.2.1. Let $P_\infty(x) = (1 - x^2)(x^2 - a)$, and let V_∞ be the Châtelet surface defined by $y^2 - az^2 = P_\infty(x)$. By the argument in the proof of Proposition 3.2.1, two degree-2 irreducible factors of $P_0(x)$ are prime to $x^2 - a$ in $K[x]$. So, the polynomials $P_0(x)$ and $P_\infty(x)$ are coprime in $K[x]$.

Secondly, we will construct a Châtelet surface bundle over a curve. Let $\tilde{P}_\infty(x_0, x_1)$ and $\tilde{P}_0(x_0, x_1)$ be the homogenizations of P_∞ and P_0 . Let $(u_0 : u_1) \times (x_0 : x_1)$ be the coordinates of $\mathbb{P}^1 \times \mathbb{P}^1$, and let $s' = u_0^2 \tilde{P}_\infty(x_0, x_1) + u_1^2 \tilde{P}_0(x_0, x_1) \in \Gamma(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(1, 2)^{\otimes 2})$. For $P_0(x)$ and $P_\infty(x)$ are coprime in $K[x]$, by Jacobian criterion, the locus Z' defined by $s' = 0$ in $\mathbb{P}^1 \times \mathbb{P}^1$ is smooth. Then the branch locus of the composition $Z' \hookrightarrow \mathbb{P}^1 \times \mathbb{P}^1 \xrightarrow{pr_1} \mathbb{P}^1$, denoted by R , is finite over K . By the assumption that Conjecture 4.0.1 holds over K , and Lemma 4.0.4, we can take a curve C defined over K such that the triple (C, K, L) is of type I. By Lemma 6.1.3, we can choose a K -morphism $\gamma: C \rightarrow \mathbb{P}^1$ such that $\gamma(C(L) \setminus C(K)) = \{0\} \subset \mathbb{P}^1(K)$, $\gamma(C(K)) = \{\infty\} \subset \mathbb{P}^1(K)$, and that γ is étale over R . Let $B = C \times \mathbb{P}^1$. Let $\mathcal{L} = (\gamma, id)^* \mathcal{O}(1, 2)$, and let $s = (\gamma, id)^*(s') \in \Gamma(B, \mathcal{L}^{\otimes 2})$. For γ is étale over the branch locus R , the locus Z defined by $s = 0$ in B is smooth. Since Z is defined by the support of the global section s , it is an effective divisor. The invertible sheaf $\mathcal{L}(Z')$ on $\mathbb{P}^1 \times \mathbb{P}^1$ is isomorphic to $\mathcal{O}(2, 4)$, which is a very ample sheaf on $\mathbb{P}^1 \times \mathbb{P}^1$. And (γ, id) is a finite morphism, so the pull back of this ample sheaf is again ample, which implies that the invertible sheaf $\mathcal{L}(Z)$ on $C \times \mathbb{P}^1$ is ample. By [Har97, Chapter III Corollary 7.9], the curve Z is geometrically connected. So the curve Z is smooth, projective, and geometrically connected. By Lemma 6.1.4, the Brauer group $\text{Br}(\bar{B}) = 0$. Let X be the zero locus of $(1, -a, -s) \in \Gamma(B, \mathcal{O}_B \oplus \mathcal{O}_B \oplus \mathcal{L}^{\otimes 2}) \subset \Gamma(B, \text{Sym}^2 \mathcal{E})$ in the projective space bundle $\text{Proj}(\mathcal{E})$ with the natural projection $\alpha: X \rightarrow B$. Using Proposition 5.0.2, the variety X is smooth, projective, and geometrically connected. Let $\beta: X \xrightarrow{\alpha} B = C \times \mathbb{P}^1 \xrightarrow{pr_1} C$ be the composition of α and pr_1 . Then β is a Châtelet surface bundle over the curve C .

At last, we will check that X has the properties.

We will show that X has a K -rational point. For any $P \in C(K)$, the fiber $\beta^{-1}(P) \cong V_\infty$. The surface V_∞ has a K -rational point $(x, y, z) = (0, 0, 1)$, so the set $X(K) \neq \emptyset$.

We will show that X satisfies weak approximation with Brauer-Manin obstruction off ∞_K . By Remark 3.0.4, the surface V_∞ satisfies weak approximation. So, for the morphism β , Assumption (3) of Lemma 6.1.1 holds. Since Conjecture 4.0.1 holds for the curve C , using Lemma 6.1.1 for the morphism β , the variety X satisfies weak approximation with Brauer-Manin obstruction off ∞_K .

We will show that X_L does not satisfy weak approximation with Brauer-Manin obstruction off T . By Proposition 5.0.2, the map $\alpha_L^*: \text{Br}(B_L) \rightarrow \text{Br}(X_L)$ is an isomorphism, so $\beta_L^*: \text{Br}(C_L) \rightarrow \text{Br}(X_L)$ is an isomorphism. By the choice of the curve C and morphism β , for any $Q \in C(L) \setminus C(K)$, the fiber $\beta^{-1}(Q) \cong V_{0L}$. By Proposition 3.2.3, the surface V_{0L} does not satisfy weak approximation off $T \cup \infty_L$. For $V_0(L) \neq \emptyset$, by Lemma 6.1.2, the variety X_L does not satisfy weak approximation with Brauer-Manin obstruction off $T \cup \infty_L$. So it does not satisfy weak approximation with Brauer-Manin obstruction off T . \square

6.3. Non-invariance of the failures of the Hasse principle explained by the Brauer-Manin obstruction. For extensions L/K of the following two cases, assuming Conjecture 4.0.1, we construct Châtelet surface bundles over curves to give negative answers to Question 1.2.2.

Theorem 6.3.1. For any number field K , and any nontrivial field extension L of odd degree over K , assuming that Conjecture 4.0.1 holds over K , there exists a Châtelet surface bundle over a curve: $X \rightarrow C$ defined over K such that

- X is a counterexample to the Hasse principle, and its failure of the Hasse principle is explained by the Brauer-Manin obstruction,

- X_L is a counterexample to the Hasse principle, but its failure of the Hasse principle cannot be explained by the Brauer-Manin obstruction.

Proof. Firstly, we will construct two Châtelet surfaces. By Theorem 2.0.2, we can take two different finite places $v_1, v_2 \in \Omega_K^f \setminus 2_K$ splitting completely in L . For the extension L/K , and $S = \{v_1, v_2\}$, we choose an element a as in Subsubsection 3.0.1. For the extension L/K , and $S_1 = \{v_1\}$, by Remark 3.0.7, choosing other parameters for the equation (1) as in Subsubsection 3.3.1, we have a Châtelet surface V_0 defined by $y^2 - az^2 = P_0(x)$ over K having the properties of Proposition 3.3.1. For the extension L/K , and $S_2 = \{v_2\}$, by Remark 3.0.7, choosing other parameters for the equation (1) as in Subsubsection 3.1.1, we have a Châtelet surface V_∞ defined by $y^2 - az^2 = P_\infty(x)$ over K having the properties of Proposition 3.1.1. By Remark 3.1.2, the polynomial $P_\infty(x)$ is irreducible over K . For $P_0(x)$ is a product of two degree-2 irreducible factors over K , the polynomials $P_0(x)$ and $P_\infty(x)$ are coprime in $K[x]$.

Secondly, we will construct a Châtelet surface bundle over a curve. Let $\tilde{P}_\infty(x_0, x_1)$ and $\tilde{P}_0(x_0, x_1)$ be the homogenizations of P_∞ and P_0 . Let $(u_0 : u_1) \times (x_0 : x_1)$ be the coordinates of $\mathbb{P}^1 \times \mathbb{P}^1$, and let $s' = u_0^2 \tilde{P}_\infty(x_0, x_1) + u_1^2 \tilde{P}_0(x_0, x_1) \in \Gamma(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(1, 2)^{\otimes 2})$. For $P_0(x)$ and $P_\infty(x)$ are coprime in $K[x]$, by Jacobian criterion, the locus Z' defined by $s' = 0$ in $\mathbb{P}^1 \times \mathbb{P}^1$ is smooth. Then the branch locus of the composition $Z' \hookrightarrow \mathbb{P}^1 \times \mathbb{P}^1 \xrightarrow{pr_1} \mathbb{P}^1$, denoted by R , is finite over K . By the assumption that Conjecture 4.0.1 holds over K , and Lemma 4.0.4, we can take a curve C defined over K such that the triple (C, K, L) is of type I. By Lemma 6.1.3, we can choose a K -morphism $\gamma: C \rightarrow \mathbb{P}^1$ such that $\gamma(C(L) \setminus C(K)) = \{0\} \subset \mathbb{P}^1(K)$, $\gamma(C(K)) = \{\infty\} \subset \mathbb{P}^1(K)$, and that γ is étale over R . Let $B = C \times \mathbb{P}^1$. Let $\mathcal{L} = (\gamma, id)^* \mathcal{O}(1, 2)$, and let $s = (\gamma, id)^*(s') \in \Gamma(B, \mathcal{L}^{\otimes 2})$. By the same argument as in the proof of Theorem 6.2.1, the locus Z defined by $s = 0$ in B is smooth, projective, and geometrically connected; the Brauer group $\text{Br}(\overline{B}) = 0$. Let X be the zero locus of $(1, -a, -s) \in \Gamma(B, \mathcal{O}_B \oplus \mathcal{O}_B \oplus \mathcal{L}^{\otimes 2}) \subset \Gamma(B, \text{Sym}^2 \mathcal{E})$ in the projective space bundle $\text{Proj}(\mathcal{E})$ with the natural projection $\alpha: X \rightarrow B$. By Proposition 5.0.2, the variety X is smooth, projective, and geometrically connected. Let $\beta: X \rightarrow C$ be the composition of α and pr_1 . Then β is a Châtelet surface bundle over the curve C .

At last, we will check that X has the properties.

We will show $X(\mathbb{A}_K) \neq \emptyset$. For any $P \in C(K)$, the fiber $\beta^{-1}(P) \cong V_\infty$. By Proposition 3.1.1, the set $V_\infty(\mathbb{A}_K^{\{v_2\}}) \neq \emptyset$. So $X(\mathbb{A}_K^{\{v_2\}}) \neq \emptyset$. For v_2 splits completely in L , take a place $v'_2 \in \Omega_L^f$ above v_2 , i.e. $v'_2 | v_2$ in L . Then $K_{v_2} = L_{v'_2}$. By Proposition 3.3.1, the set $V_0(\mathbb{A}_L) \neq \emptyset$. Take a point $Q \in C(L) \setminus C(K)$, then the fiber $\beta^{-1}(Q) \cong V_{0L}$. We have $X(K_{v_2}) = X_L(L_{v'_2}) \supset \beta^{-1}(Q)(L_{v'_2}) \cong V_0(L_{v'_2}) \neq \emptyset$. So $X(\mathbb{A}_K) \neq \emptyset$.

We will show $X(\mathbb{A}_K)^{\text{Br}} = \emptyset$. By Conjecture 4.0.1, the set $C(K)$ is finite, and $C(K) = pr^{\infty K}(C(\mathbb{A}_K)^{\text{Br}})$. By the functoriality of Brauer-Manin pairing, we have $pr^{\infty K}(X(\mathbb{A}_K)^{\text{Br}}) \subset \bigsqcup_{P \in C(K)} \beta^{-1}(P)(\mathbb{A}_K^{\infty K})$. But by Proposition 3.1.1, the set $V_\infty(K_{v_2}) = \emptyset$, so we have $pr^{\infty K}(X(\mathbb{A}_K)^{\text{Br}}) \subset \bigsqcup_{P \in C(K)} \beta^{-1}(P)(\mathbb{A}_K^{\infty K}) \cong V_\infty(\mathbb{A}_K^{\infty K}) \times C(K) = \emptyset$, which implies that $X(\mathbb{A}_K)^{\text{Br}} = \emptyset$.

So, the variety X is a counterexample to the Hasse principle, and its failure of the Hasse principle is explained by the Brauer-Manin obstruction.

We will show $X_L(\mathbb{A}_L)^{\text{Br}} \neq \emptyset$. By Proposition 5.0.2, the map $\alpha_L^*: \text{Br}(B_L) \rightarrow \text{Br}(X_L)$ is an isomorphism, so $\beta_L^*: \text{Br}(C_L) \rightarrow \text{Br}(X_L)$ is an isomorphism. By the functoriality of Brauer-Manin pairing, the set $X_L(\mathbb{A}_L)^{\text{Br}}$ contains $\bigsqcup_{Q \in C(L) \setminus C(K)} \beta^{-1}(Q)(\mathbb{A}_L) \cong V_0(\mathbb{A}_L) \times (C(L) \setminus C(K))$, which is nonempty.

We will show $X(L) = \emptyset$. By the assumption that the degree $[L : K]$ is odd, and v_1 splitting completely in L , the number $\sharp S_{1L}$ is odd. By Proposition 3.3.1 and the global reciprocity law explained in Remark 3.3.2, the set $V_0(L) = \emptyset$. By Proposition 3.1.1, the set $V_\infty(\mathbb{A}_L) = \emptyset$. Since each L -rational fiber of β is isomorphic to V_{0L} or $V_{\infty L}$, the set $X(L) = \emptyset$.

So, the variety X_L is a counterexample to the Hasse principle, but its failure of the Hasse principle cannot be explained by the Brauer-Manin obstruction. \square

In the case when both fields K and L have real places, making use of these real place information, and assuming the Stoll's conjecture, we have the following theorem to give a negative answer to Question 1.2.2

Theorem 6.3.2. For any number field K having a real place, and any nontrivial field extension L/K having a real place, assuming that Conjecture 4.0.1 holds over K , there exists a Châtelet surface bundle over a curve: $X \rightarrow C$ defined over K such that

- X is a counterexample to the Hasse principle, and its failure of the Hasse principle is explained by the Brauer-Manin obstruction,
- X_L is a counterexample to the Hasse principle, but its failure of the Hasse principle cannot be explained by the Brauer-Manin obstruction.

Proof. Firstly, we will construct two Châtelet surfaces. Take a real place v'_0 of L , and let $v_0 \in \infty_K$ be the restriction of v'_0 on K . By Theorem 2.0.2, we can take a finite place $v_1 \in \Omega_K^f \setminus 2_K$ splitting completely in L . For the extension L/K , and $S = \{v_0, v_1\}$, we choose an element a as in Subsubsection 3.0.1. For the trivial extension K/K (respectively the extension L/K), and $S_1 = \{v_0\}$ (respectively $S_2 = \{v_1\}$), by Remark 3.0.7, choosing other parameters for the equation (1) as in Subsubsection 3.1.1, we have a Châtelet surface V_0 (respectively V_∞) defined by $y^2 - az^2 = P_0(x)$ (respectively by $y^2 - az^2 = P_\infty(x)$) over K having the properties of Proposition 3.1.1. By Remark 3.1.2, the polynomial $P_\infty(x)$ is irreducible over K_{v_1} . If the polynomials $P_0(x)$ and $P_\infty(x)$ are not coprime, then $P_0(x) = \lambda P_\infty(x)$ for some $\lambda \in K^\times$. By Remark 3.1.2, we choose another $P_\infty(x)'$ to replace $P_\infty(x)$ so that the new polynomial $P_\infty(x)$ is prime to $P_0(x)$.

Secondly, we will construct a Châtelet surface bundle over a curve. Let $\tilde{P}_\infty(x_0, x_1)$ and $\tilde{P}_0(x_0, x_1)$ be the homogenizations of P_∞ and P_0 . Let $(u_0 : u_1) \times (x_0 : x_1)$ be the coordinates of $\mathbb{P}^1 \times \mathbb{P}^1$, and let $s' = u_0^2 \tilde{P}_\infty(x_0, x_1) + u_1^2 \tilde{P}_0(x_0, x_1) \in \Gamma(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(1, 2)^{\otimes 2})$. For $P_0(x)$ and $P_\infty(x)$ are coprime in $K[x]$, by Jacobian criterion, the locus Z' defined by $s' = 0$ in $\mathbb{P}^1 \times \mathbb{P}^1$ is smooth. Then the branch locus of the composition $Z' \hookrightarrow \mathbb{P}^1 \times \mathbb{P}^1 \xrightarrow{pr_1} \mathbb{P}^1$, denoted by R , is finite over K . By the assumptions that Conjecture 4.0.1 holds over K , and that the field L has a real place, using Lemma 4.0.4, we can take a curve C defined over K such that the triple (C, K, L) is of type II. By Lemma 6.1.3, we can choose a K -morphism $\gamma: C \rightarrow \mathbb{P}^1$ such that $\gamma(C(L) \setminus C(K)) = \{0\} \subset \mathbb{P}^1(K)$, $\gamma(C(K)) = \{\infty\} \subset \mathbb{P}^1(K)$, and that γ is étale over R . Let $B = C \times \mathbb{P}^1$. Let $\mathcal{L} = (\gamma, id)^* \mathcal{O}(1, 2)$, and let $s = (\gamma, id)^*(s') \in \Gamma(B, \mathcal{L}^{\otimes 2})$. By the same argument as in the proof of Theorem 6.2.1, the locus Z defined by $s = 0$ in B is smooth, projective, and geometrically connected; the Brauer group $\text{Br}(\overline{B}) = 0$. Let X be the zero locus of $(1, -a, -s) \in \Gamma(B, \mathcal{O}_B \oplus \mathcal{O}_B \oplus \mathcal{L}^{\otimes 2}) \subset \Gamma(B, \text{Sym}^2 \mathcal{E})$ in the projective space bundle $\text{Proj}(\mathcal{E})$ with the natural projection $\alpha: X \rightarrow B$. By the same argument as in the proof of Theorem 6.3.1, the variety X is smooth, projective, and geometrically connected. Let $\beta: X \rightarrow C$ be the composition of α and pr_1 . Then β is a Châtelet surface bundle over the curve C .

At last, we will check that X has the properties.

We will show $X(\mathbb{A}_K) \neq \emptyset$. For any $P \in C(K)$, the fiber $\beta^{-1}(P) \cong V_\infty$. By Proposition 3.1.1, the set $V_\infty(\mathbb{A}_K^{\{v_1\}}) \neq \emptyset$. So $X(\mathbb{A}_K^{\{v_1\}}) \neq \emptyset$. For v_1 splits completely in L , take a place $v'_1 \in \Omega_L^f$ above v_1 , i.e. $v'_1 | v_1$ in L . Then $K_{v_1} = L_{v'_1}$. By Proposition 3.1.1, the set $V_0(K_{v_1}) \neq \emptyset$. Take a point $Q \in C(L) \setminus C(K)$, then the fiber $\beta^{-1}(Q) \cong V_0 L$. We have $X(K_{v_1}) = X_L(L_{v'_1}) \supset \beta^{-1}(Q)(L_{v'_1}) \cong V_0(L_{v'_1}) \neq \emptyset$. So $X(\mathbb{A}_K) \neq \emptyset$.

By the same argument as in the proof of Theorem 6.3.1, the set $X(\mathbb{A}_K)^{\text{Br}} = \emptyset$. So, the variety X is a counterexample to the Hasse principle, and its failure of the Hasse principle is explained by the Brauer-Manin obstruction.

We will show $X_L(\mathbb{A}_L)^{\text{Br}} \neq \emptyset$. By Proposition 3.1.1, the set $V_0(\mathbb{A}_K^{\{v_0\}}) \neq \emptyset$ and $V_\infty(K_{v_0}) \neq \emptyset$. By Proposition 5.0.2, the map $\beta_L^*: \text{Br}(C_L) \rightarrow \text{Br}(X_L)$ is an isomorphism. By our choice, the space $C(K_{v_0}) \cong C(L_{v'_0}) \cong C(\mathbb{R})$ is connected. For any $A \in \text{Br}(C_L)$, since the evaluation of A on $C(L_{v'})$ is locally constant for all $v' \in \Omega_L$, it is constant on $C(K_{v_0})$, so it is constant on $C(L_{v'})$ for all $v' \in S_{1L}$. Take points $P \in C(K)$ and $Q \in C(L) \setminus C(K)$. By the

functoriality of Brauer-Manin pairing and isomorphism of $\beta_L^*: \text{Br}(C_L) \rightarrow \text{Br}(X_L)$, the set $X_L(\mathbb{A}_L)^{\text{Br}} \supset \beta^{-1}(Q)(\mathbb{A}_L^{S_{1L}}) \times \prod_{v' \in S_{1L}} \beta^{-1}(P)(L_{v'}) \cong V_0(\mathbb{A}_L^{S_{1L}}) \times \prod_{v' \in S_{1L}} V_\infty(L_{v'}) \neq \emptyset$.

We will show $X(L) = \emptyset$. By Proposition 3.1.1, the set $V_0(K_{v_0}) = \emptyset$, so the set $V_0(L_{v'_0}) = V_0(K_{v_0}) = \emptyset$. By Proposition 3.1.1, the set $V_\infty(\mathbb{A}_L) = \emptyset$. Since each L -rational fiber of β is isomorphic to V_{0L} or $V_{\infty L}$, the set $X(L) = \emptyset$.

So, the variety X is a counterexample to the Hasse principle, but its failure of the Hasse principle cannot be explained by the Brauer-Manin obstruction. \square

7. EXPLICIT UNCONDITIONAL EXAMPLES

Firstly, we will give explicit examples without assuming Conjecture 4.0.1 for Theorem 6.2.1, Theorem 6.3.1 and Theorem 6.3.2. Secondly, when $K = \mathbb{Q}$ and $L = \mathbb{Q}(i)$, besides cases of Theorem 6.3.1 and Theorem 6.3.2, we construct an explicit Châtelet surface bundle over a curve in Subsection 7.4 to give a negative answer to Question 1.2.2.

7.1. An explicit unconditional example for Theorem 6.2.1. In the subsection, let $K = \mathbb{Q}$ and $L = \mathbb{Q}(\sqrt{3})$. We will construct an explicit Châtelet surface bundle over a curve having properties of Theorem 6.2.1.

7.1.1. Choosing an elliptic curve. Let E be an elliptic curve defined over \mathbb{Q} by a homogeneous equation:

$$w_1^2 w_2 = w_0^3 - 16w_2^3$$

with homogeneous coordinates $(w_0 : w_1 : w_2) \in \mathbb{P}^2$. This is an elliptic curve with complex multiplication. Its quadratic twist $E^{(3)}$ is isomorphic to an elliptic curve defined by a homogeneous equation: $w_1^2 w_2 = w_0^3 - 432w_2^3$ with homogeneous coordinates $(w_0 : w_1 : w_2) \in \mathbb{P}^2$. These elliptic curves E and $E^{(3)}$ defined over \mathbb{Q} , are of analytic rank 0. Then the Tate-Shafarevich group $\text{III}(E, \mathbb{Q})$ is finite, so E satisfies weak approximation with Brauer-Manin obstruction off ∞_K . The Mordell-Weil groups $E(K)$ and $E^{(3)}(K)$ are finite, so $E(L)$ is finite. Indeed, the Mordell-Weil groups $E(K) = \{(0 : 1 : 0)\}$ and $E(L) = \{(4 : \pm 4\sqrt{3} : 1), (0 : 1 : 0)\}$. So the triple (E, K, L) is of type I .

7.1.2. Choosing a dominant morphism. Let $\mathbb{P}^2 \setminus \{(0 : 1 : 0)\} \rightarrow \mathbb{P}^1$ be a morphism over \mathbb{Q} given by $(w_0 : w_1 : w_2) \mapsto (w_0 - 4w_2 : w_2)$. Composite with the natural inclusion $E \setminus \{(0 : 1 : 0)\} \hookrightarrow \mathbb{P}^2 \setminus \{(0 : 1 : 0)\}$. We get a morphism $E \setminus \{(0 : 1 : 0)\} \rightarrow \mathbb{P}^1$, which can be extended to a dominant morphism $\gamma: E \rightarrow \mathbb{P}^1$ of degree 2. The morphism γ maps $E(K)$ to $\{\infty\} = \{(1 : 0)\}$, and maps $(4 : \pm 4\sqrt{3} : 1)$ to 0 point: $(0 : 1)$. By Bézout's Theorem [Har97, Chapter I. Corollary 7.8] or Hurwitz's Theorem [Har97, Chapter IV. Corollary 2.4], the branch locus of γ is $\{(1 : 0), (2\sqrt[3]{2} - 4 : 1), (2\sqrt[3]{2}e^{2\pi i/3} - 4 : 1), (2\sqrt[3]{2}e^{-2\pi i/3} - 4 : 1)\}$.

7.1.3. Construction of a Châtelet surface bundle. Let $P_\infty(x) = (1 - x^2)(x^2 - 73)$, and let $P_0(x) = (99x^2 + 1)(5428x^2/5329 + 1/5329)$. Notice that these polynomials P_∞ and P_0 are separable. Let V_∞ be the Châtelet surface given by $y^2 - 73z^2 = P_\infty(x)$. As mentioned in Example 3.2.5, let V_0 be the Châtelet surface given by $y^2 - 73z^2 = P_0(x)$. Let $\tilde{P}_\infty(x_0, x_1)$ and $\tilde{P}_0(x_0, x_1)$ be the homogenizations of P_∞ and P_0 . Let $(u_0 : u_1) \times (x_0 : x_1)$ be the coordinates of $\mathbb{P}^1 \times \mathbb{P}^1$, and let $s' = u_0^2 \tilde{P}_\infty(x_0, x_1) + u_1^2 \tilde{P}_0(x_0, x_1) \in \Gamma(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(1, 2)^{\otimes 2})$. For $P_0(x)$ and $P_\infty(x)$ are coprime in $K[x]$, by Jacobian criterion, the locus Z' defined by $s' = 0$ in $\mathbb{P}^1 \times \mathbb{P}^1$ is smooth. Then the branch locus of the composition $Z' \hookrightarrow \mathbb{P}^1 \times \mathbb{P}^1 \xrightarrow{\text{pr}_1} \mathbb{P}^1$, denoted by R , is finite, and contained in $\mathbb{P}^1 \setminus \{(1 : 0)\}$. Let $B = E \times \mathbb{P}^1$. Let $\mathcal{L} = (\gamma, id)^* \mathcal{O}(1, 2)$, and let $s = (\gamma, id)^*(s') \in \Gamma(B, \mathcal{L}^{\otimes 2})$. With these notations, we have the following lemma.

Lemma 7.1.1. The curve Z defined by $s = 0$ in B is smooth, projective, and geometrically connected.

Proof. For smoothness of Z , we need to check that the branch locus R does not intersect with the branch locus of $\gamma: E \rightarrow \mathbb{P}^1$. For R is contained in $\mathbb{P}^1 \setminus \{(1 : 0)\}$, we can assume the homogeneous coordinate $u_1 = 1$, then the point in R satisfies one of the following equations: $5329u_0^2 - 537372 = 0$, $389017u_0^2 - 1 = 0$, $27625536u_0^4 + 157730624u_0^2 + 5329 = 0$. The

polynomials of these equations are irreducible over \mathbb{Q} . By comparing the degree $[\mathbb{Q}(u_0) : \mathbb{Q}]$ with the branch locus of γ , we get the conclusion that these two branch loci do not intersect. The same argument as in the proof of Theorem 6.2.1, the locus Z defined by $s = 0$ in B is geometrically connected. So it is smooth, projective, and geometrically connected. \square

Let X be the zero locus of $(1, -a, -s) \in \Gamma(B, \mathcal{O}_B \oplus \mathcal{O}_B \oplus \mathcal{L}^{\otimes 2}) \subset \Gamma(B, \text{Sym}^2 \mathcal{E})$ in the projective space bundle $\text{Proj}(\mathcal{E})$ with the natural projection $\alpha: X \rightarrow B$. By the same argument as in the proof of Theorem 6.3.1, the variety X is smooth, projective, and geometrically connected. Let $\beta: X \rightarrow E$ be the composition of α and pr_1 . Then it is a Châtelet surface bundle over the curve E . For this X , we have the following proposition.

Proposition 7.1.2. For $K = \mathbb{Q}$ and $L = \mathbb{Q}(\sqrt{3})$, the 3-fold X has the following properties.

- X has a K -rational point, and satisfies weak approximation with Brauer-Manin obstruction off ∞_K .
- X_L does not satisfy weak approximation with Brauer-Manin obstruction off ∞_L .

Proof. This is the same as in the proof of Theorem 6.2.1. \square

The 3-fold X that we constructed, has an affine open subvariety defined by the following equations, which is a closed subvariety of \mathbb{A}^5 with affine coordinates (x, y, z, x', y') .

$$\begin{cases} y^2 - 73z^2 = (1 - x^2)(x^2 - 73)(x' - 4)^2 + (99x^2 + 1)(5428x^2/5329 + 1/5329) \\ y'^2 = x'^3 - 16 \end{cases}$$

7.2. An explicit unconditional example for Theorem 6.3.1. In the subsection, let $K = \mathbb{Q}$, and let ζ_7 be a primitive 7-th root of unity. Let $\alpha = \zeta_7 + \zeta_7^{-1}$ with the minimal polynomial $x^3 + x^2 - 2x - 1$. Let $L = \mathbb{Q}(\alpha)$. Then the degree $[L : K] = 3$. We will construct an explicit Châtelet surface bundle over a curve having properties of Theorem 6.3.1.

7.2.1. Choosing an elliptic curve. Let E be an elliptic curve defined over \mathbb{Q} by a homogeneous equation:

$$w_1^2 w_2 = w_0^3 - 343w_0 w_2^2 - 2401w_2^3$$

with homogeneous coordinates $(w_0 : w_1 : w_2) \in \mathbb{P}^2$. This elliptic curve defined over \mathbb{Q} , is of analytic rank 0, so it satisfies weak approximation with Brauer-Manin obstruction off ∞_K . By computer calculation, we have the Mordell-Weil groups $E(K) = \{(0 : 1 : 0)\}$ and $E(L) = \{(7\alpha^2 + 14\alpha - 7 : 0 : 1), (7\alpha^2 - 7\alpha - 14 : 0 : 1), (-14\alpha^2 - 7\alpha + 21 : 0 : 1), (0 : 1 : 0)\}$. So the triple (E, K, L) is of type I .

7.2.2. Choosing a dominant morphism. Let $\mathbb{P}^2 \setminus \{(1 : 0 : 0)\} \rightarrow \mathbb{P}^1$ be a morphism over \mathbb{Q} given by $(w_0 : w_1 : w_2) \mapsto (w_1 : w_2)$. Composite with the natural inclusion $E \hookrightarrow \mathbb{P}^2 \setminus \{(1 : 0 : 0)\}$. We get a morphism $\gamma: E \rightarrow \mathbb{P}^1$, which is a dominant morphism of degree 3. The morphism γ maps $E(K)$ to $\{\infty\} = \{(1 : 0)\}$, and maps $E(L) \setminus E(K)$ to $\{0\} = \{(0 : 1)\}$. By Bézout's Theorem [Har97, Chapter I. Corollary 7.8] or Hurwitz's Theorem [Har97, Chapter IV. Corollary 2.4], the branch locus of γ is $\{(1 : 0)\} \cup \{(u_0 : 1) \mid 27u_0^4 + 129654u_0^2 - 5764801 = 0\}$.

7.2.3. Construction of a Châtelet surface bundle. Let $P_\infty(x) = 14(x^4 - 89726)$, and let $P_0(x) = (x^2 - 878755181)(5x^2 - 4393775906)$. Notice that these polynomials P_∞ and P_0 are separable. As mentioned in Example 3.1.3 and Example 3.3.5, let V_∞ be the Châtelet surface given by $y^2 - 377z^2 = P_\infty(x)$, and let V_0 be the Châtelet surface given by $y^2 - 377z^2 = P_0(x)$. Let $\tilde{P}_\infty(x_0, x_1)$ and $\tilde{P}_0(x_0, x_1)$ be the homogenizations of P_∞ and P_0 . Let $(u_0 : u_1) \times (x_0 : x_1)$ be the coordinates of $\mathbb{P}^1 \times \mathbb{P}^1$, and let $s' = u_0^2 \tilde{P}_\infty(x_0, x_1) + u_1^2 \tilde{P}_0(x_0, x_1) \in \Gamma(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(1, 2)^{\otimes 2})$. For $P_0(x)$ and $P_\infty(x)$ are coprime in $K[x]$, by Jacobian criterion, the locus Z' defined by $s' = 0$ in $\mathbb{P}^1 \times \mathbb{P}^1$ is smooth. Then the branch locus of the composition $Z' \hookrightarrow \mathbb{P}^1 \times \mathbb{P}^1 \xrightarrow{pr_1} \mathbb{P}^1$, denoted by R , is finite and contained in $\mathbb{P}^1 \setminus \{(1 : 0)\}$. Let $B = E \times \mathbb{P}^1$. Let $\mathcal{L} = (\gamma, id)^* \mathcal{O}(1, 2)$, and let $s = (\gamma, id)^*(s') \in \Gamma(B, \mathcal{L}^{\otimes 2})$. With these notations, we have the following lemma.

Lemma 7.2.1. The curve Z defined by $s = 0$ in B is smooth, projective, and geometrically connected.

Proof. For smoothness of Z , we need to check that the branch locus R does not intersect with the branch locus of $\gamma: E \rightarrow \mathbb{P}^1$. For R is contained in $\mathbb{P}^1 \setminus \{(1:0)\}$, we can assume the homogeneous coordinate $u_1 = 1$, then the point in R satisfies one of the following equations: $14u_0^2 + 5 = 0$, $44863u_0^2 - 137894762198231040 = 0$, $70345184u_0^4 - 216218987126801139936u_0^2 + 1 = 0$. The polynomials of these equations are irreducible over \mathbb{Q} . By comparing these irreducible polynomials with the branch locus of γ , we get the conclusion that these two branch loci do not intersect. The same argument as in the proof of Theorem 6.2.1, the locus Z defined by $s = 0$ in B is geometrically connected. So it is smooth, projective, and geometrically connected. \square

Let X be the zero locus of $(1, -a, -s) \in \Gamma(B, \mathcal{O}_B \oplus \mathcal{O}_B \oplus \mathcal{L}^{\otimes 2}) \subset \Gamma(B, \text{Sym}^2 \mathcal{E})$ in the projective space bundle $\text{Proj}(\mathcal{E})$ with the natural projection $\alpha: X \rightarrow B$. By the same argument as in the proof of Theorem 6.3.1, the variety X is smooth, projective, and geometrically connected. Let $\beta: X \rightarrow E$ be the composition of α and pr_1 . Then it is a Châtelet surface bundle over the curve E . For this X , we have the following proposition.

Proposition 7.2.2. For $K = \mathbb{Q}$ and $L = \mathbb{Q}(\zeta_7 + \zeta_7^{-1})$, the 3-fold X has the following properties.

- X is a counterexample to the Hasse principle, and its failure of the Hasse principle is explained by the Brauer-Manin obstruction.
- X_L is a counterexample to the Hasse principle, but its failure of the Hasse principle cannot be explained by the Brauer-Manin obstruction.

Proof. This is the same as in the proof of Theorem 6.3.1. \square

The 3-fold X that we constructed, has an affine open subvariety defined by the following equations, which is closed subvariety of \mathbb{A}^5 with affine coordinates (x, y, z, x', y') .

$$\begin{cases} y^2 - 377z^2 = 14(x^4 - 89726)y'^2 + (x^2 - 878755181)(5x^2 - 4393775906) \\ y'^2 = x'^3 - 343x' - 2401 \end{cases}$$

7.3. An explicit unconditional example for Theorem 6.3.2. In the subsection, let $K = \mathbb{Q}$ and $L = \mathbb{Q}(\sqrt{3})$. We will construct an explicit Châtelet surface bundle over a curve having properties of Theorem 6.3.2.

7.3.1. Choosing an elliptic curve and a dominant morphism. Let E and $\gamma: E \rightarrow \mathbb{P}^1$ be the same as in Subsection 7.1. For $E(\mathbb{R})$ is connected, the triple (E, K, L) is of type II.

7.3.2. Construction of a Châtelet surface bundle. Let $P_\infty(x) = 5(x^4 + 805)$, and let $P_0(x) = -5(x^4 + 115)$. Notice that these polynomials P_∞ and P_0 are irreducible. As mentioned in Example 3.1.4 and Example 3.1.5, let V_∞ be the Châtelet surface given by $y^2 + 23z^2 = P_\infty(x)$, and let V_0 be the Châtelet surface given by $y^2 + 23z^2 = P_0(x)$. Let $\tilde{P}_\infty(x_0, x_1)$ and $\tilde{P}_0(x_0, x_1)$ be the homogenizations of P_∞ and P_0 . Let $(u_0 : u_1) \times (x_0 : x_1)$ be the coordinates of $\mathbb{P}^1 \times \mathbb{P}^1$, and let $s' = u_0^2 \tilde{P}_\infty(x_0, x_1) + u_1^2 \tilde{P}_0(x_0, x_1) \in \Gamma(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(1, 2)^{\otimes 2})$. For $P_0(x)$ and $P_\infty(x)$ are coprime in $K[x]$, by Jacobian criterion, the locus Z' defined by $s' = 0$ in $\mathbb{P}^1 \times \mathbb{P}^1$ is smooth. Then the branch locus of the composition $Z' \hookrightarrow \mathbb{P}^1 \times \mathbb{P}^1 \xrightarrow{pr_1} \mathbb{P}^1$, denoted by R , is finite, and contained in $\mathbb{P}^1 \setminus \{(1:0)\}$. Let $B = E \times \mathbb{P}^1$. Let $\mathcal{L} = (\gamma, id)^* \mathcal{O}(1, 2)$, and let $s = (\gamma, id)^*(s') \in \Gamma(B, \mathcal{L}^{\otimes 2})$. With these notations, we have the following lemma.

Lemma 7.3.1. The curve Z defined by $s = 0$ in B is smooth, projective, and geometrically connected.

Proof. For smoothness of Z , we need to check that the branch locus R does not intersect with the branch locus of $\gamma: E \rightarrow \mathbb{P}^1$. For R is contained in $\mathbb{P}^1 \setminus \{(1:0)\}$, we can assume the homogeneous coordinate $u_1 = 1$, then the point in R satisfies one of the following equations:

$u_0^2 - 1 = 0$, $7u_0^2 - 1 = 0$. By comparing this locus with the branch locus of γ , these two branch loci do not intersect. The same argument as in the proof of Theorem 6.2.1, the locus Z defined by $s = 0$ in B is geometrically connected. So it is smooth, projective, and geometrically connected. \square

Let X be the zero locus of $(1, -a, -s) \in \Gamma(B, \mathcal{O}_B \oplus \mathcal{O}_B \oplus \mathcal{L}^{\otimes 2}) \subset \Gamma(B, \text{Sym}^2 \mathcal{E})$ in the projective space bundle $\text{Proj}(\mathcal{E})$ with the natural projection $\alpha: X \rightarrow B$. By the same argument as in the proof of Theorem 6.3.1, the variety X is smooth, projective, and geometrically connected. Let $\beta: X \rightarrow E$ be the composition of α and pr_1 . Then it is a Châtelet surface bundle over the curve E . For this X , we have the following proposition.

Proposition 7.3.2. For $K = \mathbb{Q}$ and $L = \mathbb{Q}(\sqrt{3})$, the 3-fold X has the following properties.

- X is a counterexample to the Hasse principle, and its failure of the Hasse principle is explained by the Brauer-Manin obstruction.
- X_L is a counterexample to the Hasse principle, but its failure of the Hasse principle cannot be explained by the Brauer-Manin obstruction.

Proof. This is the same as in the proof of Theorem 6.3.2. \square

The 3-fold X that we constructed, has an affine open subvariety defined by the following equations, which is closed subvariety of \mathbb{A}^5 with affine coordinates (x, y, z, x', y') .

$$\begin{cases} y^2 + 23z^2 = 5(x^4 + 805)(x' - 4)^2 - 5(x^4 + 115) \\ y'^2 = x'^3 - 16 \end{cases}$$

7.4. Exceptions. For Question 1.2.2, when the degree $[L : K]$ is even and L has no real place, besides cases of Theorem 6.3.1 and Theorem 6.3.2, we can give some unconditional examples, case by case, to give negative answers to Question 1.2.2, although we do not have a uniform way to construct them. In this subsection, we give an explicit example to explain how it works for the case that $K = \mathbb{Q}$ and $L = \mathbb{Q}(i)$.

7.4.1. Choosing an elliptic curve. Let E be an elliptic curve defined over \mathbb{Q} by a homogeneous equation:

$$w_1^2 w_2 = w_0^3 - 16w_2^3$$

with homogeneous coordinates $(w_0 : w_1 : w_2) \in \mathbb{P}^2$. This is an elliptic curve with complex multiplication. Its quadratic twist $E^{(-1)}$ is isomorphic to an elliptic curve defined by a homogeneous equation: $w_1^2 w_2 = w_0^3 + 16w_2^3$ with homogeneous coordinates $(w_0 : w_1 : w_2) \in \mathbb{P}^2$. These elliptic curves E and $E^{(-1)}$ defined over \mathbb{Q} , are of analytic rank 0. Then the Tate-Shafarevich group $\text{III}(E, \mathbb{Q})$ is finite, so E satisfies weak approximation with Brauer-Manin obstruction off ∞_K . The Mordell-Weil groups $E(K)$ and $E^{(-1)}(K)$ are finite, so $E(L)$ is finite. Indeed, the Mordell-Weil group $E(K) = \{(0 : 1 : 0)\}$ and $E(L) = \{(0 : \pm 4i : 1), (0 : 1 : 0)\}$. So the triple (E, K, L) is of type I .

7.4.2. Choosing a dominant morphism. Let $\mathbb{P}^2 \setminus \{(1 : 0 : 0)\} \rightarrow \mathbb{P}^1$ be a morphism over \mathbb{Q} given by $(w_0 : w_1 : w_2) \mapsto (w_1 : 4w_2)$. Composite with the natural inclusion $E \hookrightarrow \mathbb{P}^2 \setminus \{(1 : 0 : 0)\}$, then we get a morphism $\gamma: E \rightarrow \mathbb{P}^1$, which is a dominant morphism of degree 3. The dominant morphism γ maps $E(K)$ to $\{\infty\} = \{(1 : 0)\}$, and maps $(0 : \pm 4i : 1)$ to $(\pm i : 1)$. By Bézout's Theorem [Har97, Chapter I. Corollary 7.8] or Hurwitz's Theorem [Har97, Chapter IV. Corollary 2.4], the branch locus of γ is $\{(1 : 0), (\pm i : 1)\}$.

7.4.3. Properties of Châtelet surfaces. Let $P_\infty(x) = 2(x^4 - 10x^2 + 15)$, $P_0(x) = -2(5x^4 - 39x^2 + 75)$, and let $P_1(x) = P_\infty(x) + iP_0(x)$. Notice that all those polynomials P_∞, P_0, P_1 are separable, and $P_1(x) = -2[x^2 - (5+i)][(-1+5i)x^2 - 15i]$. The two polynomials $x^2 - (5+i)$ and $(-1+5i)x^2 - 15i$ are irreducible over $\mathbb{Q}(i)$ (indeed, they are irreducible over $\mathbb{Q}(i)_3$). Let V_∞ be the Châtelet surface over \mathbb{Q} given by $y^2 + 15z^2 = P_\infty(x)$, and let V_1 be the Châtelet surface over $\mathbb{Q}(i)$ given by $y^2 + 15z^2 = P_1(x)$. With these notations, we have the following lemmas.

Lemma 7.4.1. The Châtelet surface V_∞ given by $y^2 + 15z^2 = 2(x^4 - 10x^2 + 15)$, has a \mathbb{Q}_v -point for all $v \in \Omega_{\mathbb{Q}} \setminus \{5\}$, but no \mathbb{Q}_5 -point.

Proof. Suppose that $v = \infty_{\mathbb{Q}}$. Let $x_0 = 0$. Then $(-15, P_\infty(x_0))_v = (-15, 30)_v = 1$, which implies that the surface V_∞^0 admits a \mathbb{R} -point with $x = 0$.

Suppose that $v = 2$. For $-15 \equiv 1 \pmod{8}$, by Hensel's lemma, the element -15 is a square in \mathbb{Q}_2 . By Remark 3.0.1, the surface V_∞ admits a \mathbb{Q}_2 -point.

Suppose that $v = 3$. Let $x_0 = 2$. Then $(-15, P_\infty(x_0))_v = (-15, -18)_v = (-15, 9)_v(-15, -2)_v = 1$, which implies that the surface V_∞^0 admits a \mathbb{Q}_3 -point with $x = 2$.

Suppose that $v \in \Omega_{\mathbb{Q}}^f \setminus \{2, 3, 5\}$. Take $x_0 \in \mathbb{Q}_v$ such that the valuation $v(x_0) < 0$. Then by Lemma 2.2.2, we have $(-15, P_\infty(x_0))_v = (-15, 2)_v = 1$, which implies that V_∞^0 admits a \mathbb{Q}_v -point with $x = x_0$.

Suppose that $v = 5$. Then $(-15, 2)_v = -1$. Let $x \in \mathbb{Q}_5$. If $v(x) \leq 0$, then by Lemma 2.2.2, we have $(-15, P_\infty(x))_v = (-15, 2x^4)_v = -1$. If $v(x) > 0$, then by Lemma 2.2.2, we have $(-15, P_\infty(x))_v = (-15, 30)_v = -1$. In each case, we have $(-15, P_\infty(x))_v = -1$, which implies that V_∞^0 has no \mathbb{Q}_5 -point. By Remark 3.0.2, we have $V_\infty(\mathbb{Q}_5) = \emptyset$. \square

Lemma 7.4.2. For any $v' \in \Omega_{\mathbb{Q}(i)}$, the Châtelet surface V_1 given by $y^2 + 15z^2 = -2[x^2 - (5+i)][(-1+5i)x^2 - 15i]$, has a $\mathbb{Q}(i)_{v'}$ -point.

Proof. For the only archimedean place is complex, we only need to consider finite places. Suppose that v' is a 2-adic place. For $-15 \in \mathbb{Q}_2^{\times 2}$, by Remark 3.0.1, the surface V_1 admits a $\mathbb{Q}(i)_{v'}$ -point.

Suppose that $v' = 3$. For $-2 \in \mathbb{Q}_3^{\times 2}$, we have $(-15, -2)_{v'} = 1$. By Lemma 2.2.2, we have $(-15, (-4-i)(-1-10i))_{v'} = (-15, (1+i)^2)_{v'}$. Let $x_0 = 1$. Then $(-15, P_1(x_0))_{v'} = (-15, -2(-4-i)(-1-10i))_{v'} = (-15, (1+i)^2)_{v'} = 1$, which implies that V_1^0 admits a $\mathbb{Q}(i)_{v'}$ -point with $x = 1$.

Suppose that $v'|5$, then $\mathbb{Q}(i)_{v'} \cong \mathbb{Q}_5$. By Lemma 2.2.2, we have $(-15, -2(-5+i)(-10-17i))_{v'} = (-15, -34)_{v'}$. Let $x_0 = 1+i$. Then $(-15, P_1(x_0))_{v'} = (-15, -2(-5+i)(-10-17i))_{v'} = (-15, -34)_{v'} = 1$, which implies that V_1^0 admits a $\mathbb{Q}(i)_{v'}$ -point with $x = 1+i$.

Suppose that $v'|13$, then $\mathbb{Q}(i)_{v'} \cong \mathbb{Q}_{13}$. Let $x_0 = 1$. By Lemma 2.2.1, we have $(-15, P_1(x_0))_{v'} = (-15, -2(-4-i)(-1-10i))_{v'} = 1$, which implies that V_1^0 admits a $\mathbb{Q}(i)_{v'}$ -point with $x = 1$.

Suppose that $v' \in \Omega_{\mathbb{Q}(i)}^f \setminus \{2, 3, 5, 13\}$. Take $x_0 \in \mathbb{Q}_{v'}$ such that the valuation $v'(x_0) < 0$. By Lemma 2.2.2, we have $(-15, P_\infty(x_0))_{v'} = (-15, -2(-1+5i)x_0^4)_{v'}$. By Lemma 2.2.1, we have $(-15, -2(-1+5i))_{v'} = 1$, so $(-15, P_\infty(x_0))_{v'} = 1$, which implies that V_1^0 admits a $\mathbb{Q}(i)_{v'}$ -point with $x = x_0$. \square

For $P_1(x)$ is a product of two degree-2 irreducible factors, according to [Sko01, Proposition 7.1.1], the Brauer group $\text{Br}(V_1)/\text{Br}(\mathbb{Q}(i)) \cong \mathbb{Z}/2\mathbb{Z}$. Furthermore, by Proposition 7.1.2 in loc. cit, we take the quaternion algebra $A = (-15, (-1+5i)x^2 - 15i) \in \text{Br}(V_1)$ as a generator element of this group. Then we have the equality $A = (-15, (-1+5i)x^2 - 15i) = (-15, -2(x^2 - (5+i)))$ in $\text{Br}(V_1)$. With these notations, we have the following lemmas.

Lemma 7.4.3. For any $v' \in \Omega_{\mathbb{Q}(i)}$ and any $P_{v'} \in V_1(\mathbb{Q}(i)_{v'})$,

$$\text{inv}_{v'}(A(P_{v'})) = \begin{cases} 0 & \text{if } v' \neq 3 \\ 1/2 & \text{if } v' = 3 \end{cases}$$

Proof. By Remark 3.0.2, it suffices to compute the local invariant $\text{inv}_{v'}(A(P_{v'}))$ for all $P_{v'} \in V_1^0(\mathbb{Q}(i)_{v'})$.

Suppose that v' is an archimedean place or a 2-adic place or a 181-adic place. Then $-15 \in \mathbb{Q}(i)_{v'}^{\times 2}$, so $\text{inv}_{v'}(A(P_{v'})) = 0$ for all $P_{v'} \in V_1(\mathbb{Q}(i)_{v'})$.

Suppose that $v'|5$. Let $x \in \mathbb{Q}(i)_{v'}$. If $v'(x) \leq 0$, then by Lemma 2.2.2, we have $(-15, (-1+5i)x^2 - 15i)_{v'} = (-15, -x^2)_{v'} = 1$. If $v'(x) > 0$, then by Lemma 2.2.2, we have $(-15, -2(x^2 - (5+i)))_{v'} = (-15, 2i)_{v'} = 1$. So $\text{inv}_{v'}(A(P_{v'})) = 0$ for all $P_{v'} \in V_1^0(\mathbb{Q}(i)_{v'})$.

Suppose that $v' \in \Omega_{\mathbb{Q}(i)}^f \setminus \{3, \text{all 2-adic, 5-adic and 181-adic places}\}$. Take an arbitrary $P_{v'} \in V_1^0(\mathbb{Q}(i)_{v'})$. If $\text{inv}_{v'}(A(P_{v'})) = 1/2$, then $(-15, (-1+5i)x^2 - 15i)_{v'} = -1 = (-15, -2(x^2 -$

$(5+i))_{v'}$ at $P_{v'}$. By Lemma 2.2.1, the first and last equalities imply that $v'((-1+5i)x^2 - 15i)$ and $v'(x^2 - (5+i))$ are odd, so they are positive. Hence $v'((-1+5i)x^2 - 15i - (-1+5i)(x^2 - (5+i))) = v'(-10+9i) > 0$. But $v' \nmid 181$, which is a contradiction. So $\text{inv}_{v'}(A(P_{v'})) = 0$.

Suppose that $v' = 3$. Take an arbitrary $P_{v'} \in V_1^0(\mathbb{Q}(i)_{v'})$. If $\text{inv}_{v'}(A(P_{v'})) = 1$, then $(-15, (-1+5i)x^2 - 15i)_{v'} = 1 = (-15, -2(x^2 - (5+i)))_{v'}$ at $P_{v'}$. For $(-15, -1+5i)_{v'} = -1$, the first equality implies that $v'(x) > 0$. Then by Lemma 2.2.2, we have $(-15, x^2 - (5+i))_{v'} = (-15, -(5+i))_{v'} = -1$, so $(-15, -2(x^2 - (5+i)))_{v'} = (-15, -2)_{v'}(-15, -5-i)_{v'} = -1$, which is a contradiction. So $\text{inv}_{v'}(A(P_{v'})) = 1/2$. \square

Lemma 7.4.4. The Châtelet surface V_1 has no $\mathbb{Q}(i)$ -rational point.

Proof. If there exists $\mathbb{Q}(i)$ -rational point P , by the global reciprocity law $\sum_{v \in \Omega_{\mathbb{Q}(i)}} \text{inv}_v(A(P)) = 0$ in \mathbb{Q}/\mathbb{Z} . But from Lemma 7.4.3, this sum is $1/2$, which is nonzero in \mathbb{Q}/\mathbb{Z} . So V_1 has no $\mathbb{Q}(i)$ -rational point. \square

7.4.4. *Construction of a Châtelet surface bundle.* Let $\tilde{P}_\infty(x_0, x_1)$ and $\tilde{P}_0(x_0, x_1)$ be the homogenizations of P_∞ and P_0 . Let $(u_0 : u_1) \times (x_0 : x_1)$ be the coordinates of $\mathbb{P}^1 \times \mathbb{P}^1$, and let $s' = (u_0^2 + 2u_1^2)\tilde{P}_\infty(x_0, x_1) + u_0u_1\tilde{P}_0(x_0, x_1) \in \Gamma(\mathbb{P}^1 \times \mathbb{P}^1, \mathcal{O}(1, 2)^{\otimes 2})$. By Jacobian criterion, the locus Z' defined by $s' = 0$ in $\mathbb{P}^1 \times \mathbb{P}^1$ is smooth. Then the branch locus of the composition $Z' \hookrightarrow \mathbb{P}^1 \times \mathbb{P}^1 \xrightarrow{pr_1} \mathbb{P}^1$ is finite, and contained in $\mathbb{P}^1 \setminus \{(1 : 0), (\pm i, 1)\}$. Let $B = E \times \mathbb{P}^1$. Let $\mathcal{L} = (\gamma, id)^*\mathcal{O}(1, 2)$, and let $s = (\gamma, id)^*(s') \in \Gamma(B, \mathcal{L}^{\otimes 2})$. With these notations, we have the following lemma.

Lemma 7.4.5. The curve Z defined by $s = 0$ in B is smooth, projective, and geometrically connected.

Proof. For the branch locus of the composition $Z' \hookrightarrow \mathbb{P}^1 \times \mathbb{P}^1 \xrightarrow{pr_1} \mathbb{P}^1$ is contained in $\mathbb{P}^1 \setminus \{(1 : 0), (\pm i, 1)\}$, and the branch locus of $\gamma: E \rightarrow \mathbb{P}^1$ is $\{(1 : 0), (\pm i, 1)\}$, they do not intersect, which implies the smoothness of Z . The same argument as in the proof of Theorem 6.2.1, the locus Z defined by $s = 0$ in B is geometrically connected. So it is smooth, projective, and geometrically connected. \square

Let X be the zero locus of $(1, -a, -s) \in \Gamma(B, \mathcal{O}_B \oplus \mathcal{O}_B \oplus \mathcal{L}^{\otimes 2}) \subset \Gamma(B, \text{Sym}^2 \mathcal{E})$ in the projective space bundle $\text{Proj}(\mathcal{E})$ with the natural projection $\alpha: X \rightarrow B$. By the same argument as in the proof of Theorem 6.3.1, the variety X is smooth, projective, and geometrically connected. Let $\beta: X \rightarrow E$ be the composition of α and pr_1 . Then it is a Châtelet surface bundle over the curve E . For this X , we have the following proposition.

Proposition 7.4.6. For $K = \mathbb{Q}$ and $L = \mathbb{Q}(i)$, the 3-fold X has the following properties.

- X is a counterexample to the Hasse principle, and its failure of the Hasse principle is explained by the Brauer-Manin obstruction.
- X_L is a counterexample to the Hasse principle, but its failure of the Hasse principle cannot be explained by the Brauer-Manin obstruction.

Proof. Let σ be the generator element of Galois group $\text{Gal}(L/K)$. We will show $X(\mathbb{A}_K) \neq \emptyset$. By our construction, each K -rational fiber of β is isomorphic to V_∞ . By Lemma 7.4.1, the set $V_\infty(\mathbb{A}_K^{\{5\}}) \neq \emptyset$. So $X(\mathbb{A}_K^{\{5\}}) \neq \emptyset$. For 5 splits completely in L , take a place $v' \in \Omega_L$ above 5, i.e. $v' \mid 5$ in L . Then $\mathbb{Q}_5 \cong L_{v'}$. By Lemma 7.4.2, the set $V_1(\mathbb{A}_L) \neq \emptyset$. Since $V_1 \sqcup \sigma(V_1) \cong \bigsqcup_{P \in E(L) \setminus E(K)} \beta^{-1}(P) \subset X_L$, the set $X(\mathbb{Q}_5) = X(L_{v'}) \supset V_1(L_{v'}) \neq \emptyset$. So $X(\mathbb{A}_K) \neq \emptyset$.

We will show $X(\mathbb{A}_K)^{\text{Br}} = \emptyset$. For $E(K) = pr^{\infty\kappa}(C(\mathbb{A}_K)^{\text{Br}})$, the functoriality of Brauer-Manin pairing implies $pr^{\infty\kappa}(X(\mathbb{A}_K)^{\text{Br}}) \subset \bigcup_{P \in E(K)} \beta^{-1}(P)(\mathbb{A}_K^{\infty\kappa})$. By Lemma 7.4.1, the set $V_\infty(\mathbb{Q}_5) = \emptyset$, so $pr^{\infty\kappa}(X(\mathbb{A}_K)^{\text{Br}}) \subset \bigcup_{P \in E(K)} \beta^{-1}(P)(\mathbb{A}_K^{\infty\kappa}) \cong V_\infty(\mathbb{A}_K^{\infty\kappa}) = \emptyset$, which implies that $X(\mathbb{A}_K)^{\text{Br}} = \emptyset$.

So, the variety X is a counterexample to the Hasse principle, and its failure of the Hasse principle is explained by the Brauer-Manin obstruction.

We will show $X_L(\mathbb{A}_L)^{\text{Br}} \neq \emptyset$. By Proposition 5.0.2, the map $\alpha_L^*: \text{Br}(C_L) \rightarrow \text{Br}(X_L)$ is an isomorphism. By the functoriality of Brauer-Manin pairing, the set $X_L(\mathbb{A}_L)^{\text{Br}}$ contains $V_1(\mathbb{A}_L)$, which is nonempty.

We will show $X(L) = \emptyset$. By Lemma 7.4.4, the set $V_1(L) = \emptyset$, so also $\sigma(V_1)(L) = \emptyset$. For 5 splits completely in L , the emptiness of $V_\infty(\mathbb{Q}_5)$ implies $V_\infty(\mathbb{A}_L) = \emptyset$. Since $X(L) \subset \bigsqcup_{P \in E(L)} \beta^{-1}(P)(L) \cong V_\infty(L) \cup V_1(L) \cup \sigma(V_1)(L)$, the set $X(L) = \emptyset$.

So, the variety X_L is a counterexample to the Hasse principle, but its failure of the Hasse principle cannot be explained by the Brauer-Manin obstruction. \square

The 3-fold X that we constructed, has an affine open subvariety defined by the following equations, which is closed subvariety of \mathbb{A}^5 with affine coordinates (x, y, z, x', y') .

$$\begin{cases} y^2 + 15z^2 = (x^4 - 10x^2 + 15)(y'^2 + 32)/8 - (5x^4 - 39x^2 + 75)y'/2 \\ y'^2 = x'^3 - 16 \end{cases}$$

Acknowledgements. The author would like to thank my thesis advisor Y. Liang for proposing the related problems, papers and many fruitful discussions. The author was partially supported by NSFC Grant No. 12071448.

REFERENCES

- [CF67] J. Cassels and A. Fröhlich, *Algebraic number theory*, Academic Press, 1967. $\uparrow 2$
- [CT10] J.-L. Colliot-Thélène, *Zéro-cycles de degré 1 sur les solides de Poonen*, Bull. Soc. Math. France **138** (2010), 249–257 (French). $\uparrow 5$
- [CTPS16] J.-L. Colliot-Thélène, A. Pál, and A. Skorobogatov, *Pathologies of the Brauer-Manin obstruction*, Math. Z. **282** (2016), 799–817. $\uparrow 1.1, 3.0.3$
- [CTSSD87a] J.-L. Colliot-Thélène, J.-J. Sansuc, and S. Swinnerton-Dyer, *Intersections of two quadrics and Châtelet surfaces I*, J. Reine Angew. Math. **373** (1987), 37–107. $\uparrow 1.1, 3.0.3, 3.2.1, 3.3.2$
- [CTSSD87b] ———, *Intersections of two quadrics and Châtelet surfaces II*, J. Reine Angew. Math. **374** (1987), 72–168. $\uparrow 1.1, 3.0.3, 3.2.1, 3.3.2$
- [Gro68] A. Grothendieck, *Le groupe de Brauer III: Exemples et compléments*. In: *Dix exposés sur la cohomologie des schémas*, Advanced Studies in Pure Mathematics, vol. 3, North-Holland, 1968 (French). pp. 88–188. $\uparrow 6.1$
- [GZ86] B. Gross and D. Zagier, *Heegner points and derivatives of L-series*, Invent. Math. **84** (1986), 225–320. $\uparrow 4.0.2$
- [Har97] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics, vol. 52, Springer-Verlag, 1997. $\uparrow 6.2, 7.1.2, 7.2.2, 7.4.2$
- [HS14] Y. Harpaz and A. Skorobogatov, *Singular curves and the étale Brauer-Manin obstruction for surfaces*, Ann. Sci. Éc. Norm. Supér. **47** (2014), 765–778. $\uparrow 1.1$
- [Isk71] V. Iskovskikh, *A counterexample to the Hasse principle for systems of two quadratic forms in five variables*, Mat. Zametki **10** (1971), 253–257. $\uparrow 1.3, 3.3$
- [Kol90] V. Kolyvagin, *Euler systems*. In: *The Grothendieck festschrift II*, Progress in Mathematics, vol. 87, Birkhäuser, 1990. pp. 435–483. $\uparrow 4.0.2$
- [Kol91] ———, *On the Mordell-Weil and the Shafarevich-Tate group of modular elliptic curves*. In: *Proceedings of the international congress of mathematicians*, Vol. I, Springer-Verlag, 1991. pp. 429–436. $\uparrow 4.0.2$
- [Lia18] Y. Liang, *Non-invariance of weak approximation properties under extension of the ground field*, Preprint, arXiv:1805.08851v1 [math.NT] (2018). $\uparrow 1.3, 1.4.1, 3.2, 5, 6.2$
- [Man71] Y. Manin, *Le groupe de Brauer-Grothendieck en géométrie diophantienne*. In: *Actes du Congrès International des Mathématiciens*, Vol. 1, Gauthier-Villars, 1971 (French). pp. 401–411. $\uparrow 1.1$
- [Neu99] J. Neukirch, *Algebraic number theory*, Springer-Verlag, 1999. $\uparrow 2$
- [Poo09] B. Poonen, *Existence of rational points on smooth projective varieties*, J. Eur. Math. Soc. **11** (2009), 529–543. $\uparrow 1.3, 3, 3.2, 3.3$
- [Poo10] ———, *Insufficiency of the Brauer-Manin obstruction applied to étale covers*, Ann. of Math. **171** (2010), 2157–2169. $\uparrow 1.1, 1.4.3, 5, 5.0.2, 6.2$
- [Sch99] V. Scharaschkin, *Local-global problems and the Brauer-Manin obstruction*, Thesis, University of Michigan (1999). $\uparrow 1.1, 4$
- [Ser73] J.-P. Serre, *A course in arithmetic*, Graduate Texts in Mathematics, vol. 7, Springer-Verlag, 1973. $\uparrow 2.2$
- [Ser79] ———, *Local fields*, Graduate Texts in Mathematics, vol. 67, Springer-Verlag, 1979. $\uparrow 2.1$
- [Sil09] J. Silverman, *The arithmetic of elliptic curves*, Graduate Texts in Mathematics, vol. 106, Springer-Verlag, 2009. $\uparrow 4.0.5$
- [Sko01] A. Skorobogatov, *Torsors and rational points*, Cambridge Tracts in Mathematics, vol. 144, Cambridge University Press, 2001. $\uparrow 1.1, 3, 3.0.2, 3.0.3, 3.2.1, 3.3, 3.3.1, 4, 7.4.3$

- [Sko99] ———, *Beyond the Manin obstruction*, *Invent. Math.* **135** (1999), 399–424. ↑1.1
- [Sto07] M. Stoll, *Finite descent obstructions and rational points on curves*, *Algebra Number Theory* **1** (2007), 349–391. ↑1.1, 4, 4.0.1
- [Wan96] L. Wang, *Brauer-Manin obstruction to weak approximation on abelian varieties*, *Israel J. Math.* **94** (1996), 189–200. ↑1.1

UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA, SCHOOL OF MATHEMATICAL SCIENCES, NO.96,
JINZHAI ROAD, BAOHE DISTRICT, HEFEI, ANHUI, 230026. P.R.CHINA.

Email address: wuhan90@mail.ustc.edu.cn