

HOPF FORMS AND HOPF-GALOIS THEORY

TIMOTHY KOHL AND ROBERT UNDERWOOD

ABSTRACT. Let K be a finite field extension of \mathbb{Q} and let N be a finite group with automorphism group $F = \text{Aut}(N)$. R. Haggenmüller and B. Pareigis have shown that there is a bijection

$$\Theta : \text{Gal}(K, F) \rightarrow \text{Hopf}(K[N])$$

from the collection of F -Galois extensions of K to the collection of Hopf forms of the group ring $K[N]$. For $N = C_n$, $n \geq 1$, C_p^m , p prime, $m \geq 1$, and $N = D_3, D_4, Q_8$, we show that $\mathbb{Q}[N]$ admits an absolutely semisimple Hopf form H and find L for which $\Theta(L) = H$. Moreover, if H is the Hopf algebra given by a Hopf-Galois structure on a Galois extension E/K , we show how to construct the preimage of H under Θ assuming certain conditions.

1. INTRODUCTION

Let K be a finite field extension of \mathbb{Q} and let N be a finite group with automorphism group $F = \text{Aut}(N)$. By a result of R. Haggenmüller and B. Pareigis there is a bijection

$$\Theta : \text{Gal}(K, F) \rightarrow \text{Hopf}(K[N])$$

from the collection of F -Galois extensions of K to the collection of Hopf forms of the group ring $K[N]$ [9, Theorem 5]. This paper concerns the map Θ and its inverse Θ^{-1} .

The map Θ can be used to classify Hopf forms of $K[N]$ for certain N . For instance let $N = C_n$ denote the cyclic group of order n . If $n = 2$ then F is trivial and so K is the only F -Galois extension of K . Consequently $\Theta(K) = K[C_2]$ is the only Hopf form of $K[C_2]$. For the cases $n = 3, 4, 6$,

$$F = \text{Aut}(C_n) = \mathbb{Z}_n^* = C_2.$$

The C_2 -Galois extensions of K are completely classified as the quadratic extensions $L = K[x]/(x^2 - b)$, where $b \in K^\times$ [14]. Thus the map Θ yields a complete classification of all Hopf forms of $K[C_n]$ for the cases $n = 3, 4, 6$ [9, Theorem 6].

For $n \geq 2$, two special Hopf forms of $K[C_n]$ can be identified. The first is the trivial Hopf form $K[C_n]$. If L is the trivial F -Galois extension $\text{Map}(F, K)$ of K , then $\Theta(L) = K[C_n]$. The second is $(K[C_n])^*$, the linear dual of $K[C_n]$, which is the “absolutely semisimple” Hopf form of $K[C_n]$. If $L = K[x]/(\Phi_n(x))$, where $\Phi_n(x)$ is the n th cyclotomic polynomial, then L is a \mathbb{Z}_n^* -Galois extension of K and $\Theta(L) = (K[C_n])^*$.

In [12, p. 91], B. Pareigis posed the following problem: for which finite groups N does $K[N]$, $K = \mathbb{Q}$, admit an absolutely semisimple Hopf form? As noted above $\mathbb{Q}[C_n]$ admits an absolutely semisimple Hopf form. We show that $\mathbb{Q}[N]$ admits such a form for the abelian

Date: February 2, 2022.

group C_p^m , $m \geq 1$, and non-abelian groups D_3 , D_4 and Q_8 . In each case we compute the F -Galois extension L of \mathbb{Q} for which $\Theta(L)$ is the absolutely semisimple Hopf form.

There is a natural application of Θ to Hopf-Galois theory. Let (H, \cdot) be a Hopf-Galois structure of type N on the Galois extension of fields E/K . Then H is a Hopf form of $K[N]$ and thus $\Theta(L) = H$ for some F -Galois extension L of K , $F = \text{Aut}(N)$. We show how to construct the F -Galois extension L for which $\Theta(L) = H$ under certain conditions.

2. GALOIS EXTENSIONS AND HOPF FORMS

Let F be a finite group. An F -Galois extension of K is a commutative K -algebra L that satisfies

- (i) F is a subgroup of $\text{Aut}_K(L)$,
- (ii) L is a finitely generated, projective K -module,
- (iii) $F \subseteq \text{End}_K(L)$ is a free generating system over K .

The notion of F -Galois extension generalizes the usual definition of a Galois extension of fields. The K -algebra of maps $\text{Map}(F, K)$ is the *trivial F -Galois extension* of K where the action of F on $\text{Map}(F, K)$ is given as $g(\phi)(h) = \phi(g^{-1}h)$ for $g, h \in F$, $\phi \in \text{Map}(F, K)$. We let $\mathcal{G}al(K, F)$ denote the collection of all F -Galois extensions of K .

B. Pareigis [12, Theorem 4.2] has completely characterized F -Galois extensions.

Theorem 2.1 (Pareigis). *Let F be a finite group. Then L is an F -Galois extension of K if and only if*

$$L = \underbrace{M \times M \times \cdots \times M}_n$$

where M is a U -Galois field extension of K for some subgroup U of F of index n .

Proof. (Sketch) Let L be an F -Galois extension. Then L is a commutative, separable K -algebra and hence

$$L = M_1 \times M_2 \times \cdots \times M_n$$

where each M_i is a separable field extension of K . Let f_1, f_2, \dots, f_n be the minimal orthogonal idempotents. We have $M_i \cong M_j, \forall i, j$, hence

$$L = \underbrace{M \times M \times \cdots \times M}_n,$$

where $M = M_1$. Let U be the stabilizer of f_1 in F . Then M is Galois over K with group U and $[L : U] = n$.

Conversely, let $U \leq F$, $n = [F : U]$, and let M/K be U -Galois. Let $g_1, g_2 \cdots g_n$ be a left transversal for U in F and let $\rho : G \rightarrow S_n$ be defined as $\rho(g)(i) = j$ if $gg_iU = g_jU$. Let

$$L = \underbrace{M \times M \times \cdots \times M}_n$$

with minimal orthogonal idempotents f_1, f_2, \dots, f_n , and let the action of F on L be defined on each component as

$$g(mf_i) = (g_{\rho(g)(i)}^{-1} g g_i)(m) f_{\rho(g)(i)},$$

for $m \in M$, $1 \leq i \leq n$. Then L is an F -Galois extension of K . (For details see [12, Theorem 4.2].) □

Let N be a finite group. Then the group ring $K[N]$ is a K -Hopf algebra. Let L be a faithfully flat K -algebra. An L -Hopf form of $K[N]$ is a K -Hopf algebra H for which

$$L \otimes_K H \cong L \otimes_K K[N] \cong L[N]$$

as L -Hopf algebras. A Hopf form of $K[N]$ is a K -Hopf algebra H for which there exists a faithfully flat K -algebra L with $L \otimes_K H \cong L \otimes_K K[N] \cong L[N]$ as L -Hopf algebras. The trivial Hopf form of $K[N]$ is $K[N]$. Let $\mathcal{H}opf(K[N])$ denote the collection of all Hopf forms of $K[N]$.

R. Haggemüller and B. Pareigis [9, Theorem 5] have classified all Hopf forms of $K[N]$.

Theorem 2.2 (Haggemüller and Pareigis). *Let N be a finite group and let $F = \text{Aut}(N)$. There is a bijection $\Theta : \mathcal{G}al(K, F) \rightarrow \mathcal{H}opf(K[N])$ which associates to each F -Galois extension L of K , the Hopf form $H = \Theta(L)$ of $K[N]$ defined as the fixed ring*

$$H = (L[N])^F,$$

where the action of F on N is through the automorphism group $F = \text{Aut}(N)$ and the action of F on L is the Galois action. The Hopf form H is an L -Hopf form of $K[N]$ with isomorphism $\psi : L \otimes_K H \rightarrow L[N]$ defined as $\psi(x \otimes h) = xh$.

Proposition 2.3. *Let N be a finite group, let $F = \text{Aut}(N)$, and let $L = \text{Map}(F, K)$ denote the trivial F -Galois extension of K . Then $\Theta(L) = (L[N])^F \cong K[N]$.*

Proof. (Sketch) $H = (L[N])^F$ has a K -basis consisting of group-like elements. Hence, $H = K[N']$ for some finite group N' . Since $L \otimes_K H = L \otimes_K K[N'] \cong L[N]$ as Hopf algebras, we conclude that $N' \cong N$. □

We illustrate Proposition 2.3 in the case that $N = C_3 = \langle \sigma \rangle = \{1, \sigma, \sigma^2\}$, the cyclic group of order 3. Here, $F = \text{Aut}(C_3) = C_2 = \{1, g\}$ with action given as $1(\sigma) = \sigma$, $g(\sigma) = \sigma^2$. A K -basis for $L = \text{Map}(C_2, K)$ is $\{e_1, e_g\}$, the basis dual to the basis $\{1, g\}$ for C_2 . We have $e_1(1) = 1$, $e_1(g) = 0$, $e_g(1) = 0$, $e_g(g) = 1$; $\{e_1, e_g\}$ is a basis of mutually orthogonal idempotents: $e_1^2 = e_1$, $e_g^2 = e_g$, $e_1 e_g = e_g e_1 = 0$, $e_1 + e_g = 1$, thus,

$$L = Ke_1 \oplus Ke_g.$$

The C_2 -Galois action on L is given as

$$1 \cdot e_1 = e_1, \quad 1 \cdot e_g = e_g, \quad g \cdot e_1 = e_g, \quad g \cdot e_g = e_1.$$

We compute $(L[C_3])^{C_2}$ directly. A typical element of $L[C_3]$ is $\sum_{i=0}^2 (\alpha_i e_1 + \beta_i e_g) \sigma^i$ for $\alpha_i, \beta_i \in K$. We require that

$$\begin{aligned}
g\left(\sum_{i=0}^2(\alpha_i e_1 + \beta_i e_g)\sigma^i\right) &= \sum_{i=0}^2 g(\alpha_i e_1 + \beta_i e_g)\sigma^{2i} \\
&= \sum_{i=0}^2 (\alpha_i e_g + \beta_i e_1)\sigma^{2i} \\
&= \sum_{i=0}^2 (\alpha_i e_1 + \beta_i e_g)\sigma^i.
\end{aligned}$$

Thus $\alpha_0 = \beta_0$, $\alpha_1 = \beta_2$, $\alpha_2 = \beta_1$. So a typical element of $(L[C_3])^{C_2}$ is

$$\begin{aligned}
&(\alpha_0 e_1 + \beta_0 e_g)1 + (\alpha_1 e_1 + \beta_1 e_g)\sigma + (\alpha_2 e_1 + \beta_2 e_g)\sigma^2 \\
&= \alpha_0(e_1 + e_g)1 + (\alpha_1 e_1 + \alpha_2 e_g)\sigma + (\alpha_2 e_1 + \alpha_1 e_g)\sigma^2 \\
&= \alpha_0 1 + \alpha_1(e_1\sigma + e_g\sigma^2) + \alpha_2(e_g\sigma + e_1\sigma^2).
\end{aligned}$$

Thus a K -basis for $(L[C_3])^{C_2}$ is $\{1, e_1\sigma + e_g\sigma^2, e_g\sigma + e_1\sigma^2\}$. We have

$$\begin{aligned}
\Delta(e_1\sigma + e_g\sigma^2) &= e_1(\sigma \otimes \sigma) + e_g(\sigma^2 \otimes \sigma^2) \\
&= (e_1\sigma + e_g\sigma^2) \otimes (e_1\sigma + e_g\sigma^2),
\end{aligned}$$

and so, $e_1\sigma + e_g\sigma^2$ is group-like. Likewise $e_g\sigma + e_1\sigma^2$ is group-like. Thus $(L[C_3])^{C_2} \cong K[C_3]$.

3. THE ABSOLUTELY SEMISIMPLE HOPF FORM OF $K[C_n]$

Let N be any finite group. For $n \geq 1$ let ζ_n denote a primitive n th root of unity. By Maschke's theorem, $K[N]$ is semisimple. Extending scalars to \mathbb{C} yields the Wedderburn-Artin decomposition

$$\mathbb{C}[N] \cong \text{Mat}_{n_1}(\mathbb{C}) \times \text{Mat}_{n_2}(\mathbb{C}) \times \cdots \times \text{Mat}_{n_l}(\mathbb{C})$$

for integers $n_i \geq 1$, $1 \leq i \leq l$. Let L be a faithfully flat K -algebra. Then any L -Hopf form of $K[N]$ is also semisimple. An L -Hopf form H of $K[N]$ is *absolutely semisimple* if

$$H \cong \text{Mat}_{n_1}(K) \times \text{Mat}_{n_2}(K) \times \cdots \times \text{Mat}_{n_l}(K).$$

Proposition 3.1. *Let $N = C_n = \langle \sigma \rangle$, $n \geq 1$. Then the linear dual $(K[C_n])^*$ is an absolutely semisimple Hopf form of $K[C_n]$.*

Proof. It is well-known that $(K[C_n])^*$ is a K -Hopf algebra. Let $\{p_j\}_{j=0}^{n-1}$ be the basis for $(K[C_n])^*$ dual to the basis $\{\sigma^i\}_{i=0}^{n-1}$ of $K[C_n]$. Let E/K be any field extension containing a primitive n th root of unity. Then E is a faithfully flat K -algebra. Let $\{\chi_j\}_{j=0}^{n-1}$ denote the set of n irreducible characters of C_n , which all are of degree 1 (since C_n is abelian). The χ_j take values in E . There exists an isomorphism of E -Hopf algebras

$$\varphi : E \otimes_K (K[C_n])^* \rightarrow E \otimes K[C_n] \cong E[C_n]$$

defined by

$$p_j \mapsto \frac{1}{n} \sum_{i=0}^{n-1} \chi_j(\sigma^{-i}) \sigma^i,$$

$0 \leq j \leq n-1$ (cf. [13, Exercise 6.4]). Thus $(K[C_n])^*$ is an E -form of $K[C_n]$. Since

$$\mathbb{C}[C_n] \cong \underbrace{\mathbb{C} \times \mathbb{C} \times \cdots \times \mathbb{C}}_n$$

and

$$(K[C_n])^* \cong \underbrace{K \times K \times \cdots \times K}_n,$$

$(K[C_n])^*$ is an absolutely semisimple Hopf form of $K[C_n]$. □

B. Pareigis has shown that $(K[C_n])^*$ is the only absolutely semisimple Hopf form of $K[C_n]$ [12, Theorem 4.3].

Theorem 3.2 (Pareigis). *$K[C_n]$ has a uniquely determined absolutely semisimple Hopf form $H = (K[C_n])^*$, where $(K[C_n])^*$ is the linear dual of $K[C_n]$.*

As a Hopf form of $K[C_n]$, $(K[C_n])^*$ comes from some F -Galois extension L , that is, $\Theta(L) = (K[C_n])^*$ for some L . In fact, if $L = K[x]/(\Phi_n(x))$, where $\Phi(x)$ is the n th cyclotomic polynomial, then L is an F -Galois extension of K and $\Theta(L) = (L[C_n])^F = (K[C_n])^*$.

Note that $K[x]/(\Phi_n(x))$ is not necessarily a field. For example, if $n = 15$ and $K = \mathbb{Q}(\zeta_3)$, then

$$K[x]/(\Phi_{15}(x)) \cong K(\zeta_{15}) \times K(\zeta_{15}).$$

In this case

$$F = \text{Aut}(C_{15}) = \mathbb{Z}_{15}^* = C_2 \times C_4.$$

The faithfully flat K -algebra $K(\zeta_{15}) \times K(\zeta_{15})$ is a $(C_2 \times C_4)$ -Galois extension of K with

$$\Theta(K(\zeta_{15}) \times K(\zeta_{15})) = (K[C_{15}])^*.$$

If $K = \mathbb{Q}$, then $\mathbb{Q}[x]/(\Phi_n(x))$ is a field, equal to $\mathbb{Q}(\zeta_n)$ where ζ_n denotes a primitive n th root of unity; $\mathbb{Q}(\zeta_n)$ is a \mathbb{Z}_n^* -Galois extension of \mathbb{Q} . In the case that $n = p^m$, for p an odd prime, $m \geq 1$, the first author has shown that $(K[C_{p^m}])^*$ is the fixed ring [11, Proposition 3.1].

Proposition 3.3 (Kohl). *Let p be an odd prime and let $L = \mathbb{Q}(\zeta_{p^m})$, $m \geq 1$. Then*

$$\Theta(L) = (L[C_{p^m}])^F = (\mathbb{Q}[C_{p^m}])^*,$$

where $F = \text{Aut}(C_{p^m}) = \mathbb{Z}_{p^m}^*$.

Proof. Let $C_{p^m} = \langle \sigma \rangle$. Then the elements

$$e_j = \frac{1}{p^m} \sum_{i=0}^{p^m-1} \zeta_{p^m}^{-ij} \sigma^i \in L[C_{p^m}],$$

$0 \leq j \leq p^m - 1$, are in the fixed ring $(L[C_{p^m}])^F$ and are mutually orthogonal idempotents. It follows that they form a \mathbb{Q} -basis for $(L[C_{p^m}])^F$. Thus $(L[C_{p^m}])^F \cong (\mathbb{Q}[C_{p^m}])^*$ as \mathbb{Q} -Hopf algebras. □

4. A PROBLEM OF PAREIGIS

B. Pareigis has stated the following problem in [12, p. 91].

Problem 4.1 (Pareigis). *For which finite groups N does $\mathbb{Q}[N]$ admit an absolutely semisimple Hopf form?*

We review solutions to Problem 4.1 for various N .

4.1. N Abelian.

Example 4.2. *Let $N = C_n$, $n \geq 1$. By Theorem 3.2, $\mathbb{Q}[C_n]$ admits the absolutely semisimple Hopf form $(\mathbb{Q}[C_n])^*$. If $L = \mathbb{Q}(\zeta_n)$, then $\Theta(L) = (\mathbb{Q}[C_n])^*$.*

Example 4.3. *Let $N = C_p^m$ be the elementary abelian group of order p^m , $p \geq 2$, $m \geq 1$. Let $E = \mathbb{Q}(\zeta_p)$. Then*

$$E \otimes_{\mathbb{Q}} (\mathbb{Q}[C_p^m])^* \cong E \otimes_{\mathbb{Q}} \mathbb{Q}[C_p^m],$$

as E -Hopf algebras. Moreover,

$$\mathbb{C}[C_p^m] \cong \underbrace{\mathbb{C} \times \mathbb{C} \times \cdots \times \mathbb{C}}_{p^m}$$

and

$$(\mathbb{Q}[C_p^m])^* \cong \underbrace{\mathbb{Q} \times \mathbb{Q} \times \cdots \times \mathbb{Q}}_{p^m}.$$

Thus $\mathbb{Q}[C_p^m]$ admits the absolutely semisimple Hopf form $(\mathbb{Q}[C_p^m])^*$.

By Theorem 2.2 there is an $\text{Aut}(C_p^m)$ -Galois extension L/\mathbb{Q} for which

$$\Theta(L) = (\mathbb{Q}[C_p^m])^*.$$

What is the structure of L ?

We have $\text{Aut}(C_p^m) = \text{GL}_m(\mathbb{F}_p)$ with $|\text{GL}_m(\mathbb{F}_p)| = \prod_{i=0}^{m-1} (p^m - p^i)$. There is a subgroup of $\text{GL}_m(\mathbb{F}_p)$ defined as

$$U = \{aI_m : a \in \mathbb{F}_p^*\} \cong \mathbb{F}_p^*$$

of index $l = (\prod_{i=0}^{m-1} (p^m - p^i))/(p - 1)$; $\mathbb{Q}(\zeta_p)$ is a U -Galois extension of fields. Thus by Theorem 2.1 there exists a $\text{GL}_m(\mathbb{F}_p)$ -Galois extension of \mathbb{Q}

$$L = \underbrace{\mathbb{Q}(\zeta_p) \times \mathbb{Q}(\zeta_p) \times \cdots \times \mathbb{Q}(\zeta_p)}_l.$$

Computing the fixed ring yields

$$\Theta(L) = (L[C_p^m])^{\text{GL}_m(\mathbb{F}_p)} = (\mathbb{Q}[C_p^m])^*.$$

4.2. **N Non-abelian.** We first prove a lemma.

Lemma 4.4. *Let D_3, D_4 denote the 3rd and 4th dihedral groups, respectively. Then*

- (i) $\text{Aut}(D_3) = D_3,$
- (ii) $\text{Aut}(D_4) = D_4.$

Proof. Let D_n denote the n th order dihedral group. Then $\text{Aut}(D_n) \cong \text{Hol}(C_n)$ since the cyclic subgroup of D_n is characteristic. To prove (i) note that

$$\text{Hol}(C_3) = C_3 \rtimes \text{Aut}(C_3) \cong C_3 \rtimes C_2 \cong D_3.$$

To prove (ii), observe that

$$\text{Hol}(C_4) = C_4 \rtimes \text{Aut}(C_4) \cong C_4 \rtimes C_2 \cong D_4,$$

where C_2 acts on C_4 by inversion. □

Example 4.5. *Let $N = D_3$. We have*

$$\mathbb{C}[D_3] \cong \mathbb{C} \times \mathbb{C} \times \text{Mat}_2(\mathbb{C})$$

and by [5, Example (7.39)]

$$\mathbb{Q}[D_3] \cong \mathbb{Q} \times \mathbb{Q} \times \text{Mat}_2(\mathbb{Q}).$$

Thus $\mathbb{Q}[D_3]$ is an absolutely semisimple Hopf form of itself. By Lemma 4.4(i), $\text{Aut}(D_3) = D_3$ and so

$$\Theta(\text{Map}(D_3, \mathbb{Q})) = \mathbb{Q}[D_3].$$

Example 4.6. *Let $N = D_4$. We have*

$$\mathbb{C}[D_4] \cong \mathbb{C} \times \mathbb{C} \times \mathbb{C} \times \mathbb{C} \times \text{Mat}_2(\mathbb{C})$$

and by [5, Example (7.39)]

$$\mathbb{Q}[D_4] \cong \mathbb{Q} \times \mathbb{Q} \times \mathbb{Q} \times \mathbb{Q} \times \text{Mat}_2(\mathbb{Q}).$$

Thus $\mathbb{Q}[D_4]$ is an absolutely semisimple Hopf form of itself. By Lemma 4.4(ii), $\text{Aut}(D_4) = D_4$ and so

$$\Theta(\text{Map}(D_4, \mathbb{Q})) = \mathbb{Q}[D_4].$$

4.2.1. *The case $N = Q_8$.* Let $N = Q_8$ denote the quaternion group

$$Q_8 = \{\pm 1, \pm i, \pm j, \pm k\},$$

with $ij = k, ji = -k, jk = i, kj = -i, ki = j, ik = -j$. We have

$$\mathbb{C}[Q_8] \cong \mathbb{C} \times \mathbb{C} \times \mathbb{C} \times \mathbb{C} \times \text{Mat}_2(\mathbb{C}),$$

yet

$$\mathbb{Q}[Q_8] \cong \mathbb{Q} \times \mathbb{Q} \times \mathbb{Q} \times \mathbb{Q} \times \mathbb{H},$$

where \mathbb{H} is the rational quaternions. Thus $\mathbb{Q}[Q_8]$ is not an absolutely semisimple form of itself. Does $\mathbb{Q}[Q_8]$ admit an absolutely semisimple Hopf form? The answer is “yes” as we will see below. The solution is due to C. Greither [4, §3].

We first identify a collection of Hopf forms of $\mathbb{Q}[Q_8]$. Let $\zeta = \zeta_4$. Then $\mathbb{Q}(\zeta)$ is a Galois extension of \mathbb{Q} with group $C_2 = \{1, g\}$. Let $\text{Aut}_{\text{Hopf}}(\mathbb{Q}[Q_8])$ denote the group of automorphisms of $\mathbb{Q}[Q_8]$ in the category of \mathbb{Q} -Hopf algebras. By [16, §17.6, Theorem], the $\mathbb{Q}(\zeta)$ -forms of $\mathbb{Q}[Q_8]$ are classified by the 1st cohomology set

$$H^1(\mathbb{Q}(\zeta)/\mathbb{Q}, \text{Aut}_{\text{Hopf}}(\mathbb{Q}[Q_8])).$$

Translating to C_2 -actions, we obtain

$$H^1(\mathbb{Q}(\zeta)/\mathbb{Q}, \text{Aut}_{\text{Hopf}}(\mathbb{Q}[Q_8])) = H^1(C_2, \text{Aut}_{\text{Hopf}}(\mathbb{Q}(\zeta) \otimes_{\mathbb{Q}} \mathbb{Q}[Q_8])).$$

by [16, §17.7, Theorem]. Since

$$\text{Aut}_{\text{Hopf}}(\mathbb{Q}(\zeta) \otimes_{\mathbb{Q}} \mathbb{Q}[Q_8]) = \text{Aut}_{\mathbb{Q}(\zeta)\text{-Hopf}}(\mathbb{Q}(\zeta)[Q_8]) = \text{Aut}(Q_8),$$

the $\mathbb{Q}(\zeta)$ -Hopf forms of $\mathbb{Q}[Q_8]$ are given by the cohomology set

$$H^1(C_2, \text{Aut}(Q_8)).$$

The action of C_2 on $\text{Aut}(Q_8)$ is trivial and so a 1-cocycle in $H^1(C_2, \text{Aut}(Q_8))$ is a homomorphism

$$\varrho : C_2 \rightarrow \text{Aut}(Q_8).$$

Given a homomorphism $\varrho \in H^1(C_2, \text{Aut}(Q_8))$, the corresponding $\mathbb{Q}(\zeta)$ -Hopf form is the fixed ring

$$H(\varrho) = (\mathbb{Q}(\zeta)[Q_8])^{C_2},$$

where $g \in C_2$ acts on $\mathbb{Q}(\zeta)$ as an element of the Galois group and on Q_8 through $\varrho(g)$.

Let $\theta : C_2 \rightarrow \text{Aut}(Q_8)$ be the homomorphism with $\theta(1) = 1$ and where $\theta(g)$ is conjugation by k .

Proposition 4.7 (Greither). *Let $H(\theta)$ be the Hopf form of $\mathbb{Q}[Q_8]$ corresponding to θ . Then the Wedderburn-Artin decomposition of $H(\theta)$ is*

$$H(\theta) = \mathbb{Q} \times \mathbb{Q} \times \mathbb{Q} \times \mathbb{Q} \times \text{Mat}_2(\mathbb{Q}).$$

Thus $H(\theta)$ is the absolutely semisimple Hopf form of $\mathbb{Q}[Q_8]$.

Proof. We compute the fixed ring

$$H(\theta) = (\mathbb{Q}(\zeta)[Q_8])^{C_2}.$$

Since the abelian part of $H(\theta)$ is equal to that of $\mathbb{Q}[Q_8]$, we have

$$H(\theta) = \mathbb{Q} \times \mathbb{Q} \times \mathbb{Q} \times \mathbb{Q} \times (\mathbb{Q}(\zeta) \otimes_{\mathbb{Q}} \mathbb{H})^{C_2}.$$

Writing

$$\mathbb{H} = (-1, -1)_{\mathbb{Q}} = \mathbb{Q} \oplus \mathbb{Q}u \oplus \mathbb{Q}v \oplus \mathbb{Q} \oplus w$$

with $u^2 = -1$, $v^2 = -1$, $uv = w$, $vu = -w$, we have that $g \in C_2$ acts on $\mathbb{Q}(\zeta)$ as an element of the Galois group and on \mathbb{H} through conjugation by w .

We claim that

$$(\mathbb{Q}(\zeta) \otimes_{\mathbb{Q}} \mathbb{H})^{C_2} \cong (1, 1)_{\mathbb{Q}} \cong \text{Mat}_2(\mathbb{Q}).$$

To this end, let

$$x = (a_0 + a_1\zeta)1 + (b_0 + b_1\zeta)u + (c_0 + c_1\zeta)v + (d_0 + d_1\zeta)w$$

be an element in

$$\mathbb{Q}(\zeta) \oplus \mathbb{Q}(\zeta)u \oplus \mathbb{Q}(\zeta)v \oplus \mathbb{Q}(\zeta)w = \mathbb{Q}(\zeta) \otimes_{\mathbb{Q}} \mathbb{H}$$

for $a_0, a_1, b_0, b_1, c_0, c_1, d_0, d_1 \in \mathbb{Q}$. Then

$$\begin{aligned} g(x) &= g((a_0 + a_1\zeta)1 + (b_0 + b_1\zeta)u + (c_0 + c_1\zeta)v + (d_0 + d_1\zeta)w) \\ &= (a_0 - a_1\zeta)1 + (b_0 - b_1\zeta)www^{-1} + (c_0 - c_1\zeta)wvw^{-1} + (d_0 - d_1\zeta)www^{-1} \\ &= (a_0 - a_1\zeta)1 + (b_0 - b_1\zeta)(-u) + (c_0 - c_1\zeta)(-v) + (d_0 - d_1\zeta)w \\ &= (a_0 - a_1\zeta)1 + (-b_0 + b_1\zeta)u + (-c_0 + c_1\zeta)v + (d_0 - d_1\zeta)w. \end{aligned}$$

Thus $g(x) = x$ if and only if $a_1 = b_0 = c_0 = d_1 = 0$. It follows that

$$B = \{1, \zeta v, \zeta u, w\}$$

is a \mathbb{Q} -basis for $(\mathbb{Q}(\zeta) \otimes_{\mathbb{Q}} \mathbb{H})^{C_2}$. Now, $(\zeta v)^2 = 1$, $(\zeta u)^2 = 1$, $(\zeta v)(\zeta u) = -1vu = w$, $(\zeta u)(\zeta v) = -1uv = -w$, and thus B is a \mathbb{Q} -basis for the quaternion algebra $(1, 1)_{\mathbb{Q}}$, which by [6, Theorem 4.3] is isomorphic to $\text{Mat}_2(\mathbb{Q})$. Consequently,

$$H(\theta) = \mathbb{Q} \times \mathbb{Q} \times \mathbb{Q} \times \mathbb{Q} \times \text{Mat}_2(\mathbb{Q})$$

as claimed.

One could also conclude that

$$(\mathbb{Q}(\zeta) \otimes_{\mathbb{Q}} \mathbb{H})^{C_2} \cong \text{Mat}_2(\mathbb{Q})$$

by noting that $\zeta u - w$ is a non-zero nilpotent element of $(\mathbb{Q}(\zeta) \otimes_{\mathbb{Q}} \mathbb{H})^{C_2}$ of index 2. □

By a standard result $\text{Aut}(Q_8) = S_4$ and so by Theorem 2.2 there is an S_4 -Galois extension L/\mathbb{Q} for which

$$\Theta(L) = H(\theta).$$

We compute L/\mathbb{Q} as follows. Let $U = \theta(C_2) \cong C_2$. Then $\theta(C_2)$ is a subgroup of S_4 of index $[S_4 : \theta(C_2)] = 12$. Thus by Theorem 2.1, there exists an S_4 -Galois extension of \mathbb{Q}

$$L = \underbrace{\mathbb{Q}(\zeta) \times \mathbb{Q}(\zeta) \times \cdots \times \mathbb{Q}(\zeta)}_{12}.$$

Moreover

$$\Theta(L) = (L[Q_8])^{S_4} = (\mathbb{Q}(\zeta)[Q_8])^{C_2} = H(\theta).$$

Remark 4.8. As C. Greither has noted there exist finite groups N for which $\mathbb{Q}[N]$ has no absolutely semisimple Hopf forms. For instance, take N to be any non-abelian group of odd order p^3 and exponent p^2 [4, §3].

5. CONNECTION TO HOPF-GALOIS THEORY

There is a natural application of the map Θ to Hopf-Galois theory.

5.1. Review of Greither-Pareigis theory. Let E/K be a Galois extension with group G . Let H be a finite dimensional, cocommutative K -Hopf algebra with comultiplication $\Delta : H \rightarrow H \otimes_R H$, counit $\varepsilon : H \rightarrow K$, and coinverse $S : H \rightarrow H$. Suppose there is a K -linear action \cdot of H on E that satisfies

$$h \cdot (xy) = \sum_{(h)} (h_{(1)} \cdot x)(h_{(2)} \cdot y), \quad h \cdot 1 = \varepsilon(h)1$$

for all $h \in H$, $x, y \in E$, where $\Delta(h) = \sum_{(h)} h_{(1)} \otimes h_{(2)}$ is Sweedler notation. Suppose also that the K -linear map

$$j : E \otimes_K H \rightarrow \text{End}_K(E), \quad j(x \otimes h)(y) = x(h \cdot y)$$

is an isomorphism of vector spaces over K . Then H together with this action, denoted as (H, \cdot) , provides a *Hopf-Galois structure* on E/K . Two Hopf-Galois structures (H_1, \cdot_1) , (H_2, \cdot_2) on E/K are *isomorphic* if there is a Hopf algebra isomorphism $f : H_1 \rightarrow H_2$ for which $h \cdot_1 x = f(h) \cdot_2 x$ for all $x \in E$, $h \in H$ (see [7, Introduction]).

C. Greither and B. Pareigis [8] have given a complete classification of Hopf-Galois structures up to isomorphism. Denote by $\text{Perm}(G)$ the group of permutations of G . A subgroup $N \leq \text{Perm}(G)$ is *regular* if $|N| = |G|$ and $\eta(g) \neq g$ for all $\eta \neq 1_N$, $g \in G$. Let $\lambda : G \rightarrow \text{Perm}(G)$, $\lambda(g)(h) = gh$, denote the left regular representation. A subgroup $N \leq \text{Perm}(G)$ is *normalized* by $\lambda(G) \leq \text{Perm}(G)$ if $\lambda(G)$ is contained in the normalizer of N in $\text{Perm}(G)$.

Theorem 5.1 (Greither and Pareigis). *Let E/K be a Galois extension with group G . There is a one-to-one correspondence between isomorphism classes of Hopf Galois structures on E/K and regular subgroups of $\text{Perm}(G)$ that are normalized by $\lambda(G)$.*

One direction of this correspondence works by Galois descent: Let N be a regular subgroup of $\text{Perm}(G)$ normalized by $\lambda(G)$. Then G acts on the group algebra $E[N]$ through the Galois action on E and conjugation by $\lambda(G)$ on N , i.e.,

$$g(x\eta) = g(x)(\lambda(g)\eta\lambda(g^{-1})), \quad g \in G, \quad x \in E, \quad \eta \in N.$$

We denote the conjugation action of $\lambda(g) \in \lambda(G)$ on $\eta \in N$ by ${}^g\eta$. Let H denote the fixed ring

$$(E[N])^G = \{x \in E[N] : g(x) = x, \forall g \in G\}.$$

Then H is an n -dimensional E -Hopf algebra, $n = [E : K]$, and E/K admits the Hopf Galois structure (H, \cdot) [8, p. 248, proof of 3.1 (b) \implies (a)], [2, Theorem 6.8, pp. 52-54]. The action of H on E/K is given as

$$\left(\sum_{\eta \in N} r_\eta \eta \right) \cdot x = \sum_{\eta \in N} r_\eta \eta^{-1} [1_G](x),$$

see [3, Proposition 1]. By [8, p. 249, proof of 3.1, (a) \implies (b)],

$$E \otimes_K H \cong E \otimes_K K[N] \cong E[N],$$

as E -Hopf algebras, so H is an E -form of $K[N]$.

If N is isomorphic to the abstract group N' , then we say that the Hopf-Galois structure (H, \cdot) on E/K is of *type* N' .

5.2. Connection to the map Θ . If (H, \cdot) is a Hopf-Galois structure on E/K of type N , then the Hopf algebra H is a Hopf form of $K[N]$. Thus H can be recovered via Theorem 2.2. In other words, with $F = \text{Aut}(N)$, there is an F -Galois extension L of K with

$$\Theta(L) = H = (L[N])^F.$$

We seek a method to construct L . We first prove a lemma.

Lemma 5.2. *Let E/K be a Galois extension with group G . Let (H, \cdot) be a Hopf-Galois structure corresponding to regular subgroup N . Let*

$$W = \{g \in \lambda(G) : {}^g\eta = \eta, \forall \eta \in N\}.$$

Then W is a normal subgroup of $\lambda(G)$.

Proof. Certainly $W \leq \lambda(G)$. Let N^{opp} denote the centralizer of N in $\text{Perm}(G)$ and note that

$$W = \lambda(G) \cap N^{opp}.$$

The normalizer of N in $\text{Perm}(G)$ equals the normalizer of N^{opp} in $\text{Perm}(G)$. Therefore since $\lambda(G)$ normalizes N , $\lambda(G)$ also normalizes N^{opp} . Moreover, $\lambda(G)$ normalizes itself and so $\lambda(G)$ normalizes W , i.e., $W \triangleleft \lambda(G)$. □

The quotient group $\lambda(G)/W$ can be viewed as a subgroup of $F = \text{Aut}(N)$.

Proposition 5.3. *Let E/K be a Galois extension with group G . Let (H, \cdot) be a Hopf-Galois structure corresponding to regular subgroup N . Let $F = \text{Aut}(N)$, let*

$$W = \{g \in \lambda(G) : {}^g\eta = \eta, \forall \eta \in N\},$$

and let $L = E^W$. If $\lambda(G)/W \cong F$, then $\Theta(L) = H$. Moreover, L is the smallest extension of K for which $L \otimes_K H \cong L[N]$.

Proof. By the Fundamental theorem of Galois theory, $L = E^W$ is Galois with group $F \cong \lambda(G)/W$, so L is an F -Galois extension. Now,

$$H = (E[N])^G = (L[N])^F,$$

and so, $\Theta(L) = H$. The second statement follows from [8, Corollary 3.2]. □

Example 5.4. We consider the splitting field E of the polynomial $x^4 - 10x^2 + 1$ over \mathbb{Q} . One has $E = \mathbb{Q}(\sqrt{2}, \sqrt{3})$; E/\mathbb{Q} is Galois with group $C_2 \times C_2 = \{1, \sigma, \tau, \tau\sigma\}$ with Galois action

$$\sigma(\sqrt{2}) = \sqrt{2}, \quad \sigma(\sqrt{3}) = -\sqrt{3}, \quad \tau(\sqrt{2}) = -\sqrt{2}, \quad \tau(\sqrt{3}) = \sqrt{3}.$$

By [1], there are three Hopf-Galois structures on E/\mathbb{Q} of type C_4 , each of which is determined by a regular subgroup $N \cong C_4$ normalized by $\lambda(C_2 \times C_2)$. One such N is given as

$$N = \{(1), (1, 3, 2, 4), (1, 2)(3, 4), (1, 4, 2, 3)\},$$

where $1 := 1$, $2 := \sigma$, $3 := \tau$, $4 := \tau\sigma$, and

$$\lambda(C_2 \times C_2) = \{(1), (1, 2)(3, 4), (1, 3)(2, 4), (1, 4)(2, 3)\}.$$

N is a regular subgroup of $\text{Perm}(C_2 \times C_2)$ normalized by $\lambda(C_2 \times C_2)$ with $N \cong C_4$.

Let (H, \cdot) be the corresponding Hopf-Galois extension with $H = (E[N])^G$. As one can check

$$W = \{g \in \lambda(C_2 \times C_2) : {}^g\eta = \eta, \forall \eta \in N\} = \{(1), (1, 2)(3, 4)\} = \{1, \sigma\}.$$

We have $\lambda(C_2 \times C_2)/W \cong F = \text{Aut}(C_4) \cong C_2$, and the fixed field $L = E^W = \mathbb{Q}(\sqrt{2})$ is an F -Galois extension of \mathbb{Q} . So by Proposition 5.3

$$\Theta(L) = H.$$

Example 5.5. Let $E = \mathbb{Q}(\zeta_3, \sqrt[3]{2})$ be the splitting field of $x^3 - 2$ over \mathbb{Q} , so that E/\mathbb{Q} is Galois with group S_3 . By [10, Lemma 1], there are three Hopf-Galois structures on E/\mathbb{Q} of type C_6 , all of which have isomorphic Hopf algebras H by [10, Proposition 3]. We have

$$W = \{g \in \lambda(S_3) : {}^g\eta = \eta, \forall \eta \in C_6\} \cong C_3.$$

Thus $\lambda(S_3)/W \cong C_2 = \text{Aut}(C_6)$ and $E^W = \mathbb{Q}(\zeta_3)$. Consequently,

$$\Theta(\mathbb{Q}(\zeta_3)) = H$$

by Proposition 5.3. In this case $H = (\mathbb{Q}[C_6])^*$ is the absolutely semisimple Hopf form of $\mathbb{Q}[C_6]$.

Example 5.6. Let E/K be a Galois extension with group G where G is a non-abelian complete group (i.e., G has trivial center and $G \cong \text{Aut}(G)$). For instance, $G = S_n$, $n > 6$, is a non-abelian complete group.

The subgroup $N = \lambda(G)$ is a regular subgroup of $\text{Perm}(G)$ normalized by itself and corresponds to the canonical non-classical Hopf-Galois structure with Hopf algebra H_λ . In this case W is trivial since the center of G is trivial, and so $E^W = E$ and

$$\lambda(G)/W \cong \lambda(G) \cong \text{Aut}(N) = F.$$

Thus by Proposition 5.3

$$\Theta(E) = H_\lambda.$$

5.3. **When $\lambda(G)/W \not\cong F$.** In Example 5.4, Example 5.5, and Example 5.6 we have $\lambda(G)/W \cong F$ and Proposition 5.3 can be applied to find L so that $\Theta(L) = H$. We next consider the case where $\lambda(G)/W$ is a proper subgroup of F .

Let E/\mathbb{Q} be Galois with quaternion group $Q_8 = \{\pm 1, \pm i, \pm j, \pm k\}$. Then $\{\pm 1\}$ is the unique order 2 subgroup of Q_8 . Let $K = E^{\{\pm 1\}}$. Since $\{\pm 1\}$ is normal in Q_8 , K/\mathbb{Q} is Galois with group

$$Q_8/\{\pm 1\} = \{\bar{1}, \bar{i}, \bar{j}, \bar{k}\} \cong C_2 \times C_2.$$

Thus K/\mathbb{Q} is the unique biquadratic subfield of E . For $s, t \in \{i, j, k\}$ with $s \neq t$, there exist elements α, β in K satisfying $\alpha^2 \in \mathbb{Q}$, $\beta^2 \in \mathbb{Q}$ with $K = \mathbb{Q}(\alpha, \beta)$. We have

$$s(\alpha) = \alpha, \quad t(\alpha) = -\alpha, \quad s(\beta) = -\beta, \quad t(\beta) = \beta.$$

Note that

$$E^{(t)} = \mathbb{Q}(\beta).$$

S. Taylor and P. J. Truman [15, Lemma 2.3] have shown that there are 6 Hopf-Galois structures of type $N = C_8$ on E/\mathbb{Q} . The 6 regular subgroups are given as the 6 subgroups

$$\{C_{s,t} \mid s, t \in \{i, j, k\}, s \neq t\},$$

where

$$C_{s,t} = \langle \eta_{s,t} \rangle \cong C_8.$$

Here the generator $\eta_{s,t}$ denotes the permutation in cycle notation

$$\eta_{s,t} = (1, s, t, t^{-1}s^{-1}, s^2, s^{-1}, t^{-1}, st).$$

The action of G on $C_{s,t}$ is given as

$${}^s\eta_{s,t} = \lambda(s)\eta_{s,t}\lambda(s^{-1}) = \eta_{s,t}^3,$$

$${}^t\eta_{s,t} = \lambda(t)\eta_{s,t}\lambda(t^{-1}) = \eta_{s,t}.$$

(See [15, proof of Lemma 2.3].)

Let $s, t \in \{i, j, k\}$, $s \neq t$. Then by Theorem 5.1 the fixed ring

$$H_{s,t} = (E[C_{s,t}])^G$$

is the Hopf algebra attached to the Hopf-Galois structure on E/\mathbb{Q} ; $H_{s,t}$ is an E -form of $\mathbb{Q}[C_8]$. Note that

$$W = \{g \in \lambda(Q_8) : {}^g\eta = \eta, \forall \eta \in C_{s,t}\} = \langle t \rangle,$$

with $\lambda(Q_8)/W \cong C_2$. Thus $\lambda(Q_8)/W$ is a proper subgroup of $F = \text{Aut}(C_8) \cong C_2 \times C_2$.

5.4. **Finding L so that $\Theta(L) = H_{s,t}$.** We want recover $H_{s,t}$ using Theorem 2.2, i.e., with $F = \text{Aut}(C_8)$, we want to find an F -Galois extension L for which

$$\Theta(L) = (L[C_8])^F = H_{s,t}.$$

We have

$$F = \mathbb{Z}_8^* = \{1, 3, 5, 7\} \cong C_2 \times C_2.$$

Let $U = \langle 3 \rangle \leq F$. We identify 3 with the element $s \in \{i, j, k\}$. Then $\mathbb{Q}(\beta)$ is a Galois extension of \mathbb{Q} with group U ; $3(\beta) = -\beta$. (Recall that $E^{(t)} = \mathbb{Q}(\beta)$.) Let $\{1, 5\}$ be a left transversal for U in F , i.e., the left cosets of U in F are

$$\langle 3 \rangle, \quad 5\langle 3 \rangle.$$

Let

$$L = \mathbb{Q}(\beta)f_1 \oplus \mathbb{Q}(\beta)f_2$$

where f_1, f_2 are mutually orthogonal idempotents. A typical element of L can be written as

$$x = (a_0 + a_1\beta)f_1 + (b_0 + b_1\beta)f_2,$$

where $a_0, a_1, b_0, b_1 \in \mathbb{Q}$. Following [12, proof of Theorem 4.2], L is an F -Galois extension of \mathbb{Q} with action of F on L given as

$$\begin{aligned} 1 \cdot x &= x, \\ 3 \cdot x &= (a_0 - a_1\beta)f_1 + (b_0 - b_1\beta)f_2, \\ 5 \cdot x &= (b_0 + b_1\beta)f_1 + (a_0 + a_1\beta)f_2, \\ 7 \cdot x &= (b_0 - b_1\beta)f_1 + (a_0 - a_1\beta)f_2. \end{aligned}$$

Proposition 5.7. *For $s, t \in \{i, j, k\}$, $s \neq t$, let $H_{s,t}$ be the Hopf algebra for the Hopf-Galois structure on E/\mathbb{Q} corresponding to regular subgroup $C_{s,t}$. Let L be as above. Then*

$$\Theta(L) = (L[C_8])^F = H_{s,t}.$$

Proof. We compute a \mathbb{Q} -basis for the fixed ring directly. Let $\eta = \eta_{s,t}$ and let

$$\sum_{m=0}^7 ((a_{m,0} + a_{m,1}\beta)f_1 + (b_{m,0} + b_{m,1}\beta)f_2)\eta^m$$

be a typical element of $L[C_{s,t}]$. Then

$$\begin{aligned} &g \left(\sum_{m=0}^7 ((a_{m,0} + a_{m,1}\beta)f_1 + (b_{m,0} + b_{m,1}\beta)f_2)\eta^m \right) \\ &= \sum_{m=0}^7 ((a_{m,0} + a_{m,1}\beta)f_1 + (b_{m,0} + b_{m,1}\beta)f_2)\eta^m \end{aligned}$$

for all $g \in F = C_2 \times C_2$ if and only if

$$a_{0,0} = b_{0,0}, \quad a_{0,1} = b_{0,1} = 0,$$

$$a_{1,0} = a_{3,0} = b_{5,0} = b_{7,0}, \quad a_{1,1} = -a_{3,1} = b_{5,1} = -b_{7,1},$$

$$b_{1,0} = b_{3,0} = a_{5,0} = a_{7,0}, \quad b_{1,1} = -b_{3,1} = a_{5,1} = -a_{7,1},$$

$$a_{2,0} = a_{6,0} = b_{2,0} = b_{6,0}, \quad a_{2,1} = -a_{6,1} = b_{2,1} = -b_{6,1},$$

$$a_{4,0} = b_{4,0}, \quad a_{4,1} = b_{4,1} = 0.$$

It follows that a \mathbb{Q} -basis for $(L[C_8])^F$ is

$$1, \quad \eta^4, \quad \eta^2 + \eta^6, \quad \beta(\eta^2 - \eta^6),$$

$$f_1(\eta + \eta^3) + f_2(\eta^5 + \eta^7), \quad f_2(\eta + \eta^3) + f_1(\eta^5 + \eta^7),$$

$$\beta(f_1(\eta - \eta^3) + f_2(\eta^5 - \eta^7)), \quad \beta(f_2(\eta - \eta^3) + f_1(\eta^5 - \eta^7)).$$

Thus a \mathbb{Q} -basis for $(L[C_8])^F$ can be written as

$$\frac{1}{8} (1 + \eta + \eta^2 + \eta^3 + \eta^4 + \eta^5 + \eta^6 + \eta^7),$$

$$\frac{1}{8} (1 - \eta + \eta^2 - \eta^3 + \eta^4 - \eta^5 + \eta^6 - \eta^7),$$

$$\frac{1}{4} (1 - \eta^2 + \eta^4 - \eta^6), \quad \frac{1}{4} \beta((\eta - \eta^3) + (\eta^5 - \eta^7)),$$

$$\frac{1}{2} (1 - \eta^4), \quad \frac{1}{2} \beta(\eta^2 - \eta^6),$$

together with two other basis elements

$$\begin{aligned} & \frac{1}{2} (f_2(\eta + \eta^3) + f_1(\eta^5 + \eta^7)) - \frac{1}{2} (f_1(\eta + \eta^3) + f_2(\eta^5 + \eta^7)) \\ &= \frac{1}{2} (f_2 - f_1)((\eta^3 - \eta^7) + (\eta - \eta^5)), \end{aligned}$$

and

$$\begin{aligned} & \frac{1}{2} \beta(f_1(\eta - \eta^3) + f_2(\eta^5 - \eta^7)) - \frac{1}{2} \beta(f_2(\eta - \eta^3) + f_1(\eta^5 - \eta^7)) \\ &= \frac{1}{2} \beta(f_2 - f_1)((\eta^3 - \eta^7) - (\eta - \eta^5)). \end{aligned}$$

Note that

$$(f_2 - f_1)^2 = f_2 + f_1 = 1.$$

Considering the \mathbb{Q} -basis for $H_{s,t}$ given in [15, page 231], we see that there is an isomorphism of \mathbb{Q} -algebras

$$\psi : (L[C_8])^F \rightarrow H_{s,t}$$

defined by

$$\frac{1}{8} (1 + \eta + \eta^2 + \eta^3 + \eta^4 + \eta^5 + \eta^6 + \eta^7) \mapsto \frac{1}{8} (1 + \eta + \eta^2 + \eta^3 + \eta^4 + \eta^5 + \eta^6 + \eta^7),$$

$$\frac{1}{8} (1 - \eta + \eta^2 - \eta^3 + \eta^4 - \eta^5 + \eta^6 - \eta^7) \mapsto \frac{1}{8} (1 - \eta + \eta^2 - \eta^3 + \eta^4 - \eta^5 + \eta^6 - \eta^7),$$

$$\frac{1}{4} (1 - \eta^2 + \eta^4 - \eta^6) \mapsto \frac{1}{4} (1 - \eta^2 + \eta^4 - \eta^6),$$

$$\frac{1}{4} \beta ((\eta - \eta^3) + (\eta^5 - \eta^7)) \mapsto \frac{1}{4} \beta ((\eta - \eta^3) + (\eta^5 - \eta^7)),$$

$$\frac{1}{2} (1 - \eta^4) \mapsto \frac{1}{2} (1 - \eta^4),$$

$$\frac{1}{2} \beta (\eta^2 - \eta^6) \mapsto \frac{1}{2} \beta (\eta^2 - \eta^6),$$

$$\frac{1}{2} (f_2 - f_1) ((\eta^3 - \eta^7) + (\eta - \eta^5)) \mapsto \frac{1}{2} ((\eta^3 - \eta^7) + (\eta - \eta^5)),$$

$$\frac{1}{2} \beta (f_2 - f_1) ((\eta^3 - \eta^7) - (\eta - \eta^5)) \mapsto \frac{1}{2} \beta ((\eta^3 - \eta^7) - (\eta - \eta^5)).$$

Moreover, since there is an embedding of L -Hopf algebras

$$L[(f_2 - f_1)(1 - \eta^4)] \rightarrow L[1 - \eta^4] \cong L[\langle \eta^4 \rangle] \cong L[C_2]$$

given as $(f_2 - f_1)(1 - \eta^4) \mapsto 1 - \eta^4$, we conclude that the \mathbb{Q} -algebra isomorphism ψ is also an isomorphism of \mathbb{Q} -Hopf algebras. □

As a byproduct of Proposition 5.7, we can recover [15, Lemma 3.3].

Proposition 5.8. *Let $s, s', t, t' \in \{i, j, k\}$ with $s \neq t$, $s' \neq t'$. Then $H_{s,t} \cong H_{s',t'}$ as Hopf algebras if and only if $t = t'$.*

Proof. If $t = t'$ then the fixed fields $E^{(t)}$ and $E^{(t')}$ are both equal to $\mathbb{Q}(\beta)$. Let

$$L = \mathbb{Q}(\beta)f_1 \oplus \mathbb{Q}(\beta)f_2.$$

Then $\Theta(L) = H_{s,t} = H_{s',t'}$ so that $H_{s,t} \cong H_{s',t'}$.

For the converse, suppose that $H_{s,t} \cong H_{s',t'}$. Then there is an F -Galois extension L of \mathbb{Q} with $\Theta(L) = H_{s,t}$ and an F -Galois extension L' of \mathbb{Q} with $\Theta(L') = H_{s',t'}$. Necessarily, $L \cong \mathbb{Q}(\beta)f_1 \oplus \mathbb{Q}(\beta)f_2$ where $E^{(t)} = \mathbb{Q}(\beta)$ and $L' \cong \mathbb{Q}(\gamma)f_1 \oplus \mathbb{Q}(\gamma)f_2$ where $E^{(t')} = \mathbb{Q}(\gamma)$. Now $H_{s,t} \cong H_{s',t'}$ implies that $L \cong L'$, which says that $\mathbb{Q}(\beta) \cong \mathbb{Q}(\gamma)$, hence $t = t'$. \square

REFERENCES

- [1] N. P. Byott, Integral Hopf-Galois structures on degree p^2 extensions of p -adic fields, *J. Algebra*, **248**, 2002, 334-365.
- [2] L. N. Childs, *Taming Wild Extensions: Hopf Algebras and Local Galois Module Theory*, AMS: Mathematical Surveys and Monographs, **80**, 2000.
- [3] L. N. Childs, Hopf Galois structures on Kummer extensions of prime power degree, *New York J. Math.*, **17**, 2011, 51-74.
- [4] L. N. Childs, C. Greither, K. Keating, A. Koch, T. Kohl, P. Truman, R. Underwood, *Hopf Algebras and Galois Module Theory*, AMS: submitted Aug. 2020.
- [5] C. W. Curtis and I. Reiner, *Methods of Representation Theory, vol. 1*, Wiley, 1981.
- [6] K. Conrad, Quaternion algebras, <https://kconrad.math.uconn.edu/blurbs/ringtheory/quaternionalg.pdf>
- [7] T. Crespo, A. Rio, M. Vela, Non-isomorphic Hopf-Galois structures with isomorphic underlying Hopf algebras, *J. Algebra*, **455**, (2015), 270-276.
- [8] C. Greither and B. Pareigis, Hopf Galois theory for separable field extensions, *J. Algebra*, **106**, 1987, 239-258.
- [9] R. Hagenmüller, B. Pareigis, Hopf algebra forms on the multiplicative group and other groups, *manuscripta math.*, **55**, (1986), 121-135.
- [10] A. Koch, T. Kohl, P. Truman, R. Underwood, The Structure of Hopf Algebras Acting on Dihedral Extensions. In: Feldvoss J., Grimley L., Lewis D., Pavelescu A., Pillen C. (eds) *Advances in Algebra. SRAC 2017. Springer Proceedings in Mathematics & Statistics*, vol 277. Springer, Cham, (2019).
- [11] T. Kohl, A Class of profinite Hopf-Galois extensions over \mathbb{Q} , *Comm. Alg.*, **46**(6), 2018, 2534-2550.
- [12] B. Pareigis, Forms of Hopf algebras and Galois theory, *Topics in Algebra*, Banach Center Publications, **26**, Part 1, PWN Polish Scientific Publishers, 1990.
- [13] J. P. Serre, *Linear Representations of Finite Groups*, GTM 42, Springer-Verlag, New York, 1977.
- [14] C. Small, The Group of quadratic extensions, *J. Pure Appl. Alg.*, **2**, (1972), 83-105.
- [15] S. Taylor, P. J. Truman, The Structure of Hopf algebras giving Hopf-Galois structures on quaternionic extensions, *New York J. Math.*, **25**, 2019, 219-237.
- [16] W. Waterhouse, *Introduction to Affine Group Schemes*, Springer-Verlag, New York, 1979.

DEPARTMENT OF MATHEMATICS AND STATISTICS, BOSTON UNIVERSITY, 111 CUMMINGTON MALL,
BOSTON, MA 02215 USA

Email address: tkohl@math.bu.edu

DEPARTMENT OF MATHEMATICS AND DEPARTMENT OF COMPUTER SCIENCE, AUBURN UNIVERSITY
AT MONTGOMERY, MONTGOMERY, AL, 36124 USA

Email address: runderwo@aum.edu