

ULAM STABILITY OF JENSEN FUNCTIONAL INEQUALITY IN GROUPS

GANG LU*, LULU QU, YUANFENG JIN*, AND CHOONKIL PARK

ABSTRACT. In this paper we establish the Ulam stability of Jensen functional inequality in some classes of groups.

1. INTRODUCTION AND PRELIMINARIES

Ulam [22] raised a stability problem for a homomorphism in metric groups. In 1941, Hyers [11] proved that if f is a mapping from a normed vector space into a Banach space and satisfies $\|f(x+y) - f(x) - f(y)\| \leq \varepsilon$, then there exists an additive function A such that $\|f(x) - A(x)\| \leq \varepsilon$. Hyers' theorem was generalized by Aoki [3] for additive mappings and by Rassias [21] for linear mappings by considering an unbounded Cauchy difference. A generalization of the Rassias theorem was obtained by Găvruta [10] by replacing the unbounded Cauchy difference by a general control function in the spirit of Rassias' approach. The stability problems for several functional equations or inequalities have been extensively investigated by a number of authors and there are many interesting results concerning this problem (see [2, 4, 5, 6, 7, 9, 12, 13, 14, 15, 17, 18, 19, 20, 23]).

In this paper we investigate the stability of the Jensen functional inequality

$$\|f(xy) + f(xy^{-1}) - 2f(x)\| \leq \|\rho(f(xy) - f(x) - f(y))\|$$

in some classes of noncommutative groups. The Jensen functional equation was studied in [1, 8, 16].

2. MAIN RESULTS

Suppose that X is an arbitrary group and Y is an arbitrary real Banach space. In this section, assume that X is an arbitrary multiplicative group and e is the identity element of X .

Definition 2.1. We say that a function $f : X \rightarrow Y$ is a ρ -Jensen function if the function satisfies

$$\|f(xy) + f(xy^{-1}) - 2f(x)\| \leq \|\rho(f(xy) - f(x) - f(y))\|. \quad (2.1)$$

for all $x, y \in X$.

2010 *Mathematics Subject Classification.* Primary 39B62, 39B52, 46B25.

Key words and phrases. Jensen functional inequality; Ulam stability; group.

*Corresponding authors: lvgang1234@163.com (G. Lu); yfkim@ybu.edu.cn (Y. Jin).

We denote the set of all ρ -Jensen functions by $J\rho(X; Y)$. Denote by $J\rho_0(X; Y)$ the subset of $J\rho(X; Y)$ consisting of functions f such that $f(e) = 0$. Obviously, $J\rho_0(X; Y)$ is a subspace of $J\rho(X; Y)$ and $J\rho(X; Y) = J\rho_0(X; Y) \oplus Y$.

Definition 2.2. We say that a function $f : X \rightarrow Y$ is an (X, Y) -quasi ρ -Jensen function if there is $c > 0$ such that

$$\|f(xy) + f(xy^{-1}) - 2f(x)\| \leq \|\rho(f(xy) - f(x) - f(y))\| + c \quad (2.2)$$

for all $x, y \in X$.

It is clear that the set of (X, Y) -quasi ρ -Jensen functions is a real linear space. Denote it by $KJ_\rho(X; Y)$. From (2.2), we have

$$\|f(y) + f(y^{-1}) - 2f(e)\| \leq |\rho|\|f(e)\| + c. \quad (2.3)$$

Therefore

$$\|f(y) + f(y^{-1})\| \leq c_1, \quad (2.4)$$

where $c_1 = c + (2 + |\rho|)\|f(e)\|$. Now letting $x = y$ in (2.2), we get

$$\|f(x^2) + f(e) - 2f(x)\| \leq |\rho|\|f(x^2) - 2f(x)\| + c. \quad (2.5)$$

Hence

$$\|f(x^2) - 2f(x)\| \leq c_2, \quad (2.6)$$

where $c_2 = \frac{c + \|f(e)\|}{1 - |\rho|}$. Again letting $y = x^2$ in (2.2), we get

$$\|f(x^3) + f(x^{-1}) - 2f(x)\| \leq |\rho|\|f(x^3) - f(x^2) - f(x)\| + c. \quad (2.7)$$

By (2.4) and (2), we obtain

$$\|f(x^3) - 3f(x)\| \leq c_3, \quad (2.8)$$

where $c_3 = \frac{|\rho|c_2 + c_1 + c}{1 - |\rho|}$.

Let c be as in (2.2) and define the set C as follow: $C = \{c_n | n \in \mathbb{N}\}$, where $c_1 = (|\rho| + 2)\|f(e)\| + c$, $c_2 = \frac{c + \|f(e)\|}{1 - |\rho|}$, $c_3 = \frac{|\rho|c_2 + c_1 + c}{1 - |\rho|}$ and $c_n = \frac{|\rho|c_{n-1} + c_{n-2} + c_1 + c}{1 - |\rho|}$ if $n > 3$.

Lemma 2.3. Let $f \in KJ_\rho(X; Y)$ such that

$$\|f(xy) + f(xy^{-1}) - 2f(x)\| \leq \|f(xy) - f(x) - f(y)\| + c.$$

Then for any $x \in X$ and any $m \in \mathbb{N}$ with $m \geq 2$ the following relation holds:

$$\|f(x^m) - mf(x)\| \leq c_m. \quad (2.9)$$

Proof. The proof is by induction on m . For $m = 2$, the lemma is established in (2.6). Suppose that for m the lemma has been already established, let us verify it for $m + 1$. Letting $y = x^m$ in (2.2), we have

$$\|f(x^{m+1}) + f(x^{-(m-1)}) - 2f(x)\| \leq |\rho|\|f(x^m) - f(x^{m-1}) - f(x)\| + c. \quad (2.10)$$

Then

$$\begin{aligned}
 & \|f(x^{m+1}) - (m+1)f(x)\| - \|f(x^{-(m-1)}) + f(x^{m-1})\| \\
 & - \|f(x^{m-1}) - (m-1)f(x)\| \\
 & \leq |\rho| \|f(x^{m+1}) - (m+1)f(x)\| + |\rho| \|f(x^m) - mf(x)\| + c
 \end{aligned} \tag{2.11}$$

and hence

$$\|f(x^{m+1}) - (m+1)f(x)\| \leq \frac{|\rho|c_m + c_{m-1} + c_1 + c}{1 - |\rho|} = c_{m+1}. \tag{2.12}$$

Now the lemma is proved. \square

Lemma 2.4. *Let $f \in KJ_\rho(X; Y)$. For any $m > 1$, $k \in \mathbb{N}$ and $x \in X$ we have*

$$\|f(x^{m^k}) - m^k f(x)\| \leq c_m(1 + m + m^2 + \cdots + m^{k-1}) \tag{2.13}$$

and

$$\left\| \frac{1}{m^k} f(x^{m^k}) - f(x) \right\| \leq c_m. \tag{2.14}$$

Proof. The proof will be based on mathematical induction for k . If $k = 1$, then (2.13) follows from (2.9). Suppose that (2.13) for k is true, let us verify it for $k + 1$. Replacing x by x^m in (2.13), we get

$$\|f(x^{m^{k+1}}) - m^k f(x^m)\| \leq c_m(1 + m + m^2 + \cdots + m^{k-1}). \tag{2.15}$$

Now, by (2.9) we get

$$\|m^k f(x^m) - m^{k+1} f(x)\| \leq c_m m^k \tag{2.16}$$

and hence

$$\|f(x^{m^{k+1}}) - m^{k+1} f(x)\| \leq c_m(1 + m + \cdots + m^k). \tag{2.17}$$

This implies

$$\left\| \frac{1}{m^{k+1}} f(x^{m^{k+1}}) - f(x) \right\| \leq c_m(1 + m + \cdots + m^k) \frac{1}{m^{k+1}} \leq c_m. \tag{2.18}$$

This completes the proof of the lemma. \square

From (2.14) it follows that the set

$$\left\{ \frac{1}{m^k} f(x^{m^k}) \mid k \in \mathbb{N} \right\}$$

is bounded.

Replacing x by x^{m^n} in (2.14), we obtain

$$\left\| \frac{1}{m^k} f(x^{m^{n+k}}) - f(x^{m^n}) \right\| \leq c_m,$$

$$\left\| \frac{1}{m^{n+k}} f(x^{m^{n+k}}) - \frac{1}{m^n} f(x^{m^n}) \right\| \leq \frac{c_m}{m^n} \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Then, the sequence

$$\left\{ \frac{1}{m^k} f(x^{m^k}) \mid k \in \mathbb{N} \right\}$$

is a Cauchy sequence. Since the space Y is complete, the Cauchy sequence has a limit and denoted by $\varphi_m(x)$. That is,

$$\varphi_m(x) = \lim_{k \rightarrow \infty} \frac{1}{m^k} f(x^{m^k}).$$

By (2.14),

$$\|\varphi_m(x) - f(x)\| \leq c_m, \quad \forall x \in X.$$

Lemma 2.5. *Let $f \in KJ_\rho(X; Y)$ such that*

$$\|f(xy) + f(xy^{-1}) - 2f(x)\| \leq \|\rho(f(xy) - f(x) - f(y))\| + c$$

for all $x, y \in X$. Then for any $m \in \mathbb{N}$, $\varphi_m \in KJ_\rho(X; Y)$.

Proof.

$$\begin{aligned} & \|\varphi_m(xy) + \varphi_m(xy^{-1}) - 2\varphi_m(x)\| \\ &= \lim_{k \rightarrow \infty} \frac{1}{m^k} \|f((xy)^{m^k}) + f((xy^{-1})^{m^k}) - 2f(x^{m^k})\| \\ &\leq \lim_{k \rightarrow \infty} \frac{1}{m^k} \|\rho(f((xy)^{m^k}) - f(x^{m^k}) - f(y^{m^k}))\| + c \\ &= \|\varphi_m(xy) - \varphi_m(x) - \varphi_m(y)\| + c \end{aligned}$$

for all $x, y \in X$. □

For all $x \in X$ we get the equation

$$\varphi_m(x^{m^k}) = m^k \varphi_m(x). \quad (2.19)$$

In fact,

$$\begin{aligned} \varphi_m(x^{m^k}) &= \lim_{l \rightarrow \infty} \frac{1}{m^l} f((x^{m^k})^{m^l}) \\ &= \lim_{l \rightarrow \infty} \frac{m^k}{m^{k+l}} f(x^{m^{k+l}}) \\ &= m^k \lim_{p \rightarrow \infty} \frac{1}{m^p} f(x^{m^p}) = m^k \varphi_m(x). \end{aligned} \quad (2.20)$$

Lemma 2.6. *If $f \in KJ_\rho(X; Y)$, then $\varphi_2 = \varphi_m$ for any $m \geq 2$.*

Proof. By the definition of φ_2 and φ_m , we can see that φ_2 and φ_m belong to $KJ_\rho(X; Y)$. We can define the function

$$g(x) = \lim_{k \rightarrow \infty} \frac{1}{m^k} \varphi_2(x^{m^k}).$$

Then $g(x^{m^k}) = m^k g(x)$ and $g(x^{2^k}) = 2^k g(x)$ for all $x \in X$ and $n \in \mathbb{N}$. Then there are $d_1, d_2 \in \mathbb{R}_+$ such that

$$\|\varphi_2(x) - g(x)\| \leq d_1 \quad \text{and} \quad \|\varphi_m(x) - g(x)\| \leq d_2.$$

Thus $\varphi_2 = \varphi_m$. □

We denote by $B(X; Y)$ the space of all bounded functions on group X that take values in Y . By $PJ_\rho(X; Y)$ the set of $(X; Y)$ -pseudo- ρ -Jensen functions, i.e., $(X; Y)$ -quasi- ρ -Jensen functions f such that $f(x^n) = n f(x)$ for all $x \in X$ and $n \in \mathbb{N}$.

Theorem 2.7. *For an arbitrary group G the following decomposition holds:*

$$KJ_\rho(X; Y) = PJ_\rho(X; Y) \oplus B(X; Y). \quad (2.21)$$

Proof. It is clear that $PJ_\rho(X; Y)$ and $B(X; Y)$ are subspaces of $KJ_\rho(X; Y)$, and

$$PJ_\rho(X; Y) \cap B(X; Y) = \{0\}.$$

Now we only need to prove that $PJ_\rho(X; Y) \oplus B(X; Y) \subseteq KJ_\rho(X; Y)$. In fact, if $f \in KJ_\rho(X; Y)$, then we can define the function $\widehat{f} := \lim_{k \rightarrow \infty} \frac{1}{2^k} f(x^{2^k})$, and it is easy to show that $\widehat{f}(x) = \varphi_m(x)$. Hence $\|\widehat{f}(x) - f(x)\| = \|\varphi_2(x) - f(x)\| \leq c_2$. Thus, we have $\widehat{f} \in PJ_\rho(X; Y)$ and $\widehat{f} - f \in B(X; Y)$. □

3. ULAM STABILITY

Suppose that G is a group and E is a real Banach space.

Definition 3.1. We say that the inequality (2.1) is stable for the pair $(X; Y)$ if for any $f : X \rightarrow Y$ satisfying functional inequality

$$\|f(xy) + f(xy^{-1}) - 2f(x)\| \leq \|f(xy) - f(x) - f(y)\| + c \quad (3.1)$$

for some $c > 0$ there is a solution j of the functional equation (2.1) such that $j(x) - f(x) \in B(X; Y)$.

Next, the inequality (2.1) is stable on X if and only if $PJ_\rho(X; Y) = J_{\rho_0}(X; Y)$.

Theorem 3.2. *The inequality (2.1) is stable if and only if $PJ_\rho(X; Y) = J_{\rho_0}(X; Y)$.*

Proof. Suppose $PJ_\rho(X; Y) = J_{\rho_0}(X; Y)$. Let f satisfy the inequality (2.1) for some c . By Lemma 2.5 and Theorem 2.7, there is a function $\widehat{f} \in PJ_\rho(X; Y)$ such that $f - \widehat{f}$ is a bounded function on X .

Now suppose $PJ_\rho(X; Y) \neq J_{\rho_0}(X; Y)$. Then we will show that the inequality (2.2) is not stable. Let $f \in PJ_\rho(X; Y) \setminus J_{\rho_0}(X; Y)$, by Theorem 2.7, there are $f \in PJ_\rho(X; Y)$ and $h \in B(X; Y)$ such that $f(x) = g(x) + h(x)$. It follows that for any $n, k \in \mathbb{N}$, we get

$$\begin{aligned} f(x^{n^k}) &= g(x^{n^k}) + h(x^{n^k}), \\ n^k f(x) &= n^k g(x) + h(x^{n^k}), \\ f(x) &= g(x) + \frac{1}{n^k} h(x^{n^k}). \end{aligned}$$

Thus, we can see that $f(x) = g(x)$ for all $x \in X$. It is a contradiction to the assumption on f . So $PJ_\rho(X; Y) = J_{\rho_0}(X; Y)$. \square

DECLARATIONS

Availability of data and materials

Not applicable.

Conflict of interest

The authors declare that they have no competing interests.

Fundings

This work was supported by National Natural Science Foundation of China (No. 11761074), Project of Jilin Science and Technology Development for Leading Talent of Science and Technology Innovation in Middle and Young and Team Project(No.20200301053RQ), and the scientific research project of Guangzhou College of Technology and Business in 2020 (No. KA202032).

Acknowledgements

Not applicable.

Authors' contributions

The authors equally conceived of the study, participated in its design and coordination, drafted the manuscript, participated in the sequence alignment, and read and approved the final manuscript.

REFERENCES

1. J. Aczel, J. Jung and C. Ng, *Symmetric second differences in product form on groups*, in: Topics in Mathematical Analysis, Th. M. Rassias (ed.), 1989, pp. 1–22.
2. L. Aiemsomboon and W. Sintunavarat, *Stability of the generalized logarithmic functional equations arising from fixed point theory*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Math. RACSAM **112** (2018), 229–238.
3. T. Aoki, *On the stability of the linear transformation in Banach spaces*, J. Math. Soc. Japan **2** (1950), 64–66.
4. L. Cădariu and V. Radu, *Fixed points and the stability of Jensen's functional equation*, J. Inequal. Pure Appl. Math. **4** (2003), no. 1, Article ID 4.
5. Y. Cho, C. Park and R. Saadati, *Functional inequalities in non-Archimedean Banach spaces*, Appl. Math. Lett. **23** (2010), 1238–1242.
6. Y. Cho, R. Saadati and Y. Yang, *Approximation of homomorphisms and derivations on Lie C^* -algebras via fixed point method*, J. Inequal. Appl. **2013**, 2013:415.
7. P. W. Cholewa, *Remarks on the stability of functional equations*, Aequationes Math. **27** (1984), 76–86.
8. J. Chung, B. Ebanks, C. Ng and P. Sahoo, *On a quadratic-trigonometric functional equation and some applications*, Trans. Am. Math. Soc. **347** (1995), 1131–1161.
9. K. Ciepliński, *Applications of fixed point theorems to the Hyers-Ulam stability of functional equations—a survey*, Ann. Funct. Anal. **3** (2012), 151–164.
10. P. Găvruta, *A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings*, J. Math. Anal. Appl. **184** (1994), 431–436.

11. D.H. Hyers, *On the stability of the linear functional equation*, Proc. Nat. Acad. Sci. USA **27** (1941), 222–224.
12. S. Jung, D. Popa and M. Th. Rassias, *On the stability of the linear functional equation in a single variable on complete metric spaces*, J. Global Optim. **59** (2014), 13–16.
13. Y. Lee, S. Jung and M. Th. Rassias, *Uniqueness theorems on functional inequalities concerning cubic-quadratic-additive equation*, J. Math. Inequal. **12** (2018), 43–61.
14. G. Lu and C. Park, *Hyers-Ulam stability of additive set-valued functional equations*, Appl. Math. Lett. **24** (2011), 1312–1316.
15. G. Lu and C. Park, *Hyers-Ulam stability of general Jensen-type mappings in Banach algebras*, Results Math. **66** (2014), 385–404.
16. C. Ng, *Jensen's functional equation on groups*, Aequationes Math. **39** (1999) 85–99.
17. C. Park, *Fixed points and Hyers-Ulam-Rassias stability of Cauchy-Jensen functional equations in Banach algebras*, Fixed Point Theory Appl. **2007**, Article ID 50175 (2007).
18. C. Park, *Additive ρ -functional inequalities and equations*, J. Math. Inequal. **9** (2015), 17–26.
19. C. Park, Y. Cho and M. Han, *Functional inequalities associated with Jordan-von Neumann type additive functional equations*, J. Inequal. Appl. **2007**, Article ID 41820 (2007).
20. C. Park, J. Lee and X. Zhang, *Additive s -functional inequality and hom-derivations in Banach algebras*, J. Fixed Point Theory Appl. **21** (2019), 2019:18.
21. Th. M. Rassias, *On the stability of the linear mapping in Banach spaces*, Proc. Am. Math. Soc. **72** (1978), 297–300.
22. S. M. Ulam, *Problems in Modern Mathematics*, Wiley, New York, 1940.
23. Z. Wang, *Stability of two types of cubic fuzzy set-valued functional equations*, Results Math. **70** (2016), 1–14.

GANG LU

DIVISION OF FOUNDATIONAL TEACHING, GUANGZHOU COLLEGE OF TECHNOLOGY AND BUSINESS,
GUANGZHOU 510850, P.R. CHINA

Email address: lvgang1234@163.com

LULU QU

ZHUHAI CITY POLYTECHNIC, ZHUHAI, 519090, P.R. CHINA

Email address: monaqu@126.com

YUANFENG JIN

DEPARTMENT OF MATHEMATICS, YANBIAN UNIVERSITY, YANJI 133001, P.R. CHINA

Email address: yfkim@ybu.edu.cn

CHOONKIL PARK

RESEARCH INSTITUTE FOR NATURAL SCIENCES, HANYANG UNIVERSITY, SEOUL 04763, KOREA

Email address: baak@hanyang.ac.kr