

The Nonexistence of Vortices for Rotating Bose-Einstein Condensates in Non-Radially Symmetric Traps

Yujin Guo*

October 28, 2021

Abstract

As a continuation of [34], we consider ground states of rotating Bose-Einstein condensates with attractive interactions in non-radially harmonic traps $V(x) = x_1^2 + \Lambda^2 x_2^2$, where $0 < \Lambda \neq 1$ and $x = (x_1, x_2) \in \mathbb{R}^2$. For any fixed rotational velocity $0 \leq \Omega < \Omega^* := 2 \min\{1, \Lambda\}$, it is known that ground states exist if and only if $a < a^*$ for some critical constant $0 < a^* < \infty$, where $a > 0$ denotes the product for the number of particles times the absolute value of the scattering length. We analyze the asymptotic expansions of ground states as $a \nearrow a^*$, which display the visible effect of Ω on ground states. As a byproduct, we further prove that ground states do not have any vortex in the region $R(a) := \{x \in \mathbb{R}^2 : |x| \leq C(a^* - a)^{-\frac{1}{2}}\}$ as $a \nearrow a^*$ for some constant $C > 0$, which is independent of $0 < a < a^*$.

Keywords: Bose-Einstein condensate; rotational velocity; nonexistence of vortices; limit profiles

1 Introduction

Bose-Einstein condensate (BEC) is a state of matter, in which atoms or particles are cooled to the sufficiently low temperature that a large fraction of them “condense” into a single quantum state. The BECs in magneto-optical traps present remarkable phenomena, once the traps are set in the rotational motion. Actually, starting from the first physical achievement of rotating BECs in the late 1990s, various interesting quantum phenomena have been observed in the experiments of rotating BECs, including the critical-mass collapse [16, 24, 37], the center-of-mass rotation [1, 26, 48], and the appearance of quantized vortices [2, 18, 26]. Therefore, numerical simulations and mathematical theories of rotating BECs have been a focus of international interest in physics and mathematics over the past two decades, see [1, 2, 6, 23, 24, 26, 38, 39, 45–47]. We also refer [2, 14, 22, 57, 58] for the mathematics of the relevant superfluids and superconductors.

The interactions between cold atoms in the condensates can be either repulsive or attractive, cf. [2, 15, 24, 26]. For the repulsive case, the complex structures, including the quantized vortices, of the rotating trapped BECs were analyzed and simulated extensively in the past few years, see [1–4, 9, 23, 24, 26, 38, 39, 50, 51] and the references therein. However, the rotating trapped BECs in the attractive case behave different extremely

*School of Mathematics and Statistics, and Hubei Key Laboratory of Mathematical Sciences, Central China Normal University, P.O. Box 71010, Wuhan 430079, P. R. China. Email: yguo@ccnu.edu.cn. Y. J. Guo is partially supported by NSFC under Grant 11931012.

from those in the well-understood repulsive case. Typically, the vortices are generally unstable in the attractive case (cf. [18, 48]), even though the vortices may form stable lattice configurations in the repulsive case, cf. [2, 26]. Because of distinct mechanisms, existing physical observations and numerical simulations show that rotating trapped BECs with attractive interactions present more complicated phenomena and structures, see [9, 18, 24, 26, 48], only few of which have however been investigated analytically so far.

As derived rigorously in [42, 43, 52, 55, 56] by a mean-field approximation, the ground state of two-dimensional attractive BECs in a rotating trap can be described equivalently by a complex constraint minimizer of the following Gross-Pitaevskii (GP) energy functional

$$F_a(u) := \int_{\mathbb{R}^2} (|\nabla u|^2 + V(x)|u|^2) dx - \frac{a}{2} \int_{\mathbb{R}^2} |u|^4 dx - \Omega \int_{\mathbb{R}^2} x^\perp \cdot (iu, \nabla u) dx, \quad u \in \mathcal{H}, \quad (1.1)$$

under the mass constraint

$$e_F(a) := \inf_{\{u \in \mathcal{H}, \|u\|_2^2=1\}} F_a(u), \quad a > 0, \quad (1.2)$$

where $x^\perp = (-x_2, x_1)$ with $x = (x_1, x_2) \in \mathbb{R}^2$, $(iu, \nabla u) = i(u\nabla\bar{u} - \bar{u}\nabla u)/2$, and the complex space \mathcal{H} is defined as

$$\mathcal{H} := \left\{ u \in H^1(\mathbb{R}^2, \mathbb{C}) : \int_{\mathbb{R}^2} V(x)|u|^2 dx < \infty \right\}. \quad (1.3)$$

The parameter $a > 0$ in $e_F(a)$ denotes the product for the number N of particles times the absolute value of the scattering length ν in the two-body interaction, and while $\Omega \geq 0$ describes the rotational velocity of the rotating trap $V(x) \geq 0$. We comment that one may impose $e_F(a)$ a different constraint $\int_{\mathbb{R}^2} |u(x)|^2 dx = N > 0$, but these two different forms can be reduced equivalently to each other, see [34]. In this paper we therefore focus on the form of $e_F(a)$ instead. It also deserves to remark that even though the mass-subcritical version of $e_F(a)$, where the nonlinear term $|u|^4$ of $F_a(u)$ is replaced by $|u|^p$ for $2 < p < 4$, was studied as early as in the pioneering work of Esteban-Lions [25], the mass-critical constraint variational problem $e_F(a)$ in the complex range was not addressed until recent years, see [7, 9, 13, 32, 34, 43] and the references therein.

The non-rotational case $\Omega = 0$ of $e_F(a)$ was studied recently in [31, 35, 36, 49, 61] and the references therein, where the existence, uniqueness, symmetry breaking and other analytical properties of complex-valued minimizers were investigated equivalently in view of the argument [21, Theorem II.1]. For a class of trapping potentials $V(x)$, above mentioned works give implicitly that $e_F(a)$ with $\Omega = 0$ admits complex-valued minimizers if and only if $a < a^*$, where $a^* = \|w\|_{L^2(\mathbb{R}^2)}^2$ and $w = w(|x|) > 0$ is the unique (cf. [41, 60]) positive solution of the following nonlinear scalar field equation

$$\Delta u - u + u^3 = 0 \quad \text{in } \mathbb{R}^2, \quad u \in H^1(\mathbb{R}^2, \mathbb{R}). \quad (1.4)$$

Starting from earlier works [9, 43], the rotational case $\Omega > 0$ of $e_F(a)$ was analyzed more recently. More precisely, considering special trapping potentials, such as typically $V(x) = |x|^2$, the existence and nonexistence, stability and some other properties of complex-valued minimizers for $e_F(a)$ with $\Omega > 0$ were studied in [7, 9, 13, 43]. Generally, if the trapping potential $0 \leq V(x) \in L_{loc}^\infty(\mathbb{R}^2)$ satisfies

$$\underline{\lim}_{|x| \rightarrow \infty} \frac{V(x)}{|x|^2} > 0, \quad (1.5)$$

then one can define as in [34] the following critical rotational velocity $\Omega^* := \Omega^*(V)$:

$$\Omega^* := \sup \left\{ \Omega > 0 : V(x) - \frac{\Omega^2}{4}|x|^2 \rightarrow \infty \text{ as } |x| \rightarrow \infty \right\}. \quad (1.6)$$

Note that if $V(x)$ satisfies the assumption (1.5), then $\Omega^* \in (0, +\infty]$ exists, and $V_\Omega(x) := V(x) - \frac{\Omega^2}{4}|x|^2 \geq 0$ holds in \mathbb{R}^2 for any $0 \leq \Omega < \Omega^*$. Under the assumption (1.5), we established in [34, Theorem 1.1] the following existence and non-existence of complex-valued minimizers:

Theorem A. (*[34, Theorem 1.1]*) *Assume $V(x) \in L^\infty_{\text{loc}}(\mathbb{R}^2)$ satisfies (1.5) such that $\Omega^* \in (0, +\infty]$ in (1.6) exists. Then we have*

1. *If $0 \leq \Omega < \Omega^*$ and $0 \leq a < a^* := \|w\|_2^2$, then there exists at least one minimizer of $e_F(a)$.*
2. *If $0 \leq \Omega < \Omega^*$ and $a \geq a^* := \|w\|_2^2$, then there is no minimizer of $e_F(a)$.*
3. *If $\Omega > \Omega^*$, then for any $a \geq 0$, there is no minimizer of $e_F(a)$.*

The proof of Theorem A needs the following Gagliardo-Nirenberg inequality

$$\int_{\mathbb{R}^2} |u(x)|^4 dx \leq \frac{2}{\|w\|_2^2} \int_{\mathbb{R}^2} |\nabla u(x)|^2 dx \int_{\mathbb{R}^2} |u(x)|^2 dx, \quad u \in H^1(\mathbb{R}^2, \mathbb{R}), \quad (1.7)$$

where the identity is attained at w , and the following diamagnetic inequality

$$|\nabla u|^2 - \Omega x^\perp \cdot (iu, \nabla u) = |(\nabla - i\mathcal{A})u|^2 - \frac{\Omega^2}{4}|x|^2|u|^2 \geq |\nabla|u||^2 - \frac{\Omega^2}{4}|x|^2|u|^2, \quad u \in H^1(\mathbb{R}^2, \mathbb{C}), \quad (1.8)$$

where $\mathcal{A} = \frac{\Omega}{2}x^\perp$, see [44, 60] for more details. By the variational theory, if $e_F(a)$ admits a minimizer u_a , then u_a is a ground state of the following Euler-Lagrange equation

$$-\Delta u_a + V(x)u_a + i\Omega(x^\perp \cdot \nabla u_a) = \mu u_a + a|u_a|^2 u_a \quad \text{in } \mathbb{R}^2, \quad \int_{\mathbb{R}^2} |u_a|^2 dx = 1, \quad (1.9)$$

where $\mu = \mu(a, \Omega, u_a) \in \mathbb{R}$ is a suitable Lagrange multiplier. We comment that there exist many interesting progresses on normalized solutions of the elliptic problem (1.9), see [8, 10–12, 17, 25, 30, 40, 49, 54] and the references therein.

By employing energy estimates and elliptic PDE theory, it was proved in [34, 43] that the minimizer u_a of $e_F(a)$ concentrates at a global minimum point of $V_\Omega(x)$ as $a \nearrow a^*$, in the sense that

$$\|u_a\|_\infty \rightarrow \infty \quad \text{and} \quad \int_{\mathbb{R}^2} V_\Omega(x)|u_a|^2 dx \rightarrow V_\Omega(x_0) := \inf_{x \in \mathbb{R}^2} V_\Omega(x) \quad \text{as } a \nearrow a^*. \quad (1.10)$$

Based on (1.10), the L^∞ uniform convergence of u_a after rescaling and translation was also obtained in [34]. By developing the method of inductive symmetry, we further proved in [34, Theorem 1.3] (see also [33]) the absence of vortices of minimizers u_a for $e_F(a)$ as $a \nearrow a^*$ in the radially symmetric case $V(x) = |x|^2$, where the rotational velocity Ω presents essentially no effect on u_a and $\text{Im}(u_a e^{i\delta_a}) \equiv 0$ as $a \nearrow a^*$ for some constant phase $\delta_a \in [0, 2\pi)$. We should emphasize that the arguments of [34, Theorem 1.3] cannot however be extended to the non-radially symmetric case of $V(x)$, since one cannot expect generally $\text{Im}(u_a e^{i\delta_a}) \equiv 0$ as $a \nearrow a^*$ for some constant phase $\delta_a \in [0, 2\pi)$, see also Remark 2.1 below.

1.1 Main Results

As a continuation of [34], it is natural to ask whether the rotational velocity Ω has some visible effect on the minimizers u_a of $e_F(a)$ as $a \nearrow a^*$, provided that the trap $V(x)$ is non-radially symmetric. The main purpose of this paper is to address the above question.

Due to the physical relevance (cf. [3, 19, 38, 50, 51]), in this paper we mainly focus on the non-radially harmonic trap $V(x) \geq 0$ of the form

$$V(x) = x_1^2 + \Lambda^2 x_2^2 \quad \text{for } x = (x_1, x_2) \in \mathbb{R}^2, \quad \text{where } 0 < \Lambda < 1, \quad (1.11)$$

since the case $\Lambda > 1$ can be established similarly. As mentioned before, the case $\Lambda = 1$ of (1.11) was already addressed in Theorem 1.3 and (1.18) of [34]. We comment that Theorems 1.1 and 1.2 below can be extended to the general case where the trap $V(x)$ is homogeneous of degree 2, see (2.1) for its definition, and however for simplicity we do not pursue such extensions in this paper.

Under the assumption (1.11), one can note from (1.6) that $e_F(a)$ admits the critical velocity $\Omega^* := 2\Lambda$. For any fixed $0 < \Omega < \Omega^* := 2\Lambda < 2$, we further have

$$(0, 0) \text{ is the unique and non-degenerate critical point of} \\ H_\Omega(y) := \int_{\mathbb{R}^2} V_\Omega(x+y)w^2(x)dx, \quad V_\Omega(x) := V(x) - \frac{\Omega^2}{4}|x|^2 \geq 0, \quad (1.12)$$

and

$$V_\Omega(x) + \frac{\Omega^2}{4}|x|^2 = V(x) = x_1^2 + \Lambda^2 x_2^2. \quad (1.13)$$

Under the assumption (1.11), we thus note from [32, Theorem 1.1] that up to the constant phase, complex-valued minimizers of $e_F(a)$ must be unique as $a \nearrow a^*$. Define

$$\lambda_0 = \left(\int_{\mathbb{R}^2} V(x)w^2 dx \right)^{\frac{1}{4}} > 0. \quad (1.14)$$

For convenience, we denote $\varphi_i(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ to be the unique solution of

$$\nabla \varphi_i(0) = 0, \quad (-\Delta + 1 - 3w^2)\varphi_i(x) = \tilde{f}_i(x) \quad \text{in } \mathbb{R}^2, \quad i = 1, 2, \quad (1.15)$$

where $\tilde{f}_i(x)$ satisfies

$$\tilde{f}_i(x) = \begin{cases} -\frac{\lambda_0^4}{a^*}w^3(x) - (x_1^2 + \Lambda^2 x_2^2)w(x), & \text{if } i = 1; \\ -(1 - \Lambda^2)x_1^2 w(x), & \text{if } i = 2. \end{cases} \quad (1.16)$$

Here the uniqueness of $\varphi_i(x)$ follows from [27, Theorem 1.2] and [59, Lemma 4.1] together with the fact $\nabla \varphi_i(0) = 0$ for $i = 1, 2$. We also denote $\varphi_I(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ to be the unique solution of

$$(-\Delta + 1 - w^2)\varphi_I(x) = -(x^\perp \cdot \nabla \varphi_2) \quad \text{in } \mathbb{R}^2, \quad \int_{\mathbb{R}^2} \varphi_I w dx = 0, \quad (1.17)$$

where φ_2 satisfies (1.15). Here the uniqueness of $\varphi_I(x)$ follows from [59, Lemma 4.1] and the fact $\int_{\mathbb{R}^2} \varphi_I w dx = 0$. Applying above notations, *the main result of this paper* is concerned with the following asymptotic expansion as $a \nearrow a^*$, which displays the visible effect of the rotational velocity Ω :

Theorem 1.1. *Suppose that the non-radially harmonic trap $V(x)$ satisfies (1.11) for some $0 < \Lambda < 1$. Then for any fixed $0 < \Omega < \Omega^* := 2\Lambda$, the complex-valued minimizer u_a of $e_F(a)$ satisfies*

$$v_a := \epsilon_a \sqrt{a^*} u_a(\epsilon_a x + x_a) e^{-i\left(\frac{\Omega}{2}\epsilon_a x \cdot x_a^\perp - \theta_a\right)} = w + \epsilon_a^4 \{\varphi_1 + C_\Lambda(w + x \cdot \nabla w)\} + i\Omega \epsilon_a^6 \varphi_I + o(\epsilon_a^6) \text{ in } L^\infty(\mathbb{R}^2, \mathbb{C}) \quad (1.18)$$

as $a \nearrow a^*$, where $\epsilon_a := \frac{(a^* - a)^{\frac{1}{4}}}{\lambda_0} > 0$, x_a is the unique global maximum point of $|u_a|$ and satisfies

$$|x_a| = o(\epsilon_a^5) \text{ as } a \nearrow a^*, \quad (1.19)$$

$\theta_a \in [0, 2\pi)$ is a suitable constant phase, and the constant $C_\Lambda \neq 0$ is independent of Ω and satisfies

$$C_\Lambda = \frac{1}{2\lambda_0^4} \left[\int_{\mathbb{R}^2} (3w^2 - 1)\varphi_1^2 - 4 \int_{\mathbb{R}^2} (x_1^2 + \Lambda^2 x_2^2) w \varphi_1 \right] \neq 0. \quad (1.20)$$

Here φ_1 and φ_I are uniquely given by (1.15)-(1.17).

Remark 1.1. (1). Under the assumption (1.11), the interesting novelty of Theorem 1.1 lies in the fact that the rotational velocity Ω affects visibly the minimizer u_a starting from the third term, which is imaginary, of v_a defined in (1.18). This is however different from the radial case $V(x) = |x|^2$ addressed earlier in [34, Theorem 1.3], where $\Omega > 0$ has essentially no effect on u_a .

(2). Specially, if $V(x) = |x|^2$, then the argument of Theorem 1.1 can yield that (1.18) holds for $\varphi_I \equiv 0$, and while the term $\varphi_1 + C_\Lambda(w + x \cdot \nabla w)$ is radially symmetric and independent of $\Omega > 0$. Following these, we guess that one may further obtain the non-existence of vortices for u_a in \mathbb{R}^2 as $a \nearrow a^*$, which approach is different slightly from that of [34]. We leave it to the interested reader.

(3). We expect that, instead of (1.19), the unique global maximum point x_a of $|u_a|$ satisfies $x_a \equiv 0$ as $a \nearrow a^*$. But our analysis does not give this conclusion, for which one needs to derive further refined estimates of u_a as $a \nearrow a^*$.

We next follow three steps to explain the general strategy of proving Theorem 1.1. As the first step, we proved in [34, Theorem 1.2] that the complex-valued minimizer u_a of $e_F(a)$ satisfies the equation (1.9) and

$$v_a(x) := \epsilon_a \sqrt{a^*} u_a(\epsilon_a x + x_a) e^{-i\left(\frac{\Omega}{2}\epsilon_a x \cdot x_a^\perp - \theta_a\right)} := [R_a(x) + w(x)] + iI_a(x) \rightarrow w(x) \quad (1.21)$$

uniformly in $L^\infty(\mathbb{R}^2, \mathbb{C})$ as $a \nearrow a^*$, where and below $\epsilon_a > 0$ and $x_a \in \mathbb{R}^2$ are as in Theorem 1.1, and $\theta_a \in [0, 2\pi)$ is chosen suitably. Under the assumption (1.11), recall from [34, Section 3] that the Lagrange multiplier μ_a of (1.9) and x_a satisfy

$$\mu_a \epsilon_a^2 \rightarrow -1 \text{ and } \frac{x_a}{\epsilon_a} \rightarrow 0 \text{ as } a \nearrow a^*. \quad (1.22)$$

Note from (1.21) that R_a and I_a satisfy

$$R_a(x) \rightarrow 0 \text{ and } I_a(x) \rightarrow 0 \text{ uniformly in } \mathbb{R}^2 \text{ as } a \nearrow a^*. \quad (1.23)$$

The second step of proving Theorem 1.1 is to establish the leading terms of $R_a(x) + iI_a(x)$, in terms of ϵ_a and $\mu_a \epsilon_a^2 + 1$, by following (1.21) and (1.22). However, we remark

that it seems difficult to reach this aim by investigating directly the Euler-Lagrange equation (1.9). To overcome this difficulty, motivated by (1.9) and (1.21) we shall consider the following coupled system of $R_a(x)$ and $I_a(x)$:

$$\begin{cases} (\mathcal{L}_a - w^2 - w(R_a + w))R_a = \epsilon_a^2 \Omega (x^\perp \cdot \nabla I_a) + \tilde{F}_a(x) & \text{in } \mathbb{R}^2, \quad \nabla R_a(0) \neq 0, \\ \mathcal{L}_a I_a = -\epsilon_a^2 \Omega (x^\perp \cdot \nabla R_a) & \text{in } \mathbb{R}^2, \quad \int_{\mathbb{R}^2} w I_a dx \equiv 0, \end{cases} \quad (1.24)$$

where $\tilde{F}_a(x)$ is an inhomogeneous term containing $\mu_a \epsilon_a^2 + 1$, and the operator \mathcal{L}_a is as in (2.23).

Even though the rest part of the second step is motivated by [31, Theorem 1.4] and (1.23), unfortunately, there appear extra difficulties. Actually, we first note from (1.24) that we have $R_a(0) \neq 0$ in \mathbb{R}^2 , which leads to a new difficulty in studying the expansion of R_a , see Lemma 2.2. Moreover, the coupled system (1.24) contains the coupled rotating terms $(x^\perp \cdot \nabla I_a)$ and $(x^\perp \cdot \nabla R_a)$. To overcome these extra difficulties, we shall proceed with the refined estimates of R_a and I_a , and make full use of the non-degeneracy of w as well. We shall finally obtain in Lemmas 2.3 and 3.2 the leading terms of I_a and R_a , respectively, in terms of ϵ_a and $\mu_a \epsilon_a^2 + 1$.

The third step of proving Theorem 1.1 is to address the refined estimate of $\mu_a \epsilon_a^2 + 1$ in terms of $\epsilon_a > 0$. We shall achieve it by taking full advantage of the mass constraint $\int_{\mathbb{R}^2} |u_a|^2 dx = 1$, as well as analytical results of the second step.

Consider the non-radially harmonic trap $V(x)$ satisfying (1.11). It then follows from [34, Theorem 1.2] that the minimizers do not have any vortex near the origin as $a \nearrow a^*$. As a byproduct of Theorem 1.1, we shall derive the following nonexistence of vortices under the assumption (1.11).

Theorem 1.2. *Suppose the non-radially harmonic trap $V(x)$ satisfies (1.11), and let u_a be a complex-valued minimizer of $e_F(a)$, where $0 < \Omega < \Omega^* := 2\Lambda$ is fixed. Then there exists a constant $C > 0$, independent of $0 < a < a^*$, such that*

$$|u_a(x)| > 0 \text{ in the region } R(a) := \{x \in \mathbb{R}^2 : |x| \leq C(a^* - a)^{-\frac{1}{12}}\} \text{ as } a \nearrow a^*, \quad (1.25)$$

i.e., u_a does not admit any vortex in the region $R(a)$ as $a \nearrow a^$.*

Remark 1.2. (1). The nonexistence of vortices for rotating trapped BECs was studied earlier by jacobian estimates, vortex ball constructions, the inductive symmetry, and some other methods, see [3–5, 19, 23, 33, 34, 38, 39, 57] and the references therein. However, as far as we know, above mentioned methods depend heavily on the radial symmetry of $V(x)$.

(2). Theorem 1.2 proves the nonexistence of vortices in a very large region $R(a)$ as $a \nearrow a^*$, including the places where $|u_a|$ is already very small. As remarked before, the argument of Theorem 1.2 holds actually for *the non-radially general case* where the trap $V(x)$ is homogeneous of degree 2, see (2.1) for its definition. This is the main contribution of Theorem 1.2, since the existing methods of [4, Theorem 1.1] and [34, Theorem 1.3] and the references therein are not applicable to the non-radially symmetric case of $V(x)$.

To prove Theorem 1.2 with the non-radially symmetric trap $V(x)$, we shall make full use of Theorem 1.1 to establish the following optimal estimate

$$|v_a(x) - w(x)| \leq C_1 \epsilon_a^4 |x|^{\frac{5}{2}} e^{-\sqrt{1-C_2 \epsilon_a^4} |x|} \text{ uniformly in } \mathbb{R}^2 \text{ as } a \nearrow a^*, \quad (1.26)$$

where $v_a(x)$ is as in (1.18), and the constants $C_1 > 0$ and $C_2 > 0$ are independent of $0 < a < a^*$. Applying (1.26), Theorem 1.2 then follows from the fact that $|v_a| \geq |w| - |w - v_a|$ holds in \mathbb{R}^2 , due to the exact exponential decay (2.5) of w . We note from above that the idea of proving Theorem 1.2 is different from those of [34, Theorem 1.3] and [34, Theorem 1.3], since the latter ones proved the nonexistence of vortices in the following approach: $v_a(x)$ is essentially a real-valued minimizer of $e_F(a)$ with $\Omega = 0$, and $|v_a(x)| > 0$ then holds in a standard way, see [35, 36].

This paper is organized as follows: In Section 2 we study refined estimates of minimizers u_a for $e_F(a)$ as $a \nearrow a^*$. Applying analytical results of the previous section, in Section 3 we first analyze the refined estimate of $\mu_a \epsilon_a^2 + 1$ in terms of $\epsilon_a > 0$, based on which we then finish the proofs of Theorems 1.1 and 1.2 in Subsection 3.1. In Appendix A we shall address the proof of the claims (3.19) and (3.36) used in Section 3.

2 Refined Estimates of Minimizers

Suppose the rotating speed $\Omega \in (0, \Omega^*)$ is fixed, the purpose of this section is to address the refined estimates of complex-valued minimizers u_a for $e_F(a)$ as $a \nearrow a^*$. To clarify the general idea, in this section we focus on the homogeneous trapping potential $V(x)$, in the sense that a function $h(x) : \mathbb{R}^2 \mapsto \mathbb{R}$ is called homogeneous of degree $p > 0$ (about the origin), if

$$h(tx) = t^p h(x) \text{ for any } t \in \mathbb{R}^+ \text{ and } x \in \mathbb{R}^2. \quad (2.1)$$

We note from above that if $0 \leq V(x) \in C^2(\mathbb{R}^2)$ is homogeneous of degree 2 and satisfies $\lim_{|x| \rightarrow \infty} V(x) = \infty$, such as (1.11), then $x = 0$ is the unique minimum point of $V(x)$, and

$$0 \leq V_\Omega(x) := V(x) - \frac{\Omega^2}{4}|x|^2 \in C^2(\mathbb{R}^2) \text{ is also homogeneous of degree 2} \quad (2.2)$$

for any fixed $0 < \Omega < \Omega^*$, where $0 < \Omega^* < \infty$ is defined as in (1.6).

Recall from [60] that the unique positive radial solution w of (1.4) is an optimizer of the following Gagliardo-Nirenberg inequality

$$\int_{\mathbb{R}^2} |u(x)|^4 dx \leq \frac{2}{a^*} \int_{\mathbb{R}^2} |\nabla u(x)|^2 dx \int_{\mathbb{R}^2} |u(x)|^2 dx, \quad u \in H^1(\mathbb{R}^2, \mathbb{R}). \quad (2.3)$$

Note also from [20, Lemma 8.1.2] and [28, Proposition 4.1] that $w = w(|x|) > 0$ satisfies

$$\int_{\mathbb{R}^2} |\nabla w|^2 dx = \int_{\mathbb{R}^2} w^2 dx = \frac{1}{2} \int_{\mathbb{R}^2} w^4 dx, \quad (2.4)$$

and

$$w(x), |\nabla w(x)| = O(|x|^{-\frac{1}{2}} e^{-|x|}) \text{ as } |x| \rightarrow \infty. \quad (2.5)$$

Define for $0 < a < a^*$,

$$\epsilon_a := \frac{(a^* - a)^{\frac{1}{4}}}{\lambda} > 0, \quad \lambda = \left[\int_{\mathbb{R}^2} \left(V_\Omega(x + y_0) + \frac{\Omega^2}{4}|x|^2 \right) w^2(x) dx \right]^{\frac{1}{4}} > 0, \quad (2.6)$$

where $V_\Omega(x) \geq 0$ is defined by (2.2), and $y_0 \in \mathbb{R}^2$ denotes a unique global minimum point of $H_\Omega(y) := \int_{\mathbb{R}^2} V_\Omega(x + y) w^2(x) dx$. Define

$$v_a(x) := \epsilon_a \sqrt{a^*} u_a(\epsilon_a x + x_a) e^{-i(\frac{\epsilon_a \Omega}{2} x \cdot x_a^\perp - \theta_a)} = \tilde{R}_a(x) + iI_a(x), \quad (2.7)$$

where x_a is a global maximal point of $|u_a(x)|$, $\tilde{R}_a(x)$ and $I_a(x)$ denote the real and imaginary parts of $v_a(x)$, respectively, and the constant phase $\theta_a \in [0, 2\pi)$ is chosen such that

$$\|v_a - w\|_{L^2(\mathbb{R}^2)} = \min_{\theta \in [0, 2\pi)} \|e^{i\theta} \tilde{v}_a - w\|_{L^2(\mathbb{R}^2)}, \quad (2.8)$$

where $\tilde{v}_a := \varepsilon_a \sqrt{a^*} u_a(\varepsilon_a x + x_a) e^{-i \frac{\varepsilon_a \Omega}{2} x \cdot x_a^\perp}$. The above property gives the following orthogonality condition on $I_a(x)$:

$$\int_{\mathbb{R}^2} w(x) I_a(x) dx = 0. \quad (2.9)$$

Using above notations, we proved in [34] the following L^∞ -uniform convergence as $a \nearrow a^*$.

Proposition 2.1. (*[34, Theorem 1.2]*) *Assume $0 \leq V(x) \in C^2(\mathbb{R}^2)$ satisfying $\lim_{|x| \rightarrow \infty} V(x) = +\infty$ is homogeneous of degree 2, and let 0 be a unique global minimum point of $H_\Omega(y) := \int_{\mathbb{R}^2} V_\Omega(x+y) w^2(x) dx$. For any fixed $0 < \Omega < \Omega^*$, where $\Omega^* > 0$ is defined as in (1.6), suppose u_a is a minimizer of $e_F(a)$. Then we have*

$$v_a(x) := \varepsilon_a \sqrt{a^*} u_a(\varepsilon_a x + x_a) e^{-i \left(\frac{\Omega}{2} \varepsilon_a x \cdot x_a^\perp - \theta_a \right)} \rightarrow w(x) \quad (2.10)$$

uniformly in $L^\infty(\mathbb{R}^2, \mathbb{C})$ as $a \nearrow a^*$, where $\varepsilon_a > 0$ is as in (2.6), $\theta_a \in [0, 2\pi)$ is chosen such that (2.8) holds true, and the unique global maximal point $x_a \in \mathbb{R}^2$ of $|u_a|$ satisfies

$$\lim_{a \nearrow a^*} \frac{x_a}{\varepsilon_a} = 0. \quad (2.11)$$

By the variational theory, the minimizer u_a of $e_F(a)$ solves the following Euler-Lagrange equation:

$$-\Delta u_a + V(x) u_a + i \Omega (x^\perp \cdot \nabla u_a) = \mu_a u_a + a |u_a|^2 u_a \quad \text{in } \mathbb{R}^2, \quad (2.12)$$

where $\mu_a := \mu_a(u_a) \in \mathbb{R}$ is a suitable Lagrange multiplier satisfying

$$\mu_a = e_F(a) - \frac{a}{2} \int_{\mathbb{R}^2} |u_a|^4 dx. \quad (2.13)$$

Note from (2.12) that the function v_a satisfies the following elliptic equation

$$\begin{aligned} & -\Delta v_a + i \varepsilon_a^2 \Omega (x^\perp \cdot \nabla v_a) + \left[\frac{\varepsilon_a^4 \Omega^2 |x|^2}{4} + \varepsilon_a^2 V_\Omega(\varepsilon_a x + x_a) \right] v_a \\ & = \mu_a \varepsilon_a^2 v_a + \frac{a}{a^*} |v_a|^2 v_a \quad \text{in } \mathbb{R}^2, \end{aligned} \quad (2.14)$$

where $V_\Omega(x) \geq 0$ is as in (2.2). The analysis of [34, Theorem 1.2] gives that the above Lagrange multiplier μ_a satisfies

$$\lim_{a \nearrow a^*} \mu_a \varepsilon_a^2 = -1. \quad (2.15)$$

In order to obtain the refined estimate of u_a as $a \nearrow a^*$, one can note from (2.10) and (2.14) that a more refined estimate than (2.15) is needed for μ_a as $a \nearrow a^*$, which is one of the main difficulties in this paper. Towards this purpose, it however seems difficult to handle directly with the single equation (2.14), instead of which we shall consider the coupled system of $Re(v_a)$ and $Im(v_a)$ in the coming subsection.

2.1 Refined estimates of v_a as $a \nearrow a^*$

Based on Proposition 2.1, in this subsection we shall derive refined estimates of v_a defined in (2.7) as $a \nearrow a^*$. Towards this purpose, we introduce the following linear operator

$$\mathcal{L} := -\Delta + 1 - w^2 \quad \text{in } \mathbb{R}^2. \quad (2.16)$$

It then obtains from [44, Theorem 11.8] and [44, Corollary 11.9] that

$$\ker \mathcal{L} = \{w\} \quad \text{and} \quad \langle \mathcal{L}v, v \rangle \geq 0 \quad \text{for all } v \in L^2(\mathbb{R}^2). \quad (2.17)$$

We also define the linearized operator $\tilde{\mathcal{L}}$ of (1.4) around $w > 0$ by

$$\tilde{\mathcal{L}} := -\Delta + 1 - 3w^2 \quad \text{in } \mathbb{R}^2. \quad (2.18)$$

It follows from [41, 53, 59] that

$$\ker \tilde{\mathcal{L}} = \left\{ \frac{\partial w}{\partial x_1}, \frac{\partial w}{\partial x_2} \right\}. \quad (2.19)$$

For convenience, we denote for $0 < a < a^*$,

$$\varepsilon_a := \frac{(a^* - a)^{\frac{1}{4}}}{\lambda} > 0, \quad \alpha_a := (\lambda \varepsilon_a)^4 = a^* - a > 0, \quad \text{and} \quad \beta_a := 1 + \mu_a \varepsilon_a^2, \quad (2.20)$$

where $\lambda > 0$ is defined by (2.6) with $y_0 = (0, 0)$ and the Lagrange multiplier μ_a satisfies (2.15). We then have

$$\alpha_a \rightarrow 0 \quad \text{and} \quad \beta_a \rightarrow 0 \quad \text{as } a \nearrow a^*.$$

Following (2.7), we now rewrite v_a as

$$v_a(x) := \tilde{R}_a(x) + iI_a(x) = [R_a(x) + w(x)] + iI_a(x), \quad (2.21)$$

so that

$$R_a(x) \rightarrow 0 \quad \text{and} \quad I_a(x) \rightarrow 0 \quad \text{uniformly in } L^\infty(\mathbb{R}^2, \mathbb{R}) \quad \text{as } a \nearrow a^*,$$

due to Proposition 2.1. Since $\nabla|v_a(0)| \equiv 0$ holds for all $0 < a < a^*$, we derive from (2.21) that

$$\nabla R_a(0) = -\frac{I_a(0)\nabla I_a(0)}{w(0) + R_a(0)} \rightarrow 0 \quad \text{as } a \nearrow a^*. \quad (2.22)$$

For simplicity, we denote the operator \mathcal{L}_a by

$$\mathcal{L}_a := -\Delta + \left(\frac{\varepsilon_a^4 \Omega^2}{4} |x|^2 + \varepsilon_a^2 V_\Omega(\varepsilon_a x + x_a) - \mu_a \varepsilon_a^2 - \frac{a}{a^*} |v_a|^2 \right) \quad \text{in } \mathbb{R}^2. \quad (2.23)$$

It then follows from (2.9), (2.14) and (2.21) that I_a satisfies

$$\mathcal{L}_a I_a = -\varepsilon_a^2 \Omega (x^\perp \cdot \nabla R_a) \quad \text{in } \mathbb{R}^2, \quad \int_{\mathbb{R}^2} w I_a dx \equiv 0, \quad (2.24)$$

and while R_a satisfies (2.22) and

$$\tilde{\mathcal{L}}_a R_a := [\mathcal{L}_a - w^2 - w\tilde{R}_a] R_a = F_a(x) \quad \text{in } \mathbb{R}^2, \quad (2.25)$$

where $F_a(x)$ is defined by

$$\begin{aligned}
F_a(x) &:= \varepsilon_a^2 \Omega(x^\perp \cdot \nabla I_a) - \left[\frac{\varepsilon_a^4 \Omega^2}{4} |x|^2 + \varepsilon_a^2 V_\Omega(\varepsilon_a x + x_a) - \beta_a - \frac{a}{a^*} I_a^2 + \frac{\alpha_a}{a^*} \tilde{R}_a^2 \right] w \\
&= -\varepsilon_a^4 \left[\frac{\Omega^2}{4} |x|^2 + V_\Omega\left(x + \frac{x_a}{\varepsilon_a}\right) \right] w + \beta_a w - \frac{\alpha_a}{a^*} \tilde{R}_a^2 w \\
&\quad + \varepsilon_a^4 \Omega\left(x^\perp \cdot \frac{\nabla I_a}{\varepsilon_a^2}\right) + \frac{a}{a^*} I_a^2 w.
\end{aligned} \tag{2.26}$$

Similar to [34, Lemma 4.2 (1)], where $\varepsilon_a > 0$ is defined in a slightly different way, one can deduce from (2.14) that there exists a constant $C > 0$, independent of $0 < a < a^*$, such that as $a \nearrow a^*$,

$$|\nabla \tilde{R}_a(x)|, |\nabla I_a(x)| \leq C e^{-\frac{2}{3}|x|} \quad \text{uniformly in } \mathbb{R}^2. \tag{2.27}$$

Applying (2.22) and (2.27), the argument of [34, Lemma 4.3] then yields from (2.24) that there exists a constant $C > 0$, independent of $0 < a < a^*$, such that as $a \nearrow a^*$,

$$|\nabla I_a(x)|, |I_a(x)| \leq C \varepsilon_a^2 e^{-\frac{1}{8}|x|} \quad \text{uniformly in } \mathbb{R}^2. \tag{2.28}$$

Setting

$$\psi_2(|x|) := -\frac{1}{2}(w + x \cdot \nabla w),$$

which is independent of $0 < \Omega < \Omega^*$, one can check that ψ_2 is a unique solution of

$$\nabla \psi_2(0) = 0, \quad \tilde{\mathcal{L}} \psi_2(|x|) = w(x) \quad \text{in } \mathbb{R}^2, \tag{2.29}$$

where the operator $\tilde{\mathcal{L}}$ is defined by (2.18). Having above estimates, we next establish the following ‘‘rough’’ limit profiles in terms of ε_a and β_a .

Lemma 2.2. *Under the assumptions of Proposition 2.1, we have*

1. *There exists a constant $C > 0$, independent of $0 < a < a^*$, such that the imaginary part I_a of (2.21) satisfies*

$$|\nabla I_a(x)|, |I_a(x)| \leq C \varepsilon_a^2 (\varepsilon_a^4 + \gamma_a) e^{-\frac{1}{16}|x|} \quad \text{uniformly in } \mathbb{R}^2 \text{ as } a \nearrow a^*, \tag{2.30}$$

where $\gamma_a > 0$ satisfies $\gamma_a = o(|\beta_a|)$ as $a \nearrow a^*$.

2. *The real part R_a of (2.21) satisfies*

$$R_a(x) := \varepsilon_a^4 \psi_1(x) + \beta_a \psi_2(|x|) + o(\varepsilon_a^4 + |\beta_a|) \quad \text{in } \mathbb{R}^2 \text{ as } a \nearrow a^*, \tag{2.31}$$

where $\psi_2(|x|) := -\frac{1}{2}(w + x \cdot \nabla w)$ is radially symmetric, and $\psi_1(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ solves uniquely

$$\nabla \psi_1(0) = 0, \quad \tilde{\mathcal{L}} \psi_1(x) = -\frac{\lambda^4}{a^*} w^3(x) - V(x)w(x) \quad \text{in } \mathbb{R}^2. \tag{2.32}$$

Here ψ_1 and ψ_2 are independent of $0 < \Omega < \Omega^*$.

Proof. Denote

$$\mathcal{R}_a(x) := R_a(x) - \varepsilon_a^4 \psi_1(x) - \beta_a \psi_2(|x|), \quad (2.33)$$

where $\psi_2(|x|) := -\frac{1}{2}(w + x \cdot \nabla w)$ is radially symmetric, and $\psi_1(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ is a solution of (2.32). Note that ψ_1 and ψ_2 are independent of $0 < \Omega < \Omega^*$, and the uniqueness of $\psi_1(x)$ follows from $\nabla \psi_1(0) = 0$ and the property (2.19) (see also [59, Lemma 4.1] and [27, Theorem 1.2]). Moreover, by the comparison principle, see (2.45) below for the detailed argument, we derive from (2.5) and (2.32) that

$$|\psi_1(x)|, |\psi_2(|x|)| \leq C e^{-\delta|x|} \text{ in } \mathbb{R}^2, \text{ where } \frac{4}{5} < \delta < 1. \quad (2.34)$$

It then yields from (2.22) and (2.32) that $\mathcal{R}_a(x)$ satisfies

$$\nabla \mathcal{R}_a(0) \rightarrow 0 \text{ as } a \nearrow a^*. \quad (2.35)$$

We next carry out the proof by the following three steps:

Step 1. Note from (2.25) and (2.33) that

$$\tilde{\mathcal{L}}_a \mathcal{R}_a = \mathcal{F}_a(x) - (\tilde{\mathcal{L}}_a - \tilde{\mathcal{L}})[\varepsilon_a^4 \psi_1(x) + \beta_a \psi_2(|x|)] := N_a(x) \text{ in } \mathbb{R}^2. \quad (2.36)$$

We deduce from (2.26), (2.28) and (2.34) that the term $\mathcal{F}_a(x)$ satisfies

$$|\mathcal{F}_a(x)| := \left| F_a(x) - \beta_a w + \varepsilon_a^4 \left(\frac{\lambda^4}{a^*} w^3 + V(x)w \right) \right| \leq C \varepsilon_a^4 e^{-\frac{1}{10}|x|} \text{ uniformly in } \mathbb{R}^2$$

as $a \nearrow a^*$, and

$$\frac{|(\tilde{\mathcal{L}}_a - \tilde{\mathcal{L}})[\varepsilon_a^4 \psi_1(x) + \beta_a \psi_2(|x|)]|}{\varepsilon_a^4 + |\beta_a|} \leq C \delta_a e^{-\frac{1}{10}|x|} \text{ uniformly in } \mathbb{R}^2 \text{ as } a \nearrow a^*,$$

where $\delta_a > 0$ satisfies $\delta_a = o(1)$ as $a \nearrow a^*$. We thus derive from above that

$$\frac{|N_a(x)|}{\varepsilon_a^4 + |\beta_a|} \leq C e^{-\frac{1}{10}|x|} \text{ uniformly in } \mathbb{R}^2 \text{ as } a \nearrow a^*. \quad (2.37)$$

We claim that there exists a constant $C > 0$, independent of $0 < a < a^*$, such that

$$|\mathcal{R}_a(x)| \leq C(\varepsilon_a^4 + |\beta_a|) \text{ uniformly in } \mathbb{R}^2 \text{ as } a \nearrow a^*. \quad (2.38)$$

Instead, assume that the above claim (2.38) is false, i.e., $\lim_{a \nearrow a^*} \frac{\|\mathcal{R}_a\|_{L^\infty(\mathbb{R}^2)}}{\varepsilon_a^4 + |\beta_a|} = \infty$. Denote $\bar{\mathcal{R}}_a := \frac{\mathcal{R}_a}{\|\mathcal{R}_a\|_{L^\infty(\mathbb{R}^2)}}$, so that $\|\bar{\mathcal{R}}_a\|_{L^\infty(\mathbb{R}^2)} = 1$. Applying (2.22) and (2.28), we derive from (2.33) that

$$|\nabla \bar{\mathcal{R}}_a(0)| = \frac{|\nabla R_a(0)|}{\|\mathcal{R}_a\|_{L^\infty(\mathbb{R}^2)}} = \frac{1}{|w(0) + R_a(0)|} \frac{|I_a(0)| |\nabla I_a(0)|}{\|\mathcal{R}_a\|_{L^\infty(\mathbb{R}^2)}} \leq \frac{C(\varepsilon_a^4 + |\beta_a|)}{\|\mathcal{R}_a\|_{L^\infty(\mathbb{R}^2)}} \text{ as } a \nearrow a^*,$$

which then implies that

$$\nabla \bar{\mathcal{R}}_a(0) \rightarrow 0 \text{ as } a \nearrow a^*. \quad (2.39)$$

We also deduce from (2.36) that

$$\tilde{\mathcal{L}}_a \bar{\mathcal{R}}_a = \frac{N_a(x)}{\|\mathcal{R}_a\|_{L^\infty(\mathbb{R}^2)}} \text{ in } \mathbb{R}^2. \quad (2.40)$$

Note from (2.37) that

$$\begin{aligned} \frac{|N_a(x)|}{\|\mathcal{R}_a\|_{L^\infty(\mathbb{R}^2)}} &:= \frac{\varepsilon_a^4 + |\beta_a|}{\|\mathcal{R}_a\|_{L^\infty(\mathbb{R}^2)}} \frac{|N_a(x)|}{\varepsilon_a^4 + |\beta_a|} \\ &\leq \frac{\varepsilon_a^4 + |\beta_a|}{\|\mathcal{R}_a\|_{L^\infty(\mathbb{R}^2)}} C e^{-\frac{1}{10}|x|} \quad \text{uniformly in } \mathbb{R}^2 \text{ as } a \nearrow a^*. \end{aligned} \quad (2.41)$$

Suppose y_a is a global maximum point of $|\bar{\mathcal{R}}_a(x)|$, so that $|\bar{\mathcal{R}}_a(y_a)| = \max_{x \in \mathbb{R}^2} \frac{|\mathcal{R}_a(x)|}{\|\mathcal{R}_a\|_{L^\infty(\mathbb{R}^2)}} =$

1. In view of the definition of the operator $\tilde{\mathcal{L}}_a$, by the maximum principle one can deduce from (2.40) and (2.41) that $|y_a| \leq C$ uniformly in $0 < a < a^*$.

On the other hand, the usual elliptic regularity theory yields that there exist a subsequence, still denoted by $\{\bar{\mathcal{R}}_a\}$, of $\{\mathcal{R}_a\}$ and a function $\bar{\mathcal{R}}_0 \in H^1(\mathbb{R}^2)$ such that $\bar{\mathcal{R}}_a \rightarrow \bar{\mathcal{R}}_0$ weakly in $H^1(\mathbb{R}^2)$ and strongly in $L_{loc}^q(\mathbb{R}^2)$ for all $q \in [2, \infty)$ as $a \nearrow a^*$. By (2.37), we obtain from (2.39)–(2.41) that $\bar{\mathcal{R}}_0$ satisfies

$$\nabla \bar{\mathcal{R}}_0(0) = 0, \quad \tilde{\mathcal{L}} \bar{\mathcal{R}}_0(x) = 0 \quad \text{in } \mathbb{R}^2,$$

which yields that $\bar{\mathcal{R}}_0 = \sum_{i=1}^2 c_i \frac{\partial w}{\partial y_i}$ in view of (2.19). Since $\nabla \bar{\mathcal{R}}_0(0) = 0$, we obtain that

$$\begin{pmatrix} \frac{\partial^2 w(0)}{\partial x_i \partial x_j} \\ c_1 \\ c_2 \end{pmatrix} = 0.$$

Because $\det\left(\frac{\partial^2 w(0)}{\partial x_i \partial x_j}\right) \neq 0$, we deduce from above that $c_1 = c_2 = 0$, and hence $\bar{\mathcal{R}}_0(x) \equiv 0$ in \mathbb{R}^2 . This however contradicts to the above fact that by passing to a subsequence if necessary, $1 = \bar{\mathcal{R}}_0(y_a) \rightarrow \bar{\mathcal{R}}_0(\bar{y}_0)$ as $a \nearrow a^*$ for some $\bar{y}_0 \in \mathbb{R}^2$. Therefore, the claim (2.38) holds true.

We next claim that

$$|\mathcal{R}_a(x)|, \quad |\nabla \mathcal{R}_a(x)| \leq C(\varepsilon_a^4 + |\beta_a|) e^{-\frac{1}{11}|x|} \quad \text{uniformly in } \mathbb{R}^2 \text{ as } a \nearrow a^*. \quad (2.42)$$

Actually, we deduce from (2.36) and (2.37) that

$$\tilde{\mathcal{L}}_a \frac{\mathcal{R}_a}{\varepsilon_a^4 + |\beta_a|} = \frac{N_a(x)}{\varepsilon_a^4 + |\beta_a|} \leq C_1 e^{-\frac{1}{10}|x|} \quad \text{uniformly in } \mathbb{R}^2 \text{ as } a \nearrow a^*.$$

On the other hand, for sufficiently large $R > 1$ there exists a constant $C_2 = C_2(R) > 0$ such that

$$\frac{\mathcal{R}_a}{\varepsilon_a^4 + |\beta_a|} \leq C_2 e^{-\frac{1}{10}|x|} \quad \text{at } |x| = R \text{ as } a \nearrow a^*, \quad (2.43)$$

and

$$C_2 \tilde{\mathcal{L}}_a e^{-\frac{1}{10}|x|} \geq C_1 e^{-\frac{1}{10}|x|} \quad \text{in } \mathbb{R}^2/B_R(0) \text{ as } a \nearrow a^*.$$

We thus have

$$\tilde{\mathcal{L}}_a \left(\frac{\mathcal{R}_a}{\varepsilon_a^4 + |\beta_a|} - C_2 e^{-\frac{1}{10}|x|} \right) \leq 0 \quad \text{in } \mathbb{R}^2/B_R(0) \text{ as } a \nearrow a^*. \quad (2.44)$$

By the comparison principle, we then obtain from (2.43) and (2.44) that

$$\frac{\mathcal{R}_a}{\varepsilon_a^4 + |\beta_a|} \leq C_2 e^{-\frac{1}{10}|x|} \quad \text{in } \mathbb{R}^2/B_R(0) \text{ as } a \nearrow a^*.$$

Similarly, one can derive that there exists a constant $C_3 = C_3(R) > 0$ such that

$$\frac{\mathcal{R}_a}{\varepsilon_a^4 + |\beta_a|} \geq -C_3 e^{-\frac{1}{10}|x|} \quad \text{in } \mathbb{R}^2/B_R(0) \text{ as } a \nearrow a^*,$$

and hence

$$|\mathcal{R}_a| \leq C(\varepsilon_a^4 + |\beta_a|)e^{-\frac{1}{10}|x|} \quad \text{in } \mathbb{R}^2/B_R(0) \text{ as } a \nearrow a^*.$$

This therefore yields that \mathcal{R}_a satisfies

$$|\mathcal{R}_a(x)| \leq C(\varepsilon_a^4 + |\beta_a|)e^{-\frac{1}{10}|x|} \quad \text{in } \mathbb{R}^2, \quad (2.45)$$

due to the estimate (2.38). By gradient estimates of (3.15) in [29], we further derive from (2.36), (2.37) and (2.45) that

$$|\nabla \mathcal{R}_a| \leq C(\varepsilon_a^4 + |\beta_a|)e^{-\frac{1}{11}|x|} \quad \text{in } \mathbb{R}^2. \quad (2.46)$$

Therefore, we obtain from (2.45) and (2.46) that (2.42) holds true.

Step 2. Since $\psi_2(|x|) := -\frac{1}{2}(w + x \cdot \nabla w)$ is radially symmetric, we deduce from (2.33) that

$$x^\perp \cdot \nabla R_a = x^\perp \cdot \nabla [\mathcal{R}_a + \beta_a \psi_2 + \varepsilon_a^4 \psi_1(x)] = x^\perp \cdot \nabla [\mathcal{R}_a + \varepsilon_a^4 \psi_1(x)] \quad \text{in } \mathbb{R}^2.$$

This then implies from (2.42) that

$$|x^\perp \cdot \nabla R_a| \leq C(\varepsilon_a^4 + |\beta_a|)e^{-\frac{1}{12}|x|} \quad \text{uniformly in } \mathbb{R}^2 \text{ as } a \nearrow a^*. \quad (2.47)$$

Similar to (2.45) and (2.46), one can deduce from (2.24) and (2.47) that

$$|I_a(x)|, \quad |\nabla I_a(x)| \leq C\varepsilon_a^2(\varepsilon_a^4 + |\beta_a|)e^{-\frac{1}{14}|x|} \quad \text{uniformly in } \mathbb{R}^2 \text{ as } a \nearrow a^*, \quad (2.48)$$

where the property (2.17) is used in view of the fact that I_a satisfies $\int_{\mathbb{R}^2} w I_a dx \equiv 0$.

Step 3. Applying (2.48), the same argument of (2.37) yields that the nonhomogeneous term $N_a(x)$ of (2.36) satisfies

$$\frac{|N_a(x)|}{\varepsilon_a^4 + |\beta_a|} \leq C\delta_a e^{-\frac{1}{10}|x|} \quad \text{uniformly in } \mathbb{R}^2 \text{ as } a \nearrow a^*, \quad (2.49)$$

where $\delta_a > 0$ satisfies $\delta_a = o(1)$ as $a \nearrow a^*$. The same argument of Step 1 then gives from (2.49) that

$$|\mathcal{R}_a|, \quad |\nabla \mathcal{R}_a| \leq C(\varepsilon_a^4 + |\beta_a|)\delta_a e^{-\frac{1}{12}|x|} \quad \text{in } \mathbb{R}^2, \quad (2.50)$$

where $\delta_a > 0$ also satisfies $\delta_a = o(1)$ as $a \nearrow a^*$. This proves (2.31) in view of (2.33).

Finally, similar to (2.47), one can derive from (2.33) and (2.50) that

$$|x^\perp \cdot \nabla R_a| \leq C(\varepsilon_a^4 + \gamma_a)e^{-\frac{1}{14}|x|} \quad \text{uniformly in } \mathbb{R}^2 \text{ as } a \nearrow a^*,$$

where $\gamma_a > 0$ satisfies $\gamma_a = o(|\beta_a|)$ as $a \nearrow a^*$. Applying this refined estimate, the same argument of (2.48) then yields that (2.30) holds true, and Lemma 2.2 is thus proved. \square

Before ending this subsection, we derive the refined estimate of I_a as $a \nearrow a^*$.

Lemma 2.3. *Under the assumptions of Proposition 2.1, the imaginary part I_a of (2.21) satisfies*

$$I_a(x) := \varepsilon_a^6 \Omega \psi_I(x) + o(\varepsilon_a^6 + \varepsilon_a^2 |\beta_a|) \quad \text{in } \mathbb{R}^2 \quad \text{as } a \nearrow a^*, \quad (2.51)$$

where $\psi_I(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ solves uniquely

$$\int_{\mathbb{R}^2} \psi_I w dx = 0, \quad \mathcal{L} \psi_I(x) = -(x^\perp \cdot \nabla \psi_1) \quad \text{in } \mathbb{R}^2, \quad (2.52)$$

where the operator \mathcal{L} is defined by (2.16), and $\psi_1(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ is given by (2.32).

Proof. Following the argument of Lemma 2.2, one can derive from (2.24), (2.33) and (2.50) that $\int_{\mathbb{R}^2} I_a w dx \equiv 0$, and

$$\mathcal{L}_a I_a = -\varepsilon_a^2 \Omega (x^\perp \cdot \nabla R_a) = -\varepsilon_a^6 \Omega (x^\perp \cdot \nabla \psi_1) + o(\varepsilon_a^6 + \varepsilon_a^2 |\beta_a|) \quad \text{as } a \nearrow a^*, \quad (2.53)$$

where $\psi_1(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ is given by (2.32). Set

$$I_{1,a}(x) = I_a(x) - \varepsilon_a^6 \Omega \psi_I(x),$$

where $\psi_I(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ is defined in (2.52).

Similar to the proof of Lemma 2.2, one can deduce from (2.30) and (2.53) that $I_{1,a}(x) = o(\varepsilon_a^6 + \varepsilon_a^2 |\beta_a|)$ uniformly in \mathbb{R}^2 as $a \nearrow a^*$, which then implies directly that (2.51) holds true. Finally, since \mathcal{L} is a linear operator, the uniqueness of $\psi_I(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ defined by (2.52) follows from the constricton $\int_{\mathbb{R}^2} \psi_I w dx = 0$ and the property (2.17) (cf. [59, Lemma 4.1]). \square

Remark 2.1. Under the general assumptions on $V_\Omega(x)$ of Proposition 2.1, the upper bound (2.30) of I_a as $a \nearrow a^*$ is optimal and I_a does not vanish. However, if the trap $V_\Omega(x)$ has a better symmetry, then the leading term $\psi_I(x)$ of I_a as $a \nearrow a^*$ may vanish and hence the upper bound (2.30) of I_a as $a \nearrow a^*$ is not optimal. Specially, if the trap $V_\Omega(x)$ is radially symmetric, we have proved in [34, Theorem 1.3] that $I_a(x) \equiv 0$ as $a \nearrow a^*$.

3 Asymptotic Expansions and Applications

In this section we first address the proof of Theorem 1.1 on asymptotic expansions of minimizers for $e_F(a)$ as $a \nearrow a^*$, based on which we then establish Theorem 1.2. Towards this purpose, throughout this section we always consider the harmonic trap $V(x)$ of the form (1.11), and let $0 < \Omega < \Omega^* := 2\Lambda < 2$ be fixed, so that all estimates of previous section are applicable.

Let $R_a(x)$ and $I_a(x)$ be defined by (2.7) and (2.21), respectively, where u_a denotes a complex-valued minimizer of $e_F(a)$ as $a \nearrow a^*$. Applying (1.12), it then follows from Lemma 2.2 that the real part R_a of (2.21) satisfies

$$R_a(x) := \varepsilon_a^4 \psi_1(x) + \beta_a \psi_2(|x|) + o(\varepsilon_a^4 + |\beta_a|) \quad \text{in } \mathbb{R}^2 \quad \text{as } a \nearrow a^*, \quad (3.1)$$

where $\varepsilon_a > 0$ and $\beta_a \in \mathbb{R}$ are defined by (2.20). Here $\psi_2(|x|) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ satisfying

$$\psi_2(|x|) := -\frac{1}{2}(w + x \cdot \nabla w) \quad (3.2)$$

is radially symmetric, and $\psi_1(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ solves uniquely

$$\nabla\psi_1(0) = 0, \quad \tilde{\mathcal{L}}\psi_1(x) = -\frac{\lambda_0^4}{a^*}w^3(x) - (x_1^2 + \Lambda^2x_2^2)w(x) \text{ in } \mathbb{R}^2, \quad (3.3)$$

where $0 < \Lambda < 1$ is as in (1.11), $\lambda_0 > 0$ is as in (1.14) in view of (1.12), (1.13) and (2.6), and while the operator $\tilde{\mathcal{L}}$ is defined by (2.18).

We next denote $\psi_{11}(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ to be the unique solution of

$$\nabla\psi_{11}(0) = 0, \quad \tilde{\mathcal{L}}\psi_{11}(x) = -\frac{\lambda_0^4}{a^*}w^3(x) - \Lambda^2|x|^2w(x) \text{ in } \mathbb{R}^2, \quad (3.4)$$

and let $\psi(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ be the unique solution of

$$\nabla\psi(0) = 0, \quad \tilde{\mathcal{L}}\psi(x) = -(1 - \Lambda^2)x_1^2w(x) \text{ in } \mathbb{R}^2, \quad x = (x_1, x_2) \in \mathbb{R}^2, \quad (3.5)$$

where $0 < \Lambda < 1$ is as above. Following (3.3), we decompose ψ_1 as

$$\psi_1(x) = \psi_{11}(|x|) + \psi(x), \quad \text{where } \psi_{11}(|x|) \text{ is radially symmetric in } \mathbb{R}^2. \quad (3.6)$$

Applying (3.6), we thus conclude from Lemma 2.3 that the imaginary part I_a of (2.21) satisfies

$$I_a(x) := \varepsilon_a^6 \Omega \psi_I(x) + o(\varepsilon_a^6 + \varepsilon_a^2 |\beta_a|) \text{ in } \mathbb{R}^2 \text{ as } a \nearrow a^*, \quad (3.7)$$

where $\psi_I(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ solves uniquely

$$\int_{\mathbb{R}^2} \psi_I w dx = 0, \quad \mathcal{L}\psi_I(x) = -(x^\perp \cdot \nabla\psi_1) = -(x^\perp \cdot \nabla\psi) \text{ in } \mathbb{R}^2, \quad (3.8)$$

the operator \mathcal{L} is defined by (2.16), and $\psi(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ is given uniquely by (3.5).

Lemma 3.1. *Under the assumptions of Theorem 1.1, the unique maximum point x_a of $|u_a(x)|$ satisfies*

$$|\varepsilon_a^3 x_a| = o([\varepsilon_a^4 + |\beta_a|]^2) \text{ as } a \nearrow a^*, \quad (3.9)$$

where we denote $o([\varepsilon_a^4 + |\beta_a|]^2) = o(\varepsilon_a^8) + o(\varepsilon_a^4 |\beta_a|) + o(\beta_a^2)$.

Proof. We calculate from (2.25) and (2.26) that

$$\begin{aligned} \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \tilde{\mathcal{L}}_a R_a &= \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} F_a(x) \\ &= -\varepsilon_a^4 \left[\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} V_\Omega \left(x + \frac{x_a}{\varepsilon_a} \right) w + \frac{\lambda_0^4}{a^*} \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \tilde{R}_a^2 w \right] \\ &\quad + \varepsilon_a^4 \Omega \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \left(x^\perp \cdot \frac{\nabla I_a}{\varepsilon_a^2} \right) + \frac{a}{a^*} \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} I_a^2 w \\ &:= A_1 + A_2, \end{aligned} \quad (3.10)$$

where $V_\Omega(x) = V(x) - \frac{\Omega^2}{4}|x|^2$ is as before, the part A_2 satisfies

$$\begin{aligned} A_2 &:= \varepsilon_a^4 \Omega \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \left(x^\perp \cdot \frac{\nabla I_a}{\varepsilon_a^2} \right) + \frac{a}{a^*} \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} I_a^2 w \\ &= \varepsilon_a^4 [\varepsilon_a^4 + o(\varepsilon_a^4 + |\beta_a|)] \Omega^2 \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} (x^\perp \cdot \nabla\psi_I) \text{ as } a \nearrow a^* \end{aligned} \quad (3.11)$$

in view of Lemmas 2.2 and 2.3, and $\psi_I(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ satisfies (3.8) after simplification. As for the part A_1 , we observe from (1.12) and Proposition 2.1 that

$$\int_{\mathbb{R}^2} \frac{\partial w^2}{\partial x_1} V_\Omega(x) dx = 0, \quad \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} w^2 \psi_2(|x|) dx = 0,$$

where the radial symmetry of $\psi_2(|x|)$ is also used. We then have

$$\begin{aligned} -\frac{1}{\varepsilon_a^4} A_1 &:= \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} V_\Omega\left(x + \frac{x_a}{\varepsilon_a}\right) w + \frac{\lambda_0^4}{a^*} \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \tilde{R}_a^2 w \\ &= \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \left[V_\Omega\left(x + \frac{x_a}{\varepsilon_a}\right) - V_\Omega(x) \right] w + \frac{\lambda_0^4}{a^*} \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} (\tilde{R}_a^2 - w^2) w \\ &= \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \left(\frac{x_a}{\varepsilon_a} \cdot \nabla V_\Omega(x) \right) w + \frac{2\lambda_0^4}{a^*} \varepsilon_a^4 \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} w^2 \psi_1 \\ &\quad + o\left(\left| \frac{x_a}{\varepsilon_a} \right| + \varepsilon_a^4 + |\beta_a|\right) \quad \text{as } a \nearrow a^*, \end{aligned} \quad (3.12)$$

where the expansion of Lemma 2.2(2) is used in the last identity. We deduce from (3.10)–(3.12) that

$$\begin{aligned} &-\varepsilon_a^4 \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \left(\frac{x_a}{\varepsilon_a} \cdot \nabla V_\Omega(x) \right) w \\ &= \varepsilon_a^8 \left\{ \frac{2\lambda_0^4}{a^*} \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} w^2 \psi_1 - \Omega^2 \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} (x^\perp \cdot \nabla \psi_I) \right\} \\ &\quad + \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \tilde{\mathcal{L}}_a R_a + o(\varepsilon_a^8 + \varepsilon_a^4 |\beta_a|) \quad \text{as } a \nearrow a^*, \end{aligned} \quad (3.13)$$

where $\psi_I(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ satisfies (3.8).

On the other hand, by the definition of $\tilde{\mathcal{L}}$, we have $\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \tilde{\mathcal{L}} R_a dx = 0$. It then follows from Lemmas 2.2 and 2.3 that

$$\begin{aligned} \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \tilde{\mathcal{L}}_a R_a &= \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} (\tilde{\mathcal{L}}_a - \tilde{\mathcal{L}}) R_a \\ &= \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \left\{ \varepsilon_a^4 \left(\frac{\Omega^2}{4} |x|^2 + \frac{\lambda_0^4}{a^*} w^2 \right) + \varepsilon_a^4 V_\Omega\left(x + \frac{x_a}{\varepsilon_a}\right) - \beta_a \right. \\ &\quad \left. - w R_a - \frac{a}{a^*} (2w + R_a) R_a - \frac{a}{a^*} I_a^2 \right\} R_a \\ &= \varepsilon_a^8 \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \left(\frac{\Omega^2}{4} |x|^2 + \frac{\lambda_0^4}{a^*} w^2 \right) \psi_1 + \varepsilon_a^8 \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} V_\Omega(x) \psi_1 \\ &\quad + \varepsilon_a^4 \beta_a \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} V_\Omega(x) \psi_2 - \varepsilon_a^4 \beta_a \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \psi_1 - \frac{\varepsilon_a^8}{2} \int_{\mathbb{R}^2} \frac{\partial w^2}{\partial x_1} \psi_1^2 \\ &\quad - \varepsilon_a^4 \beta_a \int_{\mathbb{R}^2} \frac{\partial w^2}{\partial x_1} \psi_1 \psi_2 - \frac{a}{a^*} \int_{\mathbb{R}^2} \frac{\partial w^2}{\partial x_1} (\varepsilon_a^8 \psi_1^2 + 2\varepsilon_a^4 \beta_a \psi_1 \psi_2) \\ &\quad + o([\varepsilon_a^4 + |\beta_a|]^2) \quad \text{as } a \nearrow a^*, \end{aligned}$$

due to the radial symmetry of $\psi_2(|x|)$. The above estimate thus gives that

$$\begin{aligned} \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \tilde{\mathcal{L}}_a R_a &= \varepsilon_a^8 \left\{ \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \left[\frac{\lambda_0^4}{a^*} w^2 + \left(\frac{\Omega^2}{4} |x|^2 + V_\Omega(x) \right) \right] \psi_1 - \frac{3}{2} \int_{\mathbb{R}^2} \frac{\partial w^2}{\partial x_1} \psi_1^2 \right\} \\ &\quad + \varepsilon_a^4 \beta_a \left\{ \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} V_\Omega(x) \psi_2 - \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \psi_1 - 3 \int_{\mathbb{R}^2} \frac{\partial w^2}{\partial x_1} \psi_1 \psi_2 \right\} \\ &\quad + o([\varepsilon_a^4 + |\beta_a|]^2) \quad \text{as } a \nearrow a^*. \end{aligned} \quad (3.14)$$

Applying (1.13), we now derive from (3.13) and (3.14) that

$$\begin{aligned}
& -\frac{\varepsilon_a^4}{2} \int_{\mathbb{R}^2} \frac{\partial w^2}{\partial x_1} \left(\frac{x_a}{\varepsilon_a} \cdot \nabla V_\Omega(x) \right) \\
&= \varepsilon_a^8 \left\{ \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \left[\frac{3\lambda_0^4}{a^*} w^2 + (x_1^2 + \Lambda^2 x_2^2) \right] \psi_1 \right. \\
&\quad \left. - 3 \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} w \psi_1^2 - \Omega^2 \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} (x^\perp \cdot \nabla \psi_I) \right\} \\
&\quad - \varepsilon_a^4 \beta_a \left\{ \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \psi_1 + 6 \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} w \psi_1 \psi_2 - \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} V_\Omega(x) \psi_2 \right\} \\
&\quad + o([\varepsilon_a^4 + |\beta_a|]^2) \quad \text{as } a \nearrow a^*,
\end{aligned} \tag{3.15}$$

where $\psi_I(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ is given by (3.8).

Similar to (3.23) of [31], one can obtain that

$$\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \psi_1 + 6 \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} w \psi_1 \psi_2 - \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} V_\Omega(x) \psi_2 = 0. \tag{3.16}$$

Moreover, one can note from (3.3) that $\psi_1(x)$ is even in $x \in \mathbb{R}^2$, which thus implies that

$$\int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} \left[\frac{3\lambda_0^4}{a^*} w^2 + (x_1^2 + \Lambda^2 x_2^2) \right] \psi_1 - 3 \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} w \psi_1^2 = 0.$$

Applying (3.15), we then derive from above that the unique maximum point x_a of $|u_a(x)|$ satisfies

$$\begin{aligned}
& \frac{\varepsilon_a^3 x_a}{2} \cdot \frac{\partial \nabla H_\Omega(y)}{\partial y_1} \Big|_{y=0} = \frac{\varepsilon_a^4}{2} \int_{\mathbb{R}^2} w^2(x) \left(\frac{x_a}{\varepsilon_a} \right) \cdot \frac{\partial \nabla V_\Omega(x+y)}{\partial x_1} \Big|_{y=0} dx \\
&= -\frac{\varepsilon_a^4}{2} \int_{\mathbb{R}^2} \frac{\partial w^2}{\partial x_1} \left(\frac{x_a}{\varepsilon_a} \right) \cdot \nabla V_\Omega(x) dx \\
&= -\varepsilon_a^8 \Omega^2 \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} (x^\perp \cdot \nabla \psi_I) dx + o([\varepsilon_a^4 + |\beta_a|]^2) \quad \text{as } a \nearrow a^*,
\end{aligned} \tag{3.17}$$

due to the definition of $H_\Omega(y) := \int_{\mathbb{R}^2} w^2(x) V_\Omega(x+y) dx$. Similarly, one can obtain that the unique maximum point x_a of $|u_a(x)|$ also satisfies

$$\begin{aligned}
& \frac{\varepsilon_a^3 x_a}{2} \cdot \frac{\partial \nabla H_\Omega(y)}{\partial y_2} \Big|_{y=0} = \frac{\varepsilon_a^4}{2} \int_{\mathbb{R}^2} w^2(x) \left(\frac{x_a}{\varepsilon_a} \right) \cdot \frac{\partial \nabla V_\Omega(x+y)}{\partial x_2} \Big|_{y=0} dx \\
&= -\frac{\varepsilon_a^4}{2} \int_{\mathbb{R}^2} \frac{\partial w^2}{\partial x_2} \left(\frac{x_a}{\varepsilon_a} \right) \cdot \nabla V_\Omega(x) dx \\
&= -\varepsilon_a^8 \Omega^2 \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_2} (x^\perp \cdot \nabla \psi_I) dx + o([\varepsilon_a^4 + |\beta_a|]^2) \quad \text{as } a \nearrow a^*.
\end{aligned} \tag{3.18}$$

In the appendix, we shall prove that the following claim is true:

$$II_1 = \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} (x^\perp \cdot \nabla \psi_I) = 0, \quad II_2 = \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_2} (x^\perp \cdot \nabla \psi_I) = 0, \tag{3.19}$$

where $\psi_I(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ solves uniquely (3.8). Recall from (1.12) that 0 is the unique and non-degenerate critical point of $H_\Omega(y)$. Applying (3.19), we then conclude from (3.17) and (3.18) that (3.9) holds true, and the proof of Lemma 3.1 is therefore complete. \square

Since the limit estimates of Lemma 2.2 are not enough for establishing Theorem 1.1, we next employ Lemma 3.1 to derive the following more terms of R_a in terms of ε_a and β_a .

Lemma 3.2. *Under the assumptions of Theorem 1.1, the real part R_a of (2.21) satisfies*

$$\begin{aligned} R_a(x) &:= \varepsilon_a^4 \psi_1(x) + \beta_a \psi_2(x) + \varepsilon_a^8 \psi_3(x) + \beta_a^2 \psi_4(x) + \varepsilon_a^4 \beta_a \psi_5(x) \\ &\quad + o([\varepsilon_a^4 + |\beta_a|]^2) \quad \text{in } \mathbb{R}^2 \quad \text{as } a \nearrow a^*, \end{aligned} \quad (3.20)$$

where $\psi_1(x)$ and $\psi_2(x)$ are as in (3.2) and (3.3), respectively. However, $\psi_i(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ solves uniquely

$$\nabla \psi_i(0) = 0, \quad \tilde{\mathcal{L}} \psi_i(x) = f_i(x) \quad \text{in } \mathbb{R}^2, \quad i = 3, 4, 5, \quad (3.21)$$

and $f_i(x)$ satisfies

$$f_i(x) = \begin{cases} 3w\psi_1^2 - \left[\frac{3\lambda_0^4}{a^*} w^2 + (x_1^2 + \Lambda^2 x_2^2) \right] \psi_1 + \Omega(x^\perp \cdot \nabla \psi_I), & \text{if } i = 3; \\ \psi_2 + 3w\psi_2^2, & \text{if } i = 4; \\ 6w\psi_1\psi_2 + \psi_1 - \left[\frac{3\lambda_0^4}{a^*} w^2 + (x_1^2 + \Lambda^2 x_2^2) \right] \psi_2, & \text{if } i = 5; \end{cases} \quad (3.22)$$

where $\psi_I(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ is given by (3.8).

Proof. Denote

$$M_a := R_a - \varepsilon_a^4 \psi_1 - \beta_a \psi_2,$$

which then yields from (2.25) that

$$\begin{aligned} \tilde{\mathcal{L}}_a M_a &= \tilde{\mathcal{L}}_a R_a - \tilde{\mathcal{L}}_a(\varepsilon_a^4 \psi_1 + \beta_a \psi_2) \\ &= \{ \tilde{\mathcal{L}}_a R_a - \tilde{\mathcal{L}}(\varepsilon_a^4 \psi_1 + \beta_a \psi_2) \} - (\tilde{\mathcal{L}}_a - \tilde{\mathcal{L}})(\varepsilon_a^4 \psi_1 + \beta_a \psi_2) \\ &:= II_1 + II_2. \end{aligned} \quad (3.23)$$

Direct calculations give that the term II_2 of (3.23) satisfies

$$\begin{aligned} II_2 &:= -(\tilde{\mathcal{L}}_a - \tilde{\mathcal{L}})(\varepsilon_a^4 \psi_1 + \beta_a \psi_2) \\ &= -(\varepsilon_a^4 \psi_1 + \beta_a \psi_2) \left\{ \varepsilon_a^4 \left[\frac{\Omega^2}{4} |x|^2 + V_\Omega \left(x + \frac{x_a}{\varepsilon_a} \right) \right] - \beta_a - w R_a \right. \\ &\quad \left. + \frac{\alpha_a}{a^*} w^2 - \frac{a}{a^*} (2w + R_a) R_a - \frac{a}{a^*} I_a^2 \right\}. \end{aligned} \quad (3.24)$$

Applying Lemmas 2.2 and 2.3, we thus get that

$$\begin{aligned} II_2 &:= \varepsilon_a^8 \left\{ 3w\psi_1^2 - \left(\frac{\lambda_0^4}{a^*} w^2 + (x_1^2 + \Lambda^2 x_2^2) \right) \psi_1 \right\} \\ &\quad + \varepsilon_a^4 \beta_a \left\{ 6w\psi_1\psi_2 + \psi_1 - \left(\frac{\lambda_0^4}{a^*} w^2 + (x_1^2 + \Lambda^2 x_2^2) \right) \psi_2 \right\} \\ &\quad + \beta_a^2 (\psi_2 + 3w\psi_2^2) + o([\varepsilon_a^4 + |\beta_a|]^2) \quad \text{as } a \nearrow a^*. \end{aligned} \quad (3.25)$$

As for the term II_1 of (3.23), we derive from (2.25) that

$$\begin{aligned} II_1 &:= \tilde{\mathcal{L}}_a R_a - \tilde{\mathcal{L}}(\varepsilon_a^4 \psi_1 + \beta_a \psi_2) \\ &= \tilde{\mathcal{L}}_a R_a - \beta_a w + \varepsilon_a^4 \left\{ \frac{\lambda_0^4}{a^*} w^3 + (x_1^2 + \Lambda^2 x_2^2) w \right\} \\ &= -\varepsilon_a^4 \left[V_\Omega \left(x + \frac{x_a}{\varepsilon_a} \right) - V_\Omega(x) \right] w - \frac{\alpha_a}{a^*} w (2w + R_a) R_a \\ &\quad + \varepsilon_a^4 \Omega \left(x^\perp \cdot \frac{\nabla I_a}{\varepsilon_a^2} \right) + \frac{a}{a^*} I_a^2 w. \end{aligned} \quad (3.26)$$

Applying (3.7) yields that

$$\varepsilon_a^4 \Omega \left(x^\perp \cdot \frac{\nabla I_a}{\varepsilon_a^2} \right) = \varepsilon_a^8 \Omega \left(x^\perp \cdot \nabla \psi_I \right) + o(\varepsilon_a^8 + \varepsilon_a^4 |\beta_a|) \quad \text{as } a \nearrow a^*,$$

where $\psi_I(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ is given by (3.8). By Lemma 3.1, we have

$$\begin{aligned} -\varepsilon_a^4 \left[V_\Omega \left(x + \frac{x_a}{\varepsilon_a} \right) - V_\Omega(x) \right] w &= -\varepsilon_a^3 (x_a \cdot \nabla V_\Omega(x)) w(x) [1 + o(1)] \\ &= o([\varepsilon_a^4 + |\beta_a|]^2) \quad \text{as } a \nearrow a^*. \end{aligned} \quad (3.27)$$

Applying Lemmas 2.2 and 2.3, we now deduce from above that

$$\begin{aligned} II_1 &:= \varepsilon_a^8 \left\{ -\frac{2\lambda_0^4}{a^*} w^2 \psi_1 + \Omega \left(x^\perp \cdot \nabla \psi_I \right) \right\} \\ &\quad - \varepsilon_a^4 \beta_a \frac{2\lambda_0^4}{a^*} w^2 \psi_2 + o([\varepsilon_a^4 + |\beta_a|]^2) \quad \text{as } a \nearrow a^*, \end{aligned} \quad (3.28)$$

where $\psi_I(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ is given by (3.8).

Applying (3.25) and (3.28), we now calculate from (3.23) that

$$\begin{aligned} \tilde{\mathcal{L}}_a M_a &= \varepsilon_a^8 \left\{ 3w\psi_1^2 - \left[\frac{3\lambda_0^4}{a^*} w^2 + (x_1^2 + \Lambda^2 x_2^2) \right] \psi_1 + \Omega \left(x^\perp \cdot \nabla \psi_I \right) \right\} \\ &\quad + \varepsilon_a^4 \beta_a \left\{ 6w\psi_1\psi_2 + \psi_1 - \left(\frac{3\lambda_0^4}{a^*} w^2 + (x_1^2 + \Lambda^2 x_2^2) \right) \psi_2 \right\} \\ &\quad + \beta_a^2 (\psi_2 + 3w\psi_2^2) + o([\varepsilon_a^4 + |\beta_a|]^2) \quad \text{as } a \nearrow a^*. \end{aligned} \quad (3.29)$$

Following (3.29), the argument of Lemma 2.2 then yields the estimate (3.20). Moreover, the property (2.19) implies the uniqueness of ψ_i for $i = 3, 4, 5$, and the proof of Lemma 3.2 is therefore complete. \square

Lemma 3.3. *Under the assumptions of Theorem 1.1, we have*

$$\int_{\mathbb{R}^2} w\psi_1 = 0, \quad \int_{\mathbb{R}^2} w\psi_2 = 0, \quad T_1 = \int_{\mathbb{R}^2} (2w\psi_4 + \psi_2^2) = 0, \quad (3.30)$$

and

$$T_2 = 2 \int_{\mathbb{R}^2} w\psi_5 + 2 \int_{\mathbb{R}^2} \psi_1\psi_2 = -2\lambda_0^4 < 0, \quad (3.31)$$

where $\psi_1(x), \dots, \psi_5(x), \psi_I(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ are as in Lemma 3.2.

Proof. The assumptions of Theorem 1.1 imply that

$$\int_{\mathbb{R}^2} w^2 \left(x \cdot \nabla V(x) \right) = 2 \int_{\mathbb{R}^2} V(x) w^2 dx = 2\lambda_0^4 > 0. \quad (3.32)$$

Applying (3.32), Lemma 3.3 can be proved in a similar way of [31, Lemma 3.5], and the detailed proof is omitted for simplicity. \square

3.1 Proofs of Theorems 1.1 and 1.2

In this subsection we complete the proofs of Theorems 1.1 and 1.2. Under the assumption (1.11), we remark from (1.13) that $\varepsilon_a = \epsilon_a > 0$, where $\epsilon_a > 0$ is as in the statement of Theorem 1.1.

Proof of Theorem 1.1. Applying Proposition 2.1, we derive from (2.21) that R_a satisfies

$$\int_{\mathbb{R}^2} w^2 = \int_{\mathbb{R}^2} |v_a|^2 = \int_{\mathbb{R}^2} [(w + R_a)^2 + I_a^2], \text{ i.e., } 0 = 2 \int_{\mathbb{R}^2} w R_a + \int_{\mathbb{R}^2} R_a^2 + \int_{\mathbb{R}^2} I_a^2. \quad (3.33)$$

Following Lemmas 3.2 and 3.3, we then derive from (3.33) that

$$\begin{aligned} 0 &= 2 \int_{\mathbb{R}^2} w R_a + \int_{\mathbb{R}^2} R_a^2 + \int_{\mathbb{R}^2} I_a^2 \\ &= 2 \int_{\mathbb{R}^2} w(\varepsilon_a^4 \psi_1 + \beta_a \psi_2 + \varepsilon_a^8 \psi_3 + \beta_a^2 \psi_4 + \varepsilon_a^4 \beta_a \psi_5) \\ &\quad + \int_{\mathbb{R}^2} (\varepsilon_a^4 \psi_1 + \beta_a \psi_2 + \varepsilon_a^8 \psi_3 + \beta_a^2 \psi_4 + \varepsilon_a^4 \beta_a \psi_5)^2 + o([\varepsilon_a^4 + |\beta_a|]^2) \\ &= 2\varepsilon_a^4 \left(\int_{\mathbb{R}^2} w \psi_1 \right) + 2\beta_a \left(\int_{\mathbb{R}^2} w \psi_2 \right) + \beta_a^2 \left(2 \int_{\mathbb{R}^2} w \psi_4 + \int_{\mathbb{R}^2} \psi_2^2 \right) \\ &\quad + 2\varepsilon_a^4 \beta_a \left(\int_{\mathbb{R}^2} w \psi_5 + \int_{\mathbb{R}^2} \psi_1 \psi_2 \right) + \varepsilon_a^8 \left(2 \int_{\mathbb{R}^2} w \psi_3 + \int_{\mathbb{R}^2} \psi_1^2 \right) + o([\varepsilon_a^4 + |\beta_a|]^2) \\ &= -2\lambda_0^4 \varepsilon_a^4 \beta_a + \varepsilon_a^8 \left(2 \int_{\mathbb{R}^2} w \psi_3 + \int_{\mathbb{R}^2} \psi_1^2 \right) + o([\varepsilon_a^4 + |\beta_a|]^2) \text{ as } a \nearrow a^*, \end{aligned} \quad (3.34)$$

where (3.7) and (3.8) are also used. One can derive from (3.34) that

$$2 \int_{\mathbb{R}^2} w \psi_3 + \int_{\mathbb{R}^2} \psi_1^2 \neq 0,$$

and

$$-2\lambda_0^4 \beta_a + \varepsilon_a^4 \left(2 \int_{\mathbb{R}^2} w \psi_3 + \int_{\mathbb{R}^2} \psi_1^2 \right) = 0, \quad (3.35)$$

where $\psi_1(x)$ and $\psi_3(x)$ are as in (3.3) and (3.21), respectively. In the appendix, we shall prove the following claim that

$$I := 2 \int_{\mathbb{R}^2} w \psi_3 + \int_{\mathbb{R}^2} \psi_1^2 = \int_{\mathbb{R}^2} (3w^2 - 1)\psi_1^2 - 4 \int_{\mathbb{R}^2} (x_1^2 + \Lambda^2 x_2^2) w \psi_1. \quad (3.36)$$

Together with (3.36), we conclude from (3.35) that the constant β_a satisfies

$$\beta_a = C^* \varepsilon_a^4, \text{ where } C^* = \frac{1}{2\lambda_0^4} \left[\int_{\mathbb{R}^2} (3w^2 - 1)\psi_1^2 - 4 \int_{\mathbb{R}^2} (x_1^2 + \Lambda^2 x_2^2) w \psi_1 \right] \neq 0. \quad (3.37)$$

Applying (3.20), (3.7) and (3.37), we thus conclude from (2.7) and (2.21) that the refined limit profile (1.18) holds true. Moreover, the estimate (1.19) follows directly from (3.37) and Lemma 3.1. This completes the proof of Theorem 1.1. \square

As a byproduct of Theorem 1.1, we finally establish Theorem 1.2 on the nonexistence of vortices in a very large region.

Proof of Theorem 1.2. By Theorem 1.1, for any fixed and sufficiently large $R > 1$, there exists a constant $C_R := C(R) > 0$ such that

$$|v_a(x)| \geq w(x) - C_R \varepsilon_a^4 > 0 \quad \text{in } \{x \in \mathbb{R}^2 : |x| \leq R\} \quad \text{as } a \nearrow a^*, \quad (3.38)$$

where $v_a(x)$ is as in (2.7).

Setting $\tilde{w}_a = v_a(x) - w(x)$, we claim that

$$|\tilde{w}_a| \leq C_1 \varepsilon_a^4 |x|^{\frac{5}{2}} e^{-\sqrt{1-C_2 \varepsilon_a^4} |x|} \quad \text{uniformly in } \mathbb{R}^2 \quad \text{as } a \nearrow a^*, \quad (3.39)$$

where the constants $C_1 > 0$ and $C_2 > 0$ are independent of $0 < a < a^*$. To prove the claim (3.39), we note from (2.14) that \tilde{w}_a satisfies

$$(-\Delta + \hat{V}_a)(\tilde{w}_a + w) + i\varepsilon_a^2 \Omega(x^\perp \cdot \nabla \tilde{w}_a) = 0 \quad \text{in } \mathbb{R}^2,$$

i.e.,

$$(-\Delta + \hat{V}_a)\tilde{w}_a + i\varepsilon_a^2 \Omega(x^\perp \cdot \nabla \tilde{w}_a) + (-\Delta + \hat{V}_a)w = 0 \quad \text{in } \mathbb{R}^2, \quad (3.40)$$

where

$$\hat{V}_a(x) = \frac{\varepsilon_a^4 \Omega^2}{4} |x|^2 + \varepsilon_a^2 V_\Omega(\varepsilon_a x + x_a) - \mu_a \varepsilon_a^2 - \frac{a}{a^*} |v_a|^2 \quad \text{in } \mathbb{R}^2. \quad (3.41)$$

One can derive from (3.40) that

$$\begin{aligned} & -\frac{1}{2} \Delta |\tilde{w}_a|^2 + \left[\frac{\varepsilon_a^4 \Omega^2}{4} |x|^2 + \varepsilon_a^2 V_\Omega(\varepsilon_a x + x_a) - \mu_a \varepsilon_a^2 - \frac{a}{a^*} |v_a|^2 \right] |\tilde{w}_a|^2 + |\nabla \tilde{w}_a|^2 \\ & - \varepsilon_a^2 \Omega x^\perp (i\tilde{w}_a, \nabla \tilde{w}_a) + (-\Delta + \hat{V}_a)(w, \tilde{w}_a) = 0 \quad \text{in } \mathbb{R}^2. \end{aligned} \quad (3.42)$$

By the diamagnetic inequality (1.8), we have

$$|\nabla \tilde{w}_a|^2 + \frac{\varepsilon_a^4 \Omega^2}{4} |x|^2 |\tilde{w}_a|^2 - \varepsilon_a^2 \Omega x^\perp (i\tilde{w}_a, \nabla \tilde{w}_a) \geq |\nabla |\tilde{w}_a||^2 \quad \text{in } \mathbb{R}^2. \quad (3.43)$$

Since

$$\frac{1}{2} \Delta |\tilde{w}_a|^2 = |\tilde{w}_a| \Delta |\tilde{w}_a| + |\nabla |\tilde{w}_a||^2,$$

we deduce from (3.42) and (3.43) that

$$-\Delta |\tilde{w}_a| - \mu_a \varepsilon_a^2 |\tilde{w}_a| \leq |(-\Delta + \hat{V}_a)w| + \frac{a}{a^*} |v_a|^2 |\tilde{w}_a| \quad \text{in } \mathbb{R}^2. \quad (3.44)$$

The argument of [34, Proposition 3.3] gives that as $a \nearrow a^*$,

$$|v_a|^2 \leq C e^{-\frac{4}{3}|x|} \quad \text{in } \mathbb{R}^2,$$

which implies from Theorem 1.1 that as $a \nearrow a^*$,

$$\frac{a}{a^*} |v_a|^2 |\tilde{w}_a| \leq C \varepsilon_a^4 e^{-\frac{4}{3}|x|} \quad \text{in } \mathbb{R}^2. \quad (3.45)$$

Note from (2.20) and (3.37) that as $a \nearrow a^*$,

$$-\mu_a \varepsilon_a^2 = 1 - C^* \varepsilon_a^4.$$

We then calculate from (2.5) and (3.41) that as $a \nearrow a^*$,

$$\begin{aligned} |(-\Delta + \hat{V}_a)w| &= \left| \left(\frac{\varepsilon_a^4 \Omega^2}{4} |x|^2 + \varepsilon_a^2 V_\Omega(\varepsilon_a x + x_a) - 1 - \mu_a \varepsilon_a^2 + w^2 - \frac{a}{a^*} |v_a|^2 \right) w \right| \\ &\leq C \varepsilon_a^4 |x|^{\frac{3}{2}} e^{-|x|} \quad \text{in } \mathbb{R}^2 \setminus B_R(0), \end{aligned} \quad (3.46)$$

where the sufficiently large constant $R > 1$ is as in (3.38). We thus deduce from (3.44)–(3.46) that as $a \nearrow a^*$,

$$-\Delta |\tilde{w}_a| - \mu_a \varepsilon_a^2 |\tilde{w}_a| \leq C_0 \varepsilon_a^4 |x|^{\frac{3}{2}} e^{-|x|} \quad \text{in } \mathbb{R}^2 \setminus B_R(0), \quad (3.47)$$

where the constant $C_0 > 0$ is independent of $0 < a < a^*$. Since Theorem 1.1 gives that $|\tilde{w}_a| = O(\varepsilon_a^4)$ as $a \nearrow a^*$, we have as $a \nearrow a^*$,

$$|\tilde{w}_a| \leq C \varepsilon_a^4 |x|^{\frac{5}{2}} e^{-\sqrt{1-C_* \varepsilon_a^4} |x|} \quad \text{at } |x| = R > 1, \quad (3.48)$$

where $R > 1$ is as in (3.38), and $C > 0$ is large enough and independent of $0 < a < a^*$. By the comparison principle as in (2.45), we thus derive from (3.47) and (3.48) that the claim (3.39) holds true, in view of the fact that $|\tilde{w}_a| = O(\varepsilon_a^4)$ as $a \nearrow a^*$.

Applying (2.5) and (3.39), we now have as $a \nearrow a^*$,

$$\begin{aligned} |v_a| &\geq |w| - |v_a - w| \\ &\geq C_3 |x|^{-\frac{1}{2}} e^{-|x|} - C_1 \varepsilon_a^4 |x|^{\frac{5}{2}} e^{-\sqrt{1-C_2 \varepsilon_a^4} |x|} \\ &\geq |x|^{-\frac{1}{2}} e^{-|x|} (C_3 - C_1 \varepsilon_a^4 |x|^3 e^{C \varepsilon_a^4 |x|}) > 0, \quad \text{if } R \leq |x| \leq \left(\frac{C_3}{2C_1} \right)^{\frac{1}{3}} \varepsilon_a^{-\frac{4}{3}}. \end{aligned} \quad (3.49)$$

We thus conclude from (3.38) and (3.49) that as $a \nearrow a^*$,

$$|v_a(x)| = \left| \varepsilon_a \sqrt{a^*} u_a(\varepsilon_a x + x_a) e^{-i(\frac{\varepsilon_a \Omega}{2} x \cdot x_a^\perp - \theta_a)} \right| > 0 \quad \text{in } \left\{ x \in \mathbb{R}^2 : |x| \leq \left(\frac{C_3}{2C_1} \right)^{\frac{1}{3}} \varepsilon_a^{-\frac{4}{3}} \right\}.$$

Thus, there exists a small constant $C_* > 0$, independent of $0 < a < a^*$, such that as $a \nearrow a^*$,

$$|u_a(y)| > 0 \quad \text{if } |y| \leq C_4 \varepsilon_a^{-\frac{1}{3}} \leq \frac{C_*}{(a^* - a)^{\frac{1}{12}}}, \quad (3.50)$$

i.e., u_a does not admit any vortex in the region $R(a) := \{x \in \mathbb{R}^2 : |x| \leq C_*(a^* - a)^{-\frac{1}{12}}\}$ as $a \nearrow a^*$. This completes the proof of Theorem 1.2. \square

A Appendix

In this appendix, we follow those notations of Section 3 to address the proof of the claims (3.19) and (3.36).

Proof of (3.19). Consider the polar coordinate (r, θ) in \mathbb{R}^2 , where $\theta \in [0, 2\pi]$. Rewrite $w(x) = w(r)$, $\psi(x) = \psi(r, \theta)$ and $\psi_I(x) = \psi_I(r, \theta)$, where ψ and ψ_I are given by (3.5) and (3.8), respectively. We then have

$$\frac{\partial w}{\partial x_1} = w'(r) \cos \theta, \quad \frac{\partial w}{\partial x_2} = w'(r) \sin \theta. \quad (\text{A.1})$$

Since $\nabla\psi = \frac{x}{r}\psi_r + \frac{x^\perp}{r^2}\psi_\theta$, we have

$$x^\perp \cdot \nabla\psi = \frac{\partial\psi(r, \theta)}{\partial\theta}, \quad x^\perp \cdot \nabla\psi_I = \frac{\partial\psi_I(r, \theta)}{\partial\theta}. \quad (\text{A.2})$$

By the symmetry of the linear inhomogeneous equation (3.5), we deduce that

$$\psi(r, \theta) = \psi(r, 2\pi - \theta), \quad \psi(r, \theta) = \psi(r, \theta - \pi), \quad \theta \in [\pi, 2\pi], \quad (\text{A.3})$$

where $\psi(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ is given by (3.5) as before. It then yields from (A.3) that

$$\frac{\partial\psi(r, \theta)}{\partial\theta} = -\frac{\partial\psi(r, 2\pi - \theta)}{\partial(2\pi - \theta)}, \quad \frac{\partial\psi(r, \theta)}{\partial\theta} = \frac{\partial\psi(r, \theta - \pi)}{\partial(\theta - \pi)}, \quad \theta \in [\pi, 2\pi]. \quad (\text{A.4})$$

Note from (3.8) and (A.2) that $\psi_I(x) \in C^2(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$ is the unique solution of

$$\int_{\mathbb{R}^2} \psi_I w dx = 0, \quad \mathcal{L}\psi_I(x) = -(x^\perp \cdot \nabla\psi) = -\frac{\partial\psi(r, \theta)}{\partial\theta} \text{ in } \mathbb{R}^2. \quad (\text{A.5})$$

We then derive from (A.4) and (A.5) that the unique solution $\psi_I(x)$ satisfies

$$\psi_I(r, \theta) = -\psi_I(r, 2\pi - \theta), \quad \psi_I(r, \theta) = \psi_I(r, \theta - \pi), \quad \theta \in [\pi, 2\pi]. \quad (\text{A.6})$$

Applying (A.2), we now deduce that

$$\begin{aligned} II_1 &= \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} (x^\perp \cdot \nabla\psi_I) \\ &= \int_0^\infty \int_0^{2\pi} r w'(r) \cos\theta \frac{\partial\psi_I(r, \theta)}{\partial\theta} d\theta dr \\ &= \int_0^\infty \int_0^{2\pi} r w'(r) \sin\theta \psi_I(r, \theta) d\theta dr \\ &= \int_0^\infty \int_0^\pi r w'(r) \sin\theta \psi_I(r, \theta) d\theta dr + \int_0^\infty \int_\pi^{2\pi} r w'(r) \sin\theta \psi_I(r, \theta) d\theta dr \\ &:= A_1 + A_2. \end{aligned} \quad (\text{A.7})$$

As for the term A_2 , we derive from (A.6) that

$$\begin{aligned} A_2 &:= \int_0^\infty \int_\pi^{2\pi} r w'(r) \sin\theta \psi_I(r, \theta) d\theta dr \\ &= -\int_0^\infty \int_\pi^{2\pi} r w'(r) \sin(\theta - \pi) \psi_I(r, \theta - \pi) d\theta dr \\ &= -\int_0^\infty \int_0^\pi r w'(r) \sin\delta \psi_I(r, \delta) d\delta dr = -A_1, \end{aligned} \quad (\text{A.8})$$

where we denote $\delta := \theta - \pi$. We hence obtain from (A.7) and (A.8) that

$$II_1 = \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_1} (x^\perp \cdot \nabla\psi_I) = A_1 + A_2 = 0. \quad (\text{A.9})$$

Similar to (A.7), one can conclude from (A.2) and (A.6) that

$$\begin{aligned} II_2 &= \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_2} (x^\perp \cdot \nabla\psi_I) \\ &= -\int_0^\infty \int_0^\pi r w'(r) \cos\theta \psi_I(r, \theta) d\theta dr - \int_0^\infty \int_\pi^{2\pi} r w'(r) \cos\theta \psi_I(r, \theta) d\theta dr \\ &:= -(B_1 + B_2), \end{aligned} \quad (\text{A.10})$$

where the term B_2 satisfies

$$\begin{aligned}
B_2 &:= \int_0^\infty \int_\pi^{2\pi} r w'(r) \cos \theta \psi_I(r, \theta) d\theta dr \\
&= - \int_0^\infty \int_\pi^{2\pi} r w'(r) \cos(2\pi - \theta) \psi_I(r, 2\pi - \theta) d\theta dr \\
&= \int_0^\infty \int_\pi^0 r w'(r) \cos \delta \psi_I(r, \delta) d\delta dr = -B_1,
\end{aligned} \tag{A.11}$$

where we denote $\delta := 2\pi - \theta$. The above estimates yield that

$$II_2 = \int_{\mathbb{R}^2} \frac{\partial w}{\partial x_2} (x^\perp \cdot \nabla \psi_I) = -(B_1 + B_2) = 0,$$

together with (A.9), which thus implies that (3.19) holds true, and we are done. \square

In the rest of this appendix, we establish the claim (3.36) as follows.

Proof of (3.36). Since ψ_2 is radially symmetric, we first note that for $\psi_2' = \frac{d\psi_2}{dr}$,

$$\begin{aligned}
\Omega \int_{\mathbb{R}^2} \psi_2 (x^\perp \cdot \nabla \psi_I) &= \Omega \int_{\mathbb{R}^2} \psi_2 [-x_2 (\psi_I)_{x_1} + x_1 (\psi_I)_{x_2}] \\
&= \Omega \int_{\mathbb{R}^2} [\psi_I x_2 (\psi_2)_{x_1} - \psi_I x_1 (\psi_2)_{x_2}] \\
&= \Omega \int_{\mathbb{R}^2} \psi_I \frac{x_1 x_2}{r} (\psi_2' - \psi_2') = 0.
\end{aligned}$$

Following (3.3) and (3.21), we then derive from above that for $V(x) = x_1^2 + \Lambda^2 x_2^2$,

$$\begin{aligned}
I &= 2 \int_{\mathbb{R}^2} \psi_3 \tilde{\mathcal{L}} \psi_2 + \int_{\mathbb{R}^2} \psi_1^2 \\
&= 2 \int_{\mathbb{R}^2} \psi_2 \tilde{\mathcal{L}} \psi_3 - 2\Omega \int_{\mathbb{R}^2} \psi_2 (x^\perp \cdot \nabla \psi_I) + \int_{\mathbb{R}^2} \psi_1^2 \\
&= - \int_{\mathbb{R}^2} (w + x \cdot \nabla w) \left\{ 3w\psi_1^2 - \left[\frac{3\lambda_0^4}{a^*} w^2 + V(x) \right] \psi_1 \right\} + \int_{\mathbb{R}^2} \psi_1^2 \\
&:= A + B,
\end{aligned} \tag{A.12}$$

where the part A satisfies

$$\begin{aligned}
A &= - \int_{\mathbb{R}^2} w \left\{ 3w\psi_1^2 - \left[\frac{3\lambda_0^4}{a^*} w^2 + V(x) \right] \psi_1 \right\} + \int_{\mathbb{R}^2} \psi_1^2 \\
&= - \int_{\mathbb{R}^2} |\nabla \psi_1|^2 + \frac{2\lambda_0^4}{a^*} \int_{\mathbb{R}^2} w^3 \psi_1.
\end{aligned} \tag{A.13}$$

Since

$$\int_{\mathbb{R}^2} (x \cdot \nabla \psi_1) \Delta \psi_1 = -2 \int_{\mathbb{R}^2} |\nabla \psi_1|^2 - \int_{\mathbb{R}^2} (x \cdot \nabla \psi_1) \Delta \psi_1,$$

i.e., $-\int_{\mathbb{R}^2} |\nabla \psi_1|^2 = \int_{\mathbb{R}^2} (x \cdot \nabla \psi_1) \Delta \psi_1$, it follows from (A.13) that the part A can be rewritten as

$$A = \int_{\mathbb{R}^2} (x \cdot \nabla \psi_1) \Delta \psi_1 + \frac{2\lambda_0^4}{a^*} \int_{\mathbb{R}^2} w^3 \psi_1. \tag{A.14}$$

We rewrite the part B of (A.12) as

$$B = - \int_{\mathbb{R}^2} (x \cdot \nabla w) \left\{ 3w\psi_1^2 - \left[\frac{3\lambda_0^4}{a^*} w^2 + V(x) \right] \psi_1 \right\} = B_1 + B_2 + B_3, \quad (\text{A.15})$$

where the term B_2 satisfies

$$B_2 = \frac{3\lambda_0^4}{a^*} \int_{\mathbb{R}^2} (x \cdot \nabla w) w^2 \psi_1 = -\frac{2\lambda_0^4}{a^*} \int_{\mathbb{R}^2} w^3 \psi_1 - \frac{\lambda_0^4}{a^*} \int_{\mathbb{R}^2} w^3 (x \cdot \nabla \psi_1).$$

Together with (A.14), we obtain from above that

$$A + B_2 = \int_{\mathbb{R}^2} (x \cdot \nabla \psi_1) \Delta \psi_1 - \frac{\lambda_0^4}{a^*} \int_{\mathbb{R}^2} w^3 (x \cdot \nabla \psi_1). \quad (\text{A.16})$$

The term B_1 of (A.15) satisfies

$$\begin{aligned} B_1 &= - \int_{\mathbb{R}^2} (x \cdot \nabla w) 3w\psi_1^2 = 3 \int_{\mathbb{R}^2} w^2 \psi_1^2 + 3 \int_{\mathbb{R}^2} w^2 \psi_1 (x \cdot \nabla \psi_1) \\ &= \int_{\mathbb{R}^2} (3w^2 - 1) \psi_1^2 + \int_{\mathbb{R}^2} (3w^2 - 1) \psi_1 (x \cdot \nabla \psi_1), \end{aligned} \quad (\text{A.17})$$

due to the fact that

$$- \int_{\mathbb{R}^2} \psi_1 (x \cdot \nabla \psi_1) = 2 \int_{\mathbb{R}^2} \psi_1^2 + \int_{\mathbb{R}^2} \psi_1 (x \cdot \nabla \psi_1).$$

But the term B_3 of (A.15) satisfies

$$\begin{aligned} B_3 &= \int_{\mathbb{R}^2} \psi_1 V(x) (x \cdot \nabla w) \\ &= -2 \int_{\mathbb{R}^2} V(x) w \psi_1 - \int_{\mathbb{R}^2} w \psi_1 [x \cdot \nabla V(x)] - \int_{\mathbb{R}^2} w V(x) (x \cdot \nabla \psi_1) \\ &= -4 \int_{\mathbb{R}^2} V(x) w \psi_1 - \int_{\mathbb{R}^2} w V(x) (x \cdot \nabla \psi_1), \end{aligned} \quad (\text{A.18})$$

since $V(x)$ satisfies $x \cdot \nabla V(x) = 2V(x)$. Applying (A.15)–(A.18), we now obtain from (3.3) and (A.12) that

$$\begin{aligned} I &= (A + B_2) + B_1 + B_3 \\ &= \int_{\mathbb{R}^2} (3w^2 - 1) \psi_1^2 - 4 \int_{\mathbb{R}^2} V(x) w \psi_1 \\ &\quad + \int_{\mathbb{R}^2} (x \cdot \nabla \psi_1) \left\{ (\Delta - 1 + 3w^2) \psi_1 - \frac{\lambda_0^4}{a^*} w^3 - V(x) w \right\} \\ &= \int_{\mathbb{R}^2} (3w^2 - 1) \psi_1^2 - 4 \int_{\mathbb{R}^2} V(x) w \psi_1, \end{aligned} \quad (\text{A.19})$$

and the claim (3.36) is therefore proved in view of (1.11). \square

Acknowledgements: The author is very grateful to the referees for many valuable suggestions which lead to the great improvements of the present paper. The author also thanks Dr. Yong Luo very much for his fruitful discussions on the present paper.

References

- [1] J. R. Abo-Shaeer, C. Raman, J. M. Vogels and W. Ketterle, *Observation of vortex lattices in Bose-Einstein condensate*, Science **292** (2001), 476.
- [2] A. Aftalion, *Vortices in Bose-Einstein condensates*, Progress in Nonlinear Differential Equations and their Applications, 67. Birkhäuser Boston, Inc., Boston, MA, 2006.
- [3] A. Aftalion, S. Alama and L. Bronsard, *Giant vortex and breakdown of strong pinning in a rotating Bose-Einstein condensate*, Arch. Ration. Mech. Anal. **178** (2005), 247–286.
- [4] A. Aftalion, R. L. Jerrard and J. Royo-Letelier, *Non-existence of vortices in the small density region of a condensate*, J. Funct. Anal. **260** (2011), 2387–2406.
- [5] A. Aftalion, B. Noris, and C. Sourdis, *Thomas-Fermi approximation for coexisting two component Bose-Einstein condensates and nonexistence of vortices for small rotation*, Comm. Math. Phys. **336** (2015), no. 2, 509–579.
- [6] M. H. Anderson, J. R. Ensher, M. R. Matthews, C. E. Wieman and E. A. Cornell, *Observation of Bose-Einstein condensation in a dilute atomic vapor*, Science **269** (1995), 198–201.
- [7] J. Arbunich, I. Nenciu and C. Sparber, *Stability and instability properties of rotating Bose-Einstein condensates*, Lett. Math. Phys. **109** (2019), 1415–1432.
- [8] G. Arioli and A. Szulkin, *A semilinear Schrödinger equation in the presence of a magnetic field*, Arch. Ration. Mech. Anal. **170** (2003), 277–295.
- [9] W. Bao and Y. Cai, *Ground states of two-component Bose-Einstein condensates with an internal atomic Josephson junction*, East Asia J. Appl. Math. **1** (2011), 49–81.
- [10] T. Bartsch, L. Jeanjean and N. Soave, *Normalized solutions for a system of coupled cubic Schrödinger equations on \mathbb{R}^3* , J. Math. Pures Appl. **106** (2016), no. 4, 583–614.
- [11] T. Bartsch, Y. Y. Liu and Z. L. Liu, *Normalized solutions for a class of nonlinear Choquard equations*, SN Partial Differ. Equ. Appl. (2020), 1: 34.
- [12] T. Bartsch, X. Zhong and W. Zou, *Normalized solutions for a coupled Schrödinger system*, Math. Ann. **380** (2021), 1713–1740.
- [13] N. Basharat, H. Hajaiej, Y. Hu and S. J. Zheng, *Threshold for blowup and stability for nonlinear Schrödinger equation with rotation*, submitted (2020), 37 pages, <https://arxiv.org/abs/2002.04722>
- [14] F. Bethuel, H. Brezis, and F. Helein, *Ginzburg-Landau Vortices*, Progress in Nonlinear Differential Equations and their Applications **13**, Birkhauser Boston, Inc., Boston, MA, 1994.
- [15] I. Bloch, J. Dalibard and W. Zwerger, *Many-body physics with ultracold gases*, Reviews of Modern Phys. **80** (2008), 885–964.

- [16] C. C. Bradley, C. A. Sackett, J. J. Tollett and R. G. Hulet, *Evidence of Bose-Einstein condensation in an atomic gas with attractive interactions*, Phys. Rev. Lett. **75** (1995), 1687. *Erratum* Phys. Rev. Lett. **79** (1997), 1170.
- [17] D. M. Cao and Z. W. Tang, *Existence and uniqueness of multi-bump bound states of nonlinear Schrödinger equations with electromagnetic fields*, J. Differential Equations **222** (2006), no. 2, 381–424.
- [18] L. D. Carr and C. W. Clark, *Vortices in attractive Bose-Einstein condensates in two dimensions*, Phys. Rev. Lett. **97** (2006), 010403.
- [19] Y. Castin and R. Dum, *Bose-Einstein condensates with vortices in rotating traps*, European Phys. J. D **7** (1999), 399–412.
- [20] T. Cazenave, *Semilinear Schrödinger equations*, Courant Lecture Notes in Mathematics Vol. 10, Courant Institute of Mathematical Science/AMS, New York, 2003.
- [21] T. Cazenave and P. L. Lions, *Orbital stability of standing waves for some nonlinear Schrödinger equations*, Comm. Math. Phys. **85** (1982), no. 4, 549–561.
- [22] M. Correggi, F. Pinsker, N. Rougerie, and J. Yngvason, *Rotating superfluids in anharmonic traps: From vortex lattices to giant vortices*, Phys. Rev. A **84** (2011), p. 053614.
- [23] M. Correggi and N. Rougerie, *Boundary behavior of the Ginzburg-Landau order parameter in the surface superconductivity regime*, Arch. Ration. Mech. Anal. **219** (2016), 553–606.
- [24] F. Dalfovo, S. Giorgini, L. P. Pitaevskii and S. Stringari, *Theory of Bose-Einstein condensation in trapped gases*, Reviews of Modern Phys. **71** (1999), 463–512.
- [25] M. J. Esteban and P. L. Lions, *Stationary solutions of nonlinear Schrödinger equations with an external magnetic field*, Partial differential equations and the calculus of variations, Vol. I, 401–449, Progr. Nonlinear Differential Equations Appl. 1, Birkhuser Boston, Boston, MA, 1989.
- [26] A. L. Fetter, *Rotating trapped Bose-Einstein condensates*, Reviews of Modern Phys. **81** (2009), 647–691.
- [27] R. L. Frank, *Ground states of semi-linear PDEs*, Lecture notes from “Summer school on Current Topics in Mathematical Physics”, CIRM Marseille, 2013.
- [28] B. Gidas, W. Ni and L. Nirenberg, *Symmetry of positive solutions of nonlinear elliptic equations in \mathbb{R}^n* , Mathematical analysis and applications Part A, Adv. in Math. Suppl. Stud. Vol. **7**, Academic Press, New York (1981), 369–402.
- [29] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer, 1997.
- [30] T. X. Gou, Z. T. Zhang, *Normalized solutions to the Chern-Simons-Schrödinger system*, J. Funct. Anal. **280** (2021), 108894, 65 pp.

- [31] Y. J. Guo, C. Lin and J. C. Wei, *Local uniqueness and refined spike profiles of ground states for two-dimensional attractive Bose-Einstein condensates*, SIAM J. Math. Anal. **49** (2017), 3671–3715.
- [32] Y. J. Guo, Y. Luo and S. J. Peng, *Local uniqueness of ground states for rotating Bose-Einstein condensates with attractive interactions*, Calc. Var. Partial Differential Equations **60** (2021), Paper No. 237, 27pp.
- [33] Y. J. Guo, Y. Luo and S. J. Peng, *Existence and asymptotic behavior of ground states of rotating Bose-Einstein condensates*, submitted, (2021), 31 pages, arxiv.org/abs/2106.14369
- [34] Y. J. Guo, Y. Luo and W. Yang, *The nonexistence of vortices for rotating Bose-Einstein condensates with attractive interactions*, Arch. Rational Mech. Anal. **238** (2020), 1231–1281.
- [35] Y. J. Guo and R. Seiringer, *On the mass concentration for Bose-Einstein condensates with attractive interactions*, Lett. Math. Phys. **104** (2014), 141–156.
- [36] Y. J. Guo, Z. Q. Wang, X. Y. Zeng and H. S. Zhou, *Properties of ground states of attractive Gross-Pitaevskii equations with multi-well potentials*, Nonlinearity **31** (2018), 957–979.
- [37] C. Huepe, S. Metens, G. Dewel, P. Borckmans and M.E. Brachet, *Decay rates in attractive Bose-Einstein condensates*, Phys. Rev. Lett. **82** (1999), 1616–1619.
- [38] R. Ignat and V. Millot, *The critical velocity for vortex existence in a two-dimensional rotating Bose-Einstein condensate*, J. Funct. Anal. **233** (2006), 260–306.
- [39] R. Ignat and V. Millot, *Energy expansion and vortex location for a two-dimensional rotating Bose-Einstein condensate*, Rev. Math. Phys. **18** (2006), 119–162.
- [40] L. Jeanjean, *Existence of solutions with prescribed norm for semilinear elliptic equations*, Nonlinear Anal. **28** (1997), no. 10, 1633–1659.
- [41] M. K. Kwong, *Uniqueness of positive solutions of $\Delta u - u + u^p = 0$ in \mathbb{R}^N* , Arch. Rational Mech. Anal. **105** (1989), 243–266.
- [42] M. Lewin, P. T. Nam and N. Rougerie, *The mean-field approximation and the nonlinear Schrödinger functional for trapped Bose gases*, Trans. Amer. Math. Soc. **368** (2016), 6131–6157.
- [43] M. Lewin, P. T. Nam and N. Rougerie, *Blow-up profile of rotating 2D focusing Bose gases*, Macroscopic Limits of Quantum Systems, Springer Verlag, 2018.
- [44] E. H. Lieb and M. Loss, *Analysis*, Graduate Studies in Mathematics Vol. 14. Amer. Math. Soc., Providence, RI, second edition, 2001.
- [45] E. H. Lieb and R. Seiringer, *Derivation of the Gross-Pitaevskii equation for rotating Bose gases*, Comm. Math. Phys. **264** (2006), 505–537.

- [46] E. H. Lieb, R. Seiringer, J. P. Solovej and J. Yngvason, *The mathematics of the Bose gas and its condensation*, Oberwolfach Seminars, **34** Birkhäuser Verlag, Basel, 2005.
- [47] E. H. Lieb, R. Seiringer and J. Yngvason, *Bosons in a trap: A rigorous derivation of the Gross-Pitaevskii energy functional*, Phys. Rev. A **61** (2000), 043602.
- [48] E. Lundh, A. Collin and K.-A. Suominen, *Rotational states of Bose gases with attractive interactions in anharmonic traps*, Phys. Rev. Lett. **92** (2004), 070401.
- [49] P. Luo, S. Peng, J. Wei and S. Yan, *Excited states on Bose-Einstein condensates with attractive interactions*, Calc. Var. Partial Differential Equations **60** (2021), Paper No. 155, 30pp.
- [50] K. Madison, F. Chevy, J. Dalibard and W. Wohlleben, *Vortex formation in a stirred Bose-Einstein condensate*, Phys. Rev. Lett. **84** (2000), 806.
- [51] K. Madison, F. Chevy, J. Dalibard and W. Wohlleben, *Vortices in a stirred Bose-Einstein condensate*, J. Mod. Opt. **47** (2000), 2715–2723.
- [52] P. T. Nam and N. Rougerie, *Improved stability for 2D attractive Bose gases*, J. Math. Phys. **61** (2020), 021901.
- [53] W. M. Ni and I. Takagi, *On the shape of least-energy solutions to a semilinear Neumann problem*, Comm. Pure Appl. Math. **44** (1991), 819–851.
- [54] B. Pellacci, A. Pistoia, G. Vaira and G. Verzini, *Normalized concentrating solutions to nonlinear elliptic problems*, J. Differential Equations **275** (2021), 882–919.
- [55] N. Rougerie, *Non linear Schrödinger limit of bosonic ground states, again*, Confluentes Math. **12** (2020), no. 1, 69–91.
- [56] N. Rougerie, *Scaling limits of bosonic ground states, from many-body to nonlinear Schrödinger*, EMS Surveys in Math. Sciences **7** (2020), 253–408.
- [57] E. Sandier and S. Serfaty, *Vortices in the Magnetic Ginzburg-Landau Model*, Progress in Nonlinear Differential Equations and their Applications **70**, Basel: Birkhäuser, 2007.
- [58] I. M. Sigal, *Magnetic vortices, Abrikosov lattices and automorphic functions*, Math. and comput. modeling, 19–58, Pure Appl. Math. (Hoboken), Wiley, Hoboken, NJ, 2015.
- [59] J. C. Wei, *On the construction of single-peaked solutions to a singularly perturbed semilinear Dirichlet problem*, J. Diff. Eqns. **129** (1996), no. 2, 315–333.
- [60] M. I. Weinstein, *Nonlinear Schrödinger equations and sharp interpolations estimates*, Comm. Math. Phys. **87** (1983), 567–576.
- [61] J. Zhang, *Stability of attractive Bose-Einstein condensates*, J. Stat. Phys. **101** (2000), 731–746.