

## SHADOWS OF CUBE VERTICES

YOSSI LONKE

ABSTRACT. Suppose that a finite-dimensional cube is orthogonally projected onto a central section of itself by a subspace of one dimension less. Up to dimension 9, at least one vertex is projected onto the section, but for dimension 10 or larger, there are orthogonal projections for which all the vertices are projected outside the section. In fact, this is the case for “most” orthogonal projections, as the dimension tends to infinity.

If the vertices of a square or a three-dimensional cube are projected onto a line (respectively, plane) through its center then at least one pair of opposite vertices is projected onto the intersection with the line (respectively, plane). *What is the situation for higher dimensions?* The purpose of this note is to answer this question.

For a unit vector  $u \in \mathbb{R}^n$ , let  $H_u$  denote the  $(n-1)$ -dimensional subspace of all vectors in  $\mathbb{R}^n$  orthogonal to  $u$ , and  $\pi_u$  the orthogonal projection onto  $H_u$ . Thus for each  $x \in \mathbb{R}^n$ ,

$$(1) \quad \pi_u(x) = x - \langle x, u \rangle u,$$

where  $\langle \cdot, \cdot \rangle$  is the standard inner product. The three standard norms on  $\mathbb{R}^n$  are denoted by

$$\|u\|_1 = \sum_{k=1}^n |u_k|, \quad \|u\|_2 = \left( \sum_{k=1}^n u_k^2 \right)^{1/2}, \quad \|u\|_\infty = \max_{1 \leq k \leq n} |u_k|,$$

and the cube  $[-1, 1]^n$  is denoted by  $C_n$ .

**Theorem 1.** *Let  $u \in \mathbb{R}^n$  be a unit vector not orthogonal to any of the vertices of the cube  $C_n$ , and  $\pi_u$  the orthogonal projection onto  $H_u$ . A necessary and sufficient condition that there exists a vertex  $\varepsilon \in C_n$  such that  $\pi_u(\varepsilon) \in H_u \cap C_n$  is  $\|u\|_1 \|u\|_\infty \leq 2$ .*

It turns out (see Lemma 1 below) that the maximal value of  $\|u\|_1 \|u\|_\infty$  on the unit sphere  $\mathbb{S}^{n-1}$  is  $\frac{\sqrt{n+1}}{2}$ . Thus the condition of the theorem is fulfilled for all  $u \in \mathbb{S}^{n-1}$  if and only if  $n \leq 9$ .

**Corollary 1.** *Let  $n \geq 1$  be an integer. The following are equivalent:*

- (1) *For every orthogonal projection  $\pi_u$  onto the  $(n-1)$ -dimensional subspace  $H_u$  of  $\mathbb{R}^n$ , there exists a vertex  $\varepsilon$  of the cube  $C_n$  such that  $\pi_u(\varepsilon) \in H_u \cap C_n$ .*
- (2)  $n \leq 9$ .

*Proof.* By symmetry, it can and will be assumed that the nonzero coordinates of  $u$  are positive. *The condition is necessary:* choose a vertex  $\varepsilon = (\varepsilon_1, \dots, \varepsilon_n)$  for which  $\|\pi_u(\varepsilon)\|_\infty \leq 1$ , and assume  $\langle \varepsilon, u \rangle > 0$  (otherwise, replace  $\varepsilon$  by  $-\varepsilon$ ). Let

$\{e_k\}_{k=1}^n$  denote the standard unit vectors in  $\mathbb{R}^n$ . Consider the  $k$ th coordinate of the projection,

$$(2) \quad \langle \pi_u(\varepsilon), e_k \rangle = \varepsilon_k - \langle \varepsilon, u \rangle u_k.$$

Pick an index  $k$  such that  $u_k > 0$ . If  $\varepsilon_k = -1$ , then

$$(3) \quad \langle \pi_u(\varepsilon), e_k \rangle = \varepsilon_k - \langle \varepsilon, u \rangle u_k < -1,$$

but  $\|\pi_u(\varepsilon)\|_\infty \leq 1$ , so  $u_k > 0$  implies  $\varepsilon_k = 1$ . Hence,

$$(4) \quad \langle \pi_u(\varepsilon), e_k \rangle = \begin{cases} \pm 1 & \text{if } u_k = 0, \\ 1 - u_k \sum_{i: u_i > 0} u_i & \text{if } u_k > 0. \end{cases}$$

As  $u_k \geq 0$  for all  $k$ , the sum of the nonzero coordinates of  $u$  is just  $\|u\|_1$ , and  $\|u\|_\infty = \max\{u_k : u_k > 0\}$ . Thus, (4) implies

$$(5) \quad \|\pi_u(\varepsilon)\|_\infty = \max_k |\langle \pi_u(\varepsilon), e_k \rangle| = \max\{1, |1 - \|u\|_\infty \|u\|_1|\},$$

and since  $\|\pi_u(\varepsilon)\|_\infty \leq 1$ , this implies  $\|u\|_1 \|u\|_\infty \leq 2$ .

*The condition is also sufficient*, because given a unit vector we can choose a vertex  $\varepsilon$  for which  $\varepsilon_k = 1$  for all  $k$  such that  $u_k > 0$ . If in addition we assume that  $u$  is not orthogonal to any vertex, then (4), and consequently (5), are valid. The latter, combined with the condition  $\|u\|_1 \|u\|_\infty \leq 2$ , implies that  $\|\pi_u(\varepsilon)\|_\infty \leq 1$ . The proof of the theorem is complete.  $\square$

The following lemma was mentioned above. Curiously, the bound of 2 appearing in the condition of the theorem is precisely the maximum of the corresponding function for  $n = 9$ .

**Lemma 1.** *For every  $n \geq 1$ , the maximum of the function  $f(u) = \|u\|_1 \|u\|_\infty$  on the unit sphere  $\mathbb{S}^{n-1}$  is  $\frac{\sqrt{n+1}}{2}$ .*

*Proof.* For reasons of symmetry the desired maximum coincides with the maximum of the function

$$g(u) = u_1(u_1 + u_2 + \cdots + u_n)$$

subject to the constraint  $u \in \mathbb{S}^{n-1}$ . The rest of the proof is a standard exercise in multivariable calculus, and is therefore omitted.  $\square$

**Remark 1.** *For sufficiently large  $n$ , most points  $u \in \mathbb{S}^{n-1}$ , in the sense of spherical measure, do not satisfy  $\|u\|_1 \|u\|_\infty \leq 2$ . In fact, in a sense that can be made precise using classical concentration of measure techniques, (see [1, (2.6)]) the “typical” order of magnitude of the product  $\|u\|_1 \|u\|_\infty$  in  $\mathbb{S}^{n-1}$  is  $\sqrt{\log n}$ . Moreover, those vectors that are orthogonal to some vertex of the cube consist of a union of  $2^{n-1} (n-2)$ -dimensional spheres, which do not occupy any positive measure, so for sufficiently large dimensions, “most” one-codimensional orthogonal projections carry all the cube’s vertices outside the section of the cube by the subspace being projected upon.*

#### REFERENCES

- [1] Figiel, T., Lindenstrauss, J., Milman, V.D., The dimension of almost spherical sections of convex bodies, *Acta Mathematica* **139** (1977) 53–94.

yossilonke@me.com