

QUOTIENT RINGS OF INTEGERS FROM A METRIC POINT OF VIEW

Ricardo Gallego Torromé
 Department of Mathematics
 Faculty of Mathematics, Natural Sciences and Information Technologies
 University of Primorska, Koper, Slovenia
 Email: rigato39@gmail.com

ABSTRACT. The theory of Gromov-Hausdorff convergence is applied to sequences of quotient rings of integers. It is shown the existence of limit rings (fields) as the Gromov-Hausdorff limits of sequences of metric quotient rings. The relation of these constructions with the field of the reals \mathbb{R} is discussed, showing that they are dense in \mathbb{R} but that they cannot be identified with the real field or with the rational field \mathbb{Q} , at least when \mathbb{R} and \mathbb{Q} are endowed with the usual metric structures. It is also shown that the limit rings can be endowed with an order relation.

MSC Class: 51F99; 11R20; 11R21.

1. INTRODUCTION

In this article quotient rings of integers and their limits when the number of elements goes to infinite are investigated by means of methods of metric geometry. The questions considered are originally motivated in a context of certain theory of dynamical systems. According to the general theory of non-reversible dynamics developed in [2], time parameters are described by subsets of ordered number fields. Subsets of the rational numbers \mathbb{Q} and the real numbers \mathbb{R} provide natural candidates for time parameters, but since the theory developed had a general scope, other discrete number fields were also considered. Since the prime fields $\mathbb{Z}/p\mathbb{Z}$ with p prime are endowed with a particular form of order relation, the prime fields are admissible to provide time parameter sets. Furthermore, we were interested in the case where the time parameter sets are equipped with a notion of *infinitesimal time laps*, namely a time interval which is small compared with any macroscopic time interval and that can be used in a definition of infinitesimal difference. Therefore, in the context of prime fields $\mathbb{Z}/p\mathbb{Z}$, it is natural to consider the limit $p \rightarrow +\infty$ for such purposes, since there is an infinitesimal time step given by the expression $1/(p - 1)$.

The construction of the limits $p \rightarrow +\infty$ of quotient rings $\mathbb{Z}/p\mathbb{Z}$ or, in the case when p takes values on the set of prime numbers, prime quotient fields, is investigated in this paper. The main tool used to show the existence of the limits is the theory of Gromov-Hausdorff convergence, as discussed in reference [1], chapter 7. As a result, we will show the existence of ordered rings (fields) obtained as limits in the sense of Gromov-Hausdorff of sequences composed by quotient rings (fields) endowed with metric structures. The limits provide examples of novel discrete, non-finite, ordered rings (fields). In the case of the limit of convergent sequences of prime fields, we conjecture that such limits are unique, a conjecture supported by the coincidence of the limits when applied to primes in different arithmetic progressions of Dirichlet's type.

Several questions arise on the nature and properties of the limit rings and fields that go beyond the metric point of view. In particular, the identification of the limit fields is an important open question. We show that for the case of successions of prime fields, the Gromov-Hausdorff limit field $\mathbb{Z}/\infty\mathbb{Z}$ is also a field which is dense in \mathbb{R} , but it cannot be identified with \mathbb{Q} or with \mathbb{R} .

The Gromov-Hausdorff limit fields admit arbitrarily close approximations in the Gromov-Hausdorff distance topology, whose minimal time laps is of the form $\delta t = 1/(p - 1)$. For large p , the time laps δt can be arbitrarily small compared with the other scale that appear in the theory, namely the diameter of $\mathbb{Z}/p\mathbb{Z}$, that for the metric used in the theory, is equal to 1 for every $p \in \mathbb{N} \setminus \{1\}$. This construction provides an infinitesimal time laps, necessary to consider differential relations in dynamical theories.

2. PRELIMINARY CONSIDERATIONS

Definition of the metric in $\mathbb{Z}/p\mathbb{Z}$. The quotient ring $\mathbb{Z}/p\mathbb{Z}$ is determined by the composition rules of the aggregate of equivalence classes $[k]$ module p , $[k] := \{n \in \mathbb{Z} \text{ s.t. } n - k = qp, q \in \mathbb{Z}\}$ with $k \in \{0, 1, 2, \dots, p - 1\}$. For each $p \in \mathbb{N}$ there is a natural metric function $d_p : \mathbb{Z}/p\mathbb{Z} \times \mathbb{Z}/p\mathbb{Z} \rightarrow \mathbb{R}$ defined by the expression

$$(2.1) \quad d_p([n], [m]) := |n_0 - m_0|, \quad n_0 \in [n], \quad m_0 \in [m], \quad 0 \leq n_0, m_0 \leq p - 1.$$

That the function d_p satisfies the axioms of a metric follows from the properties of the usual metric function on \mathbb{R} determined by the modulus function $|\cdot| : \mathbb{R} \rightarrow \mathbb{R}^+ \cup \{0\}$. In this way, the distance associated to the metric function (2.1) between $[n]$ and $[m]$ in $\mathbb{Z}/p\mathbb{Z}$ is $d_p([n], [m])$ and the norm of a class $[n]$ is just $\|[n]\|_p := d_p([0], [n])$.

The following properties of the distance function d_p are easily proved:

- (1) The induced topology in $\mathbb{Z}/p\mathbb{Z}$ from the metric function d_p and the discrete topology of $\mathbb{Z}/p\mathbb{Z}$ coincide.
- (2) The minimal distance $d_{min} := \min\{d_p([n], [m]), [n], [m] \in \mathbb{Z}/p\mathbb{Z}\}$ is equal to 1 for any $p \in \mathbb{N}$ greater or equal to 2.
- (3) $\mathbb{Z}/p\mathbb{Z}$ with the topology induced from d_p is Hausdorff separable. To have an intuition of this property, let us consider two points $[k_1] \neq [k_2] \in \mathbb{Z}/p\mathbb{Z}$. Then it is enough to consider the balls

$$B([k_i], 1/4) := \{[k] \in \mathbb{Z}_p \text{ s.t. } d_p([k], [k_i]) < 1/4\}, \quad i = 1, 2$$

of radii $1/4$. Then $B([k_1], 1/4) = \{[k_1]\}$, $B([k_2], 1/4) = \{[k_2]\}$ and are such that $B([k_1], 1/4) \cap B([k_2], 1/4) = \emptyset$.

- (4) For any $p \in \mathbb{N}$ greater or equal to 2 the diameter of $\mathbb{Z}/p\mathbb{Z}$ is the maximum distance between elements in $\mathbb{Z}/p\mathbb{Z}$,

$$diam(\mathbb{Z}/p\mathbb{Z}) := \max\{d_p([n], [m]), [n], [m] \in \mathbb{Z}/p\mathbb{Z}\}.$$

It follows that $diam(\mathbb{Z}/p\mathbb{Z}) = p - 1$.

Normalized metric. The *normalized metric* in $\mathbb{Z}/p\mathbb{Z}$ is defined as

$$\tilde{d}_p : \mathbb{Z}/p\mathbb{Z} \times \mathbb{Z}/p\mathbb{Z} \rightarrow \mathbb{R}, \quad ([n], [m]) \mapsto \frac{1}{diam(\mathbb{Z}/p\mathbb{Z})} d_p([n], [m]).$$

For this metric function one can prove the following fundamental properties:

- We have that $diam_{\tilde{d}_p}(\mathbb{Z}/p\mathbb{Z}) = 1, \forall p \in \mathbb{N} \setminus \{1\}$. Therefore, the spaces $\{(\mathbb{Z}/p\mathbb{Z}, \tilde{d}_p), p \in \mathbb{N}\}$ cannot be points (according to the characterization 7.4.6 in [1], since the diameter is non-zero).
- Each metric space $(\mathbb{Z}/p\mathbb{Z}, \tilde{d}_p)$ is compact with the metric topology of \tilde{d}_p .
- As a metric space, $(\mathbb{Z}/p\mathbb{Z}, \tilde{d}_p)$ cannot be identified with \mathbb{R} endowed with its usual metric structure.
- On each $\mathbb{Z}/p\mathbb{Z}$ there is a minimal distance between elements given by $1/(p - 1)$.
- We have that

$$\lim_{p \rightarrow +\infty} \tilde{d}_{min}(\mathbb{Z}/p\mathbb{Z}) = \lim_{p \rightarrow +\infty} \frac{1}{p - 1} = 0.$$

Order relation in $\mathbb{Z}/p\mathbb{Z}$. The ring \mathbb{Z}_p can be endowed with an order relation,

$$[n] < [m] \text{ iff } n_0 < m_0, \text{ with } n_0 \in [n], m_0 \in [m], 0 \leq n_0, m_0 \leq p - 1.$$

Let us remark that the order relations discussed above are not the usual order relations of ordered fields.

3. GROMOV-HAUSDORFF DISTANCE BETWEEN QUOTIENT RINGS

Let us consider the application of the Gromov-Hausdorff theory to the class of compact metric spaces

$$\mathcal{N} = \{(\mathbb{Z}/p\mathbb{Z}, \tilde{d}_p), p \in \mathbb{N} \setminus \{1\}\}.$$

The Gromov-Hausdorff distance between the metric spaces $(\mathbb{Z}/p_i\mathbb{Z}, d_{Z_{p_i}})$ and $(\mathbb{Z}/p_j\mathbb{Z}, d_{Z_{p_j}})$ is evaluated by applying the corresponding definition given in [1], section 7.3. Let us consider the isometries defining the metric functions $d_p : \mathbb{Z}_p \rightarrow \mathbb{R}$, namely the maps

$$(3.1) \quad v_p : \mathbb{Z}/p\mathbb{Z} \rightarrow \mathbb{R}, [n] \mapsto n_0, \text{ s.t. } 0 \leq n_0 \leq p-1, n_0 \in [n] \subset \mathbb{R}.$$

With the usual metric structure on \mathbb{R} provided by the module function $|\cdot| : \mathbb{R} \rightarrow \mathbb{R}^+ \cup \{0\}$, v_p is an isometric embedding of $\mathbb{Z}/p\mathbb{Z}$ in \mathbb{R} . The Hausdorff distance between $v(\mathbb{Z}/p_1\mathbb{Z}) \subset \mathbb{R}$ and $v(\mathbb{Z}/p_2\mathbb{Z}) \subset \mathbb{R}$ can be evaluated for any pair $p_1, p_2 \in \mathbb{N}$ and is given by

$$d_H(v_{p_1}(\mathbb{Z}/p_1\mathbb{Z}), v_{p_2}(\mathbb{Z}/p_2\mathbb{Z})) = |p_1 - p_2|.$$

From the geometric representation of the above isometric embedding and from the notion of Gromov-Hausdorff distance, the Gromov-Hausdorff distance between the compact, metric spaces $(\mathbb{Z}/p_1\mathbb{Z}, d_{p_1})$ and $(\mathbb{Z}/p_2\mathbb{Z}, d_{p_2})$ is

$$(3.2) \quad d_{GH}((\mathbb{Z}/p_1\mathbb{Z}, d_{p_1}), (\mathbb{Z}/p_2\mathbb{Z}, d_{p_2})) \leq |p_1 - p_2|.$$

Let us consider the analogous metric structures associated to the metric functions \tilde{d}_p on each $(\mathbb{Z}/p\mathbb{Z}, \tilde{d}_p) \in \mathcal{N}$. The corresponding Gromov-Hausdorff distance associated to \tilde{d}_p is given by the expression

$$(3.3) \quad \tilde{d}_{GH}((\mathbb{Z}/p_1\mathbb{Z}, \tilde{d}_{p_1}), (\mathbb{Z}/p_2\mathbb{Z}, \tilde{d}_{p_2})) \leq \frac{|p_1 - p_2|}{\min\{diam_{d_{p_1}}(\mathbb{Z}/p_1\mathbb{Z}), diam_{d_{p_2}}(\mathbb{Z}/p_2\mathbb{Z})\}}.$$

For the purposes of this paper, the convergence properties associated to the metrics \tilde{d}_{GH} are more adequate. Therefore, we will consider the convergence properties of \tilde{d}_{GH} instead than the properties of d_{GH} .

Let us recall that a subset \mathcal{C} of a metric space X is an ϵ -net if $d(x, \mathcal{C}) \leq \epsilon, \forall x \in X$. Then we recall too the following definition and theorem ([1], section 7.4):

Definition 3.1. *A class Υ of compact metric spaces is uniformly totally bounded if the following two conditions hold:*

- (1) *There is a constant $\delta > 0$ such that $diam(X) \leq \delta$ for all $X \in \Upsilon$.*
- (2) *For every $\epsilon > 0$ there exists a natural number $N = N(\epsilon)$ such that every $X \in \Upsilon$ contains an ϵ -net with no more than $N(\epsilon)$ points.*

Theorem 3.2. *Any uniformly totally bounded class Υ of compact metric spaces is pre-compact in the Gromov-Hausdorff topology. That is, any sequence of Υ contains a convergent sub-sequence.*

The uniformly totally bounded property and the corresponding result on convergence of subsequences can be applied to sequences of finite rings in \mathcal{N} . Let us consider the collection of metric spaces

$$\mathcal{N}_{\epsilon_0} := \{(\mathbb{Z}_p, \tilde{d}_p) \text{ s.t. } p > p_0 \text{ and } \frac{1}{p-1} < \epsilon_0\}$$

For $\epsilon < \epsilon_0$ one has that $\mathcal{N}_\epsilon \subset \mathcal{N}_{\epsilon_0}$. Then we have the following result,

Proposition 3.3. *For every $\epsilon_0 > 0$ the class \mathcal{N}_{ϵ_0} is totally uniformly bounded.*

Proof. Condition (1) holds, since $\text{diam}_{\tilde{d}_p}(\mathbb{Z}/p\mathbb{Z}) = 1$ for all $p \in \mathbb{N} \setminus \{1\}$.

In order to show that also the condition (2) is satisfied, for a given ϵ we distinguish two cases:

- (1) Let p be a natural number such that

$$\tilde{d}_p([\alpha], \mathbb{Z}/p\mathbb{Z}) \leq \frac{1}{\text{diam}_{\tilde{d}_p}(\mathbb{Z}/p\mathbb{Z})} = \frac{1}{p-1} \leq \epsilon \leq \epsilon_0,$$

for each $[\alpha] \in \mathbb{Z}/p\mathbb{Z}$. Such a choice of p satisfying $\frac{1}{p-1} \leq \epsilon$ implies that $\mathcal{C}(\epsilon) = \mathbb{Z}/p\mathbb{Z}$ is an ϵ -net with no more than p points.

- (2) If $p' > p$ we took $\mathcal{C}_{p'}$ to be the most equidistant distribution of p points in $\mathbb{Z}/p'\mathbb{Z}$. In this case, we have

$$\tilde{d}_{p'}([\alpha]', N_{p'}) \leq \frac{[p'/p]}{p'-1} = \frac{p'-r_0}{p'-1} \frac{1}{p},$$

where $[p'/p]$ is the integer part of the quotient p'/p and r_0 depends on p' and p and range in $\{0, \dots, p-1\}$. It is direct that

$$\tilde{d}_{p'}([\alpha]', N_{p'}) \leq 1/(p-1) \leq \epsilon,$$

from which follows that $\mathcal{C}_{p'}$ is an ϵ -net with p points, where p is a function of ϵ_0 .

Therefore, for each $p \in \mathbb{N}$ the ϵ_0 -net defined as discussed above contains maxima p points. □

As a consequence of *Proposition 3.3*, *Theorem 3.2* applies to the class \mathcal{N}_{ϵ_0} , since in order to find a convergent sub-sequence in \mathcal{N}_{ϵ_0} the initial terms for p small are irrelevant. Therefore,

Theorem 3.4. *For any ϵ_0 , the class of compact spaces \mathcal{N} contains convergent sequences in the Gromov-Hausdorff topology of compact metric spaces.*

By definition of the Gromov-Hausdorff convergence, if the subsequence $\{S_n\} \subset \mathcal{N}$ is convergent, then we have the relations

$$\lim_{n \rightarrow +\infty} \tilde{d}_n = \tilde{d}_\infty, \quad \lim_{n \rightarrow +\infty} (\mathbb{Z}/n\mathbb{Z}) = \mathbb{Z}/\infty\mathbb{Z}.$$

Thus the limit metric quotient ring is

$$(\mathbb{Z}/\infty\mathbb{Z}, \tilde{d}_\infty) := \lim_{n \rightarrow +\infty} (\mathbb{Z}/p_n\mathbb{Z}, \tilde{d}_{p_n}).$$

Let us consider the map

$$(3.4) \quad \text{diam}_{\tilde{d}} : \mathcal{N} \rightarrow \mathbb{R}, \quad (\mathbb{Z}/p\mathbb{Z}, \tilde{d}_p) \mapsto \text{diam}_{\tilde{d}_p}(\mathbb{Z}/p\mathbb{Z}).$$

Since the map $\text{diam}_{\tilde{d}} : \mathcal{N} \rightarrow \mathbb{R}$ is constant, for a given subsequence $\{S_n\} \subset \mathcal{N}$ it holds that

$$\text{diam}_{\tilde{d}_\infty}(\lim_{n \rightarrow +\infty} \mathbb{Z}/n\mathbb{Z}) = \lim_{n \rightarrow +\infty} \text{diam}_{\tilde{d}_n}(\mathbb{Z}/n\mathbb{Z}) = 1.$$

Then we have that

Lemma 3.5. *For any convergent subsequence $\{S_n\} \subset \mathcal{N}$ convergent to the limit space $(\mathbb{Z}/\infty\mathbb{Z}, \tilde{d}_\infty)$, the relation*

$$(3.5) \quad \text{diam}_{\tilde{d}_\infty} \mathbb{Z}/\infty\mathbb{Z} = 1$$

holds good.

Since the diameter $\text{diam}_{\tilde{d}_\infty}(\mathbb{Z}/\infty\mathbb{Z})$ is finite but it is not zero, we have

Corollary 3.6. *Given a convergent subsequence $\{S_n\} \subset \mathcal{N}$, the limit space $\mathbb{Z}/\infty\mathbb{Z}$ is compact but it is not a point.*

However, the limit space $(\mathbb{Z}/\infty\mathbb{Z}, \tilde{d}_\infty)$ depends in general on the subsequence $\{S_n\}$.

3.1. Application to prime fields. Let us consider the collection of prime fields as metric spaces,

$$(3.6) \quad \tilde{\mathcal{P}} := \{(\mathbb{Z}/p_i\mathbb{Z}, \tilde{d}_{\mathbb{Z}/p_i}), p_i \in \Pi\},$$

where $\Pi \subset \mathbb{N}$ denotes the aggregate of the increasing prime numbers. Given the collection $\tilde{\mathcal{P}}$, there exist convergent subsequences in $\tilde{\mathcal{P}}$ in the Gromov-Hausdorff sense,

Theorem 3.7. *The collection $\tilde{\mathcal{P}}$ contains a convergent subsequence of metric compact spaces.*

Proof. Since $\tilde{\mathcal{P}} \subset \mathcal{N}$, we can apply theorem 3.4. □

We would like to remark here two points. First, a direct construction of the set $\mathbb{Z}/\infty\mathbb{Z}$ as an aggregate of limits is not possible by the means considered until now, since the set $\mathbb{Z}/\infty\mathbb{Z}$ contains more elements than any $\mathbb{Z}/p\mathbb{Z}$ of the given sequence \mathcal{S} . Second, one can consider the question of uniqueness of the limit $\mathbb{Z}/\infty\mathbb{Z}$ as independent of the convergent subsequence $\{S_n\} \subset \mathcal{P}$ the sense of metric spaces. For this question, it is of relevance the following case:

Example 3.8. *An interesting example is the case of sequences $\{(\mathbb{Z}/D_n\mathbb{Z}, \tilde{d}_{D_n})\}$ where $\{D_n\}$ is of the form $\{an + b, n = 1, 2, \dots\}$ with a, b co-primes. By Dirichlet's theorem, the arithmetic progressions $\{D_n\}$ contain an infinite number of primes. One can apply the theory developed above to $\Gamma_D = \{(\mathbb{Z}/D_n\mathbb{Z}, \tilde{d}_{D_n})\}$. Intuitively, the distance between elements of convergent sequences in $\{(\mathbb{Z}/D_i\mathbb{Z}, \tilde{d}_{D_i})\}$ and $\{(\mathbb{Z}/D'_n\mathbb{Z}, \tilde{d}_{D'_n})\}$ decreases for arbitrary Dirichlet arithmetic progressions $\{D_n\}$ and $\{D'_n\}$.*

Example 3.8 provides an argument supporting the uniqueness of the limit rings and fields in this context, but this is not enough strong to support the uniqueness of the limits when considering sequences in \mathcal{N} or $\tilde{\mathcal{P}}$.

4. RING AND FIELD STRUCTURES IN \mathbb{Z}_∞

Lattices. Let us consider the product $\mathcal{Z} = \mathbb{Z}/p_1\mathbb{Z} \times \mathbb{Z}/p_2\mathbb{Z}$. Then a distance function is defined in \mathcal{Z} by the relation

$$(4.1) \quad \tilde{d}_{\mathcal{Z}} : \mathcal{Z} \times \mathcal{Z} \rightarrow \mathbb{R}, (([\alpha_1], [\alpha_2]), ([\lambda_1], [\lambda_2])) \mapsto \sqrt{\tilde{d}_{\mathbb{Z}/p_1}^2([\alpha_1], [\lambda_1]) + \tilde{d}_{\mathbb{Z}/p_2}^2([\alpha_2], [\lambda_2])}.$$

The generalization of this metric structure to product spaces $\mathcal{Z} := \prod_{i=1}^{\alpha} \mathbb{Z}/p_i\mathbb{Z}$ is direct.

The following question is on the existence of algebraic structures in the limit space $\mathbb{Z}/\infty\mathbb{Z}$:

- Is $\mathbb{Z}/\infty\mathbb{Z} = \lim_{n \rightarrow +\infty} S_n$, with $\mathcal{S} \subset \mathcal{N}$ a convergent subsequence, furnished with convenient operations, a ring?
- Is $\mathbb{Z}/\infty\mathbb{Z} = \lim_{n \rightarrow +\infty} S_n$ with $\mathcal{S} \subset \tilde{\mathcal{P}}$ a convergent subsequence, furnished with the convenient operations, a number field?

The following results answer positively the above questions.

Theorem 4.1. *Let $(\mathbb{Z}/\infty\mathbb{Z}, \tilde{d}_\infty)$ be the limit in the Gromov-Hausdorff sense of a convergent subsequence in \mathcal{N} . Then the limit space $\mathbb{Z}/\infty\mathbb{Z}$ can be furnished with a ring structure by the limiting operations $+_\infty$ and \cdot_∞ . If the convergent subsequence \mathcal{S} is in $\tilde{\mathcal{P}}$, then $(\mathbb{Z}/\infty\mathbb{Z}, +_\infty, \cdot_\infty)$ is a number field.*

Proof. Let us consider a convergent subsequence $\{(\mathbb{Z}/p\mathbb{Z}, \tilde{d}_p)\}$ of \mathcal{N} converging to $(\mathbb{Z}/\infty\mathbb{Z}, \tilde{d}_\infty)$. Then the algebraic addition operation $+_p : \mathbb{Z}/p\mathbb{Z} \times \mathbb{Z}/p\mathbb{Z} \rightarrow \mathbb{Z}/p\mathbb{Z}$ is equivalent to the set

$$\tilde{\dagger}_p := \{([n]_p, [m]_p, [n+m]_p) \in (\mathbb{Z}/p\mathbb{Z})^3, [n], [m] \in \mathbb{Z}/p\mathbb{Z}\}.$$

Since \mathcal{S} converges to $(\mathbb{Z}/\infty\mathbb{Z}, \tilde{d}_\infty)$ in the Gromov-Hausdorff sense, the sequence $\{(\tilde{\dagger}_p, \bar{d}_p)\}$, with p in the same index set than $\{\mathcal{S}_p\}$ and with

$$\bar{d}_p := \left(\tilde{d}_p^2(1) + \tilde{d}_p^2(2) + \tilde{d}_p^2(3) \right)^{1/2}$$

is convergent in the Gromov-Hausdorff sense towards the limit of the sequence $\{(\tilde{\dagger}_\infty, \bar{d}_\infty)\}$, where

$$\bar{d}_\infty = \left(\tilde{d}_\infty^2(1) + \tilde{d}_\infty^2(2) + \tilde{d}_\infty^2(3) \right)^{1/2}$$

and where $\tilde{\dagger}_\infty \subset (\mathbb{Z}/\infty\mathbb{Z})^3$. The limit $\{(\tilde{\dagger}_\infty, \bar{d}_\infty)\}$ determines an algebraic addition operation $+\infty : \mathbb{Z}/\infty\mathbb{Z} \times \mathbb{Z}/\infty\mathbb{Z} \rightarrow \mathbb{Z}/\infty\mathbb{Z}$ by the action of the canonical projections

$$pr_i : (\mathbb{Z}/\infty\mathbb{Z})^3 \rightarrow \mathbb{Z}/\infty\mathbb{Z}(i), \quad i = 1, 2, 3$$

when restricted to the subset $\{(\tilde{\dagger}_\infty, \bar{d}_\infty)\} \subset (\mathbb{Z}/\infty\mathbb{Z})^3$,

$$+\infty : pr_1(\tilde{\dagger}_\infty) \times pr_2(\tilde{\dagger}_\infty) \rightarrow pr_3(\tilde{\dagger}_\infty), \quad (pr_1(\alpha) \times pr_2(\alpha)) \mapsto pr_3(\alpha).$$

Since $\pi_i(\tilde{\dagger}_\infty) \simeq \mathbb{Z}/\infty\mathbb{Z}$, the aggregate $\tilde{\dagger}_\infty$ and the projections pr_i determine an abelian group structure in $\mathbb{Z}/\infty\mathbb{Z}$. One starts noting that since $[n]_p + [m]_p = [m]_p + [n]_p$ for all $p \in \mathbb{N} \setminus \{1\}$, then the commutativity also hold for the limits; otherwise will be two limits for the same convergent subsequence. The remaining grup properties can be proved by analogous arguments.

Similarly, the product operation $\cdot_\infty : \mathbb{Z}/\infty\mathbb{Z} \times \mathbb{Z}/\infty\mathbb{Z} \rightarrow \mathbb{Z}/\infty\mathbb{Z}$ is constructed using the the limits of subsets of the product

$$\tilde{\cdot}_p := \{([n]_p, [m]_p, [n \cdot m]_p) \in (\mathbb{Z}/p\mathbb{Z})^3, [n], [m] \in \mathbb{Z}/p\mathbb{Z}\}.$$

By taking limits in a similar way as before, one obtains $\tilde{\cdot}_\infty \subset (\mathbb{Z}/\infty\mathbb{Z})^3$. Then the projections pr_i and the corresponding limits, $\tilde{\cdot}_\infty$ determines a product operation on $\mathbb{Z}/\infty\mathbb{Z}$.

In the case of convergent sequences in $\tilde{\mathcal{P}}$, the limit inverse operation $\overset{-1}{\infty}$ is also well defined. \square

We introduce the notion of ϵ -approximation, that will be useful to prove the existence of a natural order relation in $\mathbb{Z}/\infty\mathbb{Z}$.

Definition 4.2. Let \mathcal{S} be a subsequence of \mathcal{N} convergent to $(\mathbb{Z}/\infty\mathbb{Z}, \tilde{d}_\infty)$. Then $(\mathbb{Z}/p\mathbb{Z}, \tilde{d}_p) \in \mathcal{S}$ is an ϵ -approximation to $(\mathbb{Z}/\infty\mathbb{Z}, \tilde{d}_\infty)$ if and only if

$$d_{GH}((\mathbb{Z}/p\mathbb{Z}, \tilde{d}_p), (\mathbb{Z}/\infty\mathbb{Z}, \tilde{d}_\infty)) < \epsilon.$$

It is direct that if $\mathcal{S} \rightarrow (\mathbb{Z}/\infty\mathbb{Z}, \tilde{d}_\infty)$, then for every $\epsilon > 0$ there is a $p \in \mathbb{N}$ such that $(\mathbb{Z}/p\mathbb{Z}, \tilde{d}_p) \in \mathcal{S}$ is an ϵ -approximation.

Let us consider now the order relations

$$[n]_{p_1} \leq [m]_{p_1}, [n]_{p_2} \leq [m]_{p_2}, [n]_{p_3} \leq [m]_{p_3}, \dots$$

where $n, m < p_1 < p_2 < p_3 < \dots$ are such that $(\mathbb{Z}/p_i\mathbb{Z}, \tilde{d}_{p_i}) \in \mathcal{S}$ and $[n]_{p_i}, [m]_{p_i} \in \mathbb{Z}/p_i\mathbb{Z}$. For every $\epsilon > 0$ there is an ϵ -approximation of $(\mathbb{Z}/\infty\mathbb{Z}, \tilde{d}_\infty)$ such that the relation $[n]_p \leq [m]_p$ is be valid for any $p > p_0(\epsilon)$, for a large enough p_0 . This condition is equivalent to define a function which is constant for $p > p_0(\epsilon)$ and hence, continuous for large p , that allows to define an order relation in the limit space $(\mathbb{Z}/\infty\mathbb{Z}, \tilde{d}_\infty)$ as the value of the limits of all possible characteristic functions $f(n, m)_p$ for $p \rightarrow +\infty$ and all possible $n, m \in \mathbb{N} \setminus \{1\}$, where the function $f([m], [n])_p$ is defined by

$$\begin{aligned} f([m], [n])_p &:= 1, \text{ if } [n]_p \geq [m]_p, [n]_p, [m]_p \in \mathbb{Z}/p\mathbb{Z}, \\ &0, \text{ if } [m]_p \geq [n]_p, [n]_p, [m]_p \in \mathbb{Z}/p\mathbb{Z}. \end{aligned}$$

The sequence of functions $\{f_p\}$ is convergent in the limit $p \rightarrow +\infty$. As a consequence of this construction, we have that $f : \mathbb{Z}/\infty\mathbb{Z} \times \mathbb{Z}/\infty\mathbb{Z} \rightarrow \mathbb{R}$ determines a notion of order: two classes $[m], [n] \in \mathbb{Z}/\infty\mathbb{Z}$ are related as $[m] < [n]$ iff $f([m], [n]) = 1$, that is, if there is an ϵ -approximation such that $[m]_p < [n]_p$ for any p in the sub-sequence convergent $\{S_p\}$ large enough. Therefore, we have

Theorem 4.3. *The limit number ring $\mathbb{Z}/\infty\mathbb{Z}$ has an order relation.*

Let us remark that although the limit field $\mathbb{Z}/\text{infty}\mathbb{Z}$ admits an order relation, it is not an ordered field.

5. RELATION OF THE LIMIT FIELDS WITH THE REAL FIELD \mathbb{R}

Each of the limits $\mathbb{Z}/\infty\mathbb{Z}$ is dense in the reals \mathbb{R} in the following sense. Let us consider the injective map $\varphi_p : \mathbb{Z}/p\mathbb{Z} \rightarrow \mathbb{S}^1$, $[n] \mapsto \exp(2\pi in/p)$. Let \mathcal{S} be a convergent sequence of metric spaces $(\mathbb{Z}/p\mathbb{Z}, \tilde{d}_p)$ in the Gromov-Hausdorff sense. The collection of maps $\{\varphi_p \text{ s.t. } (\mathbb{Z}/p\mathbb{Z}, \tilde{d}_p) \in \mathcal{S}\}$ determines the limit map $\varphi_\infty : \mathbb{Z}/\infty\mathbb{Z} \rightarrow \mathbb{S}^1$, which is ϵ -close to the maps $\{\varphi_p, \text{ s.t. } (\mathbb{Z}/p\mathbb{Z}, \tilde{d}_p) \in \mathcal{S}\}$. We also consider the inverse of the projection map $s^{-1} : \mathbb{R} \rightarrow \mathbb{S}^1 \setminus \{(0, 1)\}$. Then one has that $\varphi_\infty(\mathbb{Z}/\infty\mathbb{Z})$ is dense in $s^{-1}(\mathbb{R}) = \mathbb{S}^1 \setminus \{\text{North}\}$.

However, the metric space $(\mathbb{Z}/\infty\mathbb{Z}, \tilde{d}_\infty)$ cannot be identified with the metric space of the real numbers with absolute value as distance function $(\mathbb{R}, ||)$, since the first one is compact while $(\mathbb{R}, ||)$ is not compact. Therefore, we end up with the conclusion that in the case of the limits of prime fields, $(\mathbb{Z}/\infty\mathbb{Z}, \tilde{d}_\infty)$ are a new class of fields endowed with metric functions and order relations.

REFERENCES

- [1] D. Burago, Y. Burago and S. Ivanov, *A course in metric geometry*, Graduate Studies in Mathematics Vol. 33, American Mathematical Society (2001).
- [2] R. Gallego Torromé, *General theory of non-reversible local dynamics*, International Journal of Geometric Methods in Modern Physics Vol. **18**, No. 07, 2150111 (2021).