

Certain properties of contra- T^*_{12} -continuous functions

Hadi J. Mustafa

Dept. of Math., F. of Math. & Comp. Sci.

University of Kufa

Najaf, Iraq

drhadi.mustafa@gmail.com

Layth M. Alabdulsada

Institute of Mathematics

University of Debrecen

H-4002 Debrecen, P.O. Box 400 Hungary

layth.muhsin@science.unideb.hu

Abstract—The concept of contra function was introduced by Dontchev [2], in this work, we use the notion of T^*_{12} -open to study a new class of function called a contra- T^*_{12} -continuous function as generalization of contra-continuous.

Keywords: T^*_{12} -open sets; contra- T^*_{12} -continuous function; operator topological space; contra- T^*_{12} -closed graph.

I. INTRODUCTION

In 1996, Dontchev [2] introduced contra-continuous functions. In [10], the authors introduced the concept of almost contra- T^* -continuous function. In this paper, we introduce a new class of function called contra- T^*_{12} -continuous function where T_1, T_2 are operators associated with the topology τ on X . Throughout the paper, the space X and Y or (X, Y) and (Y, δ) stand for topological space, let A be a subset of X . the closure of A and the interior of A will be denoted by $Cl(A)$ and $Int(A)$, respectively.

II. PRELIMINARIES

In this section, we recall the basic facts and definitions needed in this work.

2.1 Definition: A subset A of a space X is said to be:

- i) Semi-open [6] if $A \subseteq Cl(Int(A))$,
- ii) Pre-open [7] if $A \subseteq Int(Cl(A))$,
- iii) b-open [1] if $A \subseteq Cl(Int(A) \cup Int(Cl(A)))$.

The complement of semi-open (pre-open, b-open) is said to be semi-closed (pre-closed, b-closed). The family of all semi-open (pre-open, b-open, semi-closed, pre-closed, b-closed) subset of a space X is denoted by $SO(X)$ ($PO(X)$, $BO(X)$, $SC(X)$, $PC(X)$, $BC(X)$, respectively).

2.2 Definition [4]: A function $f : X \rightarrow Y$ is called semi-continuous (pre-continuous, b-continuous) if for each $x \in X$ and each open set V of Y containing $f(x)$, there exists $U \in SO(X)$ ($U \in PO(X)$, $U \in BO(X)$) such that $f(U) \subseteq V$.

2.3 Definition: A function $f : X \rightarrow Y$ is called contra-continuous [2] (contra-semi continuous [4], contra-pre-continuous [3], contra-b-continuous [5]) if $f^{-1}(V)$ is closed (semi-closed, pre-closed, b-closed, resp.) in X for each open set V of Y .

III. OPERATOR TOPOLOGICAL SPACES

3.1 Definition [8]: Let (X, τ) be a topological space and let $T: p(X) \rightarrow p(X)$ be a function (where $p(X)$ is the power set of X) we say that T is an operator associated with the topology τ on X if $W \subseteq T(W)$ ($W \in \tau$) and the triple (X, τ, T) is called an operator topological space.

3.2 Definition [9]: Let (X, τ, T) be an operator topological space, let $A \subseteq X$

- i) A is called T -open if given $x \in A$, then there exists $V \in \tau$ there exists $x \in V \subseteq T(V) \subseteq A$.
- ii) A is called T^* -open if $A \subseteq T(A)$ (A is not necessarily open).

3.3 Remarks:

- i) Every T -open set is open.
- ii) Every open set is T^* -open, so we have the following implications:

$$T\text{-open} \rightarrow \text{open} \rightarrow T^*\text{-open}$$

- iii) Let (X, τ) be a topological space define $T: p(X) \rightarrow p(X)$ as follows: $T(A) = \text{Int Cl}(A)$ then T is an operator associated with the topology τ on X and the triple (X, τ, T) is an operator topological space.

As an example, we can suppose $X = \mathbb{R}$, $\tau = t_u =$ the usual topology on \mathbb{R} , if

$$T(A) = \text{Int Cl}(A),$$

then the triple (\mathbb{R}, t_u, T) is an operator topological space,

notice that $Q \subset \mathbb{R}$ satisfies $Q \subseteq \text{Int}(\text{Cl}(Q))$, so Q is a T^* -open (pre-open) which is not open.

3.3 Definition: Let (X, τ) be a topological space and let T_1, T_2 be two operators associated with the topology τ on X then (X, τ, T_1, T_2) is called a bi operator topological space.

3.4 Definition: Let (X, τ, T_1, T_2) be an operator topological space and let $A \subseteq X$, we say that A is a T^*_{12} -open if $A \subseteq T_1(A) \cup T_2(A)$, the complement of T^*_{12} -open is called T^*_{12} -closed for example if:

$$T_1(A) = \text{Cl}(\text{Int}(A)),$$

$T_2(A) = \text{Int}(\text{Cl}(A))$, Then

$A \subseteq \text{Cl}(\text{Int}(A)) \cup \text{Int}(A)$,

this is the definition of b-open set.

Notice that every T^*_{1} -open (T^*_{2} -open) is T^*_{12} -open because if A is T^*_{1} -open then $A \subseteq T_1(A) \subseteq T_1(A) \cup T_2(A)$, so A will be T^*_{12} -open.

IV. CONTRA- T^*_{12} -CONTINUOUS FUNCTIONS

In this section, we obtain some properties of contra- T^*_{12} -continuous functions.

4.1 Lemma [1]: Let (X, τ) be a topological space then:

- 1) The intersection of an open set and a b-open set is a b-open set.
- 2) The union of any family of b-open sets is a b-open set.

Now, we generalize Lemma 4.1 as follows:

4.2 Lemma: Let (X, τ, T_1, T_2) be a bi operator topological space assume that

$$T_1(W \cap B) = T_1(W) \cap T_1(B), W \in \tau, B \subseteq X,$$

$$T_2(W \cap B) = T_2(W) \cap T_2(B), W \in \tau, B \subseteq X, \text{ therefore:}$$

- 1) The intersection of an open set and a T^*_{12} -open set is T^*_{12} -open.
- 2) The union of any family T^*_{12} -open sets is a T^*_{12} -open set.

Proof:

1) Let $W \in X$ be an open set and let V be a T^*_{12} -open set we have to prove that $W \cap V$ is also a T^*_{12} -open set. Since W is open then:

$$W \subseteq T_1(W) \quad \dots (1)$$

$$W \subseteq T_2(W) \quad \dots (2)$$

Since V is a T^*_{12} -open then

$$V \subseteq T_1(V) \cap T_2(V) \quad \dots (3)$$

$$\begin{aligned} W \cap V &\subseteq W \cap (T_1(V) \cap T_2(V)) \\ &= (W \cap T_1(V)) \cup (W \cap T_2(V)) \\ &\subseteq (T_1(W) \cap T_1(V)) \cup (T_2(W) \cap T_2(V)) \\ &= (T_1(W \cap V)) \cup (T_2(W \cap V)) \end{aligned}$$

Then $W \cap V$ is T^*_{12} -open set.

2) Let $\mathcal{L} = \{w_\alpha \mid \alpha \in I\}$ be any family of T^*_{12} -open sets we must prove that $\bigcup_\alpha w_\alpha$ is also a T^*_{12} -open

$w_\alpha \subseteq T_1(w_\alpha) \cup T_2(w_\alpha)$ for each $\alpha \in I$

$$\begin{aligned} \bigcup_\alpha w_\alpha &\subseteq \bigcup_\alpha (T_1(w_\alpha) \cup T_2(w_\alpha)) \\ &= \bigcup_\alpha T_1(w_\alpha) \cup \bigcup_\alpha T_2(w_\alpha) \end{aligned}$$

Now $\bigcup_\alpha T_1(w_\alpha) = T_1(\bigcup_\alpha w_\alpha)$

Also $\bigcup_\alpha T_2(w_\alpha) = T_2(\bigcup_\alpha w_\alpha)$

Then $\bigcup_\alpha w_\alpha \subseteq T_1(\bigcup_\alpha w_\alpha) \cup T_2(\bigcup_\alpha w_\alpha)$ and $\bigcup_\alpha w_\alpha$ is a T^*_{12} -open.

4.3 Remarks:

i) The intersection of two T^*_{12} -open is not necessarily T^*_{12} -open, so the collection of all T^*_{12} -open sets is not necessarily a topology on X .

Let $\tau^*_{(12)}$ be the topology generated by the collection of all T^*_{12} -open sets.

ii) The intersection of any collection of T^*_{12} -closed sets is T^*_{12} -closed. Let $T^*_{12}\text{-Cl}(B)$ -intersection of all T^*_{12} -closed sets containing B .

Recall that for a function $f: X \rightarrow Y$, the subset $\{(x, f(x)) \mid x \in X\} \subseteq X \times Y$ is called the graph of f and denoted by $G(f)$.

4.4 Definition: Let $f: (X, \tau, T_1, T_2) \rightarrow (Y, \delta)$ be a function the graph $G(f)$ of f is said to be contra- T^*_{12} -closed graph if for each $(x, y) \in (X \times Y) - G(f)$ there exists U which is T^*_{12} -open containing x and a closed set V of Y containing y such that $(U \times V) \cap G(f) = \emptyset$. This implies that $f(U) \cap V = \emptyset$.

4.5 Definition: A space X is said to be contra-compact if every closed cover of X has a finite sub cover.

4.6 Theorem : Let (X, τ, T_1, T_2) be a bi operator topological space and suppose $f: (X, \tau, T_1, T_2) \rightarrow (Y, \delta)$ has a contra- T^*_{12} -closed graph, then the inverse image of a contra-compact set A of Y is T^*_{12} -closed in X .

Proof: Assume that A is contra-compact set of A and $x \notin f^{-1}(A)$ for each $a \in A$, $(x, a) \notin G(f)$. Then there exists U_a which is T^*_{12} -closed containing x and V_a which is closed in Y containing a such that

$$f(U_a) \cap V_a = \emptyset.$$

Consider $\mathcal{L} = \{A \cap V_a \mid a \in A\}$ and \mathcal{L} is a closed cover of the subspace A , but A is contra-compact then there exists $a_1, a_2, a_3, \dots, a_n$ such that

$$A \subseteq \bigcup_{i=1}^n V_{a_i}.$$

$$\text{Let } U = \bigcap_{i=1}^n U_{a_i},$$

then U is T^*_{12} -closed containing x and $f(U) \cap A = \emptyset$, therefore

$U \cap f^{-1}(A) = \emptyset$ and hence $x \notin T^*_{12}\text{-Cl}(f^{-1}(A))$, this shows that $f^{-1}(A)$ is T^*_{12} -closed.

4.7 Theorem : Let Y be contra –compact space and let $(X, \tau^*(_{12}), T_1, T_2)$ be operator topological space ,suppose $f : (X, \tau^*(_{12}), T_1, T_2) \rightarrow (Y, \delta)$ has a contra- T^*_{12} -closed graph then f is contra T^*_{12} -continuous.

Proof: First we show that an open set U of Y is contra –compact and let $\mathcal{L} = \{ V_\alpha \mid \alpha \in \Lambda \}$ be a cover of U by closed sets V_α of U for each $\alpha \in \Lambda$, then there exists a closed set K_α of Y such that $V_\alpha = K_\alpha \cap U$,then the family $\{ K_\alpha \mid \alpha \in \Lambda \} \cup \{ U^c \}$ is closed cover of Y . But Y is contra-compact then there exists $\alpha_1, \alpha_2 \dots \alpha_n$ such that

$$Y = (\bigcup_{i=1}^n K_{\alpha_i}) \cup (U^c), \text{ hence}$$

$$U = \bigcup_{i=1}^n V_{\alpha_i}.$$

This show that U is contra-compact by (theorem 4.6) $f^{-1}(U)$ is a T^*_{12} -closed in X then for f is contra T^*_{12} -continuous.

4.8 Theorem: Let $f : (X, \tau, T_1, T_2) \rightarrow (Y, \delta)$ be a function and $g : X \rightarrow X \times Y$ the graph function of f defined by $g(x) = (x, f(x))$ for every $x \in X$, if g is contra- T^*_{12} -continuous then f is contra- T^*_{12} -continuous.

Proof: Since g is contra- T^*_{12} -continuous then $f^{-1}(U) = g^{-1}(X \times U)$ is a T^*_{12} -closed in X . Then f is contra- T^*_{12} -continuous.

4.9 Theorem : If $f : (X, \tau, T_1, T_2) \rightarrow (Y, \delta)$ is contra- T^*_{12} -continuous and $g : (X, \tau, T_1, T_2) \rightarrow (Y, \delta)$ is contra-continuous and Y is Urysohn space then $E = \{ x \in X \mid f(x) = g(x) \}$ is T^*_{12} -closed in X .

Proof: Let $x \in E^c$, then $f(x) \neq g(x)$, since Y is a Urysohn then there exists open sets V and W such that $f(x) \in V, g(x) \in W$, and

$$Cl(V) \cap Cl(W) = \emptyset.$$

Since f is contra- T^*_{12} -continuous then $f^{-1}(Cl(V))$ is T^*_{12} -open in X and g is contra-continuous then $g^{-1}(Cl(W))$ is open in X , let $U = f^{-1}(Cl(V)), G = g^{-1}(Cl(W))$.

Then $x \in U \cap G = A$, where A is T^*_{12} -open in X and

$$f(A) \cap g(A) \subseteq f(U) \cap g(G) \subseteq Cl(V) \cap Cl(W) = \emptyset, \text{ hence}$$

$$f(A) \cap g(A) = \emptyset \text{ and } A \cap E = \emptyset, A \subseteq E^c,$$

where A is T^*_{12} -open there for $x \notin T^*_{12}\text{-Cl}(E)$, then E is T^*_{12} -closed in X .

4.10 Definition: A subset A of operator topological space (X, τ, T_1, T_2) is said to be T^*_{12} -dense in X if $T^*_{12}\text{-Cl}(A) = X$.

4.11 Remarks: Let (X, τ) be a topological space define:

$$T_1: p(X) \rightarrow p(X)$$

$$T_2: p(X) \rightarrow p(X) \text{ as follows}$$

$$T_1(A) = \text{Int}(Cl(A))$$

$T_2(A) = Cl(Int(A))$, then T^*_{12} -dense subset will be b -dense and $T^*_{12}-Cl(A)$ will be $b-Cl(A)$ so b -dense in X mean that $b-Cl(A) = X$.

4.12 Corollary: Let $f: (X, \tau, T_1, T_2) \rightarrow (Y, \delta)$ is contra- T^*_{12} -continuous and $g: (X, \tau, T_1, T_2) \rightarrow (Y, \delta)$ is contra continuous if Y is Urysohn and $f = g$ on T^*_{12} -dense set $A \subseteq X$ then $f = g$ on X .

Proof: since f is contra- T^*_{12} -continuous and is contra continuous and Y is Urysohn by previous Theorem $E = \{x \in X: f(x) = g(x)\}$ is a T^*_{12} -closed in X . We have $f = g$ on T^*_{12} -dense set $A \subseteq E$, then $X = T^*_{12}-Cl(A) \subseteq T^*_{12}-Cl(E) = E$. Hence $f = g$ on X .

4.13 Definition: A bi operator topological space (X, τ, T_1, T_2) is called T^*_{12} -connected if X is not the Union of two non-empty T^*_{12} -open sets.

4.14 Theorem: If $f: (X, \tau, T_1, T_2) \rightarrow (Y, \delta)$ is contra- T^*_{12} -continuous from a T^*_{12} -connected space onto Y , then Y is not a discrete space.

Proof: Suppose that Y is discrete. Let $\emptyset \neq A \subset Y$ then A is proper nonempty open and closed subset of Y . Then $f^{-1}(A)$ is a proper nonempty T^*_{12} -clopen (T^*_{12} -open and T^*_{12} -closed) subset of X such that $X = f^{-1}(A) \cup (f^{-1}(A))^c$ which means that X is T^*_{12} -disconnected which is a contradiction. Hence Y is not discrete.

REFERENCES

- [1] D. Andrijević, "On b -open sets". Mat. Vesnik 48, 1996, 59-64.
- [2] J. Dontchev, "Contra-continuous functions and strongly S -closed spaces". Internat.J. Math. Math. Sci. 19, 1996, 303-310.
- [3] S. Jafari and T. Noiri, "On contra-precontinuous functions". Bull. Malaysian Math. Sc.Soc. 25 (2002), 115-128.
- [4] E. Ekici and M. Caldas, "Slightly τ -continuous functions", Bol. Soc. Paran. Mat. 22(2) (2004), 63-74.
- [5] A.A. Nasef, "Some properties of contra-continuous functions". Chaos Solitons Fractals 24 (2005), 471-477.
- [6] N. Levine, "Semi-open sets and semi-continuity in topological spaces". Amer. Math. Monthly 70 (1963), 36-41.
- [7] A. S. Mashhour, M. E. Abd El-Monsef and S. N. El-Deeb, "On precontinuous and weak precontinuous functions". Proc. Math. Phys. Soc. Egypt 51 (1982), 47-53.
- [8] Hadi J. Mustafa, A. Lafta. "Operator topological space". Journal the college of Education, Al-Mustansiriya University. (2009)
- [9] Hadi J. Mustafa, and A. Abdul Hassan, T -open sets. M.Sc thesis. Mu'ta University Jordan. (2004)
- [10] Hadi J. Mustafa, and Layth M. Alabdulsada "On Almost contra T^* -continuous functions". Journal the college Mathematics and computer Sciences, Vol. 1, no. 6. (2012)