

AN ELEMENTARY PROOF OF A FUNDAMENTAL RESULT IN REAL PHASE RETRIEVAL

PETER G. CASAZZA AND JANET C. TREMAIN

ABSTRACT. Edidin [9] proved a fundamental result in phase retrieval: Theorem: A family of orthogonal projections $\{P_i\}_{i=1}^m$ does phase retrieval in \mathbb{R}^n if and only if for every $0 \neq x \in \mathbb{R}^n$, the family $\{P_i x\}_{i=1}^m$ spans \mathbb{R}^n . The proof of this result relies on Algebraic Geometry and so is inaccessible to many people in the field. We will give an elementary proof of this result using only frame theory. We will also solve the complex version of this result by showing that the "if" part fails and the "only if" part holds in \mathbb{C}^n . Finally, we will show that these techniques can be used to verify two classifications of norm retrieval.

1. INTRODUCTION

We start with frames.

Definition 1.1. *A family of vectors $\{x_i\}_{i=1}^m$ is a **frame** for \mathbb{H}^n if there are constants $0 < A \leq B < \infty$ so that for every $x \in \mathbb{H}^n$:*

$$A\|x\|^2 \leq \sum_{i=1}^m |\langle x, x_i \rangle|^2 \leq B\|x\|^2.$$

We have

- (1) A, B are a set of **lower** (respectively, **upper**) frame bounds.
- (2) If $A = B$ this is an **A-tight frame** and if $A = B = 1$ it is a **Parseval frame**.
- (3) The operator $T : \mathbb{H}^n \rightarrow \ell_2(m)$ given by

$$Tx = (\langle x, x_1 \rangle, \langle x, x_2 \rangle, \dots, \langle x, x_m \rangle)$$

is called the **analysis operator** of the **frame operator**. The frame operator is a positive, invertible, self-adjoint operator on the space.

Phase retrieval for Hilbert space frames was introduced in [1] and quickly became one of the most active areas of research in frame theory.

Definition 1.2. *A frame $\{x_i\}_{i=1}^m$ does **phase retrieval** in H^n if whenever $x, y \in \mathbb{H}^n$ and*

$$|\langle x, x_i \rangle| = |\langle y, x_i \rangle|, \text{ for all } i = 1, 2, \dots, m,$$

then $x = cy$ for some $|c| = 1$.

2020 *Mathematics Subject Classification.* 42C15.

The first author was supported by NSF DMS 1906725.

In some applications a signal must be reconstructed from the norms of higher dimensional components. In X-ray crystallography such a problem arises in crystal twinning [8]. In this scenario, there exists a similar phase retrieval problem [4].

Definition 1.3. A family of subspaces $\{W_i\}_{i=1}^m$ with respective projections $\{P_i\}_{i=1}^m$ does **phase retrieval** if whenever $x, y \in \mathbb{H}^n$ and

$$\|P_i x\| = \|P_i y\|, \text{ for all } i = 1, 2, \dots, m,$$

then $x = cy$ for some $|c| = 1$.

2. BACKGROUND

We need:

Definition 2.1. A family of vectors $\{x_i\}_{i=1}^n$ in \mathbb{R}^n or \mathbb{C}^n has the **complement property** if whenever we partition the set of vectors into two subsets, one of them must span the space.

A fundamental result here appeared in [1]:

Theorem 2.2. A family of vectors $\{x_i\}_{i=1}^m$ in \mathbb{R}^n does phase retrieval if and only if it has the complement property.

Corollary 2.3. If a family of vectors $\{x_i\}_{i=1}^m$ satisfies phase retrieval in \mathbb{R}^n , then $m \geq 2n - 1$.

It is easy to get families of vectors doing phase retrieval.

Definition 2.4. A frame $\{x_i\}_{i=1}^m$ is **full spark** in \mathbb{H}^n if for every $I \subset [m]$ with $|I| = n$, $\text{span } \{x_i\}_{i \in I} = \mathbb{H}^n$.

It follows that any full spark set of $\geq 2n - 1$ vectors does phase retrieval. There are full spark sets with an arbitrary number of vectors in \mathbb{H}^n . Let $\{e_i\}_{i=1}^n$ be an orthonormal basis for \mathbb{H}^n . For every $I \subset [n]$ with $|I| = n - 1$, $\text{span } \{x_i\}_{i \in I}$ is a hyperplane. Since we only have a finite number of hyperplanes, there is a vector $x_{m+1} \in \mathbb{H}^n$ so that $x_{m+1} \notin \text{span } \{x_i\}_{i \in I}$ for any $I \subset [n]$ with $|I| = n - 1$, and hence $\{x_i\}_{i=1}^{m+1}$ is full spark. Now continue this. Choose a vector $x_{m+2} \in \mathbb{H}^n$ so that $x_{m+2} \in \mathbb{H}^n$ and $x_{m+2} \notin \text{span } \{x_i\}_{i \in I}$ for all $I \subset [m+1]$ and $|I| = n - 1$. Hence, $\{x_i\}_{i=1}^{n+2}$ is full spark.

The complex case is much more complicated. It is known [2] that phase retrieval can be done in \mathbb{C}^n with $4n - 4$ vectors. It was believed this result was best possible. But Vinzant [11] showed that phase retrieval can be done in \mathbb{C}^4 with 11 vectors (in other words $4n - 5$ vectors). So the major open problem here is:

Problem 2.5. How many vectors in \mathbb{C}^n does it take to do phase retrieval?

For phase retrieval by projections the problem is still more complicated. In [4] it was shown:

Theorem 2.6. *Let $\{P_i\}_{i=1}^n$ be orthogonal projections in \mathbb{R}^n with $P_i(\mathbb{R}^n) = W_i$, for $i = 1, 2, \dots, m$. The following are equivalent:*

- (1) $\{W_i\}_{i=1}^m$ allows phase retrieval.
- (2) Whenever $\{x_{ij}\}_{j=1}^{I_i}$ is an orthonormal basis for W_i , the family $\{x_{ij}\}_{i=1, j=1}^m, I_i$ does phase retrieval.

Recently, [6] the following was shown:

Theorem 2.7. *It takes at least $2n - 2$ subspaces to do phase retrieval in \mathbb{R}^n .*

The following theorem in [13] shows a particular case of \mathbb{R}^4 in which phase retrieval is possible with $2n - 2 = 2(4) - 2 = 6$ subspaces.

Theorem 2.8. *There are 6 two-dimensional subspaces of \mathbb{R}^4 satisfying phase retrieval.*

In [3] it was shown that there are 6 hyperplanes in \mathbb{R}^4 doing phase retrieval.

Another open problem in this area is motivated by this theorem of Edidin [9]:

Theorem 2.9. *If $n = 2^k + 1$, for some k , then it takes $2n - 1$ subspaces of \mathbb{R}^n to do phase retrieval.*

So now the main open problem in this area is:

Problem 2.10. *Can we do phase retrieval in \mathbb{R}^n with $2n - 2$ subspaces whenever $n \neq 2^k + 1$ for any k .*

Again, the complex case is much more complicated.

Problem 2.11. *What is the minimum number of projections required to do phase retrieval in \mathbb{C}^n ?*

Edidin [9] proved a fundamental result in phase retrieval:

Theorem 2.12. *A family of orthogonal projections $\{P_i\}_{i=1}^m$ does phase retrieval in \mathbb{R}^n if and only if for every $0 \neq x \in \mathbb{R}^n$, the family $\{P_i x\}_{i=1}^m$ spans \mathbb{R}^n .*

Because the proof relies on Algebraic Geometry, it has been inaccessible to some people in the field. We will give an elementary proof of this result without Algebraic Geometry. We will also resolve the complex case by showing that the "if" part of the theorem fails and the "only if" part of the theorem holds in \mathbb{C}^n . We will also show that these methods verify several classifications of norm retrieval.

Remark: Edidin's Theorem also holds for symmetric matrices $\{A_i\}_{i=1}^m$ instead of projections [13]

3. PHASE RETRIEVAL

We start with the real case.

3.1. Proof of Theorem 2.12 for the real case.

Proof. \Rightarrow : We do the contrapositive. Assume there is an $x \in R^n$ with $\|x\| = 1$ so that $\text{span}\{P_i x\}_{i=1}^m \neq R^n$. Choose $y \in R^n$ with $\|y\| = 1$ and $y \perp \text{span}\{P_i x\}_{i=1}^m$. Now,

$$0 = \langle y, P_i x \rangle = \langle P_i y, P_i x \rangle, \text{ for all } i = 1, 2, \dots, m.$$

Since $P_i x \perp P_i y$,

$$\|P_i(x + y)\|^2 = \|P_i x\|^2 + \|P_i y\|^2 = \|P_i(x - y)\|^2.$$

But, $x + y \neq \pm(x - y)$. So $\{P_i\}_{i=1}^m$ fails phase retrieval.

\Leftarrow : Let $W_i = P_i(R^n)$. If $\{W_i\}_{i=1}^m$ fails phase retrieval, then there exists orthonormal bases $\{x_{ij}\}_{j=1}^{I_i}$ for W_i which can be divided into two non-spanning sets $\{x_{ij}\}_{(i,j) \in I}$ and $\{x_{ij}\}_{(i,j) \in I^c}$. Choose

$$0 \neq x \perp \text{span}\{x_{ij}\}_{(i,j) \in I}.$$

Then, $P_i x \in \text{span}\{x_{ij}\}_{(i,j) \in (I^c \cap W_i)}$ for all i . But this family of sets does not span. □

3.2. Proof of the Complex Case.

Proof. \Rightarrow : Assume there is an $x \in \mathbb{C}^n$ with $\|x\| = 1$ so that $\{P_i x\}_{i=1}^m$ does not span. Choose $y \in \mathbb{C}^n$ with $\|y\| = 1$ so that $y \perp P_i x$, for all $i = 1, 2, \dots, m$. Now

$$\begin{aligned} \|P_i(x + y)\|^2 &= \|P_i x\|^2 + \|P_i y\|^2 + 2\text{Re}\langle P_i x, P_i y \rangle \\ &= \|P_i x\|^2 + \|P_i y\|^2 + 2\text{Re}\langle P_i x, y \rangle \\ &= \|P_i x\|^2 + \|P_i y\|^2. \end{aligned}$$

Similarly,

$$\|P_i(x - y)\|^2 = \|P_i x\|^2 + \|P_i y\|^2.$$

So if $\{P_i\}_{i=1}^m$ does phase retrieval, then there is some $|c| = 1$ so that $x + y = c(x - y)$. If $c = 1$ then $x + y = x - y$ and so $y = 0$, a contradiction. Otherwise, $(1 + c)y = (c - 1)x$ and so $x = \frac{1+c}{c-1}y$. Thus, $P_i x = \frac{1+c}{c-1}P_i y$. But this is impossible since $x, y \neq 0$ and

$$\langle P_i x, P_i y \rangle = \langle P_i x, y \rangle = 0, \text{ for all } i = 1, 2, \dots, m.$$

□

It is known (see for example, [11]) that for a family of vectors $\{x_i\}_{i=1}^m$ to do phase retrieval in \mathbb{C}^n , m must be on the order of $4n$.

Example 3.1. Let $\{x_i\}_{i=1}^{2n-1}$ be a full spark family of vectors in \mathbb{C}^n . This family cannot do phase retrieval. But if $0 \neq x \in \mathbb{C}^n$, then $|\{1 \leq i \leq 2n-1 : \langle x, x_i \rangle \neq 0\}| \geq n$. For otherwise, $|\{1 \leq i \leq 2n-1 : \langle x, x_i \rangle = 0\}| \geq n$ and since this family is full spark, $x = 0$ a contradiction. By the first inequality and the fact that this family is full spark, $\{x_i : \langle x, x_i \rangle \neq 0\}$ spans the space.

Remark: There is a complete complex version of Edidin's theorem: See Theorem 2.1 (3) in [13].

A completely different approach to a complex version of Edidin's theorem was proved in [7]. In this paper they associate vectors in \mathbb{C}^n with rank 2 projections in \mathbb{R}^{2n} .

4. NORM RETRIEVAL

Norm retrieval is a weakening of phase retrieval.

Definition 4.1. A frame $\{x_i\}_{i=1}^m$ in \mathbb{H}^n does **norm retrieval** if whenever $x, y \in \mathbb{H}^n$ and

$$|\langle x, x_i \rangle| = |\langle y, x_i \rangle|, \text{ for all } i = 1, 2, \dots, m,$$

then $\|x\| = \|y\|$.

If $\{x_i\}_{i=1}^m$ does phase retrieval, then it does norm retrieval. But an orthonormal basis $\{e_i\}_{i=1}^n$ does norm retrieval in \mathbb{H}^n and fails phase retrieval.

The techniques from the previous section classify norm retrieval [10].

Theorem 4.2. A family of vectors $\{x_i\}_{i=1}^m$ does norm retrieval if and only if whenever $I \subset [m]$ and $x \perp x_i$, for all $i \in I$ and $y \perp x_i$ for all $i \in I^c$, then $x \perp y$.

Proof. \Leftarrow : Assume $|\langle x, x_i \rangle| = |\langle y, x_i \rangle|$, for all $i = 1, 2, \dots, m$. Let

$$I = \{1 \leq i \leq m : \langle x, x_i \rangle = \langle y, x_i \rangle\}.$$

Then, $x + y \perp x_i$ for all $i \in I^c$ and $x - y \perp x_i$ for all $i \in I$. By assumption, $x + y \perp x - y$. Now,

$$0 = \langle x + y, x - y \rangle = \|x\|^2 - \|y\|^2, \text{ and so } \|x\| = \|y\|.$$

So $\{x_i\}_{i=1}^m$ does norm retrieval.

\Rightarrow : Assume $\{x_i\}_{i=1}^m$ does norm retrieval and assume $I \subset [m]$ and $x \perp x_i$ for all $i \in I$ and $y \perp x_i$ for all $i \in I^c$. Then

$$|\langle x + y, x_i \rangle| = |\langle x - y, x_i \rangle|, \text{ for all } i = 1, 2, \dots, m.$$

Since $\{x_i\}_{i=1}^m$ does norm retrieval,

$$\begin{aligned} \|x + y\|^2 &= \|x\|^2 + \|y\|^2 + 2\langle x, y \rangle \\ &= \|x - y\|^2 \\ &= \|x\|^2 + \|y\|^2 - 2\langle x, y \rangle. \end{aligned}$$

Hence, $\langle x, y \rangle = 0$ and $x \perp y$. □

There is a corresponding notion for projections.

Definition 4.3. A family of projections does **norm retrieval** if whenever $x, y \in \mathbb{H}^n$ and

$$\|P_i x\| = \|P_i y\|, \text{ for all } i = 1, 2, \dots, m,$$

then $\|x\| = \|y\|$.

In general, if $\{P_i\}_{i=1}^m$ does phase retrieval, it may not be true that $\{(I - P_i)\}_{i=1}^m$ does phase retrieval. For example, if [3] $\{W_i\}_{i=1}^6$ are hyperplanes doing phase retrieval in \mathbb{R}^4 , then $\{W_i^\perp\}_{i=1}^6$ are only 6 vectors in \mathbb{R}^4 and hence cannot do phase retrieval. What is needed here is norm retrieval.

Theorem 4.4. If $\{P_i\}_{i=1}^m$ does phase retrieval, then $\{(I - P_i)\}_{i=1}^m$ does phase retrieval in \mathbb{R}^n if and only if $\{(I - P_i)\}_{i=1}^m$ does norm retrieval.

Proof. \Rightarrow : If $\{(I - P_i)\}_{i=1}^m$ does phase retrieval, it does norm retrieval.

\Leftarrow : Assume $\{P_i\}_{i=1}^m$ does phase retrieval and $\{(I - P_i)\}_{i=1}^m$ does norm retrieval. If $x, y \in \mathbb{H}^n$ and

$$\|(I - P_i)x\| = \|(I - P_i)y\|, \text{ for all } i = 1, 2, \dots, m,$$

then by assumption, $\|x\| = \|y\|$. Also,

$$\|(I - P_i)x\|^2 = \|x\|^2 - \|P_i x\|^2 = \|(I - P_i)y\|^2 = \|y\|^2 - \|P_i y\|^2.$$

Hence, $\|P_i x\| = \|P_i y\|$, for all $i = 1, 2, \dots, m$ and since $\{P_i\}_{i=1}^m$ does phase retrieval, $x = \pm y$. I.e. $\{(I - P_i)\}_{i=1}^m$ does phase retrieval. \square

There is a form of norm retrieval for projections.

Theorem 4.5. Let $\{W_i\}_{i=1}^m$ be subspaces of \mathbb{H}^n with corresponding projections $\{P_i\}_{i=1}^m$. The following are equivalent:

- (1) $\{W_i\}_{i=1}^m$ does norm retrieval.
- (2) For every orthonormal basis $\{x_{ij}\}_{j=1}^{I_i}$ of W_i the family $\{x_{ij}\}_{i=1, j=1}^m$ does norm retrieval.

Proof. (1) \Rightarrow (2): Assume $\{W_i\}_{i=1}^m$ does norm retrieval. Let $\{x_{ij}\}_{j=1}^{I_i}$ be an orthonormal basis of W_i . Assume

$$|\langle x, x_{ij} \rangle| = |\langle y, x_{ij} \rangle|, \text{ for all } (i, j).$$

Then, for all $i = 1, 2, \dots, m$,

$$\|P_i x\|^2 = \sum_{j=1}^{I_i} |\langle x, x_{ij} \rangle|^2 = \sum_{j=1}^{I_i} |\langle y, x_{ij} \rangle|^2 = \|P_i y\|^2.$$

Since $\{P_i\}_{i=1}^m$ does norm retrieval, $\|x\| = \|y\|$. So $\{x_{ij}\}_{i=1, j=1}^m$ does norm retrieval.

(2) \Rightarrow (1): Assume (2). Let $x, y \in \mathbb{H}^n$ satisfy $\|P_i x\| = \|P_i y\|$, for all $i = 1, 2, \dots, m$. If $P_i x = \pm P_i y$ then $|\langle x, x_{ij} \rangle| = |\langle y, x_{ij} \rangle|$, for all $j = 1, 2, \dots, I_i$. Otherwise, let

$$x_{i1} := \frac{P_i x + P_i y}{\|P_i x + P_i y\|}, \quad x_{i2} = \frac{P_i x - P_i y}{\|P_i x - P_i y\|}.$$

Then

$$\langle x_{i1}, x_{i2} \rangle = \frac{1}{\|P_i x + P_i y\| \|P_i x - P_i y\|} (\|P_i x\|^2 - \|P_i y\|^2 + \langle P_i x, P_i y \rangle - \langle P_i y, P_i x \rangle) = 0.$$

So we can extend $\{x_{i1}, x_{i2}\}$ to an orthonormal basis for W_i , $\{x_{ij}\}_{j=1}^{I_i}$. Note that $P_i x, P_i y \in \text{span}\{x_{i1}, x_{i2}\}$. So,

$$\langle x, x_{ij} \rangle = \langle y, x_{ij} \rangle = 0, \quad \text{for all } j = 3, 4, \dots, I_i.$$

We also have by the above

$$\begin{aligned} |\langle x, P_i x + P_i y \rangle| &= \| \|P_i x\|^2 + \langle P_i x, P_i y \rangle \| \\ &= \| \|P_i y\|^2 + \langle P_i x, P_i y \rangle \| \\ &= |\langle P_i y, P_i y + P_i x \rangle| \\ &= |\langle y, P_i x + P_i y \rangle|. \end{aligned}$$

Similarly, $|\langle x, P_i x - P_i y \rangle| = |\langle y, P_i x - P_i y \rangle|$. Hence,

$$|\langle x, x_{ij} \rangle| = |\langle y, x_{ij} \rangle|, \quad \text{for all } j = 1, 2, \dots, I_i.$$

By (2), $\|x\| = \|y\|$, and so $\{P_i\}_{i=1}^m$ does norm retrieval. \square

We have an Edidin type theorem for norm retrieval [7].

Theorem 4.6. *Given projections $\{P_i\}_{i=1}^m$ in \mathbb{R}^n , the following are equivalent:*

- (1) $\{P_i\}_{i=1}^m$ does norm retrieval.
- (2) For every $0 \neq x \in \mathbb{R}^n$, we have

$$[\text{span} \{P_i x\}]^\perp \subset x^\perp.$$

- (3) For every $x \in \mathbb{H}^n$ we have $x \in \text{span} \{P_i x\}_{i=1}^m$.

Proof. (2) \Rightarrow (1): We will prove the contrapositive. If norm retrieval fails, then there are vectors $x, y \in \mathbb{R}^n$ with $\|P_i x\| = \|P_i y\|$, for all $i = 1, 2, \dots, m$ but $\|x\| \neq \|y\|$. This implies if $v = x + y$ and $w = x - y$, then v, w are not orthogonal. But, $w \in [\text{span} \{P_i v\}_{i=1}^m]^\perp$. So (2) fails.

(2) \Leftrightarrow (3): This is immediate.

(2) \Rightarrow (1): Again by the contrapositive. Assume there exists $v, w \in \mathbb{R}^n$ which are not orthogonal (so $w \neq 0$) but $w \perp \text{span} \{P_i v\}_{i=1}^m$ and so $P_i w \perp$

$P_i v$, for all $i = 1, 2, \dots, m$. Let $v = x + y$ and $w = x - y$. Since v, w are not orthogonal, $\|x\| \neq \|y\|$. But,

$$\begin{aligned} \|P_i(v+w)\|^2 &= \langle P_i v + P_i w, P_i v + P_i w \rangle \\ &= \|P_i v\|^2 + \|P_i w\|^2 \\ &= \langle P_i v - P_i w, P_i v - P_i w \rangle \\ &= \|P_i(v-w)\|^2. \end{aligned}$$

So $\{P_i\}_{i=1}^m$ fails norm retrieval. \square

The projection form of this is:

Theorem 4.7. *Let $\{W_i\}_{i=1}^m$ be subspaces of \mathbb{H}^n with respective projections $\{P_i\}_{i=1}^m$. The following are equivalent:*

- (1) $\{W_i\}_{i=1}^m$ does norm retrieval.
- (2) If $\{x_{ij}\}_{j=1}^{I_i}$ is an orthonormal basis for W_i , and $S \subset \{(ij) : 1 \leq i \leq m, 1 \leq j \leq I_i\}$ and $x, y \in \mathbb{H}^n$ satisfy

$$x \perp x_{ij} \text{ for } (ij) \in S \text{ and } y \perp x_{ij} \text{ for } (ij) \in S^c,$$

then $x \perp y$.

Proof. (1) \Rightarrow (2): Assume $\{W_i\}_{i=1}^m$ does norm retrieval and assume $S \subset \{(i, j) : 1 \leq i \leq m, 1 \leq j \leq I_i\}$, and $x \perp x_{ij}$ for all $(i, j) \in S$ and $y \perp x_{ij}$ for all $(i, j) \in S^c$. Then

$$|\langle x + y, x_{ij} \rangle| = |\langle x - y, x_{ij} \rangle|, \text{ for all } (i, j).$$

Since $\{W_i\}_{i=1}^m$ does norm retrieval,

$$\begin{aligned} \|x + y\|^2 &= \|x\|^2 + \|y\|^2 + 2\langle x, y \rangle \\ &= \|x - y\|^2 \\ &= \|x\|^2 + \|y\|^2 - 2\langle x, y \rangle. \end{aligned}$$

Hence, $\langle x, y \rangle = 0$ and $x \perp y$.

(2) \Rightarrow (1): Assume $|\langle x, x_{ij} \rangle| = |\langle y, x_{ij} \rangle|$ for all (i, j) . Let

$$S = \{(i, j) : \langle x, x_{ij} \rangle = \langle y, x_{ij} \rangle\}.$$

Then $x + y \perp x_{ij}$ for all $(i, j) \in S^c$ and $x - y \perp x_{ij}$ for all $(i, j) \in S$. By assumption, $x + y \perp x - y$ so

$$0 = \langle x + y, x - y \rangle = \|x\|^2 - \|y\|^2, \text{ and so } \|x\| = \|y\|.$$

So $\{x_{ij}\}$ does norm retrieval. \square

REFERENCES

- [1] R. Balan, P.G. Casazza, and D. Edidin, *On Signal Reconstruction Without Phase*, Appl. Comput. Harmon. Anal., **20** (3) (2006) 345-356.
- [2] A.S. Bandeira, J. Cahill, D. Mixon and A.A. Nelson, *Saving phase: injectivity and stability for phase retrieval*, Appl. and Comput. Harmonic Anal, **37** (1) (2014) 106-125.
- [3] Sara Botelho-Andrade, Peter G. Casazza, Desai Cheng, John I. Haas, Tin T. Tran, Janet C. Tremain, and Zhiqiang Xu, *Real Phase Retrieval by orthogonal complements and hyperplanes*, Proceedings of SPIE (2017).
- [4] J. Cahill, P.G. Casazza, J. Peterson and L. Woodland, *Phase retrieval by projections*, Houston Journal of Mathematics **42** No. 2 (2016) 537-558.
- [5] P.G. Casazza and D. Cheng, *Associating vectors in \mathbb{C}^n with rank 2 projections in \mathbb{R}^{2n} : with applications*, Arxiv: 173.02657.
- [6] P.G. Casazza and D. Ghoreishi, *It takes $2n-2$ projections to do phase retrieval in \mathbb{R}^n* , Submitted JFAA.
- [7] D. Cheng and P.G. Casazza, *Associating vectors in \mathbb{C}^n with rank-2 projections in \mathbb{R}^{2n} : with applications*. arxiv: 1703.02657.
- [8] J. Drenth, *Principles of protein x-ray crystallography*, Springer, 2010.
- [9] D. Edidin, *Projections and phase retrieval*, Appl. Comput. Harmon. Anal., **42** (2)(2017) 350-359.
- [10] M. A. Hasankhani Fard, *Norm Retrievable Frames in \mathbb{R}^n* (Submitted)
- [11] C. Vinzant, *A small frame and a certificate of its injectivity*, Proceedings of the International conference on Sampling Theory and Applications, 7148879 (2015), 197-200.
- [12] Y. Wang and Z. Zu, *Generalized phase retrieval: measurement number, matrix recovery and beyond*, ArXiv 1605.08034.
- [13] Z. Xu, *The minimal measurement number for low rank matrix recovery*, Applied and Computational Harmonic Analysis (2018).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211-4100

Email address: Casazzap@missouri.edu; tremainjc@missouri.edu