

## P-BASES AND TOPOLOGICAL GROUPS

ZIQIN FENG

ABSTRACT. A topological space  $X$  is defined to have a neighborhood  $P$ -base at any  $x \in X$  from some poset  $P$  if there exists a neighborhood base  $(U_p[x])_{p \in P}$  at  $x$  such that  $U_p[x] \subseteq U_{p'}[x]$  for all  $p \geq p'$  in  $P$ . We prove that a compact space is countable, hence metrizable, if it has countable scattered height and a  $\mathcal{K}(M)$ -base for some separable metric space  $M$ . This gives a positive answer to Problem 8.6.8 in [3].

Let  $A(X)$  be the free Abelian topological group on  $X$ . It is shown that if  $Y$  is a retract of  $X$  such that the free Abelian topological group  $A(Y)$  has a  $P$ -base and  $A(X/Y)$  has a  $Q$ -base, then  $A(X)$  has a  $P \times Q$ -base. Also if  $Y$  is a closed subspace of  $X$  and  $A(X)$  has a  $P$ -base, then  $A(X/Y)$  has a  $P$ -base.

It is shown that any Fréchet-Urysohn topological group with a  $\mathcal{K}(M)$ -base for some separable metric space  $M$  is first-countable, hence metrizable. And if  $P$  is a poset with calibre  $(\omega_1, \omega)$  and  $G$  is a topological group with a  $P$ -base, then any precompact subset in  $G$  is metrizable, hence  $G$  is strictly angelic. Applications in function spaces  $C_p(X)$  and  $C_k(X)$  are discussed. We also give an example of a topological Boolean group of character  $\leq \mathfrak{d}$  such that the precompact subsets are metrizable but  $G$  doesn't have an  $\omega^\omega$ -base if  $\omega_1 < \mathfrak{d}$ . This gives a consistent negative answer to Problem 6.5 in [9].

### 1. INTRODUCTION

Let  $P$  be a partially ordered set (poset). A topological space  $X$  is defined to have a neighborhood  $P$ -base at any  $x \in X$  if there exists a neighborhood base  $(U_p[x])_{p \in P}$  at  $x$  such that  $U_p[x] \subseteq U_{p'}[x]$  for all  $p \geq p'$  in  $P$ . We say that a topological space has a  $P$ -base if it has a neighborhood  $P$ -base at each  $x \in X$ . Thus, a topological group  $G$  has a  $P$ -base if and only if it has a neighborhood  $P$ -base at the identity. Therefore, the classical metrization theorem of Birkhoff and Kakutani can be restated as ‘a Hausdorff topological group  $G$  is metrizable if and only if  $G$  has an  $\omega$ -base, i.e. is first-countable’. All the topological spaces and groups in this paper are assumed to be Tychonoff.

The concept of  $\omega^\omega$ -base with the original name ‘ $\mathfrak{G}$ -base’ was formally introduced in [7] for studying (DF)-spaces, function spaces, and spaces in the class  $\mathfrak{G}$  (see [17] for the definition). Here,  $\omega^\omega$  is the family of all sequences of natural numbers in the pointwise partial order, i.e.  $f \leq g$  if and only if  $f(n) \leq g(n)$  for all  $n \in \omega$ . Topological spaces and groups with an  $\omega^\omega$ -base were defined and studied in [9], for a systematic study see [3]. Works about topological algebra objects with an  $\omega^\omega$ -base can also be found in [4], [18], and [20].

---

*Date:* October 11, 2021.

*2010 Mathematics Subject Classification.* 22A05, 54H11, 46A50.

*Key words and phrases.* (Scattered) compact spaces, (free Abelian) topological groups, Tukey order,  $\omega^\omega$ -base,  $P$ -base, (strictly) angelic, metrizable, quotient groups.

For any  $x$  in the topological space  $X$ , the collection  $\mathcal{T}_x(X)$  of open neighborhood of  $x$  is a poset in the ordering defined by  $U \leq V$  if and only if  $U \supseteq V$ . Throughout all the note, the local base at any point is considered as a poset defined in this way. To understand this poset  $\mathcal{T}_x(X)$ , it is critical to investigate its cofinality. We use Tukey order to compare the cofinal complexity of posets. Tukey order [21] was originally introduced, early in the 20th century, as a tool to understand convergence in general topological spaces; however it was quickly seen to have broad applicability in comparing posets. Given two directed sets  $P$  and  $Q$ , we say  $Q$  is a Tukey quotient of  $P$ , denoted by  $P \geq_T Q$ , if there is a map  $\phi : P \rightarrow Q$  carrying cofinal subsets of  $P$  to cofinal subsets of  $Q$ . It is known that a topological space  $X$  has a  $P$ -base if and only if  $\mathcal{T}_x(X) \leq_T P$  for each  $x \in X$ .

This paper is organized as follows. In Section 3, we prove that a compact space is countable, hence metrizable, if it has countable scattered height and a  $\mathcal{K}(M)$ -base for some separable metric space  $M$ . Hence this is also true for a compact space with countable scattered height and an  $\omega^\omega$ -base, which gives a positive answer to Problem 8.6.8 in [3]. It is worth mentioning that in [5] the authors prove that under the assumption  $\omega_1 < \mathfrak{b}$ , any compact space with an  $\omega^\omega$ -base is metrizable and any scattered compact space with an  $\omega^\omega$ -base is countable. We also prove that  $D^2 \setminus \Delta$  is strongly dominated (see definition in Section 3) by the Bowtie space  $B$ , here  $D$  is the double arrow space. This gives negative answer to Questions 4.4-4.13 in [16].

For a Tychonoff space  $X$ , let  $A(X)$  be the free Abelian topological group over  $X$ , in the sense that there is a continuous mapping  $\sigma$  from  $X$  to  $A(X)$  such that: i)  $\sigma(X)$  topologically generates the topological group  $A(X)$ ; and ii), for every continuous mapping  $f$  from  $X$  to an Abelian group  $H$ , there is a continuous homomorphism  $\tilde{f}$  from  $A(X)$  to  $H$  such that  $\tilde{f} \circ \sigma = f$ . It is proved in [10] and independently in [4] that  $A(X)$  has an  $\omega^\omega$ -base if and only if the universal uniformity  $\mathcal{U}_X$  of  $X$  has an  $\omega^\omega$ -ordered base. In Section 4 we study the  $P$ -bases of free Abelian topological groups through quotient spaces and quotient groups. We prove that if  $Y$  is a retract of  $X$  such that  $A(Y)$  has a  $P$ -base and  $A(X/Y)$  has a  $Q$ -base, then  $A(X)$  has a  $P \times Q$ -base. We also prove that if  $Y$  is a closed subspace of  $X$  and  $A(X)$  has a  $P$ -base, then  $A(X/Y)$  has a  $P$ -base.

In Section 5 we show that any Fréchet-Urysohn topological group with a  $\mathcal{K}(M)$ -base for some separable metric space  $M$  is first-countable, hence metrizable. Also, we prove that if  $P$  is a poset with calibre  $(\omega_1, \omega)$  and  $G$  is a topological group with a  $P$ -base, then any precompact subset in  $G$  is metrizable, hence,  $G$  is strictly angelic. In this section we apply our results to function spaces  $C_p(X)$  and  $C_k(X)$ . We also give an example of a topological Boolean group of character  $\leq \mathfrak{d}$  such that the precompact subsets are metrizable and  $G$  doesn't have an  $\omega^\omega$ -base if  $\omega_1 < \mathfrak{d}$ . This gives a consistent negative answer to Problem 6.5 in [9].

## 2. PRELIMINARIES

Given that  $P$  and  $Q$  are both Dedekind complete (every bounded subset has a least upper bound),  $P \geq_T Q$  if and only if there is a map  $\phi : P \rightarrow Q$  which is order-preserving and such that  $\phi(P)$  is cofinal in  $Q$ . Throughout the note, all the posets are Dedekind complete unless otherwise stated. If  $P$  and  $Q$  are mutually Tukey quotients, we say that  $P$  and  $Q$  are Tukey equivalent, denoted by  $P =_T Q$ .

For any separable metric space  $M$ ,  $\mathcal{K}(M)$  is the collection of compact subsets of  $M$  ordered by set-inclusion. Fremlin observed that if a separable metric space  $M$  is

locally compact, then  $\mathcal{K}(M) =_T \omega$ . Its unique successor under Tukey order is the class of Polish but not locally compact spaces. For  $M$  in this class,  $\mathcal{K}(M) =_T \omega^\omega$  where  $\omega^\omega$  is ordered by  $f \leq g$  if  $f(n) \leq g(n)$  for each  $n \in \omega$ . In [11], Gartside and Mamataleshvili constructed a  $2^c$ -sized antichain in  $\mathcal{K}(\mathcal{M}) = \{\mathcal{K}(M) : M \in \mathcal{M}\}$  where  $\mathcal{M}$  is the class of separable metric spaces.

Let  $P$  be a directed poset, i.e., for any points  $p, p' \in P$ , there exists a point  $q \in P$  such that  $p \leq q$  and  $p' \leq q$ . A subset  $C$  of  $P$  is *cofinal* in  $P$  if for any  $p \in P$ , there exists a  $q \in C$  such that  $p \leq q$ . Then  $\text{cof}(P) = \min\{|C| : C \text{ is cofinal in } P\}$ . We also define  $\text{add}(P) = \min\{|Q| : Q \text{ is unbounded in } P\}$ . For any  $f, g \in \omega^\omega$ , we say that  $f \leq^* g$  if the set  $\{n \in \omega : f(n) > g(n)\}$  is finite. Then  $\mathfrak{b} = \text{add}(\omega^\omega, \leq^*)$  and  $\mathfrak{d} = \text{cof}(\omega^\omega, \leq^*)$ . See [2] for more information about small cardinals.

Let  $\kappa \geq \mu \geq \lambda$  be cardinals. We say that a poset  $P$  has *calibre*  $(\kappa, \mu, \lambda)$  if for every  $\kappa$ -sized subset  $S$  of  $P$  there is a  $\mu$ -sized subset  $S_0$  such that every  $\lambda$ -sized subset of  $S_0$  has an upper bound in  $P$ . We write *calibre*  $(\kappa, \mu, \mu)$  as *calibre*  $(\kappa, \mu)$  and *calibre*  $(\kappa, \kappa, \kappa)$  as *calibre*  $\kappa$ . It is known that  $\mathcal{K}(M)$  has *calibre*  $(\omega_1, \omega)$  for any separable metric space  $M$ , hence so does  $\omega^\omega$ . And if  $P$  has *calibre*  $(\omega_1, \omega)$  and  $Q$  is a Tukey quotient of  $P$ , then  $Q$  also has *calibre*  $(\omega_1, \omega)$ . Under the assumption  $\omega_1 = \mathfrak{b}$ ,  $\omega_1$  is a Tukey quotient of  $\omega^\omega$  (see [19] for the discussion of the spectrum of  $\omega^\omega$ ), hence  $\omega_1 + 1$  has an  $\omega^\omega$ -base. However, when  $\omega_1 < \mathfrak{b}$ ,  $\omega_1$  is not a Tukey quotient of  $\omega^\omega$ , hence the space  $\omega_1 + 1$  doesn't have an  $\omega^\omega$ -base, but it has a  $\mathcal{K}(\mathbb{Q})$ -base (see [5]), where  $\mathbb{Q}$  is the space of rational numbers.

We start with some basic properties of the property *calibre*  $(\omega_1, \omega)$  and  $P$ -bases. For any two posets  $(P, \leq_P)$  and  $(Q, \leq_Q)$ , the ordering on  $P \times Q$  is naturally defined as follows:  $(p, q) \leq (p', q')$  if and only if  $p \leq_P p'$  and  $q \leq_Q q'$ . For  $i = 1, 2$ , let  $\pi_i$  be the projection mapping from  $P \times Q$  to  $P$  or  $Q$  respectively.

**Proposition 2.1.** *If  $P$  is a poset with calibre  $(\omega_1, \omega)$  and  $Q$  is a countable poset, then  $P \times Q$  also has calibre  $(\omega_1, \omega)$ .*

*Proof.* Let  $S$  be an  $\omega_1$ -sized subset of  $P \times Q$ . Then there is a  $q \in Q$  and an  $\omega_1$ -sized subset  $S'$  of  $S$  such that  $\pi_2(x) = q$  for all  $x \in S'$ . Since  $P$  has *calibre*  $(\omega_1, \omega)$ ,  $\{\pi_1(x) : x \in S'\}$  is an  $\omega_1$ -sized subset of  $P$ , hence it has a countable subset bounded in  $P$ . Therefore, there is a countable subset of  $S$  which is bounded in  $P \times Q$ .  $\square$

**Proposition 2.2.** *Suppose that  $X$  has a  $P$ -base for some poset  $P$ . If there is an open continuous mapping  $f$  from  $X$  onto  $Y$ , then  $Y$  has a  $P$ -base.*

*Proof.* Fix  $y \in Y$ . Choose  $x \in f^{-1}(y)$ . Then there is a local  $P$ -base  $\{U_p[x] : p \in P\}$  at  $x$ . We claim that  $\{f(U_p[x]) : p \in P\}$  is a local  $P$ -base at  $y$ . Clearly  $f(U_p[x])$  is an open neighborhood of  $y$  since  $f$  is open. For any open neighborhood  $V$  of  $y$ ,  $f^{-1}(V)$  is an open neighborhood of  $x$ . Therefore, there is a  $p \in P$  such that  $U_p[x] \subseteq f^{-1}(V)$ , i.e.,  $f(U_p[x]) \subseteq V$ .  $\square$

### 3. (SCATTERED) COMPACT SPACES

We recall that a topological space  $X$  is scattered if each non-empty subspace of  $X$  has an isolated point. The complexity of a scattered space can be determined by the scattered height.

For any subspace  $A$  of a space  $X$ , let  $A'$  be the set of all non-isolated points of  $A$ . It is straightforward to see that  $A'$  is a closed subset of  $A$ . Let  $X^{(0)} = X$  and define  $X^{(\alpha)} = \bigcap_{\beta < \alpha} (X^{(\beta)})'$  for each  $\alpha > 0$ . Then a space  $X$  is scattered if  $X^{(\alpha)} = \emptyset$

for some ordinal  $\alpha$ . If  $X$  is scattered, there exists a unique ordinal  $h(x)$  such that  $x \in X^{(h(x))} \setminus X^{(h(x)+1)}$  for each  $x \in X$ . The ordinal  $h(X) = \sup\{h(x) : x \in X\}$  is called the scattered height of  $X$  and is denoted by  $h(X)$ . It is known that any compact scattered space is zero-dimensional. Also, it is straightforward to show that for any compact scattered space  $X$ ,  $X^{(h(X))}$  is a non-empty finite subset.

Let  $X$  be a compact space. If  $X$  has an  $\omega^\omega$ -base and finite scattered height, then  $X$  is proved in [3] to be countable, hence metrizable. It is proved in [5] that the same result also holds if  $X$  has finite scattered height and a  $P$ -base for some poset  $P$  with calibre  $(\omega_1, \omega)$ . The compact space  $\omega_1 + 1$  has an  $\omega^\omega$ -base under the assumption  $\omega_1 < \mathfrak{b}$ . The authors in [5] proved that if  $\omega_1 < \mathfrak{b}$  any compact scattered space is countable, hence metrizable. Hence it is natural to ask, as in [3], whether a compact space  $X$  is metrizable in ZFC if it has an  $\omega^\omega$ -base and *countable* scattered height. Next, we give a positive answer to this question by proving a more general result.

**Proposition 3.1.** *Any compact space is countable, hence metrizable, if it has a countable scattered height and a  $\mathcal{K}(M)$ -base, where  $M$  is a separable metric space.*

*Proof.* Let  $X$  be a compact space with countable scattered height  $\alpha$  and a  $\mathcal{K}(M)$ -base for some separable metric space  $X$ . We'll prove by induction on  $\alpha$ .

Assume any compact space with scattered height  $\beta$  is countable for any  $\beta < \alpha$ . Without loss of generality, we assume that  $X^{(\alpha)}$  is a singleton, denoted by  $x^*$ . For any open neighborhood  $U$  of  $x^*$ ,  $X \setminus U$  is a scattered compact space with scattered height  $< \alpha$ , hence is countable. So it is sufficient to prove that  $X$  is first-countable at  $x^*$ .

In [5], the authors prove that any compact space with countable tightness and a  $\mathcal{K}(M)$ -base is first-countable where  $M$  is a separable metric space. Hence it suffices to prove that  $X$  has countable tightness. For each  $x \in X \setminus \{x^*\}$  there is a clopen neighborhood  $U_x$  of  $x$  such that  $U_x \cap X^{(h(x))} = \{x\}$  since any scattered compact space is zero-dimensional. Clearly  $U_x$  is a compact space with scattered height  $< \alpha$ . By the inductive assumption  $U_x$  is countable. So  $X$  has countable tightness at any  $x$  in  $X \setminus \{x^*\}$ .

Next, we prove that  $X$  has countable tightness at  $x^*$ . Let  $B$  a subset of  $X \setminus \{x^*\}$  with  $x^* \in \overline{B}$ . Let  $\beta_* = \sup\{\gamma : X^{(\gamma)} \cap (\overline{B} \setminus \{x^*\}) \neq \emptyset \text{ and } \gamma < \alpha\}$ . We will show there is a countable subset  $C \subseteq B$  such that  $x^* \in \overline{C}$  in two cases: 1)  $\alpha$  is a limit ordinal; and 2)  $\alpha$  is a successor.

Firstly, we assume that  $\alpha$  is a limit ordinal. If  $\beta_* < \alpha$ , then  $\overline{B}$  is a compact space with scattered height  $\leq \beta_* + 1$ , hence it is countable. Then  $C = B$  works. Suppose  $\beta_* = \alpha$ . Then there exists a cofinal sequence  $\gamma_n$  in  $\alpha$  such that  $X^{(\gamma_n)} \cap (\overline{B} \setminus \{x^*\}) \neq \emptyset$ . For each  $n$ , pick  $x_n \in X^{(\gamma_n)} \cap (\overline{B} \setminus \{x^*\})$ . Since  $X$  is compact,  $x^*$  is in the closure of  $\{x_n : n \in \omega\}$ . For each  $n \in \omega$ , let  $C_n$  be a countable subset of  $B$  such that  $x_n \in \overline{C_n}$ . Therefore  $C = \bigcup\{C_n : n \in \omega\}$  works.

From now on, we assume that  $\alpha$  is a successor and let  $\alpha_-$  be the ordinal such that  $\alpha = \alpha_- + 1$ . Note that  $X^{(\alpha_-)}$  is a compact space with scattered height 1, hence it is countable. If  $\beta_* < \alpha_-$ , then the closure of  $B$  is a compact space with scattered height  $< \alpha$ . Hence  $\overline{B}$  is countable, so is  $B$ . Then let  $C = B$  which is clearly countable.

Suppose  $\beta_* = \alpha_-$ . If  $X^{(\alpha_-)} \cap \overline{B}$  is finite, choose a clopen neighborhood  $U$  of  $x^*$  such that  $U \cap (X^{(\alpha_-)} \cap \overline{B}) = \{x^*\}$ . Then  $U \cap \overline{B}$  is a compact subspace with

scattered height  $< \alpha$ , hence countable. Therefore,  $C = U \cap B$  works. If  $X^{(\alpha-)} \cap \overline{B}$  is infinite, list it as  $\{x_n : n \in \omega\}$ ; clearly  $x^* \in \overline{\{x_n : n \in \omega\}}$  since  $X$  is compact. By the inductive assumption, we can pick a countable subset  $C_n$  of  $B$  such that  $x_n$  is in the closure of  $C_n$ . Then  $C = \bigcup C_n$  is a countable subset of  $B$  with  $x^*$  in the closure of  $C$ . This finishes the proof.  $\square$

The corollary below answers Problem 8.6.8 in [3] affirmatively.

**Corollary 3.2.** *A compact space is countable (hence metrizable) if and only if it has countable scattered height and an  $\omega^\omega$ -base.*

We don't know the answer to the following question. The answer is proved to be positive in [5] if the space has a finite scattered height.

**Question 3.3.** *Let  $P$  be a poset with calibre  $(\omega_1, \omega)$  and  $X$  a compact space with a  $P$ -base. Is  $X$  countable if  $X$  has a countable scattered height?*

We say a space  $Z$  is dominated by  $Y$  if there exists a family  $\mathcal{D} = \{D_K : D_K \text{ is a compact subset of } Z \text{ and } K \text{ is a compact subset of } Y\}$  such that  $Z = \bigcup \mathcal{D}$  and  $D_K \subset D_L$  if  $K \subset L$ . Furthermore,  $Z$  is said to be strongly dominated by  $Y$  if the family  $\mathcal{D}$  is also confial in  $\mathcal{K}(Z)$ . It is shown that any compact space  $X$  with  $X^2 \setminus \Delta$  being dominated by  $\mathbb{Q}$  is metrizable in [6] and the same result holds for a compact space  $X$  with  $X^2 \setminus \Delta$  being dominated by  $M$  for some separable metric space  $M$ , see [15], here  $\Delta = \{(x, x) : x \in X\}$ . Questions 4.4-4.13 in [16] ask whether a compact space  $X$  is metrizable if  $X^2 \setminus \Delta$  is dominated by other topological space. The following example answers these questions negatively.

**Example 3.4. (Gartside)** *Let  $B$  denote the Bowtie space.*

*Then  $\mathcal{K}(B) =_T [\mathfrak{c}]^{<\omega}$ . Hence for every compact space,  $X$ , of weight  $\leq \mathfrak{c}$ , we have  $\mathcal{K}(B) \geq_T \mathcal{K}(X^2 \setminus \Delta)$ .*

*In particular, there is a  $\sigma$ -compact cosmic space,  $B$ , which strongly dominates  $D^2 \setminus \Delta$ , where  $D$  is the Double Arrow space – a compact, first countable but non-metrizable space.*

*Proof.* Let  $B$  be the standard Bowtie space obtained by refining the Euclidean topology on  $\mathbb{R}^2$  by declaring sets of the form  $B(x, s) = \{(x, 0)\} \cup \{(x', y) : |y| < s|x - x'|\}$  to be open for all  $x, s \in \mathbb{R}$  with  $s > 0$ .

Let  $K_0 = \{(a, b) : a \in [-1, 1], b = a^2\}$ . Observe that  $K_0$  is a compact subset of  $B$ . For each  $x$  in  $\mathbb{R}$ , let  $K_x$  be  $K_0$  translated to  $(x, 0)$ , i.e.,  $K_x = \{(x, y) : (x, y) = (x, 0) + (a, b) \text{ for some } (a, b) \in K_0\}$ .

Then  $\mathcal{C} = \{K_x : x \in \mathbb{R}\}$  has size  $\mathfrak{c}$  but no infinite subfamily has an upper bound in  $\mathcal{K}(B)$ , i.e., has union whose closure is compact. To see this, suppose for a contradiction that infinite family of  $K_x$ 's is contained in a compact set, say  $K_\infty$ . Note that the set of corresponding  $x$ 's in  $\mathbb{R}$  is infinite and has compact closure. Passing to a subset we may suppose that we have an infinite family of  $K_{x_n}$ 's where the  $x_n$ 's are (without loss of generality) an increasing sequence in  $\mathbb{R}$  converging to some  $x_\infty$ . But now, for each  $n$ ,  $K_{x_n}$  meets the line  $\{x_\infty\} \times \mathbb{R}$  in exactly one point, say  $(x_\infty, y_n)$ . And now we see that  $\{(x_\infty, y_n) : n \in \mathbb{N}\}$  is an infinite closed discrete subset of compact  $K_\infty$  – contradiction.

Hence  $\mathcal{K}(B)$  does not have calibre  $(\mathfrak{c}, \omega)$ . Then  $\mathcal{K}(B) \geq_T [\mathfrak{c}]^{<\omega}$  by [19, Lemma 11]. As  $\mathcal{K}(B)$  has size  $\mathfrak{c}$ , clearly  $\mathcal{K}(B) =_T [\mathfrak{c}]^{<\omega}$ .

Let  $X$  be a compact space with  $w(X) \leq \mathfrak{c}$ . Then  $\mathcal{K}(X^2 \setminus \Delta)$  has cofinality no more than  $\mathfrak{c}$  (because it is order isomorphic to the neighborhoods of the diagonal,

and a cofinal family of these comes from finite covers by basic open sets). Then by [19, Lemma 6],  $[c]^{<\omega} \geq_T \mathcal{K}(X^2 \setminus \Delta)$ , hence  $\mathcal{K}(B) \geq_T \mathcal{K}(X^2 \setminus \Delta)$ . Therefore  $X^2 \setminus \Delta$  is strongly dominated by the Bowtie space  $B$ .  $\square$

#### 4. FREE ABELIAN TOPOLOGICAL GROUPS

A property  $\mathcal{P}$  is said to be a *three-space property* if for every topological group  $G$  and a closed normal subgroup  $H$  of  $G$  both  $H$  and  $G/H$  having  $\mathcal{P}$  implies that  $G$  has  $\mathcal{P}$ . It is a classical result that having an  $\omega$ -base is a three-space property (see [1]). The authors ask in [9] whether having an  $\omega^\omega$ -base is a three-space property. We will start with a sufficient condition on  $(G, H)$  which guarantees that  $G$  has an  $\omega^\omega$ -base if both  $H$  and  $G/H$  have an  $\omega^\omega$ -base.

**Definition 4.1.** *Let  $G$  be a topological group and  $H$  a closed normal subgroup of  $G$ . We say that  $(G, H)$  is a good pair if there is a mapping  $p$  from a local base  $\mathcal{B}$  at the identity in  $H$  to the topology on  $G$  such that:*

- i)  $p(B) \cap H = B$  for any  $B \in \mathcal{B}$ ;
- ii) for any  $B, B' \in \mathcal{B}$ ,  $p(B)^2 \subseteq p(B')$  if  $B^2 \subseteq B'$ .
- iii) for any  $B, B' \in \mathcal{B}$ ,  $p(B)^{-1} \subseteq p(B')$  if  $B^{-1} \subseteq B'$ .

Our next result shows that if  $H$  is a metrizable closed normal subgroup of  $G$ , then  $(G, H)$  is a good pair. The proof is straightforward.

**Proposition 4.2.** *Let  $G$  be a topological group with the identity  $e$ . If  $H$  is a closed normal subgroup of  $G$  such that  $\mathcal{T}_H(e) \leq_T \omega$ , then  $(G, H)$  is a good pair.*

**Theorem 4.3.** *Let  $G$  be a topological group and  $H$  a closed normal subgroup of  $G$  such that  $(G, H)$  is a good pair. Suppose that  $P$ ,  $Q$ , and  $R$  are Dedekind-complete posets such that  $P \times Q \leq_T R$ . If  $H$  has a  $P$ -base and  $G/H$  has a  $Q$ -base, then  $G$  has an  $R$ -base.*

*Proof.* Let  $\phi$  be an order-preserving mapping from  $R$  to  $P \times Q$  such that  $\phi(R)$  is cofinal in  $P \times Q$ . Let  $e$  be the identity element in  $G$ .

Since  $(G, H)$  is a good pair and  $H$  has a  $P$ -base, there is a collection  $\{W_p : p \in P\}$  of open neighborhoods of  $e$  in  $G$  such that: i)  $\{W_p \cap H : p \in P\}$  is a  $P$ -base of  $H$ ; ii) if  $(W_p \cap H)^2 \subset W_{p'}$  then  $W_p^2 \subset W_{p'}$ ; and iii) if  $(W_p \cap H)^{-1} \subset W_{p'} \cap H$  then  $W_p^{-1} \subset W_{p'}$ . Let  $\{V_q : q \in Q\}$  be a symmetric  $Q$ -base of  $G/H$ . Let  $q$  be the quotient mapping from  $G$  to  $G/H$ . For each  $r \in R$ , define  $U_r = W_{\pi_1(\phi(r))} \cap q^{-1}(V_{\pi_2(\phi(r))})$ . By the order-preserving property of  $\phi$ , it is straightforward to verify that  $U_r \supset U_{r'}$  given  $r \leq r'$ .

We show that  $\mathcal{U} = \{U_r : r \in R\}$  is a base for  $G$ . Take an open neighborhood  $U$  of the identity element  $e$ . Then, we pick an open symmetric neighborhood  $V$  of the unit such that  $VV \subset U$ . Fix  $p_1 \in P$  such that  $W_{p_1} \cap H \subset V$ . By conditions ii) and iii) above, there exists a  $p_2 \in P$  such that  $p_2 \geq p_1$  and  $W_{p_2}^{-1}W_{p_2} \subset W_{p_1}$ . Pick  $q_1 \in Q$  such that  $q(V_{q_1}) \subset q(V \cap W_{p_2})$ . Since  $\phi(R)$  is cofinal in  $P \times Q$ , there exists an  $r \in R$  such that  $\pi_1(\phi(r)) \geq p_2$  and  $\pi_2(\phi(r)) \geq q_1$ .

We claim that  $U_r \subset U$ . Pick  $g \in U_r$ . By the definition of  $U_r$  and the choice of  $r$ ,  $g \in W_{p_2}$  and  $g \in q^{-1}(V_{\pi_2(\phi(r))}) \subset V_{q_1} \cdot H \subseteq V \cap W_{p_2} \cdot H$ . Hence  $g = ab$  for some  $a \in V \cap W_{p_2}$  and  $b \in H$ . Therefore  $b = a^{-1}g \in W_{p_2}^{-1}W_{p_2} \subset W_{p_1}$ . Finally, we obtain that  $g \in (V \cap W_{p_2}) \cdot (W_{p_1} \cap H) \subset V \cdot V \subset U$ .

Therefore,  $\mathcal{U}$  is an  $R$ -base for the topological group  $G$ .  $\square$

Therefore, if  $H$  is a metrizable closed normal subgroup of  $G$ ,  $(G, H)$  is a good pair by Proposition 4.2. Let  $M$  be any noncompact separable metric space. Note that  $\mathcal{K}(M) \geq_T \mathcal{K}(M) \times \omega$ . Then if  $G/H$  has a  $\mathcal{K}(M)$ -base, then  $G$  has a  $\mathcal{K}(M)$ -base. The case  $M = \omega^\omega$  is proved in [9, Proposition 2.9].

Next we discuss good pairs of the form  $(A(X), A(Y, X))$ , where  $A(Y, X)$  is the subgroup of  $A(X)$  generated by  $Y$ . It is shown in [1, Theorem 7.4.5] that  $A(Y, X)$  is a closed subgroup of  $A(X)$  if  $Y$  is a closed subspace of  $X$ . Also, if  $Y$  is a retract of  $X$ , then  $A(Y, X)$  is topologically isomorphic to  $A(Y)$ , in which case we consider them to be same.

**Proposition 4.4.** *Let  $G$  be a topological group and  $H$  a closed normal subgroup of  $G$ . If there is a continuous homomorphism from  $G$  to  $H$  whose restriction on  $H$  is identity mapping,  $(G, H)$  is a good pair.*

*Proof.* Let  $\phi$  be a continuous homomorphism from  $G$  to  $H$  such that  $\phi(h) = h$  given  $h \in H$ . Fix a local base  $\mathcal{B}$  of the identity in  $H$ . For each  $B \in \mathcal{B}$ , define  $p(B) = \phi^{-1}(B)$ . It is clear that  $p(B) \cap H = B$  for each  $B \in \mathcal{B}$  since  $\phi(h) = h$  for any  $h \in H$ .

Pick  $B, B' \in \mathcal{B}$  such that  $B^2 \subseteq B'$ . Choose  $g$  and  $g'$  in  $p(B)$ . We show that  $g \cdot g' \in p(B)$ . By the definition of  $p(B)$ , there exists  $h, h' \in B$  such that  $\phi(g) = h$  and  $\phi(g') = h'$ . Hence  $\phi(g \cdot g') = \phi(g) \cdot \phi(g') = h \cdot h' \in B'$ . Therefore,  $g \cdot g' \in \phi^{-1}(B') = p(B')$ .

Pick  $B, B' \in \mathcal{B}$  such that  $B^{-1} \subset B'$ . Choose  $g \in p(B)$ . Then there exists  $h \in B$  such that  $\phi(g) = h$ , then  $\phi(g^{-1}) = h^{-1}$ . Since  $B^{-1} \subset B'$ ,  $\phi(g^{-1}) = h^{-1} \in B'$ , i.e.,  $g^{-1} \in p(B')$ . This finishes the proof.  $\square$

**Proposition 4.5.** *Let  $X$  be a Tychonoff space and  $Y$  a retract of  $X$ . Then  $(A(X), A(Y, X))$  is a good pair.*

*Proof.* Since  $Y$  is a retract of  $X$ ,  $Y$  is a closed subspace of  $X$ . By Theorem 7.4.5 in [1],  $A(Y, X)$  is a closed subgroup of  $A(X)$ .

Let  $f$  be a continuous mapping from  $X$  to  $Y$  such that  $f(y) = y$  for each  $y \in Y$ . By Corollary 7.1.9 in [1],  $f$  admits an extension to a continuous homomorphism  $A(f)$  from  $A(X)$  to  $A(Y)$ . Clearly for any  $h \in A(Y)$ ,  $A(f)(h) = h$ . The identity mapping from  $A(Y)$  to  $A(Y, X)$  is a topological isomorphism. Hence, we obtain a continuous homomorphism from  $A(X)$  to  $A(Y, X)$  whose restriction on  $A(Y, X)$  is the identity mapping. By Proposition 4.4,  $(A(X), A(Y, X))$  is a good pair.  $\square$

Let  $Y$  be a closed subspace of  $X$ . Define an equivalence relation  $\sim$  on  $X$  by  $x \sim y$  if and only if  $x, y \in Y$ . We denote the quotient space  $X/\sim$  by  $X/Y$ ; there is a canonical quotient mapping from  $X$  to  $X/Y$ . We denote the equivalence class  $[y]$  for any  $y \in Y$  by  $[Y]$  which clearly is a retract of  $X/Y$ . So in the following discussion, we use  $A([Y])$  to represent the subgroup  $A([Y], X/Y)$ . Next, we investigate the relations of  $A(X)$ ,  $A(Y)$ , and  $A(X/Y)$  in terms of  $P$ -bases.

**Proposition 4.6.** *Let  $Y$  be a closed subspace of a Tychonoff space  $X$ . Then  $A(X/Y)/A([Y])$  is topologically isomorphic to  $A(X)/A(Y, X)$ .*

*Proof.* Let  $f$  be the quotient map from  $X$  to  $X/Y$ . By Corollary 7.1.9 in [1],  $f$  admits an extension to an open continuous homomorphism  $A(f)$  from  $A(X)$  to  $A(X/Y)$ . The canonical mapping  $\pi$  from  $A(X/Y)$  to  $A(X/Y)/A([Y])$  is an open continuous mapping. Hence we obtain an open continuous mapping  $\phi = \pi \circ A(f)$  from  $A(X)$  to the quotient group  $A(X/Y)/A([Y])$ .

Let  $e$  be the identity in  $A(X/Y)/A([Y])$ . Clearly,  $\phi^{-1}(e) = A(Y, X)$ . Then by Theorem 1.5.13 in [1],  $A(X)/A(Y, X)$  is topologically isomorphic to  $A(X/Y)/A([Y])$ .  $\square$

**Theorem 4.7.** *Let  $Y$  be a closed subspace of a Tychonoff space  $X$ . If the free Abelian group  $A(X)$  has a  $P$ -base, then  $A(X/Y)$  also has a  $P$ -base.*

*Proof.* Since  $Y$  is closed, the subgroup  $A(Y, X)$  is closed and normal. The canonical mapping from  $A(X)$  to  $A(X)/A(Y, X)$  is open, continuous, and onto. Then by Proposition 2.2,  $A(X)/A(Y, X)$  has a  $P$ -base. By Proposition 4.6,  $A(X/Y)/A([Y])$  has a  $P$ -base. Note that  $A([Y])$  is a discrete subspace of  $A(X/Y)$ . So  $(A(X/Y), A([Y]))$  is a good pair by Proposition 4.2. Applying Theorem 4.3,  $A(X/Y)$  has a  $1 \times P$ -base, i.e., a  $P$ -base.  $\square$

The following example is a direct application of Theorem 4.7.

**Example 4.8.** *Let  $S_n = \{x_{m,n} : m \in \omega\} \cup \{x_n\}$  be a nontrivial convergent sequence with  $x_n$  being the limit point for each  $n$  and let  $T$  be the topological sum of the family  $\{S_n : n \in \omega\}$ . Note that  $T$  is a metrizable space whose subspace of non-isolated points is  $\sigma$ -compact. Hence by Theorem 1.2 in [4], the free Abelian group  $A(T)$  has an  $\omega^\omega$ -base.*

*Let  $S_\omega$  be the quotient space of  $T$  obtained by identifying all the limit points to a singleton. Then by Theorem 4.7,  $A(S_\omega)$  has an  $\omega^\omega$ -base.*

We don't know whether the same result holds for free topological groups.

**Question 4.9.** *Let  $Y$  be a closed subspace of a Tychonoff space  $X$ . Suppose that the free group  $F(X)$  has a  $P$ -base. Does  $F(X/Y)$  have a  $P$ -base?*

Next we show that if  $Y$  is a retract of  $X$ , then the bases of  $A(Y)$  and  $A(X/Y)$  at the identity are both Turkey quotients of some base of  $A(X)$  at identity.

**Theorem 4.10.** *Suppose that  $P$ ,  $Q$ , and  $R$  are Dedekind-complete posets such that  $P \times Q \leq_T R$ . Let  $X$  be a Tychonoff space and  $Y$  a retract of  $X$ .*

*If  $A(Y)$  has a  $P$ -base and  $A(X/Y)$  have a  $Q$ -base, then  $A(X)$  has an  $R$ -base.*

*Proof.* Because  $Y$  is a retract of  $X$ ,  $(A(X), A(Y, X))$  is a good pair by Proposition 4.4. So it is sufficient to show that  $A(X)/A(Y, X)$  has a  $Q$ -base. Then  $A(X)$  has an  $R$ -base by Theorem 4.3.

Next we show that  $A(X)/A(Y, X)$  has a  $Q$ -base. Since  $A(X/Y)$  have a  $Q$ -base and the canonical mapping  $\pi$  from  $A(X/Y)$  to  $A(X/Y)/A([Y])$  is an open continuous mapping,  $A(X/Y)/A([Y])$  has a  $Q$ -base by Proposition 2.2. By Proposition 4.6,  $A(X)/A(Y, X)$  is topologically isomorphic to  $A(X/Y)/A([Y])$ . Hence  $A(X)/A(Y, X)$  has a  $Q$ -base. This finishes the proof.  $\square$

For every directed poset  $P$ , it is straightforward to verify that  $P \times P =_T P$ . Hence we obtain the following corollary.

**Corollary 4.11.** *Let  $P$  be any Dedekind-complete directed poset. Suppose that  $X$  is a Tychonoff space and  $Y$  is a retract of  $X$ . If both  $A(Y)$  and  $A(X/Y)$  have a  $P$ -base, then  $A(X)$  has a  $P$ -base.*

The result above holds clearly for all the posets in the form  $\mathcal{K}(M)$  for some topological space  $M$ . Therefore, the free Abelian topological group  $A(X)$  has an  $\omega^\omega$ -base if  $X$  is a Tychonoff space and  $Y$  is a retract of  $X$  such that both  $A(Y)$  and  $A(X/Y)$  have an  $\omega^\omega$ -base.



5. TOPOLOGICAL GROUPS

**Definition 5.1.** A space  $X$  is said to satisfy Property (AS) if the following holds:

(AS): for any family  $\{x_{m,n} : m, n \in \omega\} \subseteq X$  with  $\lim_m x_{m,n} = x$  for all  $n \in \omega$ , there exist two strictly increasing sequences of natural numbers  $\{m_k : k \in \omega\}$  and  $\{n_k : k \in \omega\}$  such that  $\lim_k x_{m_k, n_k} = x$ .

It is proved that in [17] any Fréchet-Urysohn topological group satisfies Property (AS).

**Theorem 5.2.** Let  $P = \mathcal{K}(M)$  for a separable metric space  $M$ . If  $G$  is a Fréchet-Urysohn topological group with a  $P$ -base, then  $G$  is first-countable, hence metrizable.

*Proof.* Let  $e$  be the identity element in  $G$  and  $\mathcal{U} = \{U_p : p \in P\}$  be an open  $P$ -base at the identity  $e$ .

For any separable metric space  $M$ ,  $P = \mathcal{K}(M)$  endowed with the Vietoris topology is also separable metrizable, hence second countable. Also if  $p_n$  converges to  $p$  in  $P$ , then  $p^* = p \cup (\bigcup\{p_n\})$  is also compact, hence it is an element in  $P$  with  $p_n \subset p^*$  and  $p \subset p^*$ .

Fix a countable base  $\{B_n : n \in \omega\}$  of  $P$ . For each  $p \in P$  and  $n \in \omega$  with  $p \in B_n$ , define  $D_n(p) = \bigcap\{U_q : q \in B_n\}$ . For each  $p$ , we list an increasing sequence of natural numbers  $\{n_i^p : i \in \omega\}$  such that  $p \in B_{n_i^p}$  for each  $i$  and  $\{B_{n_i^p} : i \in \omega\}$  is a decreasing base of  $P$ .

First, we claim that for each  $p \in P$  there exists  $i$  such that  $D_{n_i^p}(p)$  is a neighborhood of the identity  $e$ . Suppose for contradiction that there exists  $p \in P$  such that  $D_{n_i^p}(p)$  is not a neighborhood of  $e$  for each  $i \in \omega$ . Hence  $e$  is in the closure of  $G \setminus D_{n_i^p}(p)$  for each  $i \in \omega$ . Since  $G$  is Fréchet-Urysohn, there exists a sequence  $\{x_{m,i}\}$  in  $G \setminus D_{n_i^p}(p)$  for each  $i$  such that  $\lim_m x_{m,i} = e$ . Then we apply Property (AS) to get strictly increasing sequences of natural numbers  $\{m_k : k \in \omega\}$  and  $\{i_k : k \in \omega\}$  such that  $\lim_k x_{m_k, i_k} = e$ . For each  $k \in \omega$ , pick  $q_k \in B_{n_{i_k}^p}$  such that  $x_{m_k, i_k} \notin U_{q_k}$ . Clearly  $q_k$  converges to  $p$  in  $P$ . Hence  $p^* = p \cup \bigcup\{q_k : k \in \omega\}$  is also compact in  $M$ , hence it is in  $\mathcal{K}(M)$ . Since  $q_k \leq p^*$  for each  $k$ ,  $U_{q_k} \supset U_{p^*}$ . Therefore  $x_{m_k, i_k} \notin U_{p^*}$  for each  $k \in \omega$ . This contradicts with  $\lim_k x_{m_k, i_k} = e$  because  $U_{p^*}$  is an open neighborhood of  $e$ .

For each  $p \in U$ , fix the minimal natural number  $i_p$  such that  $D_{n_{i_p}^p}(p)$  is a neighborhood of  $e$ . The family  $\{D_{n_{i_p}^p}(p) : p \in P\}$  is clearly countable. By the definition of  $D_n(p)$ , the family  $\{\text{int}(D_{n_{i_p}^p}(p)) : p \in P\}$  is a countable base at the identity  $e$ . Hence,  $G$  is first-countable.  $\square$

**Question 5.3.** Suppose  $G$  is a Fréchet-Urysohn topological group such that a local base at the identity has Calibre  $(\omega_1, \omega)$ . Is  $G$  first-countable?

A topological space is angelic if every relatively countably compact set  $A$  in the space has a compact closure, i.e., relatively compact, and for each  $x$  in the closure of  $A$  there exists a sequence in  $A$  converging to  $x$ . Here, a subset  $A$  of a space  $X$  is relatively countably compact if every sequence in  $A$  has a cluster-point in  $X$ . A space is said to be strictly angelic if it is angelic and each separable compact subset is first-countable. It is known that any topological group is a uniform space and in uniform spaces, it is shown in [8] that any relatively countably compact subset is precompact. A subset  $B$  of a topological group  $G$  is said to be precompact in  $G$  if

for any neighbourhood  $U$  of the identity in  $G$  there is a finite subset set  $F \subset G$  such that  $B \subset FU$  and  $B \subset UF$ . It is known [1] that any precompact set is relatively compact in Raïkov complete topological groups.

Let  $K$  be a compact space. Gartide and Morgan show in [13] that  $K$  is compact if the off-diagonal subspace  $K^2 \setminus \Delta$  has a compact cover  $\mathcal{D}$  with Calibre  $(\omega_1, \omega)$  which swallows all the compact sets of  $K^2 \setminus \Delta$ , i.e.,  $\mathcal{D}$  is cofinal in  $\mathcal{K}(K^2 \setminus \Delta)$ .

The proof of the following result is very similar to Theorem 3.9 in [9].

**Theorem 5.4.** *Let  $P$  be any directed set with Calibre  $(\omega_1, \omega)$ . If  $G$  is a topological group with a  $P$ -base, then every precompact subset  $K$  in  $G$  is metrizable. Consequently,  $G$  is strictly angelic.*

*Proof.* Similar to Proposition 2.7 in [9], the Raïkov completion of any group with a  $P$ -base also has a  $P$ -base. So we can assume  $G$  is Raïkov complete. Hence it is sufficient to show that every compact subset in  $G$  is metrizable. Let  $K$  be a compact subset of  $G$  and  $\{U_p : p \in P\}$  be a  $P$ -base at the identity  $e$  in  $G$ .

For each  $p \in P$ , define  $L_p = \{(x, y) \in K \times K : xy^{-1} \notin U_p\}$ . Note that each  $L_p$  is closed, hence compact in  $K \times K$ . We show that  $\mathcal{L} = \{L_p : p \in P\}$  is a  $P$ -ordered compact cover of the off-diagonal space  $K \times K \setminus \Delta$  which swallows all its compact sets. It is straightforward to verify that  $L_p \subset L_{p'}$  given  $p \leq p'$  in  $P$ . Let  $C$  be a compact subset of  $K \times K \setminus \Delta$ . Then  $\hat{C} = \{xy^{-1} : (x, y) \in C\}$  is a compact subset of  $G$  which doesn't contain the identity  $e$ . Since  $\{U_p : p \in P\}$  is a local base at the identity  $e$ , there exists a  $p_0 \in P$  such that  $U_{p_0} \subset G \setminus \hat{C}$ , furthermore,  $C \subset L_{p_0}$ . Therefore, the collection  $\mathcal{L}$  swallows all the compact subsets of  $K \times K \setminus \Delta$ . Then by [13],  $K$  is metrizable. This finishes the proof.  $\square$

So next, we discuss some applications of Theorem 5.4 in function spaces. For any Tychonoff space  $X$ , let  $C(X)$  be the space of continuous real-valued functions. We use  $C_p(X)$  to denote  $C(X)$  equipped with the pointwise convergence topology in which a basic neighborhood of  $f \in C(X)$  is  $B(f, F, \epsilon) = \{g : |f(x) - g(x)| < \epsilon \text{ for any } x \in F\}$  where  $F$  is a finite subset of  $X$  and  $\epsilon$  is a positive real number.

The following result is proved in [14]. In fact, it is shown in [14] that the neighborhood filter at the identity element  $\mathbf{0}$  in  $C_p(X)$  is Tukey equivalent to  $[X]^{<\omega}$ .

**Proposition 5.5.** *Let  $P$  be a poset with Caliber  $(\omega_1, \omega)$  with  $P \geq_T \omega$ . Then the following are equivalent:*

- i)  $C_p(X)$  has a  $P$ -base;
- ii)  $C_p(X)$  is first-countable, i.e., has an  $\omega$ -base;
- iii)  $X$  is countable.

For any Tychonoff space  $X$ , it is known (see [17]) that  $C_p(X)$  is angelic if  $X$  is web-compact. A topological space  $X$  is said to be web-compact if there exists a nonempty subset  $\Sigma$  of  $\omega^\omega$  and a family  $\{A_\alpha\}$  of subsets of  $X$  such that the following two conditions hold:

- i)  $X = \overline{\bigcup\{A_\alpha : \alpha \in \Sigma\}}$ ;
- ii) the sequence  $(x_k)_k$  has a cluster point in  $X$  if  $x_k \in C_{\alpha|_k}$  for each  $k$ , here  $C_{\alpha|_k} = \bigcup\{A_\beta : \beta(i) = \alpha(i) \text{ for all } i < k\}$ .

We use  $C_k(X)$  to denote  $C(X)$  equipped with the compact-open topology in which a basic neighborhood of  $f \in C(X)$  is  $B(f, K, \epsilon) = \{g : |f(x) - g(x)| < \epsilon \text{ for any } x \in K\}$  where  $K$  is a compact subset of  $X$  and  $\epsilon$  is a positive real number.

Let  $M$  be a metric space. It is known (see Corollary 6.10 in [17]) that  $C_k(M)$  is angelic if and only if  $M$  is separable. Here, we get a stronger result below.

**Corollary 5.6.** *Let  $M$  be a metric space. Then the following are equivalent:*

- i)  $M$  is separable;
- ii)  $C_k(M)$  is strictly angelic;
- iii)  $C_k(M)$  is angelic.

*Proof.* The implication of ii)  $\rightarrow$  iii) is clear and iii)  $\rightarrow$  i) is proved in [17]. It remains to prove i)  $\rightarrow$  ii).

Let  $M$  be a separable metric space. Then  $\mathcal{B} = \{B(\mathbf{0}, K, 1/n) : K \in \mathcal{K}(M) \text{ and } n \in \mathbb{N}\}$  is a local base at the identity  $\mathbf{0}$  in  $C_k(M)$ . Clearly,  $\mathcal{B} =_T \mathcal{K}(M) \times \omega$  which is known to have Calibre  $(\omega_1, \omega)$ . Hence by Theorem 5.4,  $C_k(M)$  is strictly angelic.  $\square$

It is proved in [17] that  $C_k(\omega_1)$  has an  $\omega^\omega$ -base under the assumption  $\omega_1 = \mathfrak{b}$ , but not when  $\omega_1 < \mathfrak{b}$ . Note that  $\mathcal{K}(\omega_1) =_T \omega_1$ , hence it has Calibre  $(\omega_1, \omega)$ . Note that  $C_k(\omega_1)$  has a  $\mathcal{K}(\omega_1) \times \omega$ -base, hence an  $\omega_1 \times \omega$ -base. Note that  $\omega_1$  has Calibre  $(\omega_1, \omega)$ , hence so does  $\omega_1 \times \omega$ . Therefore,  $C_k(\omega_1)$  is strictly angelic by Theorem 5.4. Also, using the fact  $\omega_1 \leq_T \omega^\omega \times \omega_1 \leq_T \mathcal{K}(\mathbb{Q})$  proved in [12], it has a  $\mathcal{K}(\mathbb{Q})$ -base which clearly has Calibre  $(\omega_1, \omega)$ .

**Example 5.7.** *The function space  $C_k(\omega_1)$  has a  $\mathcal{K}(\mathbb{Q})$ -base, hence it is strictly angelic.*

Motivated by Theorem 5.4, it is natural to ask as in [9, Problem 6.5] whether it is sufficient for a topological group  $G$  to have an  $\omega^\omega$ -base if all precompact subsets in  $G$  are metrizable and the character of  $G$  is  $\leq \mathfrak{d}$ . Next we give an example which gives a consistent negative answer to this problem.

**Example 5.8.** *There is a topological Boolean group of character  $\chi(G) \leq \mathfrak{d}$  such that all the precompact subsets in  $G$  are metrizable and  $G$  doesn't have an  $\omega^\omega$ -base consistently, but it has a  $\mathcal{K}(\mathbb{Q})$ -base in ZFC.*

*Proof.* Let  $X = 2^{\omega_1}$  with the  $G_\delta$ -topology, i.e., a basic open neighborhood of any  $x \in X$  is  $B_\alpha(x) = \{y : y(\gamma) = x(\gamma) \text{ for all } \gamma < \alpha\}$  for  $\alpha < \omega_1$ . Define the operation on  $X$  as  $(x + y)(\alpha) = x(\alpha) + y(\alpha)$  for each  $\alpha < \omega_1$  following the rules:  $0 + 0 = 0$ ,  $0 + 1 = 1 + 0 = 1$ , and  $1 + 1 = 0$ . It is straightforward to verify that  $X$  is a topological Boolean group. Also, any countable subset of  $X$  is closed and discrete, hence any precompact subset in  $X$  is finite, therefore metrizable.

Let  $\mathbf{0}$  be the identity element in  $X$ , i.e.,  $\mathbf{0}(\alpha) = 0$  for each  $\alpha < \omega_1$ . For each  $\alpha < \omega_1$ , define  $B_\alpha(\mathbf{0}) = \{y : y(\gamma) = 0 \text{ for all } \gamma < \alpha\}$ . Then  $\{B_\alpha(\mathbf{0}) : \alpha < \omega_1\}$  is a local base at  $\mathbf{0}$ . Hence the topological group  $X$  has an  $\omega_1$ -base and its character is  $\leq \mathfrak{d}$ . Under the assumption  $\mathfrak{b} > \omega_1$ ,  $\omega_1$  is not a Tukey quotient of  $\omega^\omega$ , hence  $X$  doesn't have an  $\omega^\omega$ -base. Using the fact that  $\mathcal{K}(\mathbb{Q}) \geq_T \omega_1$  again,  $X$  has a  $\mathcal{K}(\mathbb{Q})$ -base in ZFC.  $\square$

The answer to the following question is still unknown.

**Question 5.9.** *Let  $G$  be a topological group of character  $\chi(G) \leq \mathfrak{d}$ . If all precompact subsets in  $G$  are metrizable, does  $G$  admit an  $\omega^\omega$ -base under the assumption  $\omega_1 = \mathfrak{b}$ ?*

**Example 5.10.** *There is a  $\sigma$ -compact Fréchet-Urysohn topological Boolean group which is not metrizable.*

*Proof.* Let  $X = \{x : x \in 2^{\omega_1} \text{ and } \text{supp}(x) \text{ is finite}\}$ , here  $\text{supp}(x) = \{\alpha : x(\alpha) \neq 0\}$ . The operation on  $X$  is same as the one defined in Example 5.8. Clearly,  $X$  is a Boolean group. Consider  $X$  as a subspace of  $2^{\omega_1}$  equipped with the product topology. Then the space  $X$  is Fréchet-Urysohn, but not metrizable.

Next we show that  $X$  is  $\sigma$ -compact. Define  $K_n = \{x : x \in X \text{ and } \text{supp}(x) \leq n\}$ . It suffices to show that  $K_n$  is a closed subset of  $2^{\omega_1}$  for each  $n \in \omega$ . Fix  $n \in \omega$  and  $x \notin K_n$ . Choose  $\{\alpha_1, \dots, \alpha_{n+1}\}$  such that  $x(\alpha_i) = 1$  for each  $i = 1, \dots, n+1$ . Let  $B(x) = \{y : y(\alpha_i) = x(\alpha_i) \text{ for each } i = 1, \dots, n+1\}$  which is an open neighborhood of  $x$ . And,  $K_n \cap B(x) = \emptyset$ . Hence  $K_n$  is compact for each  $n$ . Therefore,  $X = \bigcup\{K_n : n \in \omega\}$  is  $\sigma$ -compact.  $\square$

**Acknowledgements** I am very grateful to Professors Gary Gruenhagen and Paul Gartside for their valuable comments and suggestions which improve the paper. I would also like to thank Professor Paul Gartside for Example 3.4 and the anonymous reviewer for his/her valuable suggestions and corrections.

#### REFERENCES

- [1] A.V. Arhangel'skiĭ and M.G. Tkachenko, *Topological Groups and Related Structures*, Atlantis Press and World Sci., Paris, 2008.
- [2] E.K. van Douwen, *The Integers and Topology*, Handbook of Set-Theoretic Topology, North-Holland, Amsterdam (1984), 111–167.
- [3] Taras Banach, *Topological spaces with an  $\omega^\omega$ -base*, Dissertationes Math. **538** (2019), 141, DOI 10.4064/dm762-4-2018. MR3942223
- [4] Taras Banach and Arkady Leiderman,  *$\omega^\omega$ -dominated function spaces and  $\omega^\omega$ -bases in free objects of topological algebra*, Topology and Appl. **241** (2018), 203–241.
- [5] Alan Dow and Ziqin Feng, *Compact spaces with a  $P$ -base*, arxiv.org/abs/2008.04405.
- [6] Ziqin Feng, *Spaces with a  $\mathbb{Q}$ -diagonal*, Fund. Math. **245** (2019), no. 3, 305–320, DOI 10.4064/fm615-8-2018.
- [7] J.C. Ferrando, J. Kąkol, M. López-Pellicer, and S.A. Saxon, *Tightness and distinguished Fréchet spaces*, J. Math. Anal. Appl. **324** (2006), 862–881.
- [8] K. Floret, *Weakly compact sets*, Lecture notes in Math. 801, Springer, Berlin, 1980.
- [9] Saak Gabrielyan, Jerzy Kąkol, and A.G. Leiderman, *On topological groups with a small base and metrizability*, Fund. Math. **229** (2015), no. 2, 129–158, DOI 10.4064/fm229-2-3.
- [10] Paul Gartside, *Tukey Order and Diversity of Free Abelian Topological Groups*, Preprint.
- [11] Paul Gartside and Ana Mamatelashvili, *The Tukey order on compact subsets of separable metric spaces*, J. Symb. Log. **81** (2016), no. 1, 18–200, DOI 10.1017/jsl.2015.49. MR3471135
- [12] ———, *Tukey order, calibres and the rationals*, Ann. Pure Appl. Logic **172** (2021), no. 1.
- [13] Paul Gartside and Jeremiah Morgan, *Calibres, compacta and diagonals*, Fund. Math. **232** (2016), no. 1, 1–19.
- [14] ———, *Local networks for function spaces*, Houston J. Math. **45** (2019), no. 3, 893–923.
- [15] David Guerrero Sánchez, *Spaces with an  $M$ -diagonal*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. Racsam **114** (2020), no. 1, 16–24, DOI 10.1007/S13398-019-00745-x. MR4039696
- [16] D. Guerrero Sánchez and V. Tkachuk, *Domination by a Polish space of the complement of the diagonal of  $X$  implies that  $X$  is cosmic*, Topology and Appl. **212** (2016), 81–89.
- [17] J. Kąkol, W. Kubiś, and M. López-Pellicer, *Descriptive topology in selected topics of functional analysis*, Developments Math., Springer, 2011.
- [18] A.G. Leiderman, V. Pestov, and A.H. Tomita, *On topological groups admitting a base at the identity indexed by  $\omega^\omega$* , Fund. Math. **238** (2017), 79–100.
- [19] Ana Mamatelashvili, *Tukey order on sets of compact subsets of topological spaces*, ProQuest LLC, An Arbor, MI, 2014. Thesis (Ph.D.)—University of Pittsburgh.
- [20] Rongxin Shen and Ziqin Feng, *On  $\omega^\omega$ -bases and  $\omega^\omega$ -weak bases*, Houston J. Math. **46** (2020), no. 2, 507–518.
- [21] John W. Tukey, *Convergence and Uniformity in Topology*, Annals of Mathematics Studies, no. 2, Princeton University Press, Princeton, N. J., 1940. MR0002515

DEPARTMENT OF MATHEMATICS AND STATISTICS, AUBURN UNIVERSITY, AUBURN, AL 36849  
*Email address:* [zzf0006@auburn.edu](mailto:zzf0006@auburn.edu)