

Generalisation of Michelson contrast for operators and its properties

Sami Abdullah Abed*, Irina Nikolaeva^o, Andrei Novikov[†]

June 22, 2022

* E-mail: samialbarkish@gmail.com College of Administration and Economics, Diyala University, Baquba, Diyala Governorate, Republic of Iraq

o E-mail: i.nikolaeva@innopolis.ru Innopolis University, Universitetskaya st., 1, Innopolis, Rep. Tatarstan, 420500 Russia

† E-mail: a.hobukob@gmail.com Sobolev Institute of Mathematics, Siberian Branch of Russian Academy of Sciences, Novosibirsk, Russia

Abstract

In this article we consider generalisation of Michelson contrast for positive operators of countably decomposable W^* -algebras and find a few several of its properties.

subclass: 47B65, 47L07, 47C15

keywords: positive operator, invertible operator, tracial functional, C^* -algebra, W^* -algebra, von Neumann algebra, Michelson contrast, visibility

1 Introduction

The article was first composed under the inner motivation of functional analysis, defining and describing interesting properties of the "centrality measure", since from various points of view central and tracial elements are in some sense "better" than arbitrary elements of the algebra and functionals, respectively. The vast majority of articles dedicated to characterization of tracial functionals by inequalities can be found in the overview. [1]

The main inspiration was found 2019 article by A.M. Bikchentaev and S.A.Abed [10], that contains the following result:

For $\varphi \in \mathcal{M}_*^+$ the following conditions are equivalent:

- (i) φ trace;
- (ii) $\forall A, B \in \mathcal{M}^+$

$$\varphi(A^2 + B^2 + K(AB + BA)) \geq 0;$$

- (iii) $\exists K_0 > -1 \forall K > K_0 \forall A, B \in \mathcal{M}^+$

$$\varphi(A^2 + B^2 + K(AB + BA)) \geq 0.$$

Usually the inequalities that characterize the trace are binary, they give "yes" or "no" type answer, which concludes necessary and sufficient conditions for the element to be or not to be tracial (or central). But the latter condition of the work [10] seemed to give different variations of "closeness" to the tracial function. We worked with this conception and obtained "measure of centrality" and this article as a result.

As we see further the "measure of centrality" in case of positive bounded operators has a tight connection with the concept of invertibility and the classical results as Gelfand–Mazur theorem and Dixmier characterization of positive operators. Also, invertibility already appeared as an property in some sense similar to the commutativity in [3, 4, 5].

Surprisingly for us, after we have written the first version of the text we have discovered that the "centrality measure" that we have defined in this paper from purely algebraic motivation, eventually coincided in its form with the so called Michelson contrast (also called visibility measure), that is widely applied in image processing [7, 8], optics [6] and quantum computing [9].

2 Notations and Preliminaries

We adhere the following notation. Let \mathcal{A} denote some Banach $*$ -algebras, then $\mathcal{A}^{\text{sa}}, \mathcal{A}^+$ are its selfadjoint and positive parts. \mathcal{A}^* is the conjugate space of continuous linear functionals. If \mathcal{A} is a von Neumann algebra, then \mathcal{A}_* denotes its predual space. Also, $\mathcal{A}_*^+, \mathcal{A}^{*+}$ are the positive cones in \mathcal{A}_* and \mathcal{A}^* , respectively. By Tr we denote the canonical trace on $\mathbb{M}_n(\mathbb{C})$. By $C(H)$ and $B(H)$ we denote the ideal of compact operators and algebra of bounded operators, respectively.

The following lemma is known at least since 1969 [2, Lemma 1.6.2]

Lemma 1 (Dixmier). *Let \mathcal{A} be a unital C^* -algebra with unit $\mathbf{1}$ and its element $x \in \mathcal{A}$ be hermitian (i.e. $x = x^*$), then*

$$\| \|x\| \mathbf{1} - x \| \leq \|x\| \text{ if and only if } x \text{ is positive.}$$

3 Centrality and invertibility in $B(H)^+$

Let $\mathcal{A} = B(H)$, then the center $\mathfrak{C}(B(H))$ of $B(H)$ is equal to $\mathbb{C}\mathbf{1}$. Let us consider the function

$$\Delta(x) = \inf_{A \in \mathbb{R}^+} \left\{ \left\| \mathbf{1} - \frac{x}{A} \right\| \right\} \text{ for } x \in B(H)^+,$$

which illustrates how far the element x is from the central elements.

We see, that if $x = \mathbf{1}$, then $\Delta(\mathbf{1}) = 0$ ($A = 1$) and $\Delta(\mathbf{0}) = 1$.

Proposition 1. *Let x be positive operator ($x \in B(H)^+$), then $\Delta(x) \leq 1$.*

Proof. Straight from the Lemma 1 we got that $\left\| \mathbf{1} - \frac{x}{\|x\|} \right\| \leq 1$ for any $x \in B(H)^+$, thus the inequalities

$$\inf_{A \in \mathbb{R}^+} \left\| \mathbf{1} - \frac{x}{A} \right\| \leq \left\| \mathbf{1} - \frac{x}{\|x\|} \right\| \leq 1$$

hold. □

Proposition 2. *Let x be positive non-invertible (singular) operator, then*

$$\Delta(x) = 1.$$

Proof. We know, that x has the spectrum

$$\sigma\left(\frac{x}{A}\right) \subset \left[0, \frac{\|x\|}{A}\right], \text{ where } \inf \sigma(x) = 0, \sup \sigma(x) = \|x\|.$$

Thus,

$$\sigma\left(\mathbf{1} - \frac{x}{A}\right) \subset \left[1 - \frac{\|x\|}{A}, 1\right].$$

Note, that $\mathbf{1} - \frac{x}{A}$ always is selfadjoint, thus $\left\| \mathbf{1} - \frac{x}{A} \right\| \geq 1$ for any $A > 0$ and $\inf_{A \in \mathbb{R}^+} \left\| \mathbf{1} - \frac{x}{A} \right\| \geq \inf_{A \in \mathbb{R}^+} 1 = 1$. □

Corollary 1. *Let x be positive compact operator, then $\Delta(x) = 1$.*

The corollary characterize compact operators as the of of the kind of most non-central positive elements in sense of the function Δ in $B(H)$.

Theorem 1. *Let x be invertible positive operator ($x \in B(H)^+$), with the inverse x^{-1} , then*

$$\Delta(x) = \frac{\|x\|\|x^{-1}\| - 1}{\|x\|\|x^{-1}\| + 1}. \tag{1}$$

Proof. Let now x be invertible element of $B(H)$, then the spectrum

$$\sigma(x) \subset \left[\frac{1}{\|x^{-1}\|}, \|x\| \right] \text{ and } \inf \sigma(x) = \frac{1}{\|x^{-1}\|}, \sup \sigma(x) = \|x\|,$$

thus

$$\sigma\left(1 - \frac{x}{A}\right) \subset \left[1 - \frac{\|x\|}{A}, 1 - \frac{1}{A\|x^{-1}\|}\right],$$

$$\left\|1 - \frac{x}{A}\right\| = \max\left(\left|1 - \frac{1}{A\|x^{-1}\|}\right|, \left|1 - \frac{\|x\|}{A}\right|\right).$$

Let $\alpha = \min(\frac{1}{\|x^{-1}\|}, \|x\|)$, $\beta = \max(\frac{1}{\|x^{-1}\|}, \|x\|)$, $\lambda = \frac{1}{A} \geq 0$. Then the latter maximum is rewritten as

$$\max(|1 - \alpha\lambda|, |1 - \beta\lambda|) = \frac{|1 - \alpha\lambda| + |1 - \beta\lambda| + ||1 - \alpha\lambda| - |1 - \beta\lambda||}{2}.$$

and we need to minimize it due to $\lambda \in (0, +\infty)$

$$\max(|1 - \alpha\lambda|, |1 - \beta\lambda|) \rightarrow \min.$$

We have here three basic situations, for $\lambda \in (0, \frac{1}{\beta})$, $\lambda \in (\frac{1}{\beta}, \frac{1}{\alpha})$ and $\lambda > \frac{1}{\alpha}$.

1) Let $\lambda \in (0, \frac{1}{\beta})$, then

$$|1 - \alpha\lambda| + |1 - \beta\lambda| + ||1 - \alpha\lambda| - |1 - \beta\lambda|| = 2 - 2\alpha\lambda \geq 2\frac{\beta - \alpha}{\beta}.$$

2) Let $\lambda > \frac{1}{\alpha}$, then

$$|1 - \alpha\lambda| + |1 - \beta\lambda| + ||1 - \alpha\lambda| - |1 - \beta\lambda|| = 2\beta\lambda - 2 \geq 2\frac{\beta - \alpha}{\alpha}.$$

3) Let $\lambda \in [\frac{1}{\beta}, \frac{1}{\alpha}]$, then

$$|1 - \alpha\lambda| + |1 - \beta\lambda| + ||1 - \alpha\lambda| - |1 - \beta\lambda|| = (\beta - \alpha)\lambda + |2 - (\alpha + \beta)\lambda|.$$

Now either $\lambda > \frac{2}{\alpha + \beta}$ and then

$$(\beta - \alpha)\lambda + (\alpha + \beta)\lambda - 2 = 2(\beta\lambda - 1) \geq 2\frac{\beta - \alpha}{\alpha + \beta},$$

or $\lambda < \frac{2}{\alpha + \beta}$ and

$$(\beta - \alpha)\lambda + (\alpha + \beta)\lambda - 2 = 2(1 - \alpha\lambda) \geq 2\frac{\beta - \alpha}{\alpha + \beta}.$$

At last,

$$\inf\{\max(|1 - \alpha\lambda|, |1 - \beta\lambda|)\} = \frac{(\beta - \alpha)}{\max\{\alpha, \beta, \alpha + \beta\}} = \frac{\beta - \alpha}{\alpha + \beta}.$$

Now, note, that

$$\begin{aligned}\beta - \alpha &= \max\left(\frac{1}{\|x^{-1}\|}, \|x\|\right) - \min\left(\frac{1}{\|x^{-1}\|}, \|x\|\right) = \\ &= \left|\|x\| - \frac{1}{\|x^{-1}\|}\right| = \left|\frac{\|x\|\|x^{-1}\| - 1}{\|x^{-1}\|}\right|.\end{aligned}$$

Also, note, that $1 = \|\mathbf{1}\| = \|xx^{-1}\| \leq \|x\|\|x^{-1}\|$, thus

$$\beta - \alpha = \frac{\|x\|\|x^{-1}\| - 1}{\|x^{-1}\|}.$$

To accomplish the proof, note, that

$$\alpha + \beta = \frac{\|x\|\|x^{-1}\| + 1}{\|x^{-1}\|}.$$

□

Remark 1. Note, that from the latter theorem it follows, that for any $x \in B(H)^+$ the equality $\Delta(x) = \Delta(x^{-1})$ holds.

Remark 2. Note, that $\Delta(x) = 0$ for the invertible operator, only if $\|x^{-1}\| = \frac{1}{\|x\|}$, which would mean $\sigma(x) = [\|x\|, \|x\|]$, thus $x = \|x\|\mathbf{1}$.

Corollary 2. *Let $x \in B(H)^+$ be invertible element with the inverse element $x^{-1} \in B(H)$, then $\Delta(x) < 1$.*

Corollary 3. *Let the sequence x_n from $B(H)^+$ that converges to element $\mathbf{0} \neq x \in B(H)^+$ in terms of norm,*

$$\lim_n \Delta(x_n) = \Delta(x),$$

i.e. $\Delta : (B(H)^+ \setminus \{\mathbf{0}\}, \|\cdot\|) \mapsto [0, 1]$ is a continuous function.

Proof. It is sufficient to consider four cases:

- 1) x_n consists of invertible elements and x is invertible;
- 2) x_n consists of invertible elements and x is not invertible;
- 3) x_n consists of non-invertible elements and x is invertible;
- 4) x_n consists of non-invertible elements and x is not invertible.

The cases 1) and 4) are evident.

Consider 2), when x_n are invertible and x is not. Then

$$\Delta(x_n) = \frac{\|x_n^{-1}\| - \frac{1}{\|x_n\|}}{\|x_n^{-1}\| + \frac{1}{\|x_n\|}}.$$

The sequence $\|x_n^{-1}\|$ converges to infinity. Note, that

$$\|x_n^{-1} - x_m^{-1}\| \leq \|x_n^{-1}\| \|x_m^{-1}\| \|x_n - x_m\|,$$

thus if $\|x_n^{-1}\|$ has a bounded sub-sequence $\|x_{n_k}\|$, then $x_{n_k}^{-1}$ is fundamental, therefore there exists its limit z such that $x_{n_k}^{-1}x_{n_k} \rightarrow zx$ and $x_{n_k}x_{n_k}^{-1} \rightarrow xz$ on one hand, and $x_{n_k}^{-1}x_{n_k} \rightarrow \mathbf{1}$ on the other hand, thus $z = x^{-1}$, which is forbidden by assumption. Therefore, any sub-sequence of $\|x_n^{-1}\|$ is unbounded.

Since $\|x_n^{-1}\| \rightarrow \infty$ and $\|x_n\| \rightarrow \|x\|$, it follows, that $\Delta(x_n) \rightarrow 1$.

Consider 3) with x_n being a sequence of non-invertible positive operators converging to invertible operator x by norm. Consider $x'_n = \frac{1}{n} + x_n$, then each of x'_n is invertible and $x'_n \rightarrow x$, thus $\Delta(x'_n) \rightarrow \Delta(x)$. On the other hand, by formula 1

$$\begin{aligned} |\Delta(x_n) - \Delta(x'_n)| &= \left| 1 - \frac{\|\frac{1}{n} + x_n\| \|(\frac{1}{n} + x_n)^{-1}\| - 1}{\|\frac{1}{n} + x_n\| \|(\frac{1}{n} + x_n)^{-1}\| + 1} \right| = \\ &= \frac{2}{\|\frac{1}{n} + x_n\| \|(\frac{1}{n} + x_n)^{-1}\| + 1} = \frac{2}{2 + n\|x_n\|} \rightarrow 0, \end{aligned}$$

thus $|\Delta(x_n) - \Delta(x)| \leq |\Delta(x_n) - \Delta(x'_n)| + |\Delta(x'_n) - \Delta(x)| \rightarrow 0$. □

From the latter two corollaries we obtain the following.

Corollary 4. *If the sequence of operators is converging to a non-singular (invertible) operator, then the sequence contains not more than a finite quantity of non-invertible operators.*

Corollary 5. *For any x in $B(H)^+$ the following properties*

1)

$$\Delta(x) = \frac{\sup(\sigma(x)) - \inf(\sigma(x))}{\sup(\sigma(x)) + \inf(\sigma(x))};$$

2)

$$\Delta(x) = \frac{\sup \sigma(x)}{\sup \sigma(x) + \inf \sigma(x)} \left\| 1 - \frac{x}{\|x\|} \right\|.$$

hold.

Proof. 1) follows straight from the proof of the theorem. 2) Since invertible elements are dense in $B(H)$ and Δ is continuous, it is sufficient to consider invertible x . By Theorem 1 we have

$$\Delta(x) = \frac{\|x\|\|x^{-1}\| - 1}{\|x\|\|x^{-1}\| + 1}.$$

At the same time, $\mathbf{1} - \frac{x}{\|x\|} \geq \mathbf{0}$, thus

$$\begin{aligned} \left\| \mathbf{1} - \frac{x}{\|x\|} \right\| &= \sup \sigma \left(\mathbf{1} - \frac{x}{\|x\|} \right) = 1 - \inf \sigma \left(\frac{x}{\|x\|} \right) = \\ &= 1 - \frac{1}{\|x\|} \inf \sigma(x) = 1 - \frac{1}{\|x\|} \frac{1}{\|x^{-1}\|} = \frac{\|x\|\|x^{-1}\| - 1}{\|x\|\|x^{-1}\|}. \end{aligned}$$

At last,

$$\Delta(x) = \frac{\|x\|\|x^{-1}\|}{\|x\|\|x^{-1}\| + 1} \frac{\|x\|\|x^{-1}\| - 1}{\|x\|\|x^{-1}\|} = \frac{\|x\|}{\|x\| + \frac{1}{\|x^{-1}\|}} \left\| \mathbf{1} - \frac{x}{\|x\|} \right\|$$

□

Corollary 6. *Let $x \in B(H)^+$, then*

1)

$$\frac{1}{2} \left\| \mathbf{1} - \frac{x}{\|x\|} \right\| \leq \Delta(x) \leq \left\| \mathbf{1} - \frac{x}{\|x\|} \right\|;$$

2) $\Delta(x) = 0$ only if $x = \|x\|\mathbf{1}$.

Remark 3. Note, that algebra of compact operators $C(H)$ is dense in $B(H)$ in weak operator topology (and $C(H)^+$ in $B(H)^+$, respectively), thus Δ cannot be continuous in weak operator topology.

Now let us describe some properties of Δ .

Theorem 2. *Let $x, y \in B(H)^+$, $\lambda \in \mathbb{R}^+$ then*

1) $\Delta(\lambda x) = \Delta(x)$;

2) if x, y are invertible, then

$$\begin{aligned} &\left(\|x + y\| + \frac{1}{\|(x + y)^{-1}\|} \right) \Delta(x + y) \leq \\ &\leq \left(\|x\| + \frac{1}{\|x^{-1}\|} \right) \Delta(x) + \left(\|y\| + \frac{1}{\|y^{-1}\|} \right) \Delta(y). \end{aligned}$$

The inequality becomes an equality if $x = \lambda y$.

Proof. 1) By definition

$$\Delta(\lambda x) = \inf_{A \in \mathbb{R}^+} \left\{ \left\| \mathbf{1} - \frac{\lambda x}{A} \right\| \right\} = \inf_{A' \in \mathbb{R}^+, A' = A/\lambda} \left\{ \left\| \mathbf{1} - \frac{x}{A'} \right\| \right\} = \Delta(x).$$

2) Note, that $x + y$ is invertible, $\|x + y\| \mathbf{1} \leq \|x\| \mathbf{1} + \|y\| \mathbf{1}$, thus

$$0 \leq \|x + y\| \mathbf{1} - (x + y) \leq \|x\| \mathbf{1} - x + \|y\| \mathbf{1} - y,$$

therefore

$$\| \|x + y\| \mathbf{1} - (x + y) \| \leq \| \|x\| \mathbf{1} - x \| + \| \|y\| \mathbf{1} - y \|.$$

In terms of upper and lower bounds we see that

$$\sup \sigma(x + y) - \inf \sigma(x + y) \leq \sup \sigma(x) - \inf \sigma(x) + \sup \sigma(y) - \inf \sigma(y),$$

thus

$$\begin{aligned} \Delta(x + y) &= \frac{\sup \sigma(x + y) - \inf \sigma(x + y)}{\sup \sigma(x + y) + \inf \sigma(x + y)} \leq \\ &\leq \frac{\sup \sigma(x) - \inf \sigma(x)}{\sup \sigma(x + y) + \inf \sigma(x + y)} + \frac{\sup \sigma(y) - \inf \sigma(y)}{\sup \sigma(x + y) + \inf \sigma(x + y)} = \\ &= \frac{\sup \sigma(x) + \inf \sigma(x)}{\sup \sigma(x + y) + \inf \sigma(x + y)} \Delta(x) + \frac{\sup \sigma(y) + \inf \sigma(y)}{\sup \sigma(x + y) + \inf \sigma(x + y)} \Delta(y). \end{aligned}$$

Note, that if $x = \lambda y$, then

$$\begin{aligned} &\left(\|x + y\| + \frac{1}{\|(x + y)^{-1}\|} \right) \Delta(x + y) = \\ &= (1 + \lambda) \left(\|x\| + \frac{1}{\|x^{-1}\|} \right) \Delta((1 + \lambda)x) = \\ &= \left(\|x\| + \frac{1}{\|x^{-1}\|} \right) \Delta(x) + \lambda \left(\|x\| + \frac{1}{\|x^{-1}\|} \right) \Delta(x) = \\ &= \left(\|x\| + \frac{1}{\|x^{-1}\|} \right) \Delta(x) + \left(\|\lambda x\| + \frac{1}{\|(\lambda x)^{-1}\|} \right) \Delta(\lambda x). \end{aligned}$$

□

Corollary 7. *Let $x, y \in B(H)^+$, then from inequality iv) of the latter Theorem we obtain the following*

1) *if x, y are singular (non-invertible) and $x + y$ is invertible (non-singular), then*

$$\left(\|x + y\| + \frac{1}{\|(x + y)^{-1}\|} \right) \Delta(x + y) \leq \|x\| \Delta(x) + \|y\| \Delta(y);$$

2) if $x + y$ is singular (non-invertible), then

$$\|x + y\|\Delta(x + y) \leq \|x\|\Delta(x) + \|y\|\Delta(y);$$

3) if x is invertible and y is singular (non-invertible), then

$$\begin{aligned} & \left(\|x + y\| + \frac{1}{\|(x + y)^{-1}\|} \right) \Delta(x + y) \leq \\ & \leq \left(\|x\| + \frac{1}{\|x^{-1}\|} \right) \Delta(x) + \|y\|\Delta(y). \end{aligned}$$

Lemma 2. Let $X, Y \in \mathbb{M}_2(\mathbb{C})^+$, then $\Delta(X + Y) \leq \max(\Delta(X), \Delta(Y))$.

Proof. Using Jordan normal form decomposition we assume, that

$$X = \begin{pmatrix} \alpha_1 & 0 \\ 0 & \beta_1 \end{pmatrix} \text{ and } Y = \begin{pmatrix} \beta_2 + (\alpha_2 - \beta_2)\lambda & (\alpha_2 - \beta_2)\delta\sqrt{\lambda(1-\lambda)} \\ (\alpha_2 - \beta_2)\delta\sqrt{\lambda(1-\lambda)} & \alpha_2 + (\beta_2 - \alpha_2)\lambda \end{pmatrix};$$

where $\alpha_{1,2}, \beta_{1,2} \geq 0$, $\lambda \in [0, 1]$ and $|\delta| = 1$, $\delta \in \mathbb{C}$.

Evidently,

$$\Delta(X) = \frac{|\beta_1 - \alpha_1|}{\beta_1 + \alpha_1} \text{ and } \Delta(Y) = \frac{|\beta_2 - \alpha_2|}{\beta_2 + \alpha_2}.$$

Let us find $\Delta(X + Y)$ by finding the eigenvalues of $X + Y$. We have

$$\begin{aligned} & (\alpha_1 + \alpha_2\lambda + \beta_2 - \beta_2\lambda - t) \times \\ & \times (\beta_1 + \alpha_2 - \alpha_2\lambda + \beta_2\lambda - t) - (\alpha_2 - \beta_2)^2\lambda(1 - \lambda) = 0 \end{aligned}$$

We expand the first two brackets and regroup it into quadratic equation

$$\begin{aligned} & t^2 - t((\alpha_2 - \beta_2)\lambda + (\alpha_1 + \beta_2) + (\beta_2 - \alpha_2)\lambda + (\beta_1 + \alpha_2)) - \\ & - (\beta_2 - \alpha_2)^2\lambda^2 + (\alpha_1 + \beta_2)(\beta_2 - \alpha_2)\lambda + \\ & + (\alpha_2 + \beta_1)(\alpha_2 - \beta_2)\lambda + (\alpha_1 + \beta_2)(\beta_1 + \alpha_2) - \\ & - (\alpha_2 - \beta_2)^2\lambda + (\alpha_2 - \beta_2)^2\lambda^2 = 0. \end{aligned}$$

We simplify the equation and get the following

$$\begin{aligned} & t^2 - t(\alpha_1 + \alpha_2 + \beta_2 + \beta_1) + (\alpha_1 - \beta_1 + \beta_2 - \alpha_2)(\beta_2 - \alpha_2)\lambda + \\ & + (\alpha_1 + \beta_2)(\beta_1 + \alpha_2) - (\beta_2 - \alpha_2)^2\lambda = 0. \end{aligned}$$

At last, we obtain the equation

$$t^2 - t(\alpha_1 + \alpha_2 + \beta_2 + \beta_1) - (\beta_1 - \alpha_1)(\beta_2 - \alpha_2)\lambda + (\alpha_1 + \beta_2)(\beta_1 + \alpha_2) = 0.$$

Note, that

$$\Delta(X + Y) = \frac{|t_1 - t_2|}{t_1 + t_2} = \frac{\sqrt{D}}{2b},$$

where $D = b^2 - 4ac$,

$$a = 1, b = \alpha_1 + \alpha_2 + \beta_1 + \beta_2,$$

$$c = ((\beta_1 - \alpha_1)(\beta_2 - \alpha_2)\lambda - (\alpha_1 + \beta_2)(\beta_1 + \alpha_2)).$$

Assume $\Delta(X) \leq \Delta(Y)$, that will give us the condition $|(b_1 - a_1)(b_2 + a_2)| \leq |(b_2 - a_2)(b_1 + a_1)|$. Without loss of generality we can assume, that $a_1 \leq b_1$ and $a_2 \leq b_2$ (if it is not, then we can use renaming and changing λ to $1 - \lambda$), then $0 \leq b_2 a_1 - a_2 b_1$.

We need to prove, that

$$\text{Tr}^2(X+Y) + 4\text{Tr}(X)\text{Tr}(Y)\Delta(X)\Delta(Y)\lambda \leq 4(\alpha_1 + \beta_2)(\beta_1 + \alpha_2) + \Delta(Y)^2 \text{Tr}^2(X+Y).$$

We divide the inequality by $\text{Tr}^2(X + Y)$ and get

$$1 + 4\lambda \frac{\text{Tr}(X)\text{Tr}(Y)}{(\text{Tr}(X) + \text{Tr}(Y))^2} \Delta(X)\Delta(Y) \leq 4(1 - \theta)\theta + \Delta(Y)^2,$$

where

$$\theta_1 = \frac{\beta_1 + \alpha_2}{\text{Tr}(X + Y)} \text{ and } \theta_2 = 4 \frac{\text{Tr}(X)\text{Tr}(Y)}{(\text{Tr}(X) + \text{Tr}(Y))^2}.$$

Evidently, $\theta_1, \theta_2 \in [0, 1]$. We rewrite the inequality in the form

$$\lambda \theta_2 \Delta(X)\Delta(Y) \leq \Delta(Y)^2 - (1 - 2\theta_1)^2.$$

The latter inequality holds if and only if $\Delta(X + Y) \leq \max(\Delta(X), \Delta(Y))$. We see that λ effects only the left-hand side of inequality, thus if the inequality does not hold, then there exists a counter-example in the form

$$X = \begin{pmatrix} \alpha_1 & 0 \\ 0 & \beta_1 \end{pmatrix}; \quad Y = \begin{pmatrix} \alpha_2 & 0 \\ 0 & \beta_2 \end{pmatrix}; \quad \text{with } \alpha_1 \leq \beta_1, \alpha_2 \leq \beta_2.$$

Then

$$\Delta(X + Y) = \frac{\beta_2 + \beta_1 - \alpha_2 - \alpha_1}{\alpha_1 + \alpha_2 + \beta_1 + \beta_2} > \frac{\beta_2 - \alpha_2}{\beta_2 + \alpha_2} = \Delta(Y) \geq \Delta(X) = \frac{\beta_1 - \alpha_1}{\beta_1 + \alpha_1},$$

so

$$\begin{aligned} & (\beta_2 - \alpha_2)(\beta_2 + \alpha_2) + (\beta_1 - \alpha_1)(\beta_2 + \alpha_2) > \\ & > (\beta_2 - \alpha_2)(\beta_2 + \alpha_2) + (\beta_2 - \alpha_2)(\beta_1 + \alpha_1), \end{aligned}$$

and

$$(\beta_2 - \alpha_2)(\beta_1 + \alpha_1) \leq (\beta_2 + \alpha_2)(\beta_1 - \alpha_1)$$

at the same time, which cannot be true. □

Lemma 3. Let $X, Y \in \mathbb{M}_n(\mathbb{C})^+$, then $\Delta(X + Y) \leq \max(\Delta(X), \Delta(Y))$.

Proof. Note, that in finite dimensional spaces the unit ball is a compact set, thus

$$\begin{aligned} \sup_{\|f\|=1} \langle (X + Y)f, f \rangle &= \langle (X + Y)f_0, f_0 \rangle; \\ \sup_{\|g\|=1} \langle (X + Y)g, g \rangle &= \langle (X + Y)g_0, g_0 \rangle \end{aligned}$$

for some $f_0, g_0 \in \mathbb{C}^n$, $\|f_0\| = \|g_0\| = 1$. By H_2 we denote the complex linear span $\text{Lin}_{\mathbb{C}}\{f_0, g_0\}$ of these two vectors.

Evidently, for the restriction $(X + Y)|_{H_2}$ we have $(X + Y)|_{H_2}f_0 = (X + Y)f_0$, $(X + Y)|_{H_2}g_0 = (X + Y)g_0$ and, also,

$$\begin{aligned} \langle (X + Y)|_{H_2}f_0, f_0 \rangle &\leq \sup_{\|f\|=1, f \in H_2} (\langle (X + Y)|_{H_2}f, f \rangle) \leq \\ &\leq \sup_{\|f\|=1, f \in \mathbb{C}^n} (\langle (X + Y)f, f \rangle) = \langle (X + Y)f_0, f_0 \rangle; \\ \langle (X + Y)|_{H_2}f_0, f_0 \rangle &\geq \inf_{\|f\|=1, f \in H_2} (\langle (X + Y)|_{H_2}f, f \rangle) \geq \\ &\geq \inf_{\|f\|=1, f \in \mathbb{C}^n} (\langle (X + Y)f, f \rangle) = \langle (X + Y)f_0, f_0 \rangle. \end{aligned}$$

Now, from the latter inequalities we gain the fact, that

$$\Delta(X + Y) = \Delta((X + Y)|_{H_2}) = \Delta(X|_{H_2} + Y|_{H_2}).$$

Also, note, that

$$\begin{aligned} 0 &\leq \sup_{\|f\|=1, f \in H_2} (X|_{H_2}f, f) \leq \sup_{\|f\|=1, f \in \mathbb{C}^n} (Xf, f); \\ 0 &\leq \inf_{\|f\|=1, f \in \mathbb{C}^n} (Xf, f) \leq \inf_{\|f\|=1, f \in H_2} (X|_{H_2}f, f). \end{aligned}$$

Now, consider two cases:

- 1) If $\inf_{\|f\|=1, f \in \mathbb{C}^n} (Xf, f) = 0$, then $\Delta(X|_{H_2}) \leq 1 = \Delta(X)$
- 2) Otherwise, $\inf_{\|f\|=1, f \in \mathbb{C}^n} (Xf, f) \neq 0$ and we denote

$$t = \frac{\sup_{\|f\|=1, f \in H_2} (X|_{H_2}f, f)}{\inf_{\|f\|=1, f \in H_2} (X|_{H_2}f, f)}; T = \frac{\sup_{\|f\|=1, f \in \mathbb{C}^n} (Xf, f)}{\inf_{\|f\|=1, f \in \mathbb{C}^n} (Xf, f)}.$$

Evidently, $t \leq T$. So now we have

$$\begin{aligned} \Delta(X|_{H_2}) &= \frac{\sup_{f \in H_2, \|f\|=1} (\langle X|_{H_2}f, f \rangle) - \inf_{f \in H_2, \|f\|=1} (\langle X|_{H_2}f, f \rangle)}{\sup_{f \in H_2, \|f\|=1} (\langle X|_{H_2}f, f \rangle) + \inf_{f \in H_2, \|f\|=1} (\langle X|_{H_2}f, f \rangle)} = \\ &= \frac{t - 1}{t + 1} = 1 - \frac{2}{t + 1} \leq 1 - \frac{2}{T + 1} = \end{aligned}$$

$$= \frac{\sup_{f \in \mathbb{C}^n, \|f\|=1} (\langle Xf, f \rangle) - \inf_{f \in \mathbb{C}^n, \|f\|=1} (\langle Xf, f \rangle)}{\sup_{f \in \mathbb{C}^n, \|f\|=1} (\langle Xf, f \rangle) + \inf_{f \in \mathbb{C}^n, \|f\|=1} (\langle Xf, f \rangle)} = \Delta(X).$$

By the Lemma 2 we have, that

$$\Delta(X|_{H_2} + Y|_{H_2}) \leq \max(\Delta(X|_{H_2}), \Delta(Y|_{H_2}))$$

and we just have proved that

$$\Delta(X + Y) = \Delta(X|_{H_2} + Y|_{H_2}); \max(\Delta(X|_{H_2}), \Delta(Y|_{H_2})) \leq \max(\Delta(X), \Delta(Y)).$$

□

Theorem 3. *Let $x, y \in B(H)^+$, then $\Delta(x + y) \leq \max\{\Delta(x), \Delta(y)\}$.*

Proof. Let $\{f_i\}_{i=1}^\infty$ be the orthonormal basis of H , then let

$$p_n = \sum_{i=1}^n \langle \cdot, f_i \rangle f_i$$

be the projection to the linear span of $\{f_i\}_{i=1, \dots, n}$. By $\Delta_n(x)$ we denote $\Delta(x|_{p_n})$ in the reduced algebra $B(H)_{p_n}$ of operators on $p_n H p_n$ hilbert space, i.e.

$$\Delta_n(x) = \Delta(x|_{p_n}).$$

Then $\Delta_n(x + y) \leq \max(\Delta_n(x), \Delta_n(y))$ by Lemma 3.

Now, note that for arbitrary operator a $\sup(\sigma(a_{p_n})) \leq \sup(\sigma(a_{p_{n+1}})) \leq \sup(\sigma(a))$ and $\inf(\sigma(a_{p_n})) \geq \inf(\sigma(a_{p_{n+1}})) \geq \inf(\sigma(a))$, and, also,

$$\lim_{n \rightarrow \infty} \sup(\sigma(a_{p_n})) = \sup(\sigma(a));$$

$$\lim_{n \rightarrow \infty} \inf(\sigma(a_{p_n})) = \inf(\sigma(a)).$$

Now, if $\inf(\sigma(x)) = 0$, then $\Delta(x) = 1$ and $\Delta(x + y) \leq 1 = \max(\Delta(x), \Delta(y))$. Assume, that $\inf(\sigma(x)) \neq 0$, $\inf(\sigma(y)) \neq 0$ and thus $\inf(\sigma(x + y)) \neq 0$. For a such that $\inf \sigma(a) \neq 0$ by t_n^a we denote

$$t_n^a := \frac{\sup(\sigma(a_{p_n}))}{\inf(\sigma(a_{p_n}))}; \text{ note that } t_n^a \xrightarrow{n} t^a = \frac{\sup(\sigma(a))}{\inf(\sigma(a))}.$$

Then

$$\lim_{n \rightarrow \infty} \Delta_n(x) = \lim_{n \rightarrow \infty} \frac{t_n^x - 1}{t_n^x + 1} = \Delta(x)$$

$$\lim_{n \rightarrow \infty} \Delta_n(y) = \lim_{n \rightarrow \infty} \frac{t_n^y - 1}{t_n^y + 1} = \Delta(y)$$

$$\lim_{n \rightarrow \infty} \Delta_n(x + y) = \lim_{n \rightarrow \infty} \frac{t_n^{x+y} - 1}{t_n^{x+y} + 1} = \Delta(x + y),$$

and since $\Delta_n(x + y) \leq \max\{\Delta_n(x), \Delta_n(y)\}$, we get the proposition of the theorem. □

One can ask, when the equality holds. We do not have a comprehensive answer for that, but we assume that the following example would be interesting under these considerations.

Example 1. Let $X, Y \in \mathbb{M}_2(\mathbb{C})^+$ and commute i.e. $XY = YX$, then the equality $\Delta(X + Y) = \max(\Delta(X), \Delta(Y))$ holds only if $X = \lambda Y$, where $\lambda \in \mathbb{R}^+$.

Indeed, note that $X = a_x p + b_x p^\perp$ and $Y = a_y p + b_y p^\perp$, where p is non-zero and non-unit projection $\mathbb{M}_2(\mathbb{C})^{\text{Pf}}$. Consider

$$\frac{|a_x + a_y - b_x - b_y|}{a_x + b_x + a_y + b_y} = \frac{a_x - b_x}{a_x + b_x}.$$

Evidently, in that case $b_x \neq a_x$ (consider the contrary, then $\max\{\Delta(X), \Delta(Y)\} = \Delta(Y)$). From the equality we obtain

$$\left| 1 + \frac{a_y - b_y}{a_x - b_x} \right| = 1 + \frac{a_y + b_y}{a_x + b_x}.$$

Now, either

$$2 + \frac{a_y + b_y}{a_x + b_x} + \frac{a_y - b_y}{a_x - b_x} = 0$$

or

$$(a_y - b_y)(a_x + b_x) = (a_y + b_y)(a_x - b_x).$$

In the first case $(a_y + b_y)(a_x - b_x) + (a_y - b_y)(a_x + b_x) + 2(a_x + b_x)(a_x - b_x) = 0$;

$$a_y a_x + b_y a_x - a_y b_x - b_x b_y + a_x a_y + a_y b_x - a_x b_y - b_x b_y + 2a_x^2 - 2b_x^2 = 0;$$

$$2a_x a_y - 2b_x b_y + 2a_x^2 - 2b_x^2 = 0;$$

$$a_x(a_y + a_x) = b_x(b_x + b_y);$$

thus

$$\frac{a_x}{b_x} = \frac{b_x + b_y}{a_x + a_y}.$$

Without loss of generality we may assume that

$$a_x + a_y = 1 \text{ and } \frac{a_x}{b_x} = \frac{b_x + b_y}{a_x + a_y} = \lambda < 1;$$

then

$$\Delta(X + Y) = \Delta \begin{pmatrix} 1 & 0 \\ 0 & \lambda \end{pmatrix} = \frac{1 - \lambda}{1 + \lambda} \text{ and } \Delta(X) = \Delta \begin{pmatrix} \mu\lambda & 0 \\ 0 & \mu \end{pmatrix} = \frac{\mu - \lambda\mu}{\mu + \lambda\mu} = \frac{1 - \lambda}{1 + \lambda}.$$

with $0 \neq \mu < \lambda$, which leads to contradiction, since

$$\Delta(Y) = \Delta \begin{pmatrix} 1 - \mu\lambda & 0 \\ 0 & \lambda - \mu \end{pmatrix} = \frac{1 - \mu\lambda - \lambda + \mu}{1 - \mu\lambda + \lambda - \mu} = \frac{(1 - \lambda)(1 + \mu)}{(1 + \lambda)(1 - \mu)} > \frac{1 - \lambda}{1 + \lambda},$$

which leads to contradiction with the assumption that $\max\{\Delta(X), \Delta(Y)\} = \Delta(X)$.

In the other case

$$\begin{aligned} a_x a_y - a_x b_y + a_y b_x - b_x b_y &= a_x a_y - b_x a_y + b_y a_x - b_x b_y; \\ a_y b_x &= a_x b_y, \end{aligned}$$

and at last, $a_x/b_x = a_y/b_y$, thus if we denote $a_x/b_x = a_y/b_y$ as λ

$$\begin{aligned} \frac{a_x}{b_x} &= \frac{a_y}{b_y} = \frac{\lambda}{b_x + b_y} (b_x + b_y) = \frac{\lambda b_x}{b_x + b_y} + \frac{\lambda b_y}{b_x + b_y} = \\ &= \frac{a_x}{b_x + b_y} + \frac{a_y}{b_x + b_y} = \frac{a_x + a_y}{b_x + b_y}. \end{aligned}$$

Therefore, if by λ we denote $\lambda = a_x/b_x = a_y/b_y$

$$\begin{aligned} X &= \begin{pmatrix} a_x & 0 \\ 0 & b_x \end{pmatrix}; Y = \begin{pmatrix} a_y & 0 \\ 0 & b_y \end{pmatrix} \\ \Delta(X) &= \frac{|a_x - b_x|}{a_x + b_x} = \frac{|\lambda - 1|}{\lambda + 1}; \Delta(Y) = \frac{|a_y - b_y|}{a_y + b_y} = \frac{|\lambda - 1|}{\lambda + 1}; \\ \Delta(X + Y) &= \frac{|a_x + a_y - b_x - b_y|}{a_x + a_y + b_x + b_y} = \frac{|\lambda - 1|}{\lambda + 1}. \end{aligned}$$

Remark 4. Note, that Δ does not hold monotonicity.

For example take $x = 2p + 4p^\perp$ and $y = 3p + 9p^\perp$ with $(\mathbf{0}, \mathbf{1} \neq)p \in B(H)^{\text{pr}}$, then $x \leq y$, but

$$\Delta(x) = \frac{4-2}{4+2} = \frac{1}{3} \text{ and } \Delta(y) = \frac{9-3}{9+3} = \frac{1}{2} \text{ and } \Delta(x) = \frac{1}{3} \leq \frac{1}{2} = \Delta(y).$$

As another example take $x = p + 2p^\perp$ and $y = 2p + 3p^\perp$, then $x \leq y$ and

$$\Delta(x) = \frac{2-1}{2+1} = \frac{1}{3} \geq \frac{1}{5} = \frac{3-2}{3+2} = \Delta(y).$$

Remark 5. Note, that the inequality $\min(\Delta(x), \Delta(y)) \leq \Delta(x+y)$ does not hold, for example if $(\mathbf{1}, \mathbf{0} \neq)x = p \in B(H)^{\text{pr}}$ and $y = p^\perp$, then $\Delta(p) = \Delta(p^\perp) = 1$ and $0 = \Delta(\mathbf{1}) = \Delta(p + p^\perp)$.

Basing on Theorems 2 and 3 for $B(H)$ we are able to define a cone

$$K_c = \{x \in B(H)^+ | \Delta(x) \leq c\},$$

which would be a closed subcone with the property, that if $0 \leq c_1 \leq c_2 \leq 1$, then

$$\mathbb{R}^+ \mathbf{1} \subset K_0 \subset K_{c_1} \subset K_{c_2} \subset K_1 = \mathcal{B}^+(H).$$

The latter implies, that the cone of positive elements without a zero element has a fibration into the sets $K_c \setminus \bigcup_{0 \leq b < c} K_b$ parametrized by interval $[0, 1]$.

Theorem 4. *Let $x, y \in B(H)^+$, then the following inequalities*

- 1) $\Delta(xy) = \Delta(yx)$;
- 2) $\Delta(xy) \leq \max(\Delta(x^2), \Delta(y^2))$;
- 3) $\Delta(x^2) \leq 2\Delta(x)$;
- 4) $\Delta(x) \leq \Delta(x^2)$

hold.

Proof. *i)* Note that $\sigma(xy) \setminus \{0\} = \sigma(yx) \setminus \{0\}$. Assume that $0 \notin \sigma(x)$ and $0 \notin \sigma(y)$, then $\ker x = \ker y = \{\vec{0}\}$, thus $\ker xy = \ker yx = \{\vec{0}\}$ (without loss of generality, assume $xyf = \mathbf{0}$, then if $yf \neq \vec{0}$, then $\ker x \neq \{\vec{0}\}$ (that is impossible), or $yf = \{\vec{0}\}$, which would mean that $\ker y \neq 0$). Thus,

$$\sigma(xy) = \sigma(xy) \setminus \{0\} = \sigma(yx) \setminus \{0\} = \sigma(yx).$$

If $0 \in \sigma(x)$ and $0 \in \sigma(y)$, evidently,

$$\Delta(xy) = 1 = \Delta(yx)$$

. Now, assume $0 \in \sigma(x)$ and $0 \notin \sigma(y)$, then $\text{Im}y = H$, thus if $xg = \vec{0}$ we always have the preimage $y^{-1}g = f$ such that $xyf = 0$, thus $0 \in \sigma(xy)$ and $\Delta(xy) = 1$. Even simpler we obtain that $\Delta(yx) = 1$, thus $\Delta(xy) = \Delta(yx)$.

ii) If x or y is singular (non-invertible), then the inequality is evident.

Let x and y be invertible, then

$$\|xy\| \|(xy)^{-1}\| = \|xy\| \|y^{-1}x^{-1}\| \leq \|x\| \|y\| \|y^{-1}\| \|x^{-1}\|,$$

Assume, that $\Delta(x) \leq \Delta(y)$, then

$$\|x\| \|x^{-1}\| \leq \|y\| \|y^{-1}\|.$$

Now,

$$\begin{aligned} \Delta(xy) &= 1 - \frac{2}{\|xy\| \|y^{-1}x^{-1}\| + 1} \leq 1 - \frac{2}{\|x\| \|y\| \|y^{-1}\| \|x^{-1}\| + 1} \leq \\ &\leq 1 - \frac{2}{\|y\|^2 \|y^{-1}\|^2 + 1} = 1 - \frac{2}{\|y^2\| \|y^{-2}\| + 1} = \Delta(y^2). \end{aligned}$$

We used here the property $\|x^2\| = \|x\|^2$ of a C^* -algebra.

iii) We have

$$\begin{aligned}\Delta(x^2) &= \frac{\|x^2\|\|x^{-2}\| - 1}{\|x^2\|\|x^{-2}\| + 1} = \frac{\|x\|\|x^{-1}\| - 1}{\|x\|\|x^{-1}\| + 1} \times \frac{(\|x\|\|x^{-1}\| + 1)^2}{\|x^2\|\|x^{-2}\| + 1} = \\ &= \Delta(x) \left(1 + \frac{2\|x\|\|x^{-1}\|}{\|x\|^2\|x^{-1}\|^2 + 1} \right) \leq 2\Delta(x).\end{aligned}$$

iv) Just use the inequality

$$\Delta(x\mathbf{1}) \leq \max\{\Delta(x^2), \Delta(\mathbf{1})\} = \Delta(x^2).$$

□

4 The centrality measure

Let \mathcal{N} be a W^* -algebra with the center $\mathfrak{C}(\mathcal{N})$ and with some representation $\pi : \mathcal{N} \mapsto B(H)$.

By $\Delta_{\mathcal{N}} : x \in \mathcal{N}^+ \mapsto [0, 1]$ we denote the following functional

$$\Delta_{\mathcal{N}}(x) = \inf \left\{ \left\| z - \frac{x}{A} \right\| \mid A \in \mathbb{R}^+, z \in \mathfrak{C}(\mathcal{N}), \|z\| = 1 \right\}.$$

The Δ described in the previous section coincides with $\Delta_{B(H)}$. Note, that

$$\Delta_{\mathcal{N}}(x) \leq \Delta_{B(H)}(\pi(x)).$$

Remark 6. If algebra \mathcal{N} is simple, i.e. $\mathfrak{C}(\mathcal{N}) = \mathbb{C} \cdot \mathbf{1}$, then, evidently,

$$\Delta_{\mathcal{N}} \subset \Delta_{B(H)},$$

i.e. for any $x \in \mathcal{N}$ the equality $\Delta_{\mathcal{N}}(x) = \Delta_{B(H)}(\pi(x))$ holds, thus it holds for any W^* -factor.

However, it is easy to see, that the definitions are distinctive.

Example 2. Let $\mathcal{N} = \ell_{\infty}$ be a commutative W^* -algebra. Then $\Delta_{\mathcal{N}} = 0$ even for projections in ℓ_{∞} .

The general properties for $\Delta_{\mathcal{N}}$ are the same.

Proposition 3. *Let \mathcal{N} be a W^* -algebra and $x \in \mathcal{N}^+$, $\lambda \in \mathbb{R}^+$, then*

- 1) $\Delta_{\mathcal{N}}(\lambda x) = \Delta_{\mathcal{N}}(x)$;
- 2) $\Delta_{\mathcal{N}}(x) \leq 1$;
- 3) if x is invertible (non-singular), then $\Delta_{\mathcal{N}}(x) < 1$.

Proof. Statement 1) evidently follows from the definition, statement; 2) follows from the inequality $\Delta_{\mathcal{N}}(x) \leq \Delta_{B(H)}(\pi(x))$ and Proposition 1; 3) also follows from the same inequality and Theorem 1. \square

Theorem 5. *Let $\{\mathcal{N}_i\}_{i \in I}$ be a family W^* -algebras, and \mathcal{M} be its direct sum then for any $\oplus_{i \in I} \mathcal{N}_i$ (i.e. for any $a \in \mathcal{M}$ such that $a = \oplus_{i \in I} a_i$ with $a_i \in \mathcal{N}_i$ we have that $\sup_{i \in I} \{\|a_i\|\} < +\infty$) the equality*

$$\Delta_{\mathcal{M}}(\oplus_{i \in I} x_i) \leq \sup_{i \in I} \{\Delta_{\mathcal{N}_i}(x_i)\}$$

holds for any $a \in \mathcal{M}^+$ ($a_i \in \mathcal{N}_i^+$)

Proof. It is known that $\mathfrak{C}(\oplus_{i \in I} \mathcal{N}_i) = \oplus_{i \in I} \mathfrak{C}(\mathcal{N}_i)$. Also for any particular $x \in \oplus_{i \in I} \mathcal{N}_i$ with the proper decomposition $x = \oplus_{i \in I} x_i$ ($x_i \in \mathcal{N}_i$) we have $\|x\| = \|\oplus_{i \in I} x_i\| = \sup_{i \in I} \{\|x_i\|\}$, thus

$$\begin{aligned} & \inf \left\{ \left\| z - \frac{x}{A} \right\| \mid z \in \mathfrak{C}(\oplus_{i \in I} \mathcal{N}_i), A \in \mathbb{R}^+, \|z\| = 1 \right\} = \\ & = \inf \left\{ \left\| \oplus_{i \in I} z_i - \frac{\oplus_{i \in I} x_i}{A} \right\| \mid z_i \in \mathfrak{C}(\mathcal{N}_i), A \in \mathbb{R}^+, \sup_{i \in I} \|z_i\| = 1 \right\} = \\ & = \inf \left\{ \left\| \oplus_{i \in I} \left(z_i - \frac{x_i}{A} \right) \right\| \mid z_i \in \mathfrak{C}(\mathcal{N}_i), A \in \mathbb{R}^+, \sup_{i \in I} \|z_i\| = 1 \right\} = \\ & = \inf \left\{ \sup_{i \in I} \left\{ \left\| z_i \right\| \left\| \frac{z_i}{\|z_i\|} - \frac{x_i}{A \|z_i\|} \right\| \mid z_i \in \mathfrak{C}(\mathcal{N}_i), A \in \mathbb{R}^+, \sup_{i \in I} \|z_i\| = 1 \right\} \right\} \leq \\ & \leq \sup_{i \in I} \left\{ \inf \left\{ \left\| z_i - \frac{x_i}{A} \right\|, z_i \in \mathfrak{C}(\mathcal{N}_i), A \in \mathbb{R}^+, \|z_i\| = 1 \right\} \right\} = \sup_{i \in I} \Delta_{\mathcal{N}_i}(x_i) \end{aligned}$$

\square

Another definition that we propose and see it as more productive is the following: any countably decomposable von Neumann algebra \mathcal{N} is decomposable into direct sum of factors $\mathcal{N} = \oplus_{i \in I} F_i$.

For $x \in \mathcal{N}^+$ define

$$\Delta'_{\mathcal{N}}(x) := \sup_{i \in I} (\Delta(x_i)),$$

with $x_i \in F_i^+$.

Proposition 4. *Let \mathcal{N} be a W^* -algebra and $x \in \mathcal{N}^+$, $\lambda \in \mathbb{R}^+$, then*

$$1) \Delta'_{\mathcal{N}}(\lambda x) = \Delta'_{\mathcal{N}}(x);$$

$$2) \Delta'_{\mathcal{N}}(x) \leq 1;$$

3) if x is invertible (non-singular), then $\Delta'_{\mathcal{N}}(x) < 1$.

Remark 7. Note, that any projection p_j constructed as $p_j = \oplus_{i \in I} \delta_i^j \mathbf{1}$, where δ_i^j is Kronecker delta, actually is central element, thus Proposition 2 cannot be extended to the case of $\Delta_{\mathcal{N}}$ or $\Delta'_{\mathcal{N}}$.

However we are able to generalize Theorem 4 for $\Delta'_{\mathcal{N}}$ with \mathcal{N} being countably decomposable W^* -algebra

Theorem 6. *Let \mathcal{N} be countably decomposable von Neumann algebra and $x, y \in \mathcal{N}^+$, then*

- 1) $\Delta'_{\mathcal{N}}(xy) = \Delta'_{\mathcal{N}}(yx)$;
- 2) $\Delta'_{\mathcal{N}}(xy) \leq \max(\Delta'_{\mathcal{N}}(x^2), \Delta'_{\mathcal{N}}(y^2))$;
- 3) $\Delta'_{\mathcal{N}}(x^2) \leq 2\Delta'_{\mathcal{N}}(x)$;
- 4) $\Delta'_{\mathcal{N}}(x) \leq \Delta'_{\mathcal{N}}(x^2)$.

Proof. Simply consider two representations $x = \oplus_{i=1}^n x_i$, $y = \oplus_{i=1}^n y_i$, where $x_i, y_i \in F_i^+$, then

- 1)

$$\Delta'_{\mathcal{N}}(xy) = \sup_{i \in I} \Delta(x_i y_i) = \sup_{i \in I} \Delta(y_i x_i) = \Delta'_{\mathcal{N}}(yx);$$
- 2)

$$\begin{aligned} \Delta'_{\mathcal{N}}(xy) &= \sup_{i \in I} \Delta(x_i y_i) \leq \sup_{i \in I} \{\max\{\Delta(x_i^2), \Delta(y_i^2)\}\} = \\ &= \max\{\sup_{i \in I} \Delta(x_i^2), \sup_{i \in I} \Delta(y_i^2)\} = \max\{\Delta'_{\mathcal{N}}(x), \Delta'_{\mathcal{N}}(y)\}; \end{aligned}$$
- 3)

$$\Delta'_{\mathcal{N}}(x^2) = \sup_{i \in I} \Delta(x_i^2) \leq \sup_{i \in I} 2\Delta(x_i) = 2 \sup_{i \in I} \Delta(x_i) = 2\Delta'_{\mathcal{N}}(x);$$
- 4)

$$\Delta'_{\mathcal{N}}(x) = \sup_{i \in I} \Delta(x_i) \leq \sup_{i \in I} \Delta(x_i^2) = \Delta'_{\mathcal{N}}(x^2).$$

□

Also, we can generalize Theorem 3 for $\Delta'_{\mathcal{N}}$.

Theorem 7. *Let \mathcal{M} be countably decomposable von Neumann algebra and $x, y \in \mathcal{M}^+$, then*

$$\Delta'_{\mathcal{M}}(x + y) \leq \max\{\Delta'_{\mathcal{M}}(x), \Delta'_{\mathcal{M}}(y)\}.$$

Proof.

$$\begin{aligned}\Delta'_{\mathcal{N}}(x+y) &= \sup_{i \in I} (\Delta(x_i + y_i)) \leq \sup_{i \in I} (\max\{\Delta(x_i), \Delta(y_i)\}) = \\ &= \max \left\{ \sup_{i \in I} (\Delta(x_i)), \sup_{i \in I} (\Delta(y_i)) \right\} = \max \{ \Delta'_{\mathcal{N}}(x), \Delta'_{\mathcal{N}}(y) \}\end{aligned}$$

□

Basing on the latter Theorem and Proposition 4 for \mathcal{N} we are able to define a cone

$$K_c = \{x \in \mathcal{N}^+ | \Delta'_{\mathcal{N}}(x) \leq c\},$$

which would be a closed subcone with the property, that if $0 \leq c_1 \leq c_2 \leq 1$, then

$$\mathbb{R}^+ \mathbf{1} \subset K_0 \subset K_{c_1} \subset K_{c_2} \subset K_1 = \mathcal{M}^+.$$

5 Applications

Recall, that the Michelson contrast is defined as as

$$\frac{I_{\max} - I_{\min}}{I_{\max} + I_{\min}},$$

with I_{\max} and I_{\min} representing the highest and lowest values of some characteristic of an image, such as luminance, color channel intensity, or, in some cases even a probability. [6, 7, 8]

Assuming that the characteristic is described as a function $f(x, y) \geq 0$ we consider a multiplication operator $M_f : g \in L_2([a, b] \times [c, d]) \mapsto fg \in L_2([a, b] \times [c, d])$ we have that Michelson distance is just equal to $\Delta(M_f)$.

We must make a remark that value of Δ does not really depend on the representation of f or M_f .

5.1 Squared singular values

Let $M \in \mathbb{M}_{n \times m}(\mathbb{R})$, then evidently, $M^*M \in \mathbb{M}_m^+(\mathbb{R})$ and $MM^* \in \mathbb{M}_n^+(\mathbb{R})$. Let us denote

$$\Delta_2(M) = \min(\Delta(M^*M), \Delta(MM^*)) = \begin{cases} \Delta(M^*M) & m \leq n \\ \Delta(MM^*) & m > n \end{cases}.$$

The latter equality holds since $\max(\Delta(M^*M), \Delta(MM^*)) = 1$.

Then the following statement holds:

Corollary 8. Let $M_1, M_2 \in \mathbb{M}_{n \times m}(\mathbb{R})$, then

$$\Delta_2(M_1 + M_2) \leq \max(\Delta_2(M_1), \Delta_2(M_2), \min(\Delta(M_2^* M_1 + M_1^* M_2), \Delta(M_2 M_1^* + M_1 M_2^*)))$$

Proof. From lemma 3 and the definition we get either

$$\begin{aligned} \Delta((M_1 + M_2)^*(M_1 + M_2)) &= \Delta(M_1^* M_1 + M_2^* M_1 + M_1^* M_2 + M_2^* M_2) \leq \\ &\leq \max(\Delta(M_1^* M_1), \Delta(M_2^* M_2), \Delta(M_2^* M_1 + M_1^* M_2)) \end{aligned}$$

or

$$\begin{aligned} \Delta((M_1 + M_2)(M_1 + M_2)^*) &= \Delta(M_1 M_1^* + M_2 M_1^* + M_1 M_2^* + M_2 M_2^*) \leq \\ &\leq \max(\Delta(M_1 M_1^*), \Delta(M_2 M_2^*), \Delta(M_2 M_1^* + M_1 M_2^*)). \end{aligned}$$

□

In particular, for any orthogonal U_1 and U_2 we have

$$\Delta_2(U_1 + U_2) = \min(\Delta(U_2 U_1^* + U_1 U_2^*), \Delta(U_2^* U_1 + U_1^* U_2))$$

5.2 Multichannel images

Let M be a three dimensional array $c \times n \times m$. Then, interpreting M as an image with c channels we may consider a direct sum $\oplus_c M_c$ with $M_c \in \mathbb{M}_{n \times m}(\mathbb{R})$, thus consider $\Delta'_2(\oplus_c M_c) := \sup_c \Delta_2(M_c)$, then

Corollary 9. Let $M, N \in \mathbb{T}_{c \times n \times m}(\mathbb{R})$, then

$$\begin{aligned} \Delta'_2(M + N) &\leq \max(\Delta'_2(M), \Delta'_2(N), \\ &\min(\Delta'(N^* M + M^* N), \Delta(N M^* + M N^*))) \end{aligned}$$

Proof. $\Delta'_2(M + N) = \sup_c \Delta_2((M + N)_c) =$

$$\begin{aligned} &= \sup_c \min(\Delta((M + N)_c^* (M + N)_c), \Delta((M + N)_c (M + N)_c^*)) = \\ &= \sup_c \min(\Delta((M_c^* + N_c^*)(M_c + N_c)), \Delta((M_c + N_c)(M_c + N_c)^*)) \leq \\ &\leq \sup_c \max\{\Delta_2(M_c), \Delta_2(N_c), \min(\Delta(M_c^* N_c + N_c^* M_c), \Delta(M_c N_c^* + N_c M_c^*))\} = \\ &= \max\left\{\sup_c \left(\Delta_2(M_c), \sup_c \Delta_2(N_c), \sup_c \min(\Delta(M_c^* N_c + N_c^* M_c), \Delta(M_c N_c^* + N_c M_c^*))\right)\right\} \leq \\ &\leq \max\left\{\sup_c \Delta_2(M_c), \sup_c \Delta_2(N_c), \min(\sup_c \Delta(M_c^* N_c + N_c^* M_c), \sup_c \Delta(M_c N_c^* + N_c M_c^*))\right\} = \\ &= \max\{\Delta'_2(\oplus_c M_c), \Delta'_2(\oplus_c N_c), \min(\Delta'(\oplus_c (M_c^* N_c + N_c^* M_c)), \Delta'(\oplus_c (M_c N_c^* + N_c M_c^*)))\} \end{aligned}$$

□

Acknowledgments

The work is supported by the Mathematical Center in Akademgorodok under agreement No.075-15-2022-281 (05.04.2022) with the Ministry of Science and Higher Education of the Russian Federation.

References

- [1] A.M. Bikchentaev “Characterization of traces on C^* -algebras: a survey”, Gaspar, D. (ed.) et al., An operator theory summer. Proc. 23rd Internat. Conf. Operator Theory. Timisoara, Romania, June 29 July 4, 2010. Bucharest: The Theta Foundation. Theta Series in Advanced Math. **13**, 1–12 (2012).
- [2] J. Dixmier, *Les C^* -algebras et leurs representations* (Gauthier-Villars Editeur, Paris, 1969).
- [3] A. Novikov, “ L_1 -space for a positive operator affiliated with von Neumann algebra”, *Positivity* **21** (1) 359–375 (2017).
- [4] A. Novikov, “ C^* -algebra Positive Element Invertibility Criteria in Terms of L_1 -norms Equivalence and L_∞ -norms Equivalence”, *Lobachevskii Journal of Mathematics* **40** (5) 549–552 (2019),
- [5] A. Novikov, Z. Eskandarian, Z. Kholmatova “Limits of the Banach spaces associated with positive operator a affiliated with von Neumann algebra, which are neither purely projective nor purely inductive”, 2020 – <https://arxiv.org/abs/1901.09276>,
- [6] A. Michelson, *Studies in Optics*, U. of Chicago Press, 176 p. (1927).
- [7] H. Kukkonen, J. Rovamo, K. Tiippa, R. Näsänen, “Michelson contrast, RMS contrast and energy of various spatial stimuli at threshold”, *Vision Research* – 33(10) – 1431-1436 (1993),
- [8] J. L. Prince, *J. M. Links Medical Imaging Signals and Systems*, Pearson, 2006, 519p.
- [9] D. Oliveira, E. Giusto, B. Baheri, Q. Guan, B. Montrucchio, P. Rech “A Systematic Methodology to Compute the Quantum Vulnerability Factors for Quantum Circuits”, 2021 – <https://arxiv.org/abs/2111.07085>;

- [10] A.M. Bikchentaev, S.A. Abed, “Projections and Traces on von Neumann Algebras”, Lobachevskii Journal of Mathematics – 40 (9) – P. 1260–1267 (2019).