

Generalization of the theorems of Barndorff-Nielsen and Balakrishnan-Stepanov to Riesz spaces *

Nyasha Mushambi, Bruce A. Watson †, Bertin Zinsou
 School of Mathematics
 University of the Witwatersrand
 Private Bag 3, P O WITS 2050, South Africa

December 28, 2021

Abstract

In a Dedekind complete Riesz space, E , we show that if (P_n) is a sequence of band projections in E then

$$\limsup_{n \rightarrow \infty} P_n - \liminf_{n \rightarrow \infty} P_n = \limsup_{n \rightarrow \infty} P_n(I - P_{n+1}).$$

This identity is used to obtain conditional extensions in a Dedekind complete Riesz spaces with weak order unit and conditional expectation operator of the Barndorff-Nielsen and Balakrishnan-Stepanov generalizations of the First Borel-Cantelli Theorem.

1 Introduction

In [4, Lemma 1*] Barndorff-Nielsen gave the following generalization of the First Borel-Cantelli Theorem.

Theorem 1.1 (Barndorff-Nielsen). *Let (Ω, \mathcal{F}, P) be a probability space. If (A_n) is a sequence of events in the probability space (Ω, \mathcal{F}, P) such that $\lim_{n \rightarrow \infty} P(A_n) = 0$ and*

$$\sum_{n=1}^{\infty} P(A_n \cap A_{n+1}^c) < \infty,$$

then $P(\limsup_{n \rightarrow \infty} A_n) = 0$.

***Keywords:** Riesz spaces, vector lattices; zero-one laws, Borel-Cantelli theorem, conditional expectation operators. *Mathematics subject classification (2010):* 46B40; 60F15; 60F25.

†Supported in part by the Centre for Applicable Analysis and Number Theory and by NRF grant number IFR2011032400120.

In [5, page 51], it was shown that the assumption $\lim_{n \rightarrow \infty} P(A_n) \rightarrow 0$ could be weakened to $\liminf_{n \rightarrow \infty} P(A_n) = 0$. In [3, Lemma 3], see also [5, page 52], Balakrishnan and Stepanov further extended the Barndorff-Nielsen Theorem, as below.

Theorem 1.2 (Balakrishnan-Stepanov). *Let (Ω, \mathcal{F}, P) be a probability space and (A_n) be a sequence of events in (Ω, \mathcal{F}, P) . If $P(A_n) \rightarrow 0$ as $n \rightarrow \infty$ and*

$$\sum_{n=1}^{\infty} P(A_n^c \cap A_{n+1}^c \cap \dots \cap A_{n+m-1}^c \cap A_{n+m}) < \infty$$

for some $m \geq 1$, then $P(A_n \text{ infinitely often}) = 0$.

We show, see Theorem 3.3, that in a Dedekind complete Riesz space, E , if (P_n) is a sequence of band projections, in E , then

$$\limsup_{n \rightarrow \infty} P_n - \liminf_{n \rightarrow \infty} P_n = \limsup_{n \rightarrow \infty} P_n(I - P_{n+1}).$$

Let E be a Dedekind Riesz space with weak order unit. A conditional expectation operator on E , as defined in [7], is a positive order continuous projection T on E with range, $R(T)$, a Dedekind complete Riesz subspace of E , and having $T(e)$ a weak order unit of E for each weak order unit e of E . We refer the reader to [7] for background on conditional expectation operators on Riesz spaces and [8] for stochastic processes in Riesz spaces with recent developments in [1] and [6].

In the setting of a Dedekind complete Riesz spaces with weak order unit and conditional expectation operator we use the above band projection identity to give conditional generalizations of the theorems Barndorff-Nielsen and Balakrishnan-Stepanov to Riesz spaces.

We thank the referees for their comments and corrections.

2 Preliminaries

We refer the reader to [2, 10, 12] for general background on Riesz spaces and band projections. From [11], or [12, Theorem 12.7], we have that if $(f_n)_{n \geq 1}$ is an order bounded sequence in the Dedekind complete Riesz space E , then the sequence converges if and only if $\liminf_{n \rightarrow \infty} f_n = \limsup_{n \rightarrow \infty} f_n$ in which case this common value is the order limit of (f_n) as $n \rightarrow \infty$ and is denoted $\lim_{n \rightarrow \infty} f_n$.

For a sequence (P_n) of band projections in a Dedekind complete Riesz space, E , it is easy to verified that $\prod_{n \in \mathbb{N}} P_n = \bigwedge_{n \in \mathbb{N}} P_n$. Further for a sequence (f_n) in the positive cone,

E_+ , of such a Riesz space E , if (f_n) is order bounded and $f_n \wedge f_m = 0$ for all $n \neq m$ then $\sum_{n \in \mathbb{N}} f_n = \bigvee_{n \in \mathbb{N}} f_n$. This is the order limit of the result in [12, page 110].

The Fatou property (2.1) holds for any positive order continuous operator, T , on a Dedekind complete Riesz space, however, our interest here is only with regards to conditional expectation operators. It is well known, but included here for completeness.

Theorem 2.1 (Fatou). *Let E be a Dedekind complete Riesz space with conditional expectation operator T with weak order unit $e = Te$. If (f_n) is a bounded sequence, in E , then*

$$T(\liminf_{n \rightarrow \infty} f_n) \leq \liminf_{n \rightarrow \infty} T f_n. \quad (2.1)$$

Proof. Let $f \in E$, denote the limit inferior of f_n . For every $k \in \mathbb{N}$, define $g_k = \inf_{n \geq k} f_n$. Then the sequence (g_k) is increasing and converges to f . For $k \leq n$, we have $g_k \leq f_n$ so, as T is a positive linear operator, $Tg_k \leq T f_n$. Now taking the infimum over $n \geq k$, we have $Tg_k \leq \inf_{n \geq k} T f_n$. We know $g_k \uparrow$ with order limit f , so by the order continuity of T we have, $\lim_{k \rightarrow \infty} Tg_k = T f$. Also $\inf_{n \geq k} T f_n \uparrow \liminf_{n \rightarrow \infty} T f_n$ so taking $k \rightarrow \infty$, we have $T f \leq \liminf_{n \rightarrow \infty} T f_n$. \square

If $f_n \leq g$, for all $n \geq 1$, then applying Theorem 2.1 to $(g - f_n)$ gives following Reverse Fatou inequality.

Corollary 2.2. *Let E be a Dedekind complete Riesz space with conditional expectation operator T with weak order unit $e = Te$. If $(f_n) \subset E^+$ and $g \in E^+$ with $f_n \leq g$ for all $n \geq 1$, then*

$$T(\limsup_{n \rightarrow \infty} f_n) \geq \limsup_{n \rightarrow \infty} T f_n.$$

3 Sequences of Band projections

We present two representation lemmas for sequences of band projections in Lemmas 3.1 and 3.2. The first of these will be used in the characterization of the difference between the superior and inferior limits of a sequence of band projections in Theorem 3.3 while the latter will be used in the proof of the Balakrishnan-Stepanov Theorem, Theorem 4.3.

Lemma 3.1. *Let E be a Dedekind complete Riesz space. If (P_n) is a sequence of band projections on E , then*

$$P_n = Q + \sum_{i=1}^m Q_i, \quad (3.1)$$

where the band projections $Q = \prod_{k=n}^{n+m} P_k$ and $Q_i = \prod_{k=0}^{i-1} P_{n+k}(I - P_{n+i})$ have $QQ_i = 0 = Q_iQ_j$ for $i \neq j$.

Proof. We observe that

$$\sum_{i=1}^m \prod_{k=0}^{i-1} P_{n+k}(I - P_{n+i}) = \sum_{i=1}^m \left(\prod_{k=0}^{i-1} P_{n+k} - \prod_{k=0}^i P_{n+k} \right) = P_n - \prod_{k=n}^{n+m} P_k$$

hence proving 3.1. Now, for $1 \leq i \leq m$, we have

$$0 \leq QQ_i = \prod_{k=n}^{n+m} P_k \prod_{r=0}^{i-1} P_{n+r}(I - P_{n+i}) \leq P_{n+i} \prod_{r=0}^{i-1} P_{n+r}(I - P_{n+i}) = 0$$

and

$$0 \leq Q_jQ_i = \prod_{r=0}^{j-1} P_{n+r}(I - P_{n+j}) \prod_{k=0}^{i-1} P_{n+k}(I - P_{n+i}) \leq P_{n+i}(I - P_{n+i}) = 0,$$

for $1 \leq i < j \leq m$. □

Lemma 3.2. *Let E be a Dedekind complete Riesz space with weak order unit e . If (P_n) is a sequence of band projections acting on E then,*

$$\bigvee_{n=k}^N P_n = P_k \vee \bigvee_{n=k+1}^N (P_n \prod_{j=k}^{n-1} (I - P_j)). \quad (3.2)$$

Proof. We observe the notational convention $\prod_{j=k}^{k-1} (I - P_j) = I$. Setting $Q_n := P_n \prod_{j=k}^{n-1} (I - P_j)$

$P_j) = \prod_{j=k}^{n-1} (I - P_j) - \prod_{j=k}^n (I - P_j)$ we have that $Q_nQ_m = 0$ for $n \neq m$ and thus

$$\bigvee_{n=k}^N P_n \prod_{j=k}^{n-1} (I - P_j) = \sum_{n=k}^N \left(\prod_{j=k}^{n-1} (I - P_j) - \prod_{j=k}^n (I - P_j) \right) = I - \prod_{j=k}^N (I - P_j). \quad (3.3)$$

Here

$$\prod_{j=k}^N (I - P_j) = \bigwedge_{j=k}^N (I - P_j) = I - \bigvee_{j=k}^N P_j. \quad (3.4)$$

Combining (3.3) and (3.4) gives the required equality. □

Theorem 3.3. *Let E be a Dedekind complete Riesz space. For (P_n) a sequence of band projections in E ,*

$$\limsup_{n \rightarrow \infty} P_n - \liminf_{n \rightarrow \infty} P_n = \limsup_{n \rightarrow \infty} P_n(I - P_{n+1}). \quad (3.5)$$

Proof. Since $P_n(I - P_{n+1}) \leq P_n$ and $P_n(I - P_{n+1}) \leq I - P_{n+1}$, we have

$$\limsup_{n \rightarrow \infty} P_n(I - P_{n+1}) \leq \limsup_{n \rightarrow \infty} P_n$$

and

$$\limsup_{n \rightarrow \infty} P_n(I - P_{n+1}) \leq I - \liminf_{n \rightarrow \infty} P_n.$$

Thus, since $\liminf_{n \rightarrow \infty} P_n \leq \limsup_{n \rightarrow \infty} P_n$,

$$\limsup_{n \rightarrow \infty} P_n(I - P_{n+1}) \leq \limsup_{n \rightarrow \infty} P_n \wedge (I - \liminf_{n \rightarrow \infty} P_n) = \limsup_{n \rightarrow \infty} P_n - \liminf_{n \rightarrow \infty} P_n. \quad (3.6)$$

Taking the limit as $m \rightarrow \infty$ in Lemma 3.1, we have

$$P_n = \bigwedge_{k \geq n} P_k + \bigvee_{i \geq 1} \bigwedge_{k=0}^{i-1} P_{n+k}(I - P_{n+i}) \leq \bigwedge_{k \geq n} P_k + \bigvee_{i \geq 1} P_{n+i-1}(I - P_{n+i}).$$

Hence,

$$P_n \leq \bigwedge_{k \geq n} P_k + \bigvee_{i \geq n} P_i(I - P_{i+1}). \quad (3.7)$$

Taking the $\limsup_{n \rightarrow \infty}$ of both sides to (3.7) gives

$$\begin{aligned} \limsup_{n \rightarrow \infty} P_n &\leq \bigwedge_{r \geq 1} \bigvee_{n \geq r} \bigwedge_{k \geq n} P_k + \bigwedge_{r \geq 1} \bigvee_{n \geq r} \bigvee_{i \geq n} P_i(I - P_{i+1}) \\ &= \bigwedge_{r \geq 1} \liminf_{k \rightarrow \infty} P_k + \bigwedge_{r \geq 1} \bigvee_{i \geq r} P_i(I - P_{i+1}) \\ &= \liminf_{k \rightarrow \infty} P_k + \limsup_{i \rightarrow \infty} P_i(I - P_{i+1}). \end{aligned}$$

Since $\limsup_{n \rightarrow \infty} P_n \cdot \liminf_{k \rightarrow \infty} P_k = \liminf_{k \rightarrow \infty} P_k$, applying $I - \liminf_{n \rightarrow \infty} P_n$ to both sides of the above gives

$$\limsup_{n \rightarrow \infty} P_n - \liminf_{k \rightarrow \infty} P_k \leq \limsup_{i \rightarrow \infty} P_i(I - P_{i+1}). \quad (3.8)$$

Combining (3.6) and (3.8) concludes the proof. \square

4 Borel-Cantelli type theorems in Riesz spaces

In [9, Corollary 4.2] a generalization of the First Borel-Cantelli Lemma to Riesz spaces was given and it is quoted below. Following this we generalize the Barndorff-Nielsen Theorem, in Theorem 4.2, and the Balakrishnan-Stepanov Theorem, in Theorem 4.3, to Dedekind complete Riesz spaces with weak order unit and conditional expectation operator.

Corollary 4.1 (Borel-Cantelli Theorem in Riesz spaces). *Let E be a Dedekind complete Riesz space with strictly positive conditional expectation operator T and weak order unit $e = Te$. Let (P_j) be a sequence of band projections in E . If $\sum_{j=1}^{\infty} T(P_j e)$ is order convergent in E , then $\limsup_{j \rightarrow \infty} P_j = 0$.*

Theorem 4.2 (Barndorff-Nielsen Theorem in Riesz spaces). *Let (P_n) be a sequence of band projections on a Dedekind complete Riesz space E with strictly positive conditional expectation operator T and weak order unit $e = Te$ such that $\liminf_{n \rightarrow \infty} T(P_n e) = 0$ and $\sum_{n=1}^{\infty} T(P_n(I - P_{n+1})e)$ is convergent in E , then $\limsup_{n \rightarrow \infty} P_n = 0$.*

Proof. Let

$$Q_n = P_n(I - P_{n+1}), n \geq 1. \quad (4.1)$$

Applying Corollary 4.1 to (Q_n) , as $\sum_{n=1}^{\infty} TQ_n e$ converges in E then $\limsup_{n \rightarrow \infty} Q_n = 0$. Thus, $\limsup_{n \rightarrow \infty} P_n(I - P_{n+1}) = 0$, so, by Theorem 2.1,

$$\liminf_{n \rightarrow \infty} P_n = \limsup_{n \rightarrow \infty} P_n = \lim_{n \rightarrow \infty} P_n.$$

Hence, as T is order continuous,

$$0 = \liminf_{n \rightarrow \infty} T(P_n e) = \lim_{n \rightarrow \infty} TP_n e = T(\lim_{n \rightarrow \infty} P_n e) = T(\liminf_{n \rightarrow \infty} P_n e).$$

Now, since T is a strictly positive, we have that $\liminf_{n \rightarrow \infty} P_n e = 0$. □

Theorem 4.3 (Balakrishnan-Stepanov Theorem in Riesz spaces). *Let E be a Dedekind complete Riesz space with T a strictly positive conditional expectation operator on E and weak order unit $e = Te$. Let (P_n) be a sequence of band projections on E with $T(P_n e) \rightarrow 0$ in order, as $n \rightarrow \infty$, and*

$$\sum_{n=1}^{\infty} T\left(P_{n+m} \prod_{j=n}^{n+m-1} (I - P_j)e\right) \quad (4.2)$$

order convergent in E for some $m \geq 1$. Then $\limsup_{n \rightarrow \infty} P_n = 0$.

Proof. From definition of the limit supremum we have for each $k \in \mathbb{N}$,

$$T\left(\limsup_{n \rightarrow \infty} P_n e\right) = T\left(\bigwedge_{j \in \mathbb{N}} \bigvee_{n \geq j} P_n e\right) \leq T\left(\bigvee_{n \geq k} P_n e\right). \quad (4.3)$$

Letting $N \rightarrow \infty$ in Lemma 3.2, we get

$$\bigvee_{n \geq k} P_n e = \bigvee_{n \geq k} P_n \prod_{j=k}^{n-1} (I - P_j) e. \quad (4.4)$$

Combining (4.3) and (4.4), we obtain

$$T \left(\limsup_{n \rightarrow \infty} P_n e \right) \leq T \left(\bigvee_{n \geq k} P_n e \right) = T \left(\bigvee_{n \geq k} P_n \prod_{j=k}^{n-1} (I - P_j) e \right). \quad (4.5)$$

However, as shown in Lemma 3.2, for $n \neq m$,

$$(P_n \prod_{j=k}^{n-1} (I - P_j)) (P_m \prod_{j=k}^{m-1} (I - P_j)) = 0,$$

thus,

$$\bigvee_{n \geq k} P_n \prod_{j=k}^{n-1} (I - P_j) = \sum_{n \geq k} P_n \prod_{j=k}^{n-1} (I - P_j). \quad (4.6)$$

Combining (4.5) and (4.6) gives, for $k \in \mathbb{N}$,

$$T \left(\limsup_{n \rightarrow \infty} P_n e \right) \leq \sum_{n \geq k} T P_n \prod_{j=k}^{n-1} (I - P_j) e. \quad (4.7)$$

Now for $k \geq m + 1$ we have

$$\sum_{n \geq k} T P_n \prod_{j=k}^{n-1} (I - P_j) e = \sum_{n=k}^{k+m-1} T P_n \prod_{j=k}^{n-1} (I - P_j) e + \sum_{n \geq k} T P_{n+m} \prod_{j=k}^{n+m-1} (I - P_j) e. \quad (4.8)$$

For $n \geq k$ we have $\prod_{j=k}^{n+m-1} (I - P_j) \leq \prod_{j=n}^{n+m-1} (I - P_j)$ giving

$$\sum_{n \geq k} T P_{n+m} \prod_{j=k}^{n+m-1} (I - P_j) e \leq \sum_{n \geq k} T P_{n+m} \prod_{j=n}^{n+m-1} (I - P_j) \quad (4.9)$$

which tends to zero in order as $k \rightarrow \infty$ since (4.2) is order convergent in E . Further

$$\sum_{n=k}^{k+m-1} T P_n \prod_{j=k}^{n-1} (I - P_j) e \leq \sum_{n=k}^{k+m-1} (T P_n e) \quad (4.10)$$

which tends to zero in E as $k \rightarrow \infty$ as $T P_n e \rightarrow 0$ in order as $n \rightarrow \infty$. \square

We note that (4.2) and (4.5) with Theorem 3.3 give that $\liminf_{n \rightarrow \infty} P_n = 0$. Thus the assumption $\lim_{n \rightarrow \infty} T P_n e = 0$, as used in Theorem 4.3, cannot be replaced by $\liminf_{n \rightarrow \infty} T P_n e = 0$.

References

- [1] Y. AZOUZI, K. RAMDANE, Burkholder Inequalities in Riesz spaces, *Indag. Math.*, **28** (2017), 1076-1094.
- [2] C. D. ALIPRANTIS, K. C. BORDER, *Infinite dimensional analysis: A hitchhiker's guide*. Springer-Verlag, (1994).
- [3] N. BALAKRISHNAN, A. STEPANOV, A generalization of the Borel-Cantelli Lemma. *Math. Scientist*, **35** (2010), 61-61.
- [4] O. BARNDORFF-NIELSEN, On the rate of growth of the maxima of a sequence of independent identically distributed random variables. *Math. Scand.*, **9** (1961), 383-394.
- [5] T.K. CHANDRA, *The Borel-Cantelli Lemma.*, Springer, (2012).
- [6] J.J. GROBLER, C.C.A. LABUSCHAGNE, Girsanov's theorem in vector lattices. *Positivity*, to appear, DOI: <https://doi.org/10.1007/S11117-019-00649-5>
- [7] W.-C. KUO, C.C.A. LABUSCHAGNE, B.A. WATSON, Discrete time stochastic processes in Riesz spaces. *Indag. Math.*, **15** (2004), 15-40.
- [8] W.-C. KUO, C.C.A. LABUSCHAGNE, B.A. WATSON, Conditional Expectation on Riesz spaces. *J. Math. Anal. Appl.*, **303** (2005), 509-521.
- [9] W.-C. KUO, C.C.A. LABUSCHAGNE, B.A. WATSON, Zero-one laws for Riesz space and fuzzy random variables, *Fuzzy Logic, Soft Computing and Computational Intelligence*, Springer Verlag and Tsinghua University Press, Beijing, China, 2005, pages 393-397.
- [10] W.A.J. LUXEMBURG, A.C. ZAAANEN, *Riesz spaces I*. North Holland, (1971).
- [11] P. MEYER-NIEBERG, *Banach Lattices*, Springer Verlag, 1991.
- [12] A.C. ZAAANEN, *Introduction to Operator Theory in Riesz Spaces*. Springer-Verlag, (1997).