

# BALMER SPECTRA AND DRINFELD CENTERS

KENT B. VASHAW

**ABSTRACT.** The Balmer spectrum of a monoidal triangulated category is an important geometric construction which is closely related to the problem of classifying thick tensor ideals. We prove that the forgetful functor from the Drinfeld center of a finite tensor category  $\mathbf{C}$  to  $\mathbf{C}$  extends to a monoidal triangulated functor between their corresponding stable categories, and induces a continuous map between their Balmer spectra. In the finite-dimensional Hopf algebra setting, we give conditions under which it is injective, surjective, or a homeomorphism. We apply this general theory to prove that Balmer spectra associated to finite-dimensional cosemisimple quasitriangular Hopf algebras (in particular, group algebras in characteristic dividing the order of the group) coincide with the Balmer spectra associated to their Drinfeld doubles, and that the thick ideals of both categories are in bijection. An analogous theorem is proven for certain Benson-Witherspoon smash coproduct Hopf algebras, which are not quasitriangular in general.

## INTRODUCTION

Balmer initiated the study of tensor triangular geometry in [4] and [5]. This theory has found a wide variety of applications, for instance in modular representation theory, algebraic geometry, and algebraic topology, which are all studied through the prism of tensor triangulated categories [14, 15, 41, 6, 7]. Balmer’s original theory is a commutative geometry, in the sense that the tensor categories studied are required to have a braiding

$$X \otimes Y \cong Y \otimes X$$

for all objects. Since many interesting monoidal triangulated categories in representation theory are not necessarily braided, a noncommutative analogue of Balmer’s theory was initiated and explored in [16, 44, 45], which constructs Balmer spectra for monoidal triangulated categories without a braiding on their monoidal product. The Balmer spectrum of such a category  $\mathbf{T}$  is denoted  $\mathrm{Spc} \mathbf{T}$ ; it is defined as the collection of prime ideals of  $\mathbf{T}$ , reflecting the usual notion of prime spectrum from ring theory.

Non-braided monoidal triangulated categories arise naturally as the stable categories of finite tensor categories. Broadly speaking, if  $\mathbf{C}$  is a finite tensor category, then the stable category of  $\mathbf{C}$ , denoted  $\mathrm{st}(\mathbf{C})$ , is the category obtained by factoring out the projective objects of  $\mathbf{C}$ . One motivation for factoring out projectives comes from the theory of support varieties, where the support variety of an object only distinguishes an object up to direct sums with projective objects. The stable category is a triangulated category, and the monoidal product of  $\mathbf{C}$  extends to  $\mathrm{st}(\mathbf{C})$ ; in this way,  $\mathrm{st}(\mathbf{C})$  becomes a monoidal triangulated category.

An important tool in the study of tensor categories is the Drinfeld center, a categorical analogue of the center of a ring; it is a generalization of the quantum or Drinfeld double

---

2020 *Mathematics Subject Classification.* 18G80, 18M05, 16T05 .

*Key words and phrases.* Hopf algebra, stable module category, tensor triangulated category, thick ideal.

Research of K.B.V. was supported in part by a Board of Regents LSU fellowship, an Arthur K. Barton Superior Graduate Student Scholarship in Mathematics from LSU, NSF grant DMS-1901830, and NSF Postdoctoral Fellowship DMS-2103272.

construction for Hopf algebras, originally introduced by Drinfeld in [21]. For any tensor category  $\mathbf{C}$ , its Drinfeld center  $Z(\mathbf{C})$  is a braided tensor category equipped with a functor  $F : Z(\mathbf{C}) \rightarrow \mathbf{C}$ . The Drinfeld center satisfies the universal property: if  $G : \mathbf{D} \rightarrow \mathbf{C}$  is a strict tensor functor between strict tensor categories, and  $\mathbf{D}$  is braided, such that  $G$  is bijective on objects and surjective on morphisms, then there exists a strict tensor functor  $H : \mathbf{D} \rightarrow Z(\mathbf{C})$

$$\begin{array}{ccc} \mathbf{C} & & \\ \uparrow F & \swarrow G & \\ Z(\mathbf{C}) & \xleftarrow{H} & \mathbf{D} \end{array}$$

with  $F \circ H = G$ .

If  $\mathbf{C}$  is abelian, then  $Z(\mathbf{C})$  is automatically abelian as well. We do not see an analogue for this argument in the triangulated case: if  $\mathbf{C}$  is triangulated, it does not seem to follow immediately that  $Z(\mathbf{C})$  is triangulated. This is a reflection of the fact that the morphism given in the extension axiom for triangulated categories is not necessarily unique.

However, if  $\mathbf{C}$  is a finite tensor category, one can form its stable category  $\text{st}(\mathbf{C})$  on one hand; on the other hand,  $Z(\mathbf{C})$  is again a finite tensor category, and one can form its stable category  $\text{st}(Z(\mathbf{C}))$ . The natural question that arises is, therefore: how are the Balmer spectra between these two categories connected?

This question is of particular interest because Balmer spectra are related intimately with cohomological support varieties (as in [13]); for example, the projectivization of the spectrum of the cohomology ring of the small quantum groups  $u_\zeta(\mathfrak{b})$  of Borel subalgebras at roots of unity (as computed in [30, 8]) identifies with the Balmer spectrum of its stable category [44], which can be used to show that the support varieties for the small quantum Borel possess the tensor product property [49, 45]. In many specific cases, for instance see [25, 48, 50], the cohomology of Drinfeld doubles has been studied, and its relationship to the cohomology of the original finite tensor category explored.

Additionally, this project will provide tools to aid in thick ideal classification problems. Balmer spectra, which are defined as the collection of prime ideals of the category, are intimately related to these problems, since every thick ideal of a rigid monoidal triangulated category is equal to an intersection of prime ideals. Classifications of thick ideals in various settings have been undertaken in many different settings, for instance in various categories arising from

- (1) commutative algebra and algebraic geometry [34, 56, 41];
- (2) Lie superalgebras [14];
- (3) finite groups and finite group schemes [9, 26];
- (4) tilting modules for quantum groups and algebraic groups in positive characteristic [52, 1];
- (5) Hopf algebras which are not necessarily commutative, cocommutative, or even quasitriangular [12, 14, 44, 45].

There are examples, for instance the small quantum groups of Borel subalgebras  $u_\zeta(\mathfrak{b})$  at roots of unity, where, as mentioned above, the Balmer spectrum and thick ideals are known for its stable module category; however, it is an open question to classify the Balmer spectrum and thick ideals for the stable category of its Drinfeld center, that is, the stable module category of  $u_\zeta(\mathfrak{g}) \otimes \mathbb{k}T$ , where  $\mathbb{k}T$  is the group algebra of the group of generators  $K_i$  for  $u_\zeta(\mathfrak{g})$ . This motivates our central question, to reiterate: what relationship exists between the Balmer spectra of  $\text{st}(\mathbf{C})$  and  $\text{st}(Z(\mathbf{C}))$ ?

We answer this question by the following approach.

In Section 1, we give a brief background on tensor triangular geometry, support data, stable module categories, and Drinfeld centers, and establish notation.

Next, in Section 2, we consider directly the relationship between the Balmer spectra  $\mathrm{Spcst}(\mathbf{C})$  and  $\mathrm{Spcst}(Z(\mathbf{C}))$ . Since the prime ideals of the Balmer spectrum of a non-braided monoidal triangulated category are a categorical analogue of the prime ideals in a noncommutative ring, we are motivated by prime ideal contraction, that is, the statement that if  $\mathfrak{p}$  is a prime ideal of a noncommutative ring  $R$ , then  $\mathfrak{p} \cap Z(R)$  is a prime ideal in  $Z(R)$ , the center of  $R$ . For general background on prime ideals for noncommutative rings, see [31, Chapter 3]. Finding a categorical analogue to prime ideal contraction is complicated by the fact that we work with  $\mathrm{st}(Z(\mathbf{C}))$  rather than  $Z(\mathrm{st}(\mathbf{C}))$ ; the latter is equipped with a forgetful functor  $Z(\mathrm{st}(\mathbf{C})) \rightarrow \mathrm{st}(\mathbf{C})$ , but, as noted above, is not necessarily triangulated. Nevertheless, we verify that the forgetful functor  $F : Z(\mathbf{C}) \rightarrow \mathbf{C}$  extends to a functor  $\overline{F} : \mathrm{st}(Z(\mathbf{C})) \rightarrow \mathrm{st}(\mathbf{C})$ .

Reflecting the analogous property for rings, Balmer spectra of braided monoidal triangulated categories are functorial; but in the non-braided situation, a monoidal triangulated functor does not necessarily induce a continuous map between Balmer spectra. However, we show that  $\overline{F}$  does induce a continuous map, and we obtain an analogue of prime ideal contraction. This is summarized by the following:

**Theorem.** (See Proposition 2.1.1 and Proposition 2.1.2). *Let  $\mathbf{C}$  be a finite tensor category. There exists a monoidal triangulated functor  $\overline{F} : \mathrm{st}(Z(\mathbf{C})) \rightarrow \mathrm{st}(\mathbf{C})$  extending the forgetful functor  $F : Z(\mathbf{C}) \rightarrow \mathrm{st}(\mathbf{C})$ , which induces a continuous map  $f : \mathrm{Spcst}(\mathbf{C}) \rightarrow \mathrm{Spcst}(Z(\mathbf{C}))$ , defined by*

$$f : \mathbf{P} \mapsto \{X : \overline{F}(X) \in \mathbf{P}\},$$

for  $\mathbf{P}$  a thick prime ideal of  $\mathrm{st}(\mathbf{C})$ .

In the case that  $\mathbf{C} = \mathrm{mod}(H)$ , the category of finite-dimensional modules for a finite-dimensional Hopf algebra  $H$ , we are able to utilize the machinery of localization and colocalization functors to better describe the image of this map. The reason for moving this situation is that we have access to the “big stable module categories”  $\mathrm{St}(\mathbf{C})$  and  $\mathrm{St}(Z(\mathbf{C}))$ , defined as the stable categories of all (in other words, not necessarily finite-dimensional) modules of  $H$  and its Drinfeld double  $D(H)$ , respectively. Under certain conditions on the kernel  $\mathbf{K}$  of  $\overline{F}$ , which consists of  $D(H)$ -modules  $X$  such that  $\overline{F}(X)$  is isomorphic to 0 in  $\mathrm{St}(\mathbf{C})$ , we determine the image of  $f$ .

**Theorem.** (See Proposition 2.4.1). *Let  $H$  be a finite-dimensional Hopf algebra,  $\overline{F}$  the functor induced by the forgetful functor as in the previous theorem,  $\mathbf{K}$  the kernel of  $\overline{F}$ , and  $f : \mathrm{Spcst}(\mathbf{C}) \rightarrow \mathrm{Spcst}(Z(\mathbf{C}))$  the continuous map induced by  $\overline{F}$ . Then there are containments*

$$\begin{aligned} \{\mathbf{P} \in \mathrm{Spcstmod}(D(H)) : \mathbf{P} \supseteq \mathbf{K} \cap \mathrm{st}(Z(\mathbf{C}))\} &\supseteq \mathrm{im} f \\ &\supseteq \{\mathbf{P} \in \mathrm{Spcstmod}(D(H)) : \mathrm{Loc}(\mathbf{P}) \supseteq \mathbf{K}\}. \end{aligned}$$

Here,  $\mathrm{Loc}(\mathbf{P})$  refers to the localizing subcategory (meaning triangulated and closed under set-indexed coproducts) of  $\mathrm{StMod}(D(H))$  generated by  $\mathbf{P}$ .

This implies that if  $H$  satisfies the following property, then  $f$  is surjective.

(\*) For  $X$  in  $\mathrm{Mod}(D(H))$ , if  $F(X)$  is projective, then so is  $X$ .

Additionally, if  $\mathbf{C}$  was a braided tensor category to begin with, then we prove that  $f$  is injective. This leads to the following theorem.

**Theorem.** (See Theorem 2.5.1). Let  $\mathbf{C} = \text{mod}(H)$  for a finite-dimensional quasi-triangular Hopf algebra  $H$  satisfying (\*). Then  $f$  is a homeomorphism  $\text{Spcst}(\mathbf{C}) \xrightarrow{\cong} \text{Spcst}(Z(\mathbf{C}))$ , and there is a bijection between the thick ideals of  $\text{st}(Z(\mathbf{C}))$  and the thick ideals of  $\text{st}(\mathbf{C})$ , given by

$$\mathbf{I} \mapsto \overline{F}(X) : X \in \mathbf{I}$$

for a thick ideal  $\mathbf{I}$  of  $\text{st}(Z(\mathbf{C}))$ .

In Section 3, we illustrate the theory with concrete examples. We first consider  $\mathbf{C}$  to be alternately  $\text{mod}(\mathbb{k}G)$  and  $\text{mod}((\mathbb{k}[G]^{\text{cop}}))$ , for  $G$  a finite group and  $\mathbb{k}$  a field of characteristic  $p$  dividing the order of  $G$ , where  $\mathbb{k}G$  denotes the group algebra of  $G$  and  $\mathbb{k}[G]$  denotes the dual group algebra to  $\mathbb{k}G$ . Of these two examples, the first satisfies Property (\*) and the second does not. This allows us to classify the Balmer spectrum and classify the thick ideals for  $\text{stmod}(D(\mathbb{k}G))$ , where  $D(\mathbb{k}G)$  is the Drinfeld double of the group algebra  $\mathbb{k}G$ . We are then able to generalize this example in the following way.

**Theorem.** (See Proposition 3.1.2, Proposition 3.2.5, and Theorem 3.3.4). For the following classes of Hopf algebras  $H$ , the Balmer spectrum of  $\text{stmod}(D(H))$  is homeomorphic via the map  $f$  to the Balmer spectrum of  $\text{stmod}(H)$ , and the thick ideals of the two categories are in bijection:

- (1) finite-dimensional cosemisimple quasitriangular Hopf algebras (e.g. group algebras of finite groups  $G$  in characteristic dividing the order of  $G$ );
- (2) Benson-Witherspoon smash coproducts  $(\mathbb{k}[G] \# \mathbb{k}L)^*$ , where  $G$  and  $L$  are finite groups with dual group algebra and group algebra  $\mathbb{k}[G]$  and  $\mathbb{k}L$  respectively,  $\mathbb{k}$  a field of characteristic  $p$  dividing the order of  $G$  and not dividing the order of  $L$ , such that  $L$  acts by group automorphisms on  $G$ .

**Acknowledgements.** The author would like to thank Daniel Nakano and Milen Yakimov for many useful discussions, and also to thank Pavel Etingof, Siu-Hung Ng, Victor Ostrik, and Sean Sanford for providing valuable comments used to improve this paper.

## 1. PRELIMINARIES

**1.1. Tensor triangular geometry.** We will recall some of the background of noncommutative tensor triangular geometry. A monoidal triangulated category  $\mathbf{T}$  is a category such that the following conditions hold.

- (1)  $\mathbf{T}$  is triangulated: it is equipped with an autoequivalence  $\Sigma$ , called the *shift functor*, and a collection of distinguished triangles

$$A \rightarrow B \rightarrow C \rightarrow \Sigma A,$$

subject to the usual axioms (see [32, 47]);

- (2)  $\mathbf{T}$  is monoidal: it is equipped with a monoidal product  $- \otimes -$  and unit  $\mathbf{1}_{\mathbf{T}}$ , subject to the usual associativity and unit axioms (see [36, 3, 23]);
- (3) The triangulated and monoidal structures on  $\mathbf{T}$  are compatible: for any object  $A$  of  $\mathbf{T}$ , the functors  $A \otimes -$  and  $- \otimes A$  are triangulated (i.e. exact) functors. In other words, there exists a natural isomorphism  $\Sigma(A) \otimes B \cong \Sigma(A \otimes B) \cong A \otimes \Sigma(B)$ , such that if

$$A \rightarrow B \rightarrow C \rightarrow \Sigma A$$

is a distinguished triangle, then for any object  $D$ , the triangles

$$D \otimes A \rightarrow D \otimes B \rightarrow D \otimes C \rightarrow \Sigma(D \otimes A)$$

and

$$A \otimes D \rightarrow B \otimes D \rightarrow C \otimes D \rightarrow \Sigma(A \otimes D)$$

are distinguished.

The setting is the same as the monoidal triangulated categories in [44, 45]. The only difference between such a category and the tensor triangulated categories considered in [4, 5] is that a monoidal triangulated category is not required to have any braiding  $A \otimes B \cong B \otimes A$ .

We will recall the definition of the Balmer spectrum of a monoidal triangulated category  $\mathbf{T}$ , as in [16, 44].

- (1) A (*two-sided*) *thick ideal*  $\mathbf{I}$  of  $\mathbf{T}$  is a full subcategory such that the following hold.
- (a)  $\mathbf{I}$  is triangulated: it is closed under  $\Sigma$  and  $\Sigma^{-1}$ , and if

$$A \rightarrow B \rightarrow C \rightarrow \Sigma A$$

is a distinguished triangle, then if any two of  $A, B$ , and  $C$  are in  $\mathbf{I}$ , then so is the third;

- (b)  $\mathbf{I}$  is *thick*: if  $A \oplus B$  is in  $\mathbf{I}$ , then so are  $A$  and  $B$ ;
- (c)  $\mathbf{I}$  is an *ideal*: if  $A \in \mathbf{I}$ , then so are  $A \otimes B$  and  $B \otimes A$ , for any object  $B$ .

The collection of thick ideals of  $\mathbf{T}$  will be denoted by  $\text{ThickId}(\mathbf{T})$ , and the thick ideal generated by a collection of objects  $\mathcal{T}$  will be denoted  $\langle \mathcal{T} \rangle$ .

- (2) A thick ideal  $\mathbf{P}$  is called *prime* if  $\mathbf{I} \otimes \mathbf{J} \subseteq \mathbf{P}$  implies either  $\mathbf{I}$  or  $\mathbf{J} \subseteq \mathbf{P}$ , for all thick ideals  $\mathbf{I}$  and  $\mathbf{J}$  of  $\mathbf{T}$ ; equivalently,  $\mathbf{P}$  is prime if and only if  $A \otimes \mathbf{T} \otimes B \subseteq \mathbf{P}$  implies either  $A$  or  $B$  is in  $\mathbf{P}$ . Here  $\mathbf{I} \otimes \mathbf{J}$  refers to the collection of objects  $\{A \otimes B : A \in \mathbf{I}, B \in \mathbf{J}\}$ , and  $A \otimes \mathbf{T} \otimes B$  refers to the collection of objects  $\{A \otimes C \otimes B : C \in \mathbf{T}\}$ .
- (3) A thick ideal  $\mathbf{P}$  is called *completely prime* if  $A \otimes B \in \mathbf{P}$  implies either  $A$  or  $B \in \mathbf{P}$ , for all objects  $A$  and  $B$  of  $\mathbf{T}$ .
- (4) The *Balmer spectrum* of  $\mathbf{T}$ , denoted  $\text{Spc } \mathbf{T}$ , is the collection of prime ideals of  $\mathbf{T}$  under the Zariski topology, where closed sets are defined as the sets

$$V_{\mathbf{T}}(\mathcal{T}) = \{\mathbf{P} \in \text{Spc } \mathbf{T} : \mathcal{T} \cap \mathbf{P} = \emptyset\}$$

for all collections of objects  $\mathcal{T}$  of  $\mathbf{T}$ .

Note that every completely prime ideal is prime. If  $\mathbf{T}$  is a braided category then every prime ideal is completely prime, and so in that case the two notions coincide.

Suppose  $\mathbf{T}$  is rigid, in other words, every object is dualizable (cf. [23, Section 2.10]). We then obtain the following facts, which we recall for reference. Both follow directly from the fact that if  $A$  is dualizable with dual  $A^*$ , then  $A$  is a direct summand of  $A \otimes A^* \otimes A$ .

**Proposition 1.1.1.** *Let  $\mathbf{T}$  be a monoidal triangulated category such that every object is dualizable. Let  $A$  be an object of  $\mathbf{T}$  with dual  $A^*$ . Then*

- (1)  $\langle A \rangle = \langle A^* \rangle$ , and
- (2) every thick two-sided ideal  $\mathbf{I}$  of  $\mathbf{T}$  is *semiprime*, i.e. it is the intersection

$$\mathbf{I} = \bigcap_{\mathbf{P} \in \text{Spc } \mathbf{T}, \mathbf{I} \subseteq \mathbf{P}} \mathbf{P}$$

of prime ideals over itself. Equivalently, for every ideal  $\mathbf{I}$  of  $\mathbf{T}$ , if the set  $A \otimes \mathbf{T} \otimes A \subseteq \mathbf{I}$ , then  $A \in \mathbf{I}$ , where  $A \otimes \mathbf{T} \otimes A$  refers to the collection  $\{A \otimes B \otimes A : B \in \mathbf{T}\}$ .

For the details of the proofs, see [44, Lemma 5.1.1] and [45, Proposition 4.1.1].

**1.2. Support data.** Suppose that  $\mathbf{T}$  is a monoidal triangulated category and  $S$  a topological space. We will denote the collection of subsets of  $S$  by  $\mathcal{X}(S)$ , closed subsets of  $S$  by  $\mathcal{X}_{cl}(S)$ , and specialization-closed subsets of  $S$  by  $\mathcal{X}_{sp}(S)$ ; recall that by definition, a set is specialization-closed if it is a union of closed sets. When the underlying space is clear from context, we will denote these collections by  $\mathcal{X}$ ,  $\mathcal{X}_{cl}$ , and  $\mathcal{X}_{sp}$ .

Given a monoidal triangulated category  $\mathbf{T}$  and a topological space  $S$ , a *support datum on  $\mathbf{T}$  with value in  $S$*  is a map  $\sigma : \mathbf{T} \rightarrow \mathcal{X}(S)$  satisfying the following axioms:

- (1)  $\sigma(0) = \emptyset$  and  $\sigma(\mathbf{1}) = S$ ;
- (2)  $\sigma(A \oplus B) = \sigma(A) \cup \sigma(B)$ ,  $\forall A, B \in \mathbf{T}$ ;
- (3)  $\sigma(\Sigma A) = \sigma(A)$ ,  $\forall A \in \mathbf{T}$ ;
- (4) if  $A \rightarrow B \rightarrow C \rightarrow \Sigma A$  is a distinguished triangle, then  $\sigma(A) \subseteq \sigma(B) \cup \sigma(C)$ ;
- (5)  $\bigcup_{C \in \mathbf{T}} \sigma(A \otimes C \otimes B) = \sigma(A) \cap \sigma(B)$ ,  $\forall A, B \in \mathbf{T}$ .

See [44, Section 4] for a more in-depth discussion of support data. For any monoidal triangulated category  $\mathbf{T}$ , the map  $V_{\mathbf{T}}(A) = \{\mathbf{P} \in \mathrm{Spc} \mathbf{T} : A \notin \mathbf{P}\}$  defined above is a support datum  $\mathbf{T} \rightarrow \mathcal{X}(\mathrm{Spc} \mathbf{T})$ . By definition,  $V_{\mathbf{T}}(A)$  is a closed set in  $\mathrm{Spc} \mathbf{T}$ ; in other words,  $V_{\mathbf{T}}$  takes values in  $\mathcal{X}_{cl}(\mathrm{Spc} \mathbf{T})$ . We will refer to this support datum as the *Balmer support*.

For any support datum  $\sigma$ , we have a map

$$\Phi_{\sigma}(\mathcal{T}) := \bigcup_{A \in \mathcal{T}} \sigma(A),$$

where  $\mathcal{T}$  is any collection of objects of  $\mathbf{T}$ . If  $\sigma$  takes values in  $\mathcal{X}_{cl}$ , then by definition  $\Phi_{\sigma}$  takes values in  $\mathcal{X}_{sp}$ . The map  $\Phi_{\sigma}$  in fact only depends on thick ideals rather than arbitrary subsets, since by [44, Lemma 4.3.2] we have  $\Phi_{\sigma}(\mathcal{T}) = \Phi_{\sigma}(\langle \mathcal{T} \rangle)$ .

For a support datum  $\sigma$ , we have a second map  $\Theta_{\sigma} : \mathcal{X}_{sp}(S) \rightarrow \mathrm{ThickId}(\mathbf{T})$  defined by

$$\Theta_{\sigma}(S') := \{A \in \mathbf{T} : \sigma(A) \subseteq S'\}$$

for any specialization-closed subset  $S'$  of  $S$ . For any specialization closed set  $S'$ , the collection  $\Theta_{\sigma}(S')$  is a thick ideal of  $\mathbf{T}$ . Hence, we have the following collection of maps, given a support datum  $\sigma$  on  $\mathbf{T}$  which takes values in  $\mathcal{X}_{cl}$ :

$$\mathrm{ThickId}(\mathbf{T}) \begin{array}{c} \xrightarrow{\Phi_{\sigma}} \\ \xleftarrow{\Theta_{\sigma}} \end{array} \mathcal{X}_{sp}.$$

Classifications of thick ideals are obtained in many cases (see [4, 5, 14, 15, 44, 45] for examples) by constructing a support datum for which these maps are bijective and inverse to each other.

**1.3. Stable module categories.** The primary examples of monoidal triangulated categories that we will concern ourselves with in this paper are the stable categories of finite tensor categories (and, in particular, the stable module categories of finite-dimensional Hopf algebras). Recall that a finite tensor category (following the notation given in [24, 23]) consists of a monoidal category  $\mathbf{C}$  such that:

- (1)  $\mathbf{C}$  is abelian and  $\mathbb{k}$ -linear for some field  $\mathbb{k}$ ;
- (2) the tensor product  $- \otimes -$  is bilinear on spaces of morphisms;
- (3) every object of  $\mathbf{C}$  has finite length;
- (4)  $\mathrm{Hom}_{\mathbf{C}}(\mathbf{1}, \mathbf{1}) \cong \mathbb{k}$ ;
- (5) for any pair of objects  $A$  and  $B$ , the vector space  $\mathrm{Hom}_{\mathbf{C}}(A, B)$  is finite-dimensional over  $\mathbb{k}$ ;
- (6)  $\mathbf{C}$  has enough projectives;

- (7) there are finitely many isomorphism classes of simple objects of  $\mathbf{C}$ ;
- (8)  $\mathbf{C}$  is rigid, i.e. every object has a left and a right dual.

The stable category of a finite tensor category  $\mathbf{C}$ , which we will denote by  $\text{st}(\mathbf{C})$ , is constructed by factoring out the projective objects of  $\mathbf{C}$ . In more detail, let  $\text{PHom}_{\mathbf{C}}(X, Y)$  consist of the morphisms  $f : A \rightarrow B$  in  $\mathbf{C}$  such that  $f$  factors through a projective object. The stable category  $\text{st}(\mathbf{C})$  is the category where

- (1) objects are the same as the objects of  $\mathbf{C}$ ;
- (2) morphisms  $A \rightarrow B$  are defined to be  $\text{Hom}_{\mathbf{C}}(A, B)/\text{PHom}_{\mathbf{C}}(A, B)$ .

There is a straightforward functor  $G : \mathbf{C} \rightarrow \text{st}(\mathbf{C})$  sending objects to themselves and morphisms to their image in the quotient.

If  $P$  is a projective object of  $\mathbf{C}$ , note that the corresponding object  $G(P)$  in  $\text{st}(\mathbf{C})$  is isomorphic to 0, since  $\text{id}_P$  factors through a projective; and the converse is also true, since  $G(P) \cong 0$  in  $\text{st}(\mathbf{C})$  implies that the 0-morphism  $G(P) \rightarrow G(P)$  is equal to  $\text{id}_{G(P)}$  in  $\text{Hom}_{\text{st}(\mathbf{C})}(G(P), G(P))$ , in other words,  $\text{id}_P$  factors through a projective  $Q$  in  $\mathbf{C}$ :

$$\begin{array}{ccc} P & \xrightarrow{\text{id}_P} & P \\ & \searrow & \nearrow \\ & & Q \end{array}$$

Of course, this implies  $P$  is a summand of  $Q$ , and so  $P$  is projective.

If  $\mathbf{C}$  is a finite tensor category, then  $\text{st}(\mathbf{C})$  is a monoidal triangulated category. The monoidal product is inherited directly from  $\mathbf{C}$ : we define  $G(A) \otimes G(B) := G(A \otimes B)$ , and similarly for morphisms  $f : A \rightarrow B$  and  $g : C \rightarrow D$  we define  $G(f) \otimes G(g) := G(f \otimes g)$ . This is well-defined: if  $G(f) = G(\hat{f})$ , then  $f - \hat{f}$  factors through a projective  $P$ , and then  $f \otimes g - \hat{f} \otimes g$  factors through  $P \otimes D$ , which is projective by projectivity of  $P$  (see [23, Proposition 4.2.12]).

We recall the triangulated structure on  $\text{st}(\mathbf{C})$ , for reference. If  $A$  is an object of  $\mathbf{C}$ , denote by  $\Omega(A)$  the kernel of the projective cover of  $A$ . The functor  $\Omega$  extends to the stable module category, and this in fact gives an autoequivalence on  $\text{st}(\mathbf{C})$ . The shift  $\Sigma$  is then defined to be  $\Sigma(A) = \Omega^{-1}(A)$ . For any short exact sequence of objects in  $\mathbf{C}$ , say

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0,$$

there exists a triangle

$$A \rightarrow B \rightarrow C \rightarrow \Sigma A$$

in  $\text{st}(\mathbf{C})$ ; the distinguished triangles of  $\text{st}(\mathbf{C})$  are then defined to be all triangles which are isomorphic to triangles of this form. For more details on the triangulated structure of  $\text{st}(\mathbf{C})$ , see [32, Chapter I].

**1.4. The Drinfeld center.** Let  $\mathbf{C}$  be a strict monoidal category. Then the *Drinfeld center* or *center* of  $\mathbf{C}$ , which we will denote by  $Z(\mathbf{C})$ , is defined as the following braided monoidal category.

- (1) Objects are pairs  $(A, \gamma)$  where  $A$  is an object of  $\mathbf{T}$  and  $\gamma$  is a natural isomorphism  $\gamma_B : B \otimes A \xrightarrow{\cong} A \otimes B$  for all  $B \in \mathbf{C}$ , satisfying the diagram

$$\begin{array}{ccccc} & & \gamma_{B \otimes C} & & \\ & \searrow & \text{---} & \nearrow & \\ B \otimes C \otimes A & \xrightarrow{\text{id}_B \otimes \gamma_C} & B \otimes A \otimes C & \xrightarrow{\gamma_B \otimes \text{id}_C} & A \otimes B \otimes C \end{array}$$

for all  $B$  and  $C$ . Such a natural isomorphism  $\gamma$  is called a *half-braiding* of  $A$ .

- (2) Morphisms  $(A, \gamma) \rightarrow (A', \gamma')$  are morphisms  $f : A \rightarrow A'$  such that for all  $B$ , the diagram

$$\begin{array}{ccc} B \otimes A & \xrightarrow{\text{id}_B \otimes f} & B \otimes A' \\ \downarrow \gamma_B & & \downarrow \gamma'_B \\ A \otimes B & \xrightarrow{f \otimes \text{id}_B} & A' \otimes B \end{array}$$

commutes.

- (3) The monoidal product  $(A, \gamma) \otimes (A', \gamma')$  is defined as  $(A \otimes A', \tilde{\gamma})$  where  $\tilde{\gamma}$  is defined as

$$\begin{array}{ccc} B \otimes A \otimes A' & \xrightarrow{\gamma_B \otimes \text{id}_{A'}} & A \otimes B \otimes A' \\ \downarrow \tilde{\gamma}_B & \swarrow \text{id}_{A'} \otimes \gamma'_B & \\ A \otimes A' \otimes B & & \end{array}$$

- (4) The braiding  $c_{(A, \gamma), (A', \gamma')} : (A, \gamma) \otimes (A', \gamma') \xrightarrow{\cong} (A', \gamma') \otimes (A, \gamma)$  is defined as  $\gamma'_A$ . The map  $\gamma'_A$  being a valid map in  $\mathbf{Z}(\mathbf{C})$  amounts to checking the commutativity of the diagram

$$\begin{array}{ccc} B \otimes A \otimes A' & \xrightarrow{\text{id}_B \otimes \gamma'_A} & B \otimes A' \otimes A \\ \downarrow \gamma_B \otimes \text{id}_{A'} & & \downarrow \gamma'_B \otimes \text{id}_A \\ A \otimes B \otimes A' & & A' \otimes B \otimes A \\ \downarrow \text{id}_A \otimes \gamma'_B & & \downarrow \text{id}_{A'} \otimes \gamma_B \\ A \otimes A' \otimes B & \xrightarrow{\gamma'_A \otimes \text{id}_B} & A' \otimes A \otimes B \end{array}$$

This diagram commutes by the naturality of  $\gamma'$ , since it can be rewritten, using the defining diagram for  $\gamma'$ , as

$$\begin{array}{ccc} B \otimes A \otimes A' & \xrightarrow{\gamma'_B \otimes A} & A' \otimes B \otimes A \\ \downarrow \gamma_B \otimes \text{id}_{A'} & & \downarrow \text{id}_{A'} \otimes \gamma_B \\ A \otimes B \otimes A' & \xrightarrow{\gamma'_A \otimes B} & A' \otimes A \otimes B \end{array}$$

We will denote by  $F : \mathbf{Z}(\mathbf{C}) \rightarrow \mathbf{C}$  the forgetful functor sending  $(A, \gamma) \mapsto A$ .

If  $H$  is a Hopf algebra and  $\mathbf{C}$  is the category of  $H$ -modules, it is well-known that the Drinfeld center  $\mathbf{Z}(\mathbf{C})$  of  $\mathbf{C}$  is equivalent to the category of modules of  $D(H)$  the Drinfeld (or quantum) double of  $H$ . For the details of Drinfeld doubles, see [42, Section 10.3], [17, Section 4.2.D], [36, Section IX.4], or [23, Section 7.14]. The Drinfeld double  $D(H)$  is isomorphic as a vector space to  $(H^{\text{op}})^* \otimes H$ , and contains both  $H$  and  $(H^{\text{op}})^*$  as Hopf subalgebras. Here if  $H$  is a Hopf algebra with multiplication  $\mu$ , unit  $\eta$ , comultiplication  $\Delta$ , counit  $\epsilon$ , and antipode  $S$ , then  $(H^{\text{op}})^*$  is the Hopf algebra with multiplication  $\Delta^*$ , unit  $\epsilon^*$ , comultiplication  $(\mu^{\text{op}})^*$ , counit  $\eta^*$ , and antipode  $(S^{-1})^*$ .

The following result of Etingof-Ostrik will be important in extending the forgetful functor  $\mathbf{Z}(\mathbf{C}) \rightarrow \mathbf{C}$  to the stable categories.

**Proposition 1.4.1.** ([24]). *If  $\mathbf{C}$  is a finite tensor category, then its Drinfeld center  $\mathbf{Z}(\mathbf{C})$  is a finite tensor category, and the forgetful functor  $F$  is exact and sends projective objects to projective objects.*

The fact that  $F$  preserves projectivity is a generalization of the classical Nichols-Zoeller theorem for Hopf algebras, which states that a finite-dimensional Hopf algebra is free as a module over any Hopf subalgebra [51].

## 2. THE DRINFELD CENTER AND BALMER SPECTRA

**2.1. Construction of a continuous map between Balmer spectra.** Recall the stable categories defined in Section 1.3. For the rest of this section, let  $\mathbf{C}$  be a finite tensor category,  $\text{st}(\mathbf{C})$  its stable category,  $Z(\mathbf{C})$  its Drinfeld center, and  $\text{st}(Z(\mathbf{C}))$  the stable category of its Drinfeld center (which may be formed by Proposition 1.4.1). We have a forgetful functor  $F : Z(\mathbf{C}) \rightarrow \mathbf{C}$ , and we have functors  $G : \mathbf{C} \rightarrow \text{st}(\mathbf{C})$  and  $H : Z(\mathbf{C}) \rightarrow \text{st}(Z(\mathbf{C}))$ . We have the respective Balmer support data associated to  $\text{st}(\mathbf{C})$  and  $\text{st}(Z(\mathbf{C}))$ :

$$V_{\text{st} \mathbf{C}} : \text{st}(\mathbf{C}) \rightarrow \mathcal{X}_{cl}(\text{Spc st}(\mathbf{C}))$$

and

$$V_{\text{st}(Z(\mathbf{C}))} : \text{st}(Z(\mathbf{C})) \rightarrow \mathcal{X}_{cl}(\text{Spc st}(Z(\mathbf{C}))),$$

defined in their respective categories by sending

$$A \mapsto \{\text{primes not containing } A\}.$$

For readability, we will denote  $V_{\mathbf{C}} := V_{\text{st} \mathbf{C}}$  and  $V_Z := V_{\text{st}(Z(\mathbf{C}))}$ . The corresponding maps  $\Phi$  (recalling the construction from Section 1.2) associated to these support data will similarly be denoted  $\Phi_{\mathbf{C}}$  and  $\Phi_Z$ .

The following proposition is probably well-known to experts, but we record it for completeness.

**Proposition 2.1.1.** *There is a functor  $\overline{F} : \text{st}(Z(\mathbf{C})) \rightarrow \text{st}(\mathbf{C})$  which extends the forgetful functor  $F$ , i.e. the diagram of functors*

$$\begin{array}{ccc} \mathbf{C} & \xrightarrow{G} & \text{st}(\mathbf{C}) \\ \uparrow F & & \uparrow \overline{F} \\ Z(\mathbf{C}) & \xrightarrow{H} & \text{st}(Z(\mathbf{C})) \end{array}$$

*commutes. This functor  $\overline{F}$  is monoidal and triangulated.*

*Proof.* Since the objects of  $\text{st}(Z(\mathbf{C}))$  are in bijection with those of  $Z(\mathbf{C})$ ,  $\overline{F}$  is well-defined on objects, namely by defining

$$\overline{F}(H(X)) := G(F(X)).$$

Let  $f : X \rightarrow Y$  be a morphism in  $Z(\mathbf{C})$ . Then for  $\overline{F}(H(f)) := GF(f)$  to be well-defined, we need  $GF(g) = 0$  for each morphism  $g$  which factors through a projective in  $Z(\mathbf{C})$ . In other words, we need  $F(g)$  to factor through a projective in  $\mathbf{C}$ . Hence, to define  $\overline{F}$ , it is enough to know that  $G \circ F$  sends all projective objects of  $Z(\mathbf{C})$  to 0, which is true by Proposition 1.4.1.

Let  $H(X) \in \text{st}(Z(\mathbf{C}))$  an arbitrary object, where  $X \in Z(\mathbf{C})$ . Then  $\Sigma H(X)$  is defined as  $H(Z)$ , such that there exists a short exact sequence

$$0 \rightarrow X \rightarrow P \rightarrow Z \rightarrow 0$$

in  $Z(\mathbf{C})$ , where  $P$  is projective. The object  $\Sigma H(Z)$  is well-defined in  $\text{st}(Z(\mathbf{C}))$ , by Schanuel's Lemma. Since  $F$  is exact and sends projectives to projectives,

$$0 \rightarrow F(X) \rightarrow F(P) \rightarrow F(Z) \rightarrow 0$$

is an exact sequence in  $\mathbf{C}$  with  $F(P)$  projective; therefore,  $\Sigma(GF(X)) \cong GF(Z)$  in  $\mathbf{st}(\mathbf{C})$ , and so we have  $\overline{F}(\Sigma X) \cong \Sigma \overline{F}(X)$ .

Now, let  $X \rightarrow Y \rightarrow Z \rightarrow \Sigma X$  be a distinguished triangle in  $\mathbf{st}(\mathbf{Z}(\mathbf{C}))$ . Then it is isomorphic to a triangle of the form

$$H(X') \rightarrow H(Y') \rightarrow H(Z') \rightarrow \Sigma H(X')$$

for some short exact sequence

$$0 \rightarrow X' \rightarrow Y' \rightarrow Z' \rightarrow 0$$

in  $\mathbf{Z}(\mathbf{C})$ . Since  $F$  is exact, and  $G$  sends exact sequences to triangles, we have that the composition  $GF$  is exact and hence

$$\overline{F}H(X') \rightarrow \overline{F}H(Y') \rightarrow \overline{F}H(Z') \rightarrow \Sigma \overline{F}H(X')$$

a triangle in  $\mathbf{st}(\mathbf{C})$ . Therefore,

$$\overline{F}(X) \rightarrow \overline{F}(Y) \rightarrow \overline{F}(Z) \rightarrow \Sigma \overline{F}(X)$$

is a triangle as well, and so  $\overline{F}$  is a triangulated functor.  $\square$

For braided tensor triangulated categories, the Balmer spectrum  $\mathrm{Spc}$  is functorial, as Balmer has shown in [4, Proposition 3.6]. This is a categorical reflection the ring-theoretic fact that  $\mathrm{Spec}$  is functorial for commutative rings. On the other hand, for noncommutative rings,  $\mathrm{Spec}$  is not a functor (for an in-depth exploration of the extent of the failure of functoriality of  $\mathrm{Spec}$  for noncommutative rings, see [54]). It is not surprising, then, that for generic monoidal triangulated categories, the Balmer spectrum is also not functorial; in other words, an exact monoidal functor between monoidal triangulated categories does not necessarily induce a map between their Balmer spectra.

However, reflecting the classical prime ideal contraction for noncommutative rings, the forgetful functor  $\overline{F}$  does induce a map between the Balmer spectra of  $\mathbf{st}(\mathbf{C})$  and  $\mathbf{st}(\mathbf{Z}(\mathbf{C}))$ .

**Proposition 2.1.2.** *The functor  $\overline{F}$  induces a continuous map  $\mathrm{Spc} \mathbf{st}(\mathbf{C}) \xrightarrow{f} \mathrm{Spc} \mathbf{st}(\mathbf{Z}(\mathbf{C}))$ , defined by*

$$f(\mathbf{P}) := \{X \in \mathbf{st}(\mathbf{Z}(\mathbf{C})) : \overline{F}(X) \in \mathbf{P}\}.$$

*Proof.* Let  $\mathbf{P}$  be a prime ideal of  $\mathbf{st}(\mathbf{C})$ . We must first show that  $f(\mathbf{P})$  is a prime ideal of  $\mathbf{st}(\mathbf{Z}(\mathbf{C}))$ .

We first check that  $f(\mathbf{P})$  is a thick ideal of  $\mathbf{st}(\mathbf{Z}(\mathbf{C}))$ . This necessitates checking three properties:

(Triangulated) Suppose  $\Sigma X \in f(\mathbf{P})$ , in other words,  $\overline{F}(\Sigma X) \in \mathbf{P}$ . Since  $\overline{F}$  is exact, this is true if and only if  $\Sigma \overline{F}(X) \in \mathbf{P}$ , which is true if and only if  $\overline{F}(X) \in \mathbf{P}$ , in other words,  $X \in f(\mathbf{P})$ . Now, suppose

$$X \rightarrow Y \rightarrow Z \rightarrow \Sigma X$$

is a distinguished triangle with  $X$  and  $Y$  in  $f(\mathbf{P})$ . This means that  $\overline{F}(X)$  and  $\overline{F}(Y)$  are in  $\mathbf{P}$ . Since  $\overline{F}$  is triangulated, the triangle

$$\overline{F}(X) \rightarrow \overline{F}(Y) \rightarrow \overline{F}(Z) \rightarrow \Sigma \overline{F}(X)$$

is distinguished in  $\mathbf{st}(\mathbf{C})$ . Now since the first two objects are in  $\mathbf{P}$ , so is  $\overline{F}(Z)$ , and so  $Z \in f(\mathbf{P})$ .

(Thick) If  $X \oplus Y$  is in  $f(\mathbf{P})$ , then  $\overline{F}(X \oplus Y) \in \mathbf{P}$ ;  $\overline{F}$  is an additive functor, and so  $\overline{F}(X) \oplus \overline{F}(Y) \in \mathbf{P}$ . This implies that both  $\overline{F}(X)$  and  $\overline{F}(Y)$  are in  $\mathbf{P}$ , and so  $X$  and  $Y$  are both in  $f(\mathbf{P})$ .

(Ideal) Suppose  $X \in f(\mathbf{P})$  and  $Y \in \text{st}(\mathbf{Z}(\mathbf{C}))$ . Since  $\overline{F}$  is monoidal,  $\overline{F}(X \otimes Y) \cong \overline{F}(X) \otimes \overline{F}(Y)$ . Since  $\overline{F}(X) \in \mathbf{P}$ , so is  $\overline{F}(X) \otimes \overline{F}(Y)$ , and thus  $\overline{F}(X \otimes Y) \in \mathbf{P}$  as well. Hence  $X \otimes Y \in f(\mathbf{P})$ . The symmetric argument shows that  $Y \otimes X$  is in  $f(\mathbf{P})$  as well, so  $f(\mathbf{P})$  is a two-sided ideal.

(Prime) Let  $A \otimes B \in f(\mathbf{P})$ . Then  $\overline{F}(A) \otimes \overline{F}(B) \in \mathbf{P}$ . But  $\overline{F}(A)$  and  $\overline{F}(B)$  commute with every object of  $\text{st}(\mathbf{C})$ : by the ideal property of  $\mathbf{P}$ , we have

$$\begin{aligned} \text{st}(\mathbf{C}) \otimes \overline{F}(A) \otimes \overline{F}(B) &\subseteq \mathbf{P} \\ \Rightarrow \overline{F}(A) \otimes \text{st}(\mathbf{C}) \otimes \overline{F}(B) &\subseteq \mathbf{P} \\ \Rightarrow \overline{F}(A) \text{ or } \overline{F}(B) &\in \mathbf{P}, \end{aligned}$$

with the last step following by primeness of  $\mathbf{P}$ . This implies that either  $A$  or  $B$  is in  $f(\mathbf{P})$ , which means that  $f(\mathbf{P})$  is prime.

We can also check directly that  $f$  is continuous: an arbitrary closed set of  $\text{Spc}(\text{st}(\mathbf{Z}(\mathbf{C})))$  is of the form  $V_{\mathbf{Z}}(\mathcal{T}) = \{\mathbf{P} \in \text{Spc}(\text{st}(\mathbf{Z}(\mathbf{C}))) : \mathcal{T} \cap \mathbf{P} = \emptyset\}$  for some collection of objects  $\mathcal{T}$  of  $\text{st}(\mathbf{Z}(\mathbf{C}))$ . Then

$$\begin{aligned} f^{-1}(V_{\mathbf{Z}}(\mathcal{T})) &= \{\mathbf{P} \in \text{Spc} \text{st}(\mathbf{C}) : \mathcal{T} \cap \{X \in \text{st}(\mathbf{Z}(\mathbf{C})) : \overline{F}(X) \in \mathbf{P}\} = \emptyset\} \\ &= \{\mathbf{P} \in \text{Spc} \text{st}(\mathbf{C}) : \overline{F}(\mathcal{T}) \cap \mathbf{P} = \emptyset\} \\ &= V_{\mathbf{C}}(\overline{F}(\mathcal{T})), \end{aligned}$$

where by  $\overline{F}(\mathcal{T})$  we mean the collection  $\{\overline{F}(X) : X \in \mathcal{T}\}$ .  $\square$

**Remark 2.1.3.** If  $H$  is a finite-dimensional Hopf algebra, then  $\mathbf{Z}(\text{mod}(H)) \cong \text{mod}(D(H))$ , recall Section 1.4. In this case,  $D(H) \cong D((H^{\text{op}})^*)$ , and so we have two functors:

$$\begin{array}{ccc} \text{mod}(H) & & \text{mod}((H^{\text{op}})^*) \\ & \swarrow F_H & \searrow F_{(H^{\text{op}})^*} \\ & \mathbf{Z}(\text{mod}(H)) \cong \text{mod}(D(H)) \cong \mathbf{Z}(\text{mod}((H^{\text{op}})^*)) & \end{array}$$

which then give two maps between Balmer spectra:

$$\begin{array}{ccc} \text{Spc}(\text{stmod}(H)) & & \text{Spc}(\text{stmod}((H^{\text{op}})^*)) \\ & \searrow f_H & \swarrow f_{(H^{\text{op}})^*} \\ & \text{Spc}(\text{stmod}(D(H))) & \end{array}$$

**2.2. A support data interpretation.** We can interpret the map  $f$  in the context of support data (recalling the definition from Section 1.2), by first defining a new support datum given as the composition of the functor  $\overline{F}$  with the Balmer support  $V_{\mathbf{C}}$  on  $\text{st}(\mathbf{C})$ .

**Proposition 2.2.1.** *Define a map  $W : \text{st}(\mathbf{Z}(\mathbf{C})) \rightarrow \mathcal{X}_{\text{cl}}(\text{Spc} \text{st}(\mathbf{C}))$  by*

$$W(X) := V_{\mathbf{C}}(\overline{F}(X)) = \{\mathbf{P} \in \text{Spc} \text{st}(\mathbf{C}) : \overline{F}(X) \notin \mathbf{P}\}.$$

*This map is a support datum.*

*Proof.* The first four conditions follow directly from the facts that  $\overline{F}$  is an exact functor and  $V_{\mathbf{C}}$  is itself a support datum, since

- (1)  $\overline{F}(0_{\text{st}(\mathbf{Z}(\mathbf{C}))}) = 0_{\text{st}(\mathbf{C})}$ ,
- (2)  $\overline{F}(X \oplus Y) = \overline{F}(X) \oplus \overline{F}(Y)$ ,
- (3)  $\overline{F}(\Sigma X) \cong \Sigma \overline{F}(X)$ ,

(4) and if  $X \rightarrow Y \rightarrow Z \rightarrow \Sigma X$  is a distinguished triangle, then so is  $\overline{F}(X) \rightarrow \overline{F}(Y) \rightarrow \overline{F}(Z) \rightarrow \Sigma \overline{F}(X)$ .

To check the last condition, we need to show that

$$\bigcup_{Z \in \text{st}(\mathbf{Z}(\mathbf{C}))} W(X \otimes Z \otimes Y) = W(X) \cap W(Y).$$

By the ideal condition, if  $\mathbf{P}$  is a prime ideal which does not contain  $\overline{F}(X) \otimes \overline{F}(Z) \otimes \overline{F}(Y)$  for some object  $Z$ , then it must also not contain  $\overline{F}(X)$  or  $\overline{F}(Y)$ . Hence,

$$\bigcup_{Z \in \text{st}(\mathbf{Z}(\mathbf{C}))} W(X \otimes Z \otimes Y) \subseteq W(X) \cap W(Y)$$

is automatic.

For the reverse containment, suppose  $\mathbf{P}$  is a prime ideal which does not contain  $\overline{F}(X)$  or  $\overline{F}(Y)$ . By the prime condition, that means  $\mathbf{P}$  does not contain the entire collection of objects  $\overline{F}(X) \otimes \text{st}(\mathbf{C}) \otimes \overline{F}(Y)$ . But since  $\overline{F}(X)$  and  $\overline{F}(Y)$  commute up to isomorphism with all elements of  $\text{st}(\mathbf{C})$ , if  $\overline{F}(X) \otimes \overline{F}(Y) \in \mathbf{P}$ , that would imply  $\overline{F}(X) \otimes \overline{F}(Y) \otimes \text{st}(\mathbf{C}) \subseteq \mathbf{P}$ , which would then imply  $\overline{F}(X) \otimes \text{st}(\mathbf{C}) \otimes \overline{F}(Y) \subseteq \mathbf{P}$ , a contradiction. Hence,  $\mathbf{P} \in W(X \otimes Y)$ , and we have the claimed equality.  $\square$

By the universal property of the Balmer spectrum (see [44, Theorem 4.2.2]), the support datum  $W$  induces a continuous map  $\text{Spc st}(\mathbf{C}) \rightarrow \text{Spc st}(\mathbf{Z}(\mathbf{C}))$ . This map is defined as

$$\mathbf{P} \mapsto \{X \in \text{st}(\mathbf{Z}(\mathbf{C})) : \mathbf{P} \notin W(X)\}.$$

This map is the same as the map defined in Proposition 2.1.2. We have the following diagram, which commutes by definition:

$$\begin{array}{ccc} \text{st}(\mathbf{Z}(\mathbf{C})) & \xrightarrow{W} & \mathcal{X}_{cl}(\text{Spc}(\text{st}(\mathbf{C}))) \\ & \searrow \overline{F} & \nearrow v_{\mathbf{C}} \\ & & \text{st}(\mathbf{C}) \end{array}$$

On the level of ideals, we have the following induced maps, recall  $\Phi$  and  $\Theta$  associated to a support datum as constructed in Section 1.2:

$$\begin{array}{ccc} & \Theta_W & \\ & \curvearrowright & \\ \text{ThickId}(\text{st}(\mathbf{Z}(\mathbf{C}))) & \xrightarrow{\Phi_W} & \mathcal{X}_{sp}(\text{Spc}(\text{st}(\mathbf{C}))) \\ & \searrow \Psi & \nearrow \Phi_{\mathbf{C}} \\ & & \text{ThickId}(\text{st}(\mathbf{C})) \\ & \swarrow \Lambda & \searrow \Theta_{\mathbf{C}} \end{array}$$

Here, for thick ideals  $\mathbf{I}$  of  $\text{st}(\mathbf{Z}(\mathbf{C}))$  and  $\mathbf{J}$  of  $\text{st}(\mathbf{C})$ , the maps  $\Psi$  and  $\Lambda$  are defined by

$$\begin{aligned} \Psi : \mathbf{I} &\mapsto \langle \overline{F}(\mathbf{I}) \rangle, \\ \Lambda : \mathbf{J} &\mapsto \{X \in \text{st}(\mathbf{Z}(\mathbf{C})) : \overline{F}(X) \in \mathbf{J}\}. \end{aligned}$$

By definition, the inner and outer triangles commute: in other words,  $\Phi_W = \Phi_{\mathbf{C}} \circ \Psi$ , and  $\Theta_W = \Lambda \circ \Theta_{\mathbf{C}}$ .

**2.3. Recovering ideals from their supports.** In [44, Theorem 6.2.1], conditions were given under which an arbitrary support datum  $\sigma : \mathbf{T} \rightarrow \mathcal{X}(S)$  has the property that  $\Phi_\sigma$  is a left, right, and two-sided inverse to  $\Theta_\sigma$ . In the second case, this means that all thick ideals can be recovered from their supports; in the third case, the ideals are completely classified by the topological space  $S$ . Since the support datum  $W(-)$  defined above might not satisfy conditions under which every ideal may be recovered from their support (see Section 3 for examples), in this section we discuss precisely which ideals can be recovered in this way; this allows us to describe the image of the map  $f$  defined above.

We will use the technology of localization and colocalization functors defined in [10]. To use these functors, we must work in the context of a compactly generated monoidal triangulated category; that is, a monoidal triangulated category which is closed under arbitrary coproducts, such that it is generated, as a triangulated category, by the collection of compact objects. Recall that an object  $C$  in a category  $\mathbf{T}$  is said to be compact if  $\mathrm{Hom}_{\mathbf{T}}(C, -)$  commutes with set-indexed coproducts. For background on compactly generated triangulated categories, see [11, Section 1.3.9].

To work in this situation, we specialize to the case where  $\mathbf{C} = \mathrm{mod}(H)$  for a Hopf algebra  $H$ . In that case, we denote  $\mathrm{St}(\mathbf{C})$  for the stable module category of (not necessarily finite-dimensional) modules of  $H$ . The category  $\mathrm{St}(\mathbf{C})$  is compactly generated, and the compact objects of  $\mathrm{St}(\mathbf{C})$  are the finite-dimensional modules, i.e.  $\mathrm{st}(\mathbf{C})$ , the stable module category of finite-dimensional modules of  $H$ . In this situation, then, we have  $\mathbf{Z}(\mathbf{C}) \cong \mathrm{mod}(D(H))$ , and  $\mathrm{st}(\mathbf{Z}(\mathbf{C})) \cong \mathrm{stmod}(D(H))$  is the collection of compact objects in the compactly generated category  $\mathrm{St}(\mathbf{Z}(\mathbf{C})) \cong \mathrm{StMod}(D(H))$ .

We recall some terminology for compactly generated triangulated categories. A subcategory of a compactly generated triangulated category  $\mathbf{T}$  is said to be *localizing* if it is triangulated and closed under arbitrary set-indexed coproducts. Such a category is automatically thick, by a version of the Eilenberg swindle. Given a collection of objects  $\mathcal{T}$  in  $\mathbf{T}$ , we will denote by  $\mathrm{Loc}(\mathcal{T})$  the smallest localizing subcategory of  $\mathbf{T}$  containing  $\mathcal{T}$ .

We will need to invoke the following theorem, see [14, Theorem 3.1.1, Theorem 3.1.2]. For details on the localization and colocalization functors, see [10, Section 3].

**Theorem 2.3.1.** *Let  $\mathbf{T}$  be a compactly generated triangulated category. Given a thick subcategory  $\mathbf{S}$  of  $\mathbf{T}^c$ , there exist functors  $\Gamma_{\mathbf{S}}$  and  $L_{\mathbf{S}}$  from  $\mathbf{T} \rightarrow \mathbf{T}$ , which gives for every object  $M$  of  $\mathbf{T}$  a distinguished triangle*

$$\Gamma_{\mathbf{S}}(M) \rightarrow M \rightarrow L_{\mathbf{S}}(M) \rightarrow \Sigma(\Gamma_{\mathbf{S}}(M))$$

such that

- (1)  $L_{\mathbf{S}}$  and  $\Gamma_{\mathbf{S}}$  are unique up to isomorphism,
- (2)  $\Gamma_{\mathbf{S}}(M)$  is in  $\mathrm{Loc}(\mathbf{S})$ ,
- (3) there are no nonzero maps from  $\mathrm{Loc}(\mathbf{S}) \rightarrow L_{\mathbf{S}}(M)$ , and
- (4)  $M \in \mathrm{Loc}(\mathbf{S})$  if and only if  $\Gamma_{\mathbf{S}}(M) \cong M$ , or, equivalently,  $L_{\mathbf{S}}(M) \cong 0$ .

We now introduce some terminology, which will be useful for our reconstruction theory.

**Definition 2.3.2.** The kernel of  $\overline{F}$ , which we will denote by  $\mathbf{K}$  for the sake of brevity, is defined by

$$\mathbf{K} := \{X \in \mathrm{St}(\mathbf{Z}(\mathbf{C})) : \overline{F}(X) \cong 0 \in \mathrm{St}(\mathbf{C})\}.$$

An equivalent characterization of the kernel of  $\overline{F}$  can be given by

$$\mathbf{K} = \{H(X) : X \text{ a } D(H)\text{-module, such that } F(X) \text{ is projective as an } H\text{-module}\}.$$

**Lemma 2.3.3.** *The kernel of  $\overline{F}$  is a thick ideal of  $\mathrm{St}(\mathbf{Z}(\mathbf{C}))$ .*

*Proof.* Since  $\overline{F}$  is a monoidal triangulated functor, it is straightforward to verify that the collection of objects  $X$  such that  $\overline{F}(X) \cong 0$  is closed under taking cones, shifts, direct summands, and by tensoring on the left or right by arbitrary objects of  $\text{St}(\mathbf{Z}(\mathbf{C}))$ .  $\square$

**Lemma 2.3.4.** *An object  $A \in \text{st}(\mathbf{Z}(\mathbf{C}))$  satisfies  $W(A) = \emptyset$  if and only if  $A \in \mathbf{K}$ .*

*Proof.* First, note that if  $A \in \mathbf{K}$ , then by definition  $\overline{F}(A) \cong 0$ , and so

$$W(A) = V_{\mathbf{C}}(0) = \{\mathbf{P} \in \text{Spc}(\text{st}(\mathbf{C})) : 0 \notin \mathbf{P}\} = \emptyset.$$

For the other direction, recall that by the rigidity of  $\mathbf{C}$ , all thick ideals of  $\text{st}(\mathbf{C})$  are semiprime, i.e. intersections of prime ideals, by Proposition 1.1.1. This implies in particular that the ideal  $\langle 0 \rangle$  is semiprime; in other words, the only object contained in all prime ideals of  $\text{st}(\mathbf{C})$  is 0. By definition, this means that if  $X$  is an object of  $\text{st}(\mathbf{C})$  such that  $V_{\mathbf{C}}(X) = \emptyset$ , then  $X \cong 0$ . Hence, we have

$$\emptyset = W(A) = V_{\mathbf{C}}(\overline{F}(A)) \Rightarrow \overline{F}(A) \cong 0 \Rightarrow A \in \mathbf{K}.$$

$\square$

Using the localization and colocalization functors, we are now able to prove the following, which is the critical step in determining which ideals can be recovered from their  $W$ -support and determining the image of the map  $f : \text{Spc st}(\mathbf{C}) \rightarrow \text{Spc st}(\mathbf{Z}(\mathbf{C}))$ .

**Theorem 2.3.5.** *Let  $\mathbf{I}$  be a thick ideal of  $\text{st}(\mathbf{Z}(\mathbf{C}))$  such that  $\text{Loc}(\mathbf{I})$  contains  $\mathbf{K}$ . Suppose that  $X$  is an object of  $\text{st}(\mathbf{Z}(\mathbf{C}))$  such that  $\overline{F}(X) \in \langle \overline{F}(\mathbf{I}) \rangle$ , that is, the thick ideal of  $\text{st}(\mathbf{C})$  generated by all  $\overline{F}(Y)$  for  $Y \in \mathbf{I}$ . Then  $X$  is in  $\mathbf{I}$ .*

*Proof.* By Theorem 2.3.1, we have a distinguished triangle

$$\Gamma_{\mathbf{I}}(X) \rightarrow X \rightarrow L_{\mathbf{I}}(X) \rightarrow \Sigma \Gamma_{\mathbf{I}}(X)$$

in  $\text{St}(\mathbf{Z}(\mathbf{C}))$ , using the localization and colocalization functors associated to the thick ideal  $\mathbf{I}$ . We know that there are no morphisms from  $\mathbf{I}$  to  $L_{\mathbf{I}}(X)$ ; in other words, if  $Y \in \mathbf{I}$  and  $Z$  is any compact object in  $\text{St}(\mathbf{Z}(\mathbf{C}))$ , then

$$\begin{aligned} 0 &= \text{Hom}_{\text{St}(\mathbf{Z}(\mathbf{C}))}(Z \otimes Y, L_{\mathbf{I}}(X)) \\ &\cong \text{Hom}_{\text{St}(\mathbf{Z}(\mathbf{C}))}(Z, L_{\mathbf{I}}(X) \otimes Y^*). \end{aligned}$$

Since this holds for all compact objects  $Z$ , this implies that  $L_{\mathbf{I}}(X) \otimes Y^* \cong 0$ . Since all compact objects are rigid, and by Proposition 1.1.1 all thick ideals are closed under taking duals, we have  $L_{\mathbf{I}}(X) \otimes Y \cong 0$  for all  $Y \in \mathbf{I}$ . Since  $\overline{F}$  is a monoidal functor, this additionally implies that

$$\overline{F}(L_{\mathbf{I}}(X)) \otimes \overline{F}(Y) \cong 0$$

in  $\text{St}(\mathbf{C})$ , for all  $Y \in \mathbf{I}$ .

Now, consider the thick ideal  $\langle \overline{F}(\mathbf{I}) \rangle$ . This is formed successively by taking shifts, cones, direct summands, and tensor products with arbitrary elements of  $\text{st}(\mathbf{C})$ , starting from the collection of objects of the form  $\overline{F}(Y)$  for  $Y \in \mathbf{I}$ . This allows us to conclude inductively that  $\overline{F}(L_{\mathbf{I}}(X)) \otimes A \cong 0$  for all  $A$  in  $\langle \overline{F}(\mathbf{I}) \rangle$ , since inductively each step by which we construct  $\langle \overline{F}(\mathbf{I}) \rangle$  preserves the property that tensoring with  $\overline{F}(L_{\mathbf{I}}(X))$  gives 0. To be more explicit, if

$$A \rightarrow B \rightarrow C \rightarrow \Sigma A$$

is a distinguished triangle in  $\text{st}(\mathbf{C})$  such that  $A \otimes \overline{F}(L_{\mathbf{I}}(X)) \cong B \otimes \overline{F}(L_{\mathbf{I}}(X)) \cong 0$ , then it is straightforward that additionally  $C \otimes \overline{F}(L_{\mathbf{I}}(X)) \cong 0$  as well. Similarly, if  $A \otimes \overline{F}(L_{\mathbf{I}}(X)) \cong 0$ , then  $\Sigma(A) \otimes \overline{F}(L_{\mathbf{I}}(X)) \cong \Sigma(A \otimes \overline{F}(L_{\mathbf{I}}(X))) \cong \Sigma 0 \cong 0$ . Furthermore,

if we have  $(A \oplus B) \otimes \overline{F}(L_{\mathbf{I}}(X)) \cong 0$ , then we also have  $A \otimes \overline{F}(L_{\mathbf{I}}(X)) \cong 0 \cong B \otimes \overline{F}(L_{\mathbf{I}}(X))$ . Lastly, if  $A \otimes \overline{F}(L_{\mathbf{I}}(X)) \cong 0$  and  $B$  is an arbitrary object in  $\mathbf{st}(\mathbf{C})$ , then  $A \otimes B \otimes \overline{F}(L_{\mathbf{I}}(X)) \cong A \otimes \overline{F}(L_{\mathbf{I}}(X)) \otimes B \cong 0$  as well, using the commutativity of  $\overline{F}(L_{\mathbf{I}}(X))$ .

To reiterate, the upshot of the previous paragraph is that  $A \otimes \overline{F}(L_{\mathbf{I}}(X)) \cong 0$  for all  $A \in \langle \overline{F}(\mathbf{I}) \rangle$ . But by assumption, we have  $\overline{F}(X) \in \langle \overline{F}(\mathbf{I}) \rangle$ . Hence,

$$\overline{F}(X \otimes L_{\mathbf{I}}(X)) \cong \overline{F}(X) \otimes \overline{F}(L_{\mathbf{I}}(X)) \cong 0.$$

Therefore,  $X \otimes L_{\mathbf{I}}(X)$  is an object in  $\mathbf{K}$ , the collection of objects of  $\mathbf{St}(\mathbf{Z}(\mathbf{C}))$  mapped to 0 by  $\overline{F}$ . By assumption,  $\mathbf{Loc}(\mathbf{I})$  contains  $\mathbf{K}$ , and so  $X \otimes L_{\mathbf{I}}(X) \in \mathbf{Loc}(\mathbf{I})$ .

Now, consider the distinguished triangle obtained by tensoring the triangle

$$\Gamma_{\mathbf{I}}(X) \rightarrow X \rightarrow L_{\mathbf{I}}(X) \rightarrow \Sigma \Gamma_{\mathbf{I}}(X)$$

by  $X$ : this gives us

$$X \otimes \Gamma_{\mathbf{I}}(X) \rightarrow X \otimes X \rightarrow X \otimes L_{\mathbf{I}}(X) \rightarrow \Sigma X \otimes \Gamma_{\mathbf{I}}(X).$$

We have just finished showing that the third object of this triangle is in  $\mathbf{Loc}(\mathbf{I})$ . The first object is in  $\mathbf{Loc}(\mathbf{I})$  as well, by Theorem 2.3.1. Since  $\mathbf{Loc}(\mathbf{I})$  is triangulated, this implies  $X \otimes X$  is in  $\mathbf{Loc}(\mathbf{I})$ . But by [46, Lemma 2.2], since  $\mathbf{I}$  is a thick subcategory of compact objects, the compact objects in  $\mathbf{Loc}(\mathbf{I})$  are precisely the objects of  $\mathbf{I}$ . Thus,  $X \otimes X \in \mathbf{I}$ , and by semiprimeness of  $\mathbf{I}$  (Proposition 1.1.1) so is  $X$ ; this completes the proof.  $\square$

We can now give a condition under which an ideal  $\mathbf{I}$  can be recovered from its support  $\Phi_W(\mathbf{I})$ .

**Corollary 2.3.6.** *Let  $\mathbf{I}$  be an ideal such that  $\mathbf{Loc}(\mathbf{I})$  contains  $\mathbf{K}$ . Then  $\Theta_W \circ \Phi_W(\mathbf{I}) = \mathbf{I}$ .*

*Proof.* By definition,

$$\begin{aligned} \Theta_W \circ \Phi_W(\mathbf{I}) &= \Theta_W(\Phi_{\mathbf{C}}(\overline{F}(\mathbf{I}))) \\ &= \{X \in \mathbf{st}(\mathbf{Z}(\mathbf{C})) : W(X) \subseteq \Phi_{\mathbf{C}}(\overline{F}(\mathbf{I}))\} \\ &= \{X \in \mathbf{st}(\mathbf{Z}(\mathbf{C})) : V_{\mathbf{C}}(\overline{F}(X)) \subseteq \Phi_{\mathbf{C}}(\langle \overline{F}(\mathbf{I}) \rangle)\} \\ &= \{X \in \mathbf{st}(\mathbf{Z}(\mathbf{C})) : \forall \mathbf{P} \in \mathbf{Spc} \mathbf{st}(\mathbf{C}) \text{ with } \overline{F}(X) \notin \mathbf{P}, \langle \overline{F}(\mathbf{I}) \rangle \not\subseteq \mathbf{P}\} \\ &= \{X \in \mathbf{st}(\mathbf{Z}(\mathbf{C})) : \forall \mathbf{P} \in \mathbf{Spc} \mathbf{st}(\mathbf{C}) \text{ with } \langle \overline{F}(\mathbf{I}) \rangle \subseteq \mathbf{P}, \overline{F}(X) \in \mathbf{P}\} \\ &= \left\{ X \in \mathbf{st}(\mathbf{Z}(\mathbf{C})) : \overline{F}(X) \in \bigcap_{\mathbf{P} \in \mathbf{Spc} \mathbf{st}(\mathbf{C}), \langle \overline{F}(\mathbf{I}) \rangle \subseteq \mathbf{P}} \mathbf{P} \right\} \\ &= \{X \in \mathbf{st}(\mathbf{Z}(\mathbf{C})) : \overline{F}(X) \in \langle \overline{F}(\mathbf{I}) \rangle\}. \end{aligned}$$

The last equality follows from Proposition 1.1.1. The corollary now follows directly from Theorem 2.3.5.  $\square$

**2.4. The image of prime ideal contraction.** We now describe the relationship of the image of the map  $f$  to the kernel  $\mathbf{K}$  of  $\overline{F}$ . As in Section 2.3, we continue our assumption in this section that we are working with the category of modules for a finite-dimensional Hopf algebra, allowing us to apply Theorem 2.3.5.

**Proposition 2.4.1.** *Let  $\mathbf{C}$  be the category of modules of a finite-dimensional Hopf algebra  $H$ .*

- (1) *If  $\mathbf{P}$  is in the image of the map  $f : \mathbf{Spc} \mathbf{st}(\mathbf{C}) \rightarrow \mathbf{Spc} \mathbf{st}(\mathbf{Z}(\mathbf{C}))$ , then  $\mathbf{P}$  contains  $\mathbf{K} \cap \mathbf{st}(\mathbf{Z}(\mathbf{C}))$ , the kernel of  $\overline{F}$  restricted to compact objects.*

- (2) If  $\mathbf{P}$  is a prime ideal of  $\text{st}(\mathbf{Z}(\mathbf{C}))$  such that  $\text{Loc}(\mathbf{P})$  contains  $\mathbf{K}$ , then  $\mathbf{P}$  is in the image of  $f$ .

*Proof.* For (1), if  $\mathbf{Q}$  is a prime ideal of  $\text{st}(\mathbf{C})$ , then  $f(\mathbf{Q})$  contains  $\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C}))$ , which are by definition the finite-dimensional objects  $X$  such that  $\overline{F}(X) \cong 0$ : if  $X$  is in  $\text{st}(\mathbf{Z}(\mathbf{C}))$  and  $\overline{F}(X) \cong 0$ , then  $X \in \{Y : \overline{F}(Y) \in \mathbf{Q}\} = f(\mathbf{Q})$ , since 0 is in every prime ideal of  $\text{st}(\mathbf{C})$ .

Part (2) is an application of both Theorem 2.3.5 and [44, Theorem 3.2.3]. Let  $\mathbf{P}$  be a prime ideal of  $\text{st}(\mathbf{Z}(\mathbf{C}))$  such that  $\text{Loc}(\mathbf{P})$  contains  $\mathbf{K}$ . Consider the following two collections of objects in  $\text{st}(\mathbf{C})$ :

- (1) the ideal  $\mathbf{I} := \langle \overline{F}(X) : X \in \mathbf{P} \rangle$  of  $\text{st}(\mathbf{C})$ ;
- (2) the collection  $\mathcal{M} := \{\overline{F}(Y) : Y \notin \mathbf{P}\}$  of objects in  $\text{st}(\mathbf{C})$ .

We first claim that these two collections of objects are disjoint. If  $\overline{F}(Y) \in \mathbf{I}$  then  $Y \in \Theta_W(\Phi_W(\mathbf{P}))$ , implying that  $Y \in \mathbf{P}$  by Corollary 2.3.6. This means that in particular, if  $\overline{F}(X) \cong \overline{F}(Y)$ , then either both  $X$  and  $Y$  are in  $\mathbf{P}$ , or neither are, and so  $\mathbf{I}$  and  $\mathcal{M}$  are indeed disjoint.

Since  $\mathbf{P}$  is a proper ideal of  $\text{st}(\mathbf{Z}(\mathbf{C}))$ , it follows that  $\mathcal{M}$  is nonempty, and thus  $\mathbf{I}$  is a proper ideal of  $\text{st}(\mathbf{C})$ . We claim that  $\mathcal{M}$  is a multiplicative subset. Suppose  $\overline{F}(X)$  and  $\overline{F}(Y)$  are in  $\mathcal{M}$ . Then if  $\overline{F}(X) \otimes \overline{F}(Y) \cong \overline{F}(X \otimes Y)$  was not in  $\mathcal{M}$ , this would imply that  $X \otimes Y \in \mathbf{P}$ ; by the prime condition of  $\mathbf{P}$ , either  $X$  or  $Y$  (without loss of generality, say  $Y$ ) would then be in  $\mathbf{P}$ . This is a contradiction, since  $\overline{F}(Y) \in \mathcal{M}$  implies  $Y \notin \mathbf{P}$ , which is a consequence of the observation above that  $\mathbf{I}$  and  $\mathcal{M}$  are disjoint.

By [44, Theorem 3.2.3], given a disjoint pair consisting of a multiplicative subset and a proper ideal of any monoidal triangulated category (in this case,  $\text{st}(\mathbf{C})$ ), there exists a prime ideal  $\mathbf{Q}$  of  $\text{st}(\mathbf{C})$  such that  $\mathbf{Q} \cap \mathcal{M} = \emptyset$  and  $\mathbf{I} \subseteq \mathbf{Q}$ . We have

$$f(\mathbf{Q}) = \{X \in \text{st}(\mathbf{Z}(\mathbf{C})) : \overline{F}(X) \in \mathbf{Q}\},$$

and then since  $\mathbf{I} \subseteq \mathbf{Q}$ , it is automatic that  $\mathbf{P} \subseteq f(\mathbf{Q})$ ; and since  $\mathbf{Q}$  is disjoint from  $\mathcal{M}$ , in fact  $\mathbf{P} = f(\mathbf{Q})$ . Thus,  $f$  surjects onto the collection of prime ideals  $\mathbf{P}$  such that  $\text{Loc}(\mathbf{P})$  contains  $\mathbf{K}$ , which completes the proof.  $\square$

By Proposition 2.4.1, we have inclusions of the following subsets of  $\text{Spc st}(\mathbf{Z}(\mathbf{C}))$ :

$$(2.4.1) \quad \{\mathbf{P} : \mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C})) \subseteq \mathbf{P}\} \supseteq \text{im } f \supseteq \{\mathbf{P} : \mathbf{K} \subseteq \text{Loc}(\mathbf{P})\}.$$

We note the following lemma, which can be observed as a direct consequence of Theorem 2.3.1.

**Lemma 2.4.2.** *The following are equivalent.*

- (1) The kernel  $\mathbf{K}$  of  $\overline{F}$  is generated as a localizing category by the set  $\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C}))$ .
- (2) For every nonzero  $X$  in  $\mathbf{K}$ , there exists a finite-dimensional module  $Y$  in  $\mathbf{K}$  which has some nonzero map  $Y \rightarrow X$  in  $\text{St}(\mathbf{Z}(\mathbf{C}))$ .

If these conditions are satisfied, then we can sharpen (2.4.1), as well as Corollary 2.3.6.

**Corollary 2.4.3.** *Suppose the kernel  $\mathbf{K}$  of  $\overline{F}$  satisfies the equivalent conditions of Lemma 2.4.2.*

- (1) The image of  $f$  is precisely the collection of prime ideals of  $\text{st}(\mathbf{Z}(\mathbf{C}))$  which contain  $\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C}))$ , that is, the collection of finite-dimensional  $D(H)$ -modules  $X$  such that  $\overline{F}(X) \cong 0$ .
- (2) A thick ideal  $\mathbf{I}$  of  $\text{st}(\mathbf{Z}(\mathbf{C}))$  satisfies  $\Theta_W \circ \Phi_W(\mathbf{I}) = \mathbf{I}$  if and only if  $\mathbf{I}$  contains  $\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C}))$ .

*Proof.* Suppose  $\mathbf{K}$  satisfies the conditions of Lemma 2.4.2, in other words,  $\text{Loc}(\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C}))) = \mathbf{K}$ .

For (1), let  $\mathbf{P}$  be a prime ideal of  $\text{st}(\mathbf{Z}(\mathbf{C}))$  containing  $\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C}))$ . Then  $\text{Loc}(\mathbf{P})$  contains  $\text{Loc}(\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C}))) = \mathbf{K}$ . Hence the collection of inequalities of (2.4.1) becomes an equality, and we are done.

For (2), similarly, we have by Corollary 2.3.6 that if  $\text{Loc}(\mathbf{I})$  contains  $\mathbf{K}$ , then  $\Theta_W \circ \Phi_W(\mathbf{I}) = \mathbf{I}$ . Since  $\mathbf{K} = \text{Loc}(\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C})))$ , we have  $\mathbf{K} \subseteq \text{Loc}(\mathbf{I})$  if and only if  $\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C})) \subseteq \mathbf{I}$ . For the other direction, we note that for any ideal  $\mathbf{I}$ , we have  $\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C})) \subseteq \Theta_W \circ \Phi_W(\mathbf{I})$ , and so any thick ideal satisfying  $\Theta_W \circ \Phi_W(\mathbf{I}) = \mathbf{I}$  must have  $\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C})) \subseteq \mathbf{I}$  as well.  $\square$

**Remark 2.4.4.** Corollary 2.4.3 implies that if  $H$  satisfies the conditions of Lemma 2.4.2, then the image of  $f : \text{Spc st}(\mathbf{C}) \rightarrow \text{Spc st}(\mathbf{Z}(\mathbf{C}))$  is automatically the complement of a specialization-closed set, since we have

$$\begin{aligned} \text{im}(f) &= \{\mathbf{P} \in \text{Spc st}(\mathbf{Z}(\mathbf{C})) : \mathbf{P} \supseteq \mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C}))\} \\ &= (\Phi_{\mathbf{Z}}(\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C}))))^c. \end{aligned}$$

In other words, the image of  $f$  can be written as an intersection of open sets. If  $\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C}))$  is generated (as a thick ideal) by a finite collection of objects, say  $\{X_i\}_{i=1}^n$ , then it follows that  $\text{im}(f)$  is in fact an open subset of  $\text{Spc st}(\mathbf{Z}(\mathbf{C}))$ , namely

$$\text{im}(f) = (V_{\mathbf{Z}}(X_1 \oplus \dots \oplus X_n))^c.$$

**Remark 2.4.5.** In the situation of Corollary 2.4.3 (2), we have Corollary 2.3.6 sharpened from a one-way implication to a two-way implication. We note on the other hand that if the conditions of Lemma 2.4.2 are not satisfied, then Corollary 2.3.6 can never be an if-and-only-if, for the following reason. The collection of objects  $\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C}))$  is itself a thick ideal of  $\text{st}(\mathbf{Z}(\mathbf{C}))$ , since it is in particular the kernel of the monoidal triangulated functor  $\overline{F}$  restricted to compact objects. But now note that

$$\begin{aligned} \Theta_W \circ \Phi_W(\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C}))) &= \{X \in \text{st}(\mathbf{Z}(\mathbf{C})) : W(X) \subseteq \Phi_W(\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C})))\} \\ &= \{X \in \text{st}(\mathbf{Z}(\mathbf{C})) : W(X) \subseteq \emptyset\} \\ &= \{X \in \text{st}(\mathbf{Z}(\mathbf{C})) : \overline{F}(X) \cong 0\} \\ &= \mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C})), \end{aligned}$$

recall Lemma 2.3.4. In other words, the thick ideal  $\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C}))$  can be recovered from its support. But plainly, since we are assuming the conditions of Lemma 2.4.2 are not satisfied, we have

$$\text{Loc}(\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C}))) \not\supseteq \mathbf{K},$$

and so Corollary 2.3.6 cannot be sharpened to an if-and-only-if statement.

## 2.5. Conditions under which $f$ is injective, surjective, or a homeomorphism.

Under the same assumptions as the previous section, we now give conditions under which  $\Phi_W$  and  $\Theta_W$  are inverses, and  $f$  is surjective, injective, and a homeomorphism.

**Theorem 2.5.1.** *Suppose  $\mathbf{C} = \text{mod}(H)$  for a finite-dimensional Hopf algebra  $H$ .*

- (1) *The following conditions are equivalent.*
  - (a) *For all  $X \in \mathbf{K}$ , there exists an isomorphism  $X \cong 0$  in  $\text{St}(\mathbf{Z}(\mathbf{C}))$ .*
  - (b) *The map  $f$  is surjective and  $\mathbf{K}$  is generated as a localizing category by  $\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C}))$ .*
  - (c) *As maps  $\text{ThickId}(\text{st}(\mathbf{Z}(\mathbf{C}))) \rightarrow \text{ThickId}(\text{st}(\mathbf{Z}(\mathbf{C})))$ , we have  $\Lambda \circ \Psi = \text{id}$ .*

- (d) As maps  $\text{ThickId}(\mathbf{Z}(\mathbf{C})) \rightarrow \text{ThickId}(\mathbf{Z}(\mathbf{C}))$ , we have  $\Theta_W \circ \Phi_W = \text{id}$ .
- (2) If  $\mathbf{C}$  is braided, then the following hold.
- (a) The map  $f$  is injective.
  - (b) As maps  $\text{ThickId}(\text{st}(\mathbf{C})) \rightarrow \text{ThickId}(\text{st}(\mathbf{C}))$ , we have  $\Psi \circ \Lambda = \text{id}$ .
  - (c) If additionally  $\text{Spc st}(\mathbf{C})$  is topologically Noetherian, then  $\Phi_W \circ \Theta_W = \text{id}$ .
- (3) If  $X \cong 0$  in  $\text{St}(\mathbf{Z}(\mathbf{C}))$  for all  $X \in \mathbf{K}$  and  $\mathbf{C}$  is braided, then the following hold.
- (a)  $f$  is a homeomorphism.
  - (b) The maps  $\Psi$  and  $\Lambda$  define mutually inverse bijections between  $\text{ThickId}(\text{st}(\mathbf{Z}(\mathbf{C})))$  and  $\text{ThickId}(\text{st}(\mathbf{C}))$ .
  - (c) If additionally  $\text{Spc st}(\mathbf{C})$  is topologically Noetherian, then  $\Phi_W$  and  $\Theta_W$  are mutually inverse bijections between  $\text{ThickId}(\text{st}(\mathbf{Z}(\mathbf{C})))$  and  $\mathcal{X}_{sp}(\text{Spc}(\text{st}(\mathbf{C})))$ .

*Proof.* Suppose (1a) holds, and so  $\mathbf{K}$  consists only of objects isomorphic to 0, in other words, for all  $D(H)$ -modules  $X$ ,

$$F(X) \text{ is projective as an } H\text{-module} \Leftrightarrow X \text{ is projective as a } D(H)\text{-module.}$$

In particular this means that  $\mathbf{K}$  is generated by  $\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C}))$ , since all objects of  $\mathbf{K}$  are isomorphic to 0. Then (1c) follows from Theorem 2.3.5, and the conditions (1b) and (1d) follow directly from Corollary 2.4.3.

Now, suppose (1b) is satisfied. By Proposition 2.4.1, this means that every prime ideal of  $\text{st}(\mathbf{Z}(\mathbf{C}))$  contains  $\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C}))$ . But since every ideal is semiprime, the zero ideal is equal to the intersection of all primes of  $\text{st}(\mathbf{Z}(\mathbf{C}))$ , and so  $\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C}))$  is contained in the zero ideal. Since  $\mathbf{K}$  is generated by  $\mathbf{K} \cap \text{st}(\mathbf{Z}(\mathbf{C}))$ , i.e. the zero ideal, this implies that (1a) holds.

Note that

$$\begin{aligned} \Lambda(\Psi(\langle 0_{\text{st}(\mathbf{Z}(\mathbf{C}))} \rangle)) &= \Lambda(\langle 0_{\text{st}(\mathbf{C})} \rangle) \\ &= \{X \in \text{st}(\mathbf{Z}(\mathbf{C})) : \overline{F}(X) \in \langle 0_{\text{st}(\mathbf{C})} \rangle\} \\ &= \{X \in \text{st}(\mathbf{Z}(\mathbf{C})) : \overline{F}(X) \cong 0\} \\ &= \mathbf{K}. \end{aligned}$$

Hence (1c) implies (1a).

Lastly, suppose condition (1d) holds. This implies by Corollary 2.3.6 that  $\mathbf{K} \subseteq \text{Loc}(\mathbf{I})$  for every thick ideal  $\mathbf{I}$ ; in particular, this means that  $\mathbf{K}$  is contained in the localizing category generated by 0, which consists only of objects isomorphic to 0. Hence, (1a) holds.

To show (2), first note that if  $\mathbf{C}$  is braided with a braiding  $\gamma$ , then  $\overline{F}$  is essentially surjective, since for any object  $X$  in  $\mathbf{C}$ , the pair  $(X, \gamma_X)$  is an object of  $\mathbf{Z}(\mathbf{C})$  and  $\overline{F}$  sends

$H(X, \gamma_X)$  to  $G(X)$ . Now, we note that if  $\mathbf{P}$  and  $\mathbf{Q}$  are prime ideals of  $\text{st}(\mathbf{C})$ , then:

$$\begin{aligned}
f(\mathbf{P}) &= f(\mathbf{Q}) \\
&\Downarrow \\
\{X : \overline{F}(X) \in \mathbf{P}\} &= \{X : \overline{F}(X) \in \mathbf{Q}\} \\
&\Downarrow \\
\forall X \in \text{st}(\mathbf{Z}(\mathbf{C})), \overline{F}(X) \in \mathbf{P} &\Leftrightarrow \overline{F}(X) \in \mathbf{Q} \\
&\Downarrow \\
\forall Y \in \text{st}(\mathbf{C}), Y \in \mathbf{P} &\Leftrightarrow Y \in \mathbf{Q} \\
&\Downarrow \\
\mathbf{P} &= \mathbf{Q}.
\end{aligned}$$

Hence, if  $\mathbf{C}$  is braided then (2a) follows.

Condition (2b) also follows directly from the fact that  $\overline{F}$  is essentially surjective.

For (2c), note that by [4, Corollary 2.17],  $\text{Spc}(\text{st}(\mathbf{C}))$  is Noetherian if and only if every closed set is of the form  $V_{\mathbf{C}}(A)$  for some object  $A \in \text{st}(\mathbf{C})$ . If  $S$  is a specialization-closed set in  $\text{Spc}(\text{st}(\mathbf{C}))$ , then by definition

$$\begin{aligned}
\Phi_W(\Theta_W(S)) &= \Phi_W(\{X \in \text{st}(\mathbf{Z}(\mathbf{C})) : W(X) \subseteq S\}) \\
&= \bigcup_{X \in \Theta_W(S)} W(X) \\
&\subseteq S.
\end{aligned}$$

For the other direction, we can write  $S$  as a union of closed sets, say  $S = \bigcup_{i \in I} S_i$ , and by the Noetherianity of  $\text{Spc}(\text{st}(\mathbf{C}))$ , there exist objects  $A_i$  of  $\text{st}(\mathbf{C})$  such that  $S_i = V_{\mathbf{C}}(A_i)$ . Since  $\overline{F}$  is essentially surjective, we can pick  $X_i \in \text{st}(\mathbf{Z}(\mathbf{C}))$  with  $\overline{F}(X_i) = A_i$ . Since

$$W(X_i) = V_{\mathbf{C}}(A_i) = S_i \subseteq S,$$

we have by definition each  $X_i$  is in  $\Theta_W(S)$ . Therefore,

$$\begin{aligned}
\Phi_W(\Theta_W(S)) &\supseteq \bigcup_{i \in I} W(X_i) \\
&= \bigcup_{i \in I} S_i \\
&= S.
\end{aligned}$$

Thus  $S = \Phi_W(\Theta_W(S))$ .

Suppose the assumptions of (3). Then (3b) and (3c) follow immediately from parts (1) and (2). To show (3a), it is enough to show that  $f$  is a closed map, by (1a) and (2a). Take an arbitrary closed set  $V_{\mathbf{C}}(\mathcal{T})$  in  $\text{Spc}(\text{st}(\mathbf{C}))$ . We claim that the image of  $V_{\mathbf{C}}(\mathcal{T})$  under  $f$  is precisely  $V_{\mathbf{Z}}(\hat{\mathcal{T}})$ , where  $\hat{\mathcal{T}} = \{X \in \text{st}(\mathbf{Z}(\mathbf{C})) : \overline{F}(X) \in \mathcal{T}\}$ .

For the first direction, suppose  $\mathbf{P} \in V_{\mathbf{C}}(\mathcal{T})$ , in other words,  $\mathbf{P} \cap \mathcal{T} = \emptyset$ . Since  $f(\mathbf{P}) = \{X : \overline{F}(X) \in \mathbf{P}\}$ , this implies that for all  $X \in f(\mathbf{P})$ , we have  $X \notin \hat{\mathcal{T}}$ . Therefore  $f(\mathbf{P}) \cap \hat{\mathcal{T}} = \emptyset$ , and so  $f(\mathbf{P}) \in V_{\mathbf{Z}}(\hat{\mathcal{T}})$ . This shows  $f(V_{\mathbf{C}}(\mathcal{T})) \subseteq V_{\mathbf{Z}}(\hat{\mathcal{T}})$ .

For the other containment, suppose  $\mathbf{Q}$  is a prime ideal of  $\text{st}(\mathbf{Z}(\mathbf{C}))$  in  $V_{\mathbf{Z}}(\hat{\mathcal{T}})$ . Then  $\overline{F}(X) \notin \mathcal{T}$  for all  $X \in \mathbf{Q}$ . Since  $f$  is surjective, we can pick  $\mathbf{P} \in \text{Spc}(\text{st}(\mathbf{C}))$  with  $f(\mathbf{P}) = \mathbf{Q}$ , and for all  $\overline{F}(X) \in \mathbf{P}$ , we must have  $\overline{F}(X) \notin \mathcal{T}$ . Since  $\overline{F}$  is essentially surjective, this

implies  $A \notin \mathcal{T}$  for all  $A \in \mathbf{P}$ , and so  $\mathbf{P} \cap \mathcal{T} = \emptyset$ , i.e.  $\mathbf{P} \in V_{\mathbf{C}}(\mathcal{T})$ . This shows the other containment  $f(V_{\mathbf{C}}(\mathcal{T})) \supseteq V_{\mathbf{Z}}(\hat{\mathcal{T}})$ , and so we have equality.

Hence,  $f$  sends the closed set  $V_{\mathbf{C}}(\mathcal{T})$  to the closed set  $V_{\mathbf{Z}}(\hat{\mathcal{T}})$ , and so it is a continuous, bijective, closed map, and therefore a homeomorphism.  $\square$

### 3. APPLICATIONS

The time has come for concrete applications of our theory.

**3.1. Group algebras and dual group algebras.** Let  $G$  be a finite group,  $\mathbb{k}$  be a field of characteristic  $p$  which divides the order of  $G$ , and  $\mathbb{k}G$  the group algebra of  $G$  over  $\mathbb{k}$ . Let  $\mathbf{C} = \mathbf{mod}(\mathbb{k}G)$ , a finite tensor category. The Drinfeld double  $D(\mathbb{k}G)$  is a Hopf algebra containing  $\mathbb{k}G$  and  $(\mathbb{k}G^{\text{op}})^*$  as Hopf subalgebras. We will denote the dual of the group algebra by  $\mathbb{k}[G]$ , and in that case we can write  $(\mathbb{k}G^{\text{op}})^* = \mathbb{k}[G]^{\text{cop}}$ . The collection

$$\{p_g h : g, h \in G\}$$

is a  $\mathbb{k}$ -basis of  $D(\mathbb{k}G)$ , where the elements  $\{p_g : g \in G\}$  refer to the basis of  $\mathbb{k}[G]^{\text{cop}}$  dual to the standard basis of  $\mathbb{k}G$ . The multiplication is determined by the relations

$$hp_g = p_{hgh^{-1}}h,$$

see for instance [36, Section IX.4.3].

**Lemma 3.1.1.** *Let  $G$  and  $\mathbb{k}$  be as above and  $F : \mathbf{Mod}(D(\mathbb{k}G)) \rightarrow \mathbf{Mod}(\mathbb{k}G)$  the forgetful functor. Then if  $F(P)$  is projective as a  $\mathbb{k}G$ -module, then  $P$  is projective as a  $D(\mathbb{k}G)$ -module.*

*Proof.* A module for  $D(\mathbb{k}G)$  is a  $\mathbb{k}G$  module  $M$  which is also a  $G$ -graded vector space, such that if  $m \in M$  is a homogeneous element of degree  $g$ , then  $h.m$  is homogeneous of degree  $hgh^{-1}$ . Suppose we have a short exact sequence

$$0 \rightarrow A \rightarrow B \xrightarrow{t} C \rightarrow 0$$

of  $D(\mathbb{k}G)$ -modules such that

$$0 \rightarrow F(A) \rightarrow F(B) \rightarrow F(C) \rightarrow 0$$

is a split short exact sequence of  $G$ -modules. We claim that the original sequence splits as  $D(\mathbb{k}G)$ -modules. Pick a homogeneous basis  $\{c_i\}$  of  $C$  under the  $G$ -grading, where  $c_i$  has degree  $g_i$ . Now pick a splitting  $s : C \rightarrow B$ . Define  $\hat{s}(c_i) = p_{g_i}s(c_i)$ . This map is homogeneous with respect to the  $G$ -grading, and it is still a  $G$ -module map:

$$\begin{aligned} g\hat{s}(c_i) &= gp_{g_i}s(c_i) \\ &= p_{gg_i g^{-1}}gs(c_i) \\ &= p_{gg_i g^{-1}}s(gc_i) \\ &= \hat{s}(gc_i). \end{aligned}$$

Since on the basis  $\{c_i\}$  we have

$$\begin{aligned} t \circ \hat{s}(c_i) &= t(p_{g_i} \cdot s(c_i)) \\ &= p_{g_i} \cdot ts(c_i) \\ &= p_{g_i} \cdot c_i \\ &= c_i, \end{aligned}$$

we have that  $\hat{s}$  is a splitting of  $D(\mathbb{k}G)$ -modules.

Now, to prove the original claim, suppose  $F(P)$  is projective as a  $G$ -module. Since  $F$  is exact, this means that for every short exact sequence

$$0 \rightarrow A \rightarrow B \rightarrow P \rightarrow 0$$

in  $D(H)$ -modules, the sequence

$$0 \rightarrow F(A) \rightarrow F(B) \rightarrow F(P) \rightarrow 0$$

is split as  $G$ -modules. Therefore, the original sequences are all split, and so  $P$  is projective.  $\square$

We recall that by [4, Corollary 5.10],  $\mathrm{Spc\,stmod}(\mathbb{k}G) \cong \mathrm{Proj}\, \mathbf{H}^\bullet(G, \mathbb{k})$ , where  $\mathbf{H}^\bullet(G, \mathbb{k}) := \bigoplus_{i \geq 0} \mathrm{Ext}_{\mathbb{k}G}^i(\mathbb{k}, \mathbb{k})$  is the cohomology ring of  $G$ .

**Proposition 3.1.2.** *Let  $G$ ,  $\mathbb{k}$ , and  $\mathbf{H}^\bullet(G, \mathbb{k})$  be as above.*

- (1) *The map  $f : \mathrm{Spc\,stmod}(\mathbb{k}G) \rightarrow \mathrm{Spc\,stmod}(D(\mathbb{k}G))$  gives a homeomorphism, and so  $\mathrm{Spc\,stmod}(D(\mathbb{k}G)) \cong \mathrm{Spc\,stmod}(\mathbb{k}G) \cong \mathrm{Proj}\, \mathbf{H}^\bullet(G, \mathbb{k})$ .*
- (2) *Thick ideals of  $\mathrm{stmod}(D(\mathbb{k}G))$  are in bijection with specialization-closed sets in  $\mathrm{Proj}\, \mathbf{H}^\bullet(G, \mathbb{k})$ , which are in bijection with thick ideals of  $\mathrm{stmod}(\mathbb{k}G)$ , via the maps*

$$\mathrm{ThickId}(\mathrm{stmod}(D(\mathbb{k}G))) \begin{array}{c} \xrightarrow{\Phi_W} \\ \xleftarrow{\Theta_W} \end{array} \mathcal{X}_{sp}(\mathrm{Proj}\, \mathbf{H}^\bullet(G, \mathbb{k})) \begin{array}{c} \xrightarrow{\Theta_{\mathbb{k}G}} \\ \xleftarrow{\Phi_{\mathbb{k}G}} \end{array} \mathrm{ThickId}(\mathrm{stmod}(\mathbb{k}G)).$$

*Proof.* Since  $\mathbb{k}G$  is cocommutative,  $\mathrm{mod}(\mathbb{k}G)$  is braided symmetric. By Lemma 3.1.1, we have  $X \cong 0$  in  $\mathrm{StMod}(D(H))$  for all  $X \in \mathbf{K}$ , and so we are in the situation given of Theorem 2.5.1(3). Additionally, since cohomology rings of groups are finitely generated (for instance by the more general result of [27], in which finite generation of cohomology rings for finite-dimensional cocommutative Hopf algebras in positive characteristic was proven), we know that  $\mathrm{Proj}\, \mathbf{H}^\bullet(G, \mathbb{k})$  is a Noetherian topological space. Using Balmer's classification of thick ideals [4, Theorem 4.10], the thick ideals of  $\mathrm{stmod}(\mathbb{k}G)$  are in bijection with specialization-closed sets in  $\mathrm{Spc\,stmod}(\mathbb{k}G)$ . The rest of the theorem now follows directly as an application of Theorem 2.5.1.  $\square$

Now, note that since  $\mathbb{k}[G]^{\mathrm{cop}}$  is a semisimple algebra,  $\mathrm{stmod}(\mathbb{k}[G]^{\mathrm{cop}})$  consists only of the zero object, up to isomorphism, and so  $\mathrm{Spc}(\mathrm{stmod}(\mathbb{k}[G]^{\mathrm{cop}}))$  is the empty set. Thus, the diagram from Remark 2.1.3 becomes

$$\begin{array}{ccc} \mathrm{Spc\,stmod}(\mathbb{k}G) & \xleftarrow{\cong} & \mathrm{Proj}\, \mathbf{H}^\bullet(G, \mathbb{k}) & & \mathrm{Spc\,stmod}(\mathbb{k}[G]^{\mathrm{cop}}) = \emptyset \\ & \searrow \cong & & \swarrow & \\ & & \mathrm{Spc\,stmod}(D(\mathbb{k}G)) & & \end{array}$$

**3.2. Cosemisimple Hopf algebras.** In fact, we are able to generalize Lemma 3.1.1 and Proposition 3.1.2 from the group algebra case to the case certain finite-dimensional cosemisimple Hopf algebras. Recall that a finite-dimensional Hopf algebra is called *cosemisimple* if its Hopf dual is semisimple, as an algebra. There has been significant interest in the algebraic properties of cosemisimple Hopf algebras in the past few decades, see e.g. [39, 40, 22, 18, 19].

We first record the following straightforward lemma.

**Lemma 3.2.1.** *Let  $H$  be a finite-dimensional Hopf algebra such that  $\mathbf{1}_{D(H)}$  is a direct summand of  $D(H) \otimes_H \mathbf{1}_H$  as  $D(H)$ -modules, and let  $F : \mathrm{Mod}(D(H)) \rightarrow \mathrm{Mod}(H)$  be the forgetful functor. Then  $F(P)$  is projective in  $\mathrm{Mod}(H)$  if and only if  $P$  is projective in  $\mathrm{Mod}(D(H))$ .*

*Proof.* The functor  $D(H) \otimes_H -$  is a left adjoint to the forgetful functor  $F$ . Since  $F$  is exact, if  $Q$  is a projective  $H$ -module then

$$\mathrm{Hom}_H(Q, F(-)) \cong \mathrm{Hom}_{D(H)}(D(H) \otimes_H Q, -)$$

is an exact functor (recalling that projectives are also injective), and so  $D(H) \otimes_H -$  preserves projectivity. Therefore, if  $P$  is a  $D(H)$ -module such that  $F(P)$  is projective, then  $D(H) \otimes_H F(P)$  is a projective  $D(H)$ -module. But then, we have

$$D(H) \otimes_H F(P) \cong D(H) \otimes_H (\mathbf{1}_H \otimes_{\mathbb{k}} F(P)) \cong (D(H) \otimes_H \mathbf{1}_H) \otimes_{\mathbb{k}} P,$$

where the last isomorphism here can be seen from e.g. [28, Proposition 1.7] and the remark following it, which notes that although the proposition is stated for certain universal enveloping algebras, in fact the proof uses only the Hopf algebra structure, and so the result holds for arbitrary Hopf algebras. Note that it holds not just for finite-dimensional modules, but for arbitrary modules, which we need since in this case  $P$  may be infinite-dimensional.

Now, since  $\mathbf{1}_{D(H)}$  is a summand of  $D(H) \otimes_H \mathbf{1}_H$ , we have that  $P \cong \mathbf{1}_{D(H)} \otimes_{\mathbb{k}} P$  is a direct summand of  $(D(H) \otimes_H \mathbf{1}_H) \otimes_{\mathbb{k}} P$ , which is a projective  $D(H)$ -module, and hence  $P$  is projective as well, and the claim is proven.  $\square$

Recall that a Hopf algebra (or, more generally, a tensor category) is called *unimodular* if its spaces of left and right integrals coincide (cf. [42, Section 2.1], [23, Section 6.5]); unimodular Hopf algebras are of particular interest due to their use in constructing Hennings-Kaufman-Radford invariants for 3-manifolds [33, 37]. In light of Shimizu's result [55, Theorem 4.10] on unimodular finite tensor categories, if  $H$  satisfies the conditions of Lemma 3.2.1— that is, if  $\mathbf{1}_{D(H)}$  is a direct summand of  $D(H) \otimes_H \mathbf{1}_H$ — then  $H$  must be unimodular. The converse is not true; the dual of a finite group algebra is unimodular [55, Corollary 5.5], but  $\mathbf{1}_{D(\mathbb{k}[G])}$  is not a direct summand of  $D(\mathbb{k}[G]) \otimes_{\mathbb{k}G} \mathbf{1}_{\mathbb{k}[G]}$  (since  $F(\mathbf{1}_{D(\mathbb{k}[G])}) = \mathbf{1}_{\mathbb{k}[G]}$  is projective and  $\mathbf{1}_{D(\mathbb{k}[G])}$  is not).

**Corollary 3.2.2.** *Let  $H$  be a finite-dimensional unimodular cosemisimple Hopf algebra with Drinfeld double  $D(H)$  and forgetful functor  $F : \mathrm{Mod}(D(H)) \rightarrow \mathrm{Mod}(H)$ . Then  $F(P)$  is projective as an  $H$ -module if and only if  $P$  is projective as a  $D(H)$ -module.*

*Proof.* This follows from Lemma 3.2.1 and the proof of [23, Proposition 7.18.15]. In the course of the proof of the latter, it is shown that if  $H$  is unimodular and cosemisimple, then  $\mathbf{1}_{D(H)}$  is a direct summand of  $D(H) \otimes_H \mathbf{1}_H$  as  $D(H)$ -modules (note that here, we are reversing the roles of  $H$  and  $H^*$  given in their proof). Although this proposition assumes a stronger condition— that  $H$  itself is also semisimple— this assumption is not used for the part of the proof by which  $D(H) \otimes_H \mathbf{1}_H$  has  $\mathbf{1}_{D(H)}$  as a summand. By Lemma 3.2.1, the corollary follows.  $\square$

**Remark 3.2.3.** The condition that  $H$  is unimodular in Corollary 3.2.2 is not too restrictive. It is a long-standing conjecture of Kaplansky [35] that finite-dimensional cosemisimple Hopf algebras are involutory (e.g. the square of the antipode is the identity). In view of results of Larson [38, Corollary 4.2], a weaker form of the Kaplansky conjecture is that all finite-dimensional cosemisimple Hopf algebras are unimodular [2, Remark 3.9]. This conjecture is still open.

Corollary 3.2.2 and Theorem 2.5.1 now immediately imply the following.

**Proposition 3.2.4.** *Let  $H$  be a finite-dimensional unimodular cosemisimple Hopf algebra. Then the map  $f : \mathrm{Spcstmod}(H) \rightarrow \mathrm{Spcstmod}(D(H))$  constructed in Section 2.3 is*

surjective, and the maps  $\Lambda \circ \Psi$  and  $\Theta_W \circ \Phi_W$  (as in Section 2.2) are each the identity, as maps from the collection of thick ideals of  $\mathbf{stmod}(D(H))$  to itself.

Gelaki has shown [29, Theorem 1.3.6] that every quasitriangular cosemisimple Hopf algebra is unimodular. Hence, again by Corollary 3.2.2 and Theorem 2.5.1, we conclude:

**Proposition 3.2.5.** *Let  $H$  be a finite-dimensional quasitriangular cosemisimple Hopf algebra.*

- (1) *The map  $f$  constructed in Section 2.3 is a homeomorphism*

$$\mathrm{Spc\,stmod}(H) \xrightarrow{\cong} \mathrm{Spc\,stmod}(D(H)),$$

*and the maps  $\Psi$  and  $\Lambda$  as in Section 2.2 give inverse bijections between the thick ideals of  $\mathbf{stmod}(H)$  and  $\mathbf{stmod}(D(H))$ .*

- (2) *If  $\mathrm{Spc\,stmod}(H)$  is topologically Noetherian, then the  $\Phi_W$  and  $\Theta_W$  constructed in Section 2.2 are inverse maps, and so we have the following bijections of thick ideals:*

$$\mathrm{ThickId}(\mathbf{stmod}(D(H))) \begin{array}{c} \xrightarrow{\Phi_W} \\ \xleftarrow{\Theta_W} \end{array} \mathcal{X}_{sp}(\mathrm{Spc}(\mathbf{stmod}(H))) \begin{array}{c} \xrightarrow{\Theta_H} \\ \xleftarrow{\Phi_H} \end{array} \mathrm{ThickId}(\mathbf{stmod}(H)).$$

Of course, if  $H$  itself is also semisimple, then Proposition 3.2.4 and Proposition 3.2.5 are not particularly illuminating, since this implies that  $D(H)$  is also semisimple, and then the Balmer spectra of  $\mathbf{stmod}(H)$  and  $\mathbf{stmod}(D(H))$  are both  $\emptyset$ . It is a classical theorem of Larson-Radford [39] that in characteristic 0, all cosemisimple finite-dimensional Hopf algebras are also semisimple. Hence, Proposition 3.2.4 and Proposition 3.2.5 only provide interesting examples in positive characteristic.

**3.3. Benson-Witherspoon smash coproduct Hopf algebras.** We will now consider the Benson-Witherspoon smash coproducts which were originally studied in [12], with generalizations studied in [43] and [53]; their Balmer spectra and thick ideals were classified in [44]. We recall the general construction of these algebras. Let  $G$  and  $L$  be finite groups, such that  $L$  acts on  $G$  by group automorphisms, and let  $\mathbb{k}$  be a field of characteristic dividing the order of  $G$ . We then define  $H_{G,L}$  to be the Hopf algebra dual of the smash product  $\mathbb{k}[G] \# \mathbb{k}L$ , where  $\mathbb{k}[G]$  is the coordinate ring of  $G$ , and  $\mathbb{k}L$  is the group algebra of  $L$ .

As an algebra,  $H_{G,L}$  is isomorphic to  $\mathbb{k}G \otimes \mathbb{k}[L]$ . We will denote by  $\{p_x : x \in L\}$  the standard dual basis for  $\mathbb{k}[L]$ , as in Section 3.1. The additional Hopf algebra structures of comultiplication, counit, and antipode on  $A$  are defined by

$$\begin{aligned} \Delta(g \otimes p_x) &= \sum_{y \in L} (g \otimes p_y) \otimes (y^{-1} \cdot g \otimes p_{y^{-1}x}), \\ \epsilon(g \otimes p_x) &= \delta_{x,1}, \\ S(g \otimes p_x) &= x^{-1} \cdot (g^{-1}) \otimes p_{x^{-1}}, \end{aligned}$$

for all  $g \in G$  and  $x \in L$ .

Since as an algebra  $H_{G,L} \cong \mathbb{k}G \otimes \mathbb{k}[L]$ , an  $H_{G,L}$ -module is the same as a  $G$ -module with an  $L$ -grading, such that the action of  $G$  preserves the  $L$ -grading. In other words, every  $H_{G,L}$ -module  $M$  may be decomposed

$$M \cong \bigoplus_{x \in L} M_x \otimes \mathbb{k}_x$$

where  $M_x$  is a  $G$ -module, and  $\mathbb{k}_x$  is the 1-dimensional  $\mathbb{k}[L]$ -module on which  $p_x$  acts as the identity, and  $p_y$  acts as 0 for  $y \neq x$  (in other words, the  $\mathbb{k}[L]$ -module corresponding to a  $L$ -graded vector space of one dimension where every element is homogeneous of degree  $x$ ). The  $H_{G,L}$ -action on the component  $M_x \otimes \mathbb{k}_x$  is defined by letting  $\mathbb{k}G$  act on the first tensorand, and  $\mathbb{k}[L]$  act on the second.

Using the definition of the coproduct on  $H_{G,L}$ , Benson and Witherspoon [12, Theorem 2.1] compute the formula for the tensor product of  $H_{G,L}$ -modules:

$$(M_x \otimes \mathbb{k}_x) \otimes (N_y \otimes \mathbb{k}_y) = (M_x \otimes^x N_y) \otimes \mathbb{k}_{xy},$$

for any  $\mathbb{k}G$ -modules  $M_x$  and  $N_y$ , and for all  $x, y \in L$ , where the module  ${}^x N_y$  is defined as the twist of the module  $N_y$  by the action of  $x$ . Namely, this is the  $\mathbb{k}G$ -module which is equal to  $N_y$  as a vector space, and if we write  $g \cdot v$  for the action of  $G$  on the original module  $N_y$ , then the new action  $*$  of  $G$  on  ${}^x N_y$  is defined  $g * v = (x^{-1}g) \cdot v$ .

**Proposition 3.3.1.** *Let  $H_{G,L}$  the Benson-Witherspoon smash coproduct Hopf algebra as defined above,  $\mathbf{C}$  the category  $\text{mod}(H_{G,L})$ , and  $\mathbf{Z}(\mathbf{C})$  the category  $\text{mod}(D(H_{G,L}))$  for the Drinfeld double  $D(H_{G,L})$  of  $H_{G,L}$ .*

- (1) *The continuous map  $f : \text{Spcst}(\mathbf{C}) \rightarrow \text{Spcst}(\mathbf{Z}(\mathbf{C}))$  constructed in Section 2.3 is injective.*
- (2) *The map  $\Psi \circ \Lambda$  constructed in Section 2.2 is equal to the identity, as a map  $\text{ThickId}(\text{st}(\mathbf{C})) \rightarrow \text{ThickId}(\text{st}(\mathbf{C}))$ .*
- (3) *The map  $\Phi_W \circ \Theta_W$  constructed in Section 2.2 is equal to the identity, as a map  $\mathcal{X}_{sp}(\text{Spcst}(\mathbf{C})) \rightarrow \mathcal{X}_{sp}(\text{Spcst}(\mathbf{C}))$ .*

**Remark 3.3.2.** We note that if  $\mathbf{C}$  was braided, then Proposition 3.3.1 would follow directly from Theorem 2.5.1. However,  $H_{G,L}$  is not a quasitriangular Hopf algebra; the category of  $H_{G,L}$ -modules is not generally braided.

Proposition 3.3.1 will be proven by first showing the following intermediary lemma.

**Lemma 3.3.3.** *Suppose  $\mathbf{I}$  and  $\mathbf{J}$  are thick ideals of  $\text{st}(\mathbf{C})$  such that*

$$\{X \in \text{st}(\mathbf{Z}(\mathbf{C})) : \overline{F}(X) \in \mathbf{I}\} = \{X \in \text{st}(\mathbf{Z}(\mathbf{C})) : \overline{F}(X) \in \mathbf{J}\}.$$

*Then  $\mathbf{I} = \mathbf{J}$ . In particular, if  $M$  is an object of  $\text{st}(\mathbf{Z}(\mathbf{C}))$ , then there exists an object  $\hat{M}$  which is in the image of  $\overline{F}$ , and given any thick ideal  $\mathbf{I}$ , the object  $M$  is in  $\mathbf{I}$  if and only if  $\hat{M}$  is in  $\mathbf{I}$ .*

*Proof.* Suppose  $\mathbf{I}$  and  $\mathbf{J}$  are thick ideals satisfying the condition above. Since  $\mathbf{I}$  and  $\mathbf{J}$  are thick, it is enough to show that the indecomposable objects in  $\mathbf{I}$  are equal to the indecomposable objects in  $\mathbf{J}$ . Suppose  $M_x \otimes \mathbb{k}_x$  is an object in  $\mathbf{I}$ . Then the module

$$(M_x \otimes \mathbb{k}_x) \otimes (\mathbb{k} \otimes \mathbb{k}_{x^{-1}}) \cong M_x \otimes \mathbb{k}_{\text{id}}$$

is in  $\mathbf{I}$ . We also then have

$$(\mathbb{k} \otimes \mathbb{k}_y) \otimes (M_x \otimes \mathbb{k}_{\text{id}}) \otimes (\mathbb{k} \otimes \mathbb{k}_{y^{-1}}) \cong {}^y M_x \otimes \mathbb{k}_{\text{id}}$$

is an object of  $\mathbf{I}$  as well. The ideal  $\mathbf{I}$  then contains the direct sum

$$\hat{M} := \bigoplus_{y \in H} {}^y M_x \otimes \mathbb{k}_{\text{id}}.$$

We claim that  $\hat{M}$  is in the image of  $\overline{F}$ ; in other words,  $\hat{M}$  has a half-braiding which allows it to be lifted to the Drinfeld center. To see this, consider an  $H_{G,L}$ -module  $N_z \otimes \mathbb{k}_z$ . We

observe that

$$\begin{aligned}\hat{M} \otimes (N_z \otimes \mathbb{k}_z) &\cong \bigoplus_{y \in L} ({}^y M_x \otimes N_z) \otimes \mathbb{k}_z, \\ (N_z \otimes \mathbb{k}_z) \otimes \hat{M} &\cong \bigoplus_{y \in L} (N_z \otimes {}^{zy} M_x) \otimes \mathbb{k}_z.\end{aligned}$$

Since  $\mathbb{k}G$  is itself cocommutative (and thus  ${}^y M_x \otimes N_z \cong N_z \otimes {}^y M_x$  in a natural way), this formula can be used to observe a natural isomorphism  $\hat{M} \otimes - \cong - \otimes \hat{M}$ . This isomorphism satisfies the half-braiding condition, and so  $\hat{M}$  is in the image of  $\overline{F}$ .

Since **I** and **J** are assumed to agree on their intersections with the image of  $\overline{F}$ , we can conclude that  $\hat{M}$  is in **J** as well. But then its summand  $M_x \otimes \mathbb{k}_{\text{id}}$ , and hence

$$(M_x \otimes \mathbb{k}_{\text{id}}) \otimes (\mathbb{k} \otimes \mathbb{k}_x) \cong M_x \otimes \mathbb{k}_x,$$

is also an object of **J**. Note that we have proven generally that  $M_x \otimes \mathbb{k}_x$  is in any thick ideal if and only if  $\hat{M}$ , as constructed above, is in that ideal. Thus, the objects of **I** are a subset of the objects of **J**, and by symmetry the ideals are equal.  $\square$

We can now prove Proposition 3.3.1, as a consequence of Lemma 3.3.3:

*Proof.* The map  $f$  is defined by

$$f(\mathbf{P}) = \{X : \overline{F}(X) \in \mathbf{P}\}$$

for a given prime ideal  $\mathbf{P}$  in  $\text{Spcst}(\mathbf{C})$ . But Lemma 3.3.3 has shown that if  $\mathbf{P}$  and  $\mathbf{Q}$  are two prime ideals with  $f(\mathbf{P}) = f(\mathbf{Q})$ , then since  $\mathbf{P}$  and  $\mathbf{Q}$  are more generally examples of thick ideals, we have  $\mathbf{P} = \mathbf{Q}$ . Hence,  $f$  is injective, showing (1).

For (2), let  $S$  be an arbitrary specialization-closed set in  $\text{Spcst}(\mathbf{C})$ , in other words, a (possibly infinite) union  $S = \bigcup_{i \in I} S_i$  where each  $S_i$  is a closed set. Recall that by construction, it is automatic that  $\Phi_W(\Theta_W(S)) \subseteq S$  (the details are included above in the proof of Theorem 2.5.1).

To show the opposite containment, we note that by the classification of thick ideals and Balmer spectrum of  $\text{st}(\mathbf{C})$  as given in [44],  $\text{Spcst}(\mathbf{C})$  is a Noetherian topological space. We claim that this implies that every closed set in  $\text{Spcst}(\mathbf{C})$  has the form  $V_{\mathbf{C}}(M)$ , for some object  $M$  of  $\text{st}(\mathbf{C})$ , just as in the commutative setting [4, Corollary 2.17], using Lemma 3.3.3 as a substitute for the commutativity of the tensor product. Let  $V_{\mathbf{C}}(\mathcal{T})$  be an arbitrary closed set in  $\text{Spcst}(\mathbf{C})$ , for some collection  $\mathcal{T}$  of objects in  $\text{st}(\mathbf{C})$ . Then the complement

$$V_{\mathbf{C}}(\mathcal{T})^c = \{\mathbf{P} \in \text{Spcst}(\mathbf{C}) : \mathbf{P} \cap \mathcal{T} \neq \emptyset\}$$

has an open cover

$$V_{\mathbf{C}}(\mathcal{T})^c = \bigcup_{A \in \mathcal{T}} V_{\mathbf{C}}(A)^c = \bigcup_{A \in \mathcal{T}} \{\mathbf{P} \in \text{Spcst}(\mathbf{C}) : A \in \mathbf{P}\}.$$

By Noetherianity, this set is compact, and hence has a finite subcover

$$V_{\mathbf{C}}(\mathcal{T})^c = \bigcup_{A \in \mathcal{T}'} V_{\mathbf{C}}(A)^c,$$

where  $\mathcal{T}' \subseteq \mathcal{T}$  is some finite collection of objects. Enumerate the objects of  $\mathcal{T}'$  by  $A_1, \dots, A_n$ . Choose  $\hat{A}_1, \dots, \hat{A}_n$  as constructed in Lemma 3.3.3: they are in the image of  $\overline{F}$ ,

and for any thick ideal  $\mathbf{I}$ , we have  $A_j \in \mathbf{I}$  if and only if  $\hat{A}_j \in \mathbf{I}$ . Using this property, it is clear that  $V_{\mathbf{C}}(A_j) = V_{\mathbf{C}}(\hat{A}_j)$  for all  $j$ . Now we claim that

$$V_{\mathbf{C}}(\mathcal{T})^c = V_{\mathbf{C}}(A_1)^c \cup \dots \cup V_{\mathbf{C}}(A_n)^c = V_{\mathbf{C}}(\hat{A}_1)^c \cup \dots \cup V_{\mathbf{C}}(\hat{A}_n)^c = V_{\mathbf{C}}(\hat{A}_1 \otimes \dots \otimes \hat{A}_n)^c.$$

The last equality (more specifically, the containment  $\supseteq$ ) uses the fact that each  $\hat{A}_j$  is in the image of  $\overline{F}$ , and hence commutes with all objects of  $\text{st}(\mathbf{C})$  up to isomorphism, since this implies that

$$\hat{A}_1 \otimes \dots \otimes \hat{A}_n \in \mathbf{P} \Rightarrow A_j \in \mathbf{P} \text{ for some } j.$$

Our claim is now shown: every closed set in  $\text{Spc st}(\mathbf{C})$  is of the form  $V_{\mathbf{C}}(A)$  for some object  $A$ .

In particular, each of the closed sets  $S_i$ , for  $i \in I$ , can be written as  $V_{\mathbf{C}}(M_i)$  for some object  $M_i \in \text{st}(\mathbf{C})$ . As above, we can replace  $M_i$  by  $\hat{M}_i$ , which is in the image of  $\overline{F}$ , i.e. we can pick an object  $X_i$  in  $\text{st}(\mathbf{Z}(\mathbf{C}))$  with  $\overline{F}(X_i) = \hat{M}_i$ . Since

$$W(X_i) = V_{\mathbf{C}}(\overline{F}(X_i)) = V_{\mathbf{C}}(\hat{M}_i) = V_{\mathbf{C}}(M_i) = S_i \subseteq S,$$

we have  $X_i \in \Theta_W(S)$  by definition. Hence, we now have

$$\begin{aligned} \Phi_W(\Theta_W(S)) &\supseteq \bigcup_{i \in I} W(X_i) \\ &= \bigcup_{i \in I} S_i \\ &= S. \end{aligned}$$

Since we have both containments, we can conclude that  $\Phi_W(\Theta_W(S)) = S$  for any specialization-closed set  $S$  in  $\text{Spc st}(\mathbf{C})$ .  $\square$

We also note that if  $p$  does not divide the order of  $L$ , then we can apply the results of the previous section to obtain:

**Theorem 3.3.4.** *Let  $G, L, \mathbb{k}, H_{G,L}, \mathbf{C} = \text{mod}(H_{G,L})$ , and  $\mathbf{Z}(\mathbf{C}) = \text{mod}(D(H_{G,L}))$  be as above, and assume additionally that  $p$  does not divide the order of  $L$ . Then we have the following.*

- (1) *The map  $f$  constructed in Section 2.3 is a homeomorphism*

$$\text{Spc st}(\mathbf{C}) \xrightarrow{\cong} \text{Spc st}(\mathbf{Z}(\mathbf{C})).$$

- (2) *The maps  $\Phi_W$  and  $\Theta_W$  constructed in Section 2.2 are mutually inverse, and so we have the following bijections of thick ideals:*

$$\text{ThickId}(\text{stmod}(D(H_{G,L}))) \begin{array}{c} \xrightarrow{\Phi_W} \\ \xleftarrow{\Theta_W} \end{array} \mathcal{X}_{sp}(\text{Spc}(\text{stmod}(H_{G,L}))) \begin{array}{c} \xrightarrow{\Theta_{H_{G,L}}} \\ \xleftarrow{\Phi_{H_{G,L}}} \end{array} \text{ThickId}(\text{stmod}(H_{G,L})).$$

*Proof.* First, note that  $H_{G,L}$  is cosemisimple: its dual is the smash product  $\mathbb{k}[G] \# \mathbb{k}L$ . Since  $p$  does not divide the order of  $L$ , the group algebra  $\mathbb{k}L$  is semisimple, and by [20, Theorem 6], as the smash product of two semisimple algebras,  $\mathbb{k}[G] \# \mathbb{k}L$  is semisimple as well.

Next, we claim that  $H_{G,L}$  is unimodular. This can be observed directly, by noting that the element

$$h := \left( \sum_{g \in G} g \right) \otimes p_1$$

is both a left and a right integral in  $H_{G,L}$ .

By application of Proposition 3.3.1 and Proposition 3.2.4,  $f$  is bijective and the maps  $\Phi_W$  and  $\Theta_W$  are inverse bijections. To conclude the proof, we must just prove that  $f$  is closed, and hence a homeomorphism. This follows similarly to the proof of Theorem 2.5.1(3a), except that we must again use Lemma 3.3.3 as a substitute for commutativity of the tensor product. Let  $V_{\mathbf{C}}(M)$  an arbitrary closed set, and, just as before, we may assume (by replacing  $M$  with  $\hat{M}$  as in Lemma 3.3.3 if need be) that  $M$  is in the image of  $\overline{F}$ , and so we can pick  $X \in Z(\text{st}(\mathbf{C}))$  with  $\overline{F}(X) = M$ . We now have

$$\begin{aligned} f(V_{\mathbf{C}}(M)) &= \{f(\mathbf{P}) : \mathbf{P} \in \text{Spc st}(\mathbf{C}), M \notin \mathbf{P}\} \\ &= \{\mathbf{Q} \in \text{Spc st}(Z(\mathbf{C})) : X \notin \mathbf{Q}\} \\ &= V_Z(X). \end{aligned}$$

The second equality follows from the fact that  $f$  is bijective. Hence,  $f$  is closed, and the theorem is complete.  $\square$

#### REFERENCES

- [1] P. N. Achar, W. Hardesty, and S. Riche, *On the Humphreys conjecture on support varieties of tilting modules*, Transform. Groups **24** (2019), no. 3, 597–657.
- [2] E. Aljadeff, P. Etingof, S. Gelaki, and D. Nikshych, *On twisting of finite-dimensional Hopf algebras*, J. Algebra **256** (2002), no. 2, 484–501.
- [3] B. Bakalov and A. Kirillov Jr., *Lectures on tensor categories and modular functors*, University Lecture Series, 21. American Mathematical Society, Providence, RI, 2001.
- [4] P. Balmer, *The spectrum of prime ideals in tensor triangulated categories*, J. Reine Angew. Math. **588** (2005), 149–168.
- [5] P. Balmer, *Spectra, spectra, spectra–tensor triangular spectra versus Zariski spectra of endomorphism rings*, Algebr. Geom. Topol. **10** (2010), no. 3, 1521–1563.
- [6] P. Balmer, *A guide to tensor-triangular classification*, Handbook of Homotopy Theory. Chapman & Hall/CRC, 2019.
- [7] P. Balmer and B. Sanders, *The spectrum of the equivariant stable homotopy category of a finite group*, Invent. Math. **208** (2017), no. 1, 283–326.
- [8] C. P. Bendel, D. K. Nakano, B. J. Parshall, and C. Pillen, *Cohomology for quantum groups via the geometry of the nullcone*, Mem. Amer. Math. Soc. **229** (2014), no. 1077, x+93 pp.
- [9] D. J. Benson, J. F. Carlson, and J. Rickard, *Thick subcategories of the stable module category*, Fund. Math. **153** (1997), no. 1, 59–80.
- [10] D. J. Benson, S. B. Iyengar, and H. Krause, *Local cohomology and support for triangulated categories*, Ann. Sci. Éc. Norm. Supér. (4) **41** (2008), no. 4, 573–619.
- [11] D. J. Benson, S. B. Iyengar, and H. Krause, *Representations of finite groups: local cohomology and support*, Oberwolfach Seminars vol. 43, Birkhäuser/Springer Basel AG, Basel, 2012.
- [12] D. J. Benson and S. Witherspoon, *Examples of support varieties for Hopf algebras with noncommutative tensor products*, Arch. Math. (Basel) **102** (2014), no. 6, 512–520.
- [13] P. A. Bergh, J. Y. Plavnik, and S. Witherspoon, *Support varieties for finite tensor categories: Complexity, realization, and connectedness*, J. Pure Appl. Algebra **225** (2021), no. 9, 106705, 21 pp.
- [14] B. D. Boe, J. R. Kujawa, and D. K. Nakano, *Tensor triangular geometry for classical Lie superalgebras*, Adv. Math. **314** (2017), 228–277.
- [15] B. D. Boe, J. R. Kujawa, and D. K. Nakano, *Tensor triangular geometry for quantum groups*, arXiv:1702.01289.
- [16] A.B. Buan, H. Krause, and Ø. Solberg, *Support varieties—an ideal approach*, Homology, Homotopy Appl., **9** (2007), 45–74.
- [17] V. Chari and A. Pressley, *A guide to quantum groups*, Cambridge University Press, Cambridge, 1994. xvi+651 pp.
- [18] A. Chirvasitu, *Cosemisimple Hopf algebras are faithfully flat over Hopf subalgebras*, Algebra Number Theory **8** (2014), no. 5, 1179–1199.
- [19] A. Chirvasitu, C. Walton, and X. Wang, *Gelfand-Kirillov dimension of cosemisimple Hopf algebras*, Proc. Amer. Math. Soc. **147** (2019), no. 11, 4665–4672.
- [20] M. Cohen and D. Fishman, *Hopf algebra actions*, J. Algebra **100** (1986), no. 2, 363–379.

- [21] V. G. Drinfeld, *Quantum groups*, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Berkeley, Calif., 1986), 798–820, Amer. Math. Soc., Providence, RI, 1987.
- [22] P. Etingof and S. Gelaki, *On finite-dimensional semisimple and cosemisimple Hopf algebras in positive characteristic*, Internat. Math. Res. Notices 1998, no. 16, 851–864.
- [23] P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik, *Tensor categories*, Math. Surveys and Monographs vol. 205, Amer. Math. Soc., Providence, RI, 2015.
- [24] P. Etingof and V. Ostrik, *Finite tensor categories*, Mosc. Math. J. **4** (2004), no. 3, 627–654.
- [25] E. Friedlander and C. Negron, *Cohomology for Drinfeld doubles of some infinitesimal group schemes*, Algebra Number Theory **12** (2018), no. 5, 1281–1309.
- [26] E. M. Friedlander and J. Pevtsova,  $\Pi$ -supports for modules for finite group schemes, Duke Math. J. **139** (2007), no. 2, 317–368.
- [27] E. M. Friedlander and A. Suslin, *Cohomology of finite group schemes over a field*, Invent. Math. **127** (1997), no. 2, 209–270.
- [28] H. Garland and J. Lepowsky, *Lie algebra homology and the Macdonald-Kac formulas*, Invent. Math. **34** (1976), no. 1, 37–76.
- [29] S. Gelaki, *Quantum groups of dimension  $pq^2$* , Israel J. Math. **102** (1997), 227–267.
- [30] V. Ginzburg and S. Kumar, *Cohomology of quantum groups at roots of unity*, Duke Math. J. **69** (1993), no. 1, 179–198.
- [31] K. R. Goodearl and R. B. Warfield, Jr., *An introduction to noncommutative Noetherian rings*, Second edition. London Math. Soc. Student Texts vol. 61. Cambridge Univ. Press, Cambridge, 2004.
- [32] D. Happel, *Triangulated categories in the representation theory of finite-dimensional algebras*, London Mathematical Society Lecture Note Series, 119. Cambridge University Press, Cambridge, 1988.
- [33] M. Hennings, *Invariants of links and 3-manifolds obtained from Hopf algebras*, J. London Math. Soc. (2) **54** (1996), no. 3, 594–624.
- [34] M. J. Hopkins, *Global methods in homotopy theory*, Homotopy theory (Durham, 1985), 73–96, London Math. Soc. Lecture Note Ser., 117, Cambridge Univ. Press, Cambridge, 1987.
- [35] I. Kaplansky, *Bialgebras*, Lecture Notes in Mathematics. Department of Mathematics, University of Chicago, Chicago, Ill., 1975.
- [36] C. Kassel, *Quantum groups*, Graduate Texts in Mathematics, 155. Springer-Verlag, New York, 1995.
- [37] L. H. Kauffman and D. E. Radford, *Invariants of 3-manifolds derived from finite-dimensional Hopf algebras*, J. Knot Theory Ramifications **4** (1995), no. 1, 131–162.
- [38] R. G. Larson, *Characters of Hopf algebras*, J. Algebra **17** (1971), 352–368.
- [39] R. G. Larson and D. E. Radford, *Finite-dimensional cosemisimple Hopf algebras in characteristic 0 are semisimple*, J. Algebra **117** (1988), no. 2, 267–289.
- [40] R. G. Larson and D. E. Radford, *Semisimple cosemisimple Hopf algebras*, Amer. J. Math. **110** (1988), no. 1, 187–195.
- [41] H. Matsui and R. Takahashi, *Thick tensor ideals of right bounded derived categories*, Algebra Number Theory **11** (2017), no. 7, 1677–1738.
- [42] S. Montgomery, *Hopf algebras and their actions on rings*, CBMS Regional Conference Series in Mathematics, 82. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1993.
- [43] S. Montgomery, M. Vega, and S. Witherspoon, *Hopf automorphisms and twisted extensions*, J. Algebra Appl. **15** (2016), no. 6, 1650103, 14 pp.
- [44] D. K. Nakano, K. B. Vashaw, and M. T. Yakimov, *Noncommutative tensor triangular geometry*, arXiv:1909.04304v6, to appear in Amer. J. Math.
- [45] D. K. Nakano, K. B. Vashaw, and M. T. Yakimov, *Noncommutative tensor triangular geometry and the tensor product property for support maps*, arXiv:2008.09924, to appear in Int. Math. Res. Notices.
- [46] A. Neeman, *The connection between the K-theory localization theorem of Thomason, Trobaugh and Yao and the smashing subcategories of Bousfield and Ravenel*, Ann. Sci. École Norm. Sup. (4) **25** (1992), no. 5, 547–566.
- [47] A. Neeman, *Triangulated categories*, Annals of Mathematics Studies, 148. Princeton University Press, Princeton, NJ, 2001.
- [48] C. Negron, *Finite generation of cohomology for Drinfeld doubles of finite group schemes*, Selecta Math. (N.S.) **27** (2021), no. 2, Paper No. 26, 20 pp.
- [49] C. Negron and J. Pevtsova, *Support for integrable Hopf algebras via noncommutative hypersurfaces*, Int. Math Res. Not. IMRN. DOI: <https://doi.org/10.1093/imrn/rnab264>.

- [50] C. Negron and J. Y. Plavnik, *Cohomology of finite tensor categories: duality and Drinfeld centers*, Trans. Amer. Math. Soc. DOI: <https://doi.org/10.1090/tran/8548>.
- [51] W. D. Nichols and M. B. Zoeller, *A Hopf algebra freeness theorem*, Amer. J. Math. **111** (1989), no. 2, 381–385.
- [52] V. Ostrik, *Tensor ideals in the category of tilting modules*, Transform. Groups **2** (1997), no. 3, 279–287.
- [53] J. Y. Plavnik and S. Witherspoon, *Tensor products and support varieties for some noncocommutative Hopf algebras*, Algebr. Represent. Theory **21** (2018), no. 2, 259–276.
- [54] M. L. Reyes, *Obstructing extensions of the functor Spec to noncommutative rings*, Israel J. Math. **192** (2012), no. 2, 667–698.
- [55] K. Shimizu, *On unimodular finite tensor categories*, Int. Math. Res. Not. IMRN 2017, no. 1, 277–322.
- [56] R. W. Thomason, *The classification of triangulated subcategories*, Compositio Math. **105** (1997), no. 1, 1–27.

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139, U.S.A.

*Email address:* `kentv@mit.edu`