

Index of continuous semi-Fredholm families, regularities and semiregularities

M. Berkani

Abstract

In this paper, we define and index for continuous families of semi-Fredholm bounded linear operators. Moreover, we study various regularities and semiregularities of continuous families of bounded linear operators.

1 Introduction

In this part, we will introduce some definitions and notations that will be needed in the next sections.

In [7], the concept of regularity was defined by the following :

DEFINITION 1.1. Let A be a unital Banach algebra, with unit \mathbf{e} . A non-empty subset \mathbf{R} of A is called a regularity if it satisfies the following conditions:

- i) If $a \in A$ and $n \geq 1$, then $a \in \mathbf{R}$ if and only if $a^n \in \mathbf{R}$.
- ii) If $a, b, c, d \in L(X)$ are mutually commuting elements of A satisfying $ac + bd = \mathbf{e}$, then $ab \in \mathbf{R}$ if and only if $a, b \in \mathbf{R}$.

A regularity \mathbf{R} defines in a natural way a spectrum by $\sigma_{\mathbf{R}}(a) = \{\lambda \in \mathbb{C} : a - \lambda \mathbf{e} \notin \mathbf{R}\}$, for $a \in A$. Moreover the spectrum $\sigma_{\mathbf{R}}$ satisfies the spectral mapping theorem.

Let X be an infinite dimensional Banach space and $L(X)$ be the Banach algebra of bounded linear operators acting X . Various regularities of the Banach algebra $L(X)$ were studied in [7] and [1]. Here, we will consider similar regularities but for continuous families of bounded linear operators.

In the sequel if E and F are two vector spaces, the notation $E \simeq F$ will mean that E and F are isomorphic. If E and F are vector subspaces of the same vector space X , we will say that E is essentially included in F and write $E \subset_e F$, if there exist a finite dimensional vector subspace G of X such that $E \subset F + G$. We will say that E

2010 *Mathematics Subject Classification*: 47A53, 47A60

Key words and phrases: connected components, semi-Fredholm, homotopy, index, regularity, semiregularity,

is essentially equal to F and write $E =_e F$ if $E \subset_e F$ and $F \subset_e E$. We define also the infimum of the empty set to be ∞ .

For $T \in L(X)$, we will denote by $N(T)$ the null space of T , by $\alpha(T)$ the nullity of T , by $R(T)$ the range of T and by $\beta(T)$ its defect. If the range $R(T)$ of T is closed and $\alpha(T) < \infty$ (resp. $\beta(T) < \infty$), then T is called an upper semi-Fredholm (resp. a lower semi-Fredholm) operator. A semi-Fredholm operator is an upper or a lower semi-Fredholm operator. In the sequel $\Phi_+(X)$ (resp. $\Phi_-(X)$) will denote the set of upper (resp. lower) semi-Fredholm operators and $\mathfrak{s}\Phi(X) = \Phi_+(X) \cup \Phi_-(X)$ will denote the set of semi-Fredholm operators. If both of $\alpha(T)$ and $\beta(T)$ are finite, then T is called a Fredholm operator and the index of T is defined by $ind(T) = \alpha(T) - \beta(T)$; $\Phi(X)$ will denote the set of Fredholm operators. It is well known that $\Phi_+(X), \Phi_-(X), \mathfrak{s}\Phi(X)$ and $\Phi(X)$ are all open subset of $L(X)$, see for example [4, Chapter 4].

DEFINITION 1.2. Let $T \in L(X)$, $n \in \mathbb{N}$ and let $c_n(T) = \dim R(T^n)/R(T^{n+1})$. The descent of T is defined by $\delta(T) = \inf\{n : c_n(T) = 0\} = \inf\{n : R(T^n) = R(T^{n+1})\}$ and the essential descent of T is defined by $\delta_e(T) = \inf\{n : c_n(T) < \infty\} = \inf\{n : R(T^n) =_e R(T^{n+1})\}$.

DEFINITION 1.3. Let $T \in L(X)$, $n \in \mathbb{N}$ and let $c'_n(T) = \dim N(T^{n+1})/N(T^n)$. Then the ascent $a(T)$ of T is defined by $a(T) = \inf\{n : c'_n(T) = 0\} = \inf\{n : N(T^n) = N(T^{n+1})\}$ and the essential ascent of T is defined by $a_e(T) = \inf\{n : c'_n(T) < \infty\} = \inf\{n : N(T^{n+1}) =_e N(T^n)\}$.

DEFINITION 1.4. [5] : Let $T \in L(X)$. The sequence $(k_n(T))_n$ is defined by $k_n(T) = \dim[(R(T^n) \cap N(T))/(R(T^{n+1}) \cap N(T))]$, for all $n \in \mathbb{N}$.

DEFINITION 1.5. [5] Let $T \in L(X)$ and let $d \in \mathbb{N}$. Then T has a uniform descent for $n \geq d$ if $R(T) + N(T^n) = R(T) + N(T^d)$ for all $n \geq d$. If in addition $R(T) + N(T^d)$ is closed, then T is said to have a topological uniform descent for $n \geq d$.

Let \mathbb{X} be a compact topological space and let $\mathcal{C}(\mathbb{X}, L(X))$ be the algebra of continuous maps from the topological space \mathbb{X} into the Banach algebra $L(X)$. If $\mathcal{T} : \mathbb{X} \rightarrow L(X)$, we denote by \mathcal{T}_x the image $\mathcal{T}(x)$ of an element $x \in \mathbb{X}$.

Recall that $\mathcal{C}(\mathbb{X}, L(X))$ is a unital algebra with the usual properties of addition, scalar multiplication and multiplication defined by:

$$(\lambda\mathcal{S} + \mathcal{T})_x = \lambda\mathcal{S}_x + \mathcal{T}_x, (\mathcal{S}\mathcal{T})_x = \mathcal{S}_x\mathcal{T}_x, \forall(\mathcal{S}, \mathcal{T}) \in \mathcal{C}(\mathbb{X}, L(X))^2, \forall x \in \mathbb{X}, \forall \lambda \in \mathbb{C}.$$

The unit element \mathcal{I} of the algebra $\mathcal{C}(\mathbb{X}, L(X))$ is the map defined by: $\mathcal{I}_x = I$, for all $x \in \mathbb{X}$, where I is the identity operator. Moreover as \mathbb{X} is compact, if we set $\|\mathcal{T}\| = \sup_{x \in \mathbb{X}} \|\mathcal{T}_x\|, \forall \mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X))$, then $\mathcal{C}(\mathbb{X}, L(X))$ equipped with this norm is a unital Banach algebra.

2 Index of continuous semi-Fredholm families

In [2] and [3] respectively, we defined an analytical index for continuous Fredholm families of bounded linear operators into a Hilbert space (respectively into a Banach space). Here, in the same way as in [2] and [3], we define an analytical index for continuous semi-Fredholm families into a Banach space.

Let \mathbb{X} be a topological space. Define an equivalence relation \mathbf{c} on the space \mathbb{X} by setting that $x \mathbf{c} y$, if and only if x and y belongs to the same connected component of \mathbb{X} . Let \mathbb{X}^c be the quotient space associated to this equivalence relation. We extend here the definition [3, Definition 1.1] of the index of a continuous family of Fredholm operators to the case of a continuous family of semi-Fredholm operators.

DEFINITION 2.1. The index of a continuous family of semi-Fredholm operators $\mathcal{T} : \mathbb{X} \rightarrow L(X)$, parameterized by a topological space \mathbb{X} is defined by $ind(\mathcal{T}) = (ind(\mathcal{T}_x))_{\bar{x} \in \mathbb{X}^c}$.

Thus the index of a continuous family of semi-Fredholm operators \mathcal{T} is a sequence of elements of $\bar{\mathbb{Z}} = \mathbb{Z} \cup \{-\infty, +\infty\}$, which may be a finite sequence or infinite sequence, depending on the cardinal n_c of the connected components of \mathbb{X} , which is assumed here to have at most a countable set of connected components.

THEOREM 2.2. *The index of a continuous family of semi-Fredholm operators \mathcal{T} parameterized by a topological space \mathbb{X} is well defined as an element of $\bar{\mathbb{Z}}^{n_c}$. In particular if \mathbb{X} is reduced to a single element, then the index of \mathcal{T} is equal to the usual index of the semi-Fredholm operator \mathcal{T} .*

Proof. From the properties of the index [4, Chapter 4], it follows that two semi-Fredholm operators located in the same connected component of the set of semi-Fredholm operators have the same index. Moreover, as \mathcal{T} is continuous, the image of a connected component of the topological space \mathbb{X} , is included in a connected component of the set of semi-Fredholm operators. This shows that the index of a family of semi-Fredholm operators is well defined. Moreover, it is clear that if \mathbb{X} is reduced to a single element, the index of \mathcal{T} defined here is equal to the usual index of the single semi-Fredholm operator T .

DEFINITION 2.3. A continuous family \mathcal{K} from \mathbb{X} to $L(X)$ is said to be compact if \mathcal{K}_x is compact for all $x \in X$.

PROPOSITION 2.4. *i) Let $\mathcal{T} \in \mathcal{C}(\mathbb{X}, \mathbf{s}\Phi(X))$ be a continuous semi-Fredholm family and let \mathcal{K} be a continuous compact family. Then $\mathcal{T} + \mathcal{K}$ is a semi-Fredholm family and $ind(\mathcal{T}) = ind(\mathcal{T} + \mathcal{K})$.*

ii) Let $\mathcal{S}, \mathcal{T} \in \mathcal{C}(\mathbb{X}, \mathbf{s}\Phi(X))$ be two semi-Fredholm families, such that for all $x \in \mathbb{X}$, \mathcal{S}_x and \mathcal{T}_x are both upper semi-Fredholm or both lower semi-Fredholm. Then the family \mathcal{ST} defined by $(\mathcal{ST})_x = \mathcal{S}_x \mathcal{T}_x$ is a continuous semi-Fredholm family and $ind(\mathcal{ST}) = ind(\mathcal{S}) + ind(\mathcal{T})$.

Proof. This is clear from the usual properties of semi-Fredholm operators.

DEFINITION 2.5. Let \mathcal{S}, \mathcal{T} be in $\mathcal{C}(\mathbb{X}, \mathbf{s}\Phi(X))$. We will say that \mathcal{S} and \mathcal{T} are semi-Fredholm homotopic, if there exists a continuous map $h : [0, 1] \times X \rightarrow L(X)$ such that $h(0, x) = \mathcal{S}_x$, $h(1, x) = \mathcal{T}_x$ and for all $(t, x) \in [0, 1] \times \mathbb{X}$, $h(t, x)$ is a semi-Fredholm operator.

THEOREM 2.6. *Let \mathcal{S}, \mathcal{T} be two semi-Fredholm homotopic elements of $\mathcal{C}(\mathbb{X}, L(X))$. Then $\text{ind}(\mathcal{T}) = \text{ind}(\mathcal{S})$.*

Proof. Since \mathcal{S} and \mathcal{T} are semi-Fredholm homotopic, there exists a continuous map $h : \mathbb{X} \times [0, 1] \rightarrow \text{Fred}(X)$ such that $h(0, x) = \mathcal{S}(x)$ and $h(1, x) = \mathcal{T}(x)$ for all $x \in \mathbb{X}$. For a fixed $x \in \mathbb{X}$, the map $h_x : [0, 1] \rightarrow \mathbf{s}\Phi(X)$, defined by $h_x(t) = h(t, x)$ is a continuous path in $\mathbf{s}\Phi(X)$ linking \mathcal{S}_x to \mathcal{T}_x . Since $[0, 1]$ is connected and since the usual index is constant on connected sets of $\mathbf{s}\Phi(X)$, then $\text{ind}(\mathcal{S}_x) = \text{ind}(\mathcal{T}_x)$. Therefore $\text{ind}(\mathcal{S}) = \text{ind}(\mathcal{T})$.

THEOREM 2.7. *Let \mathbb{X} be a compact topological space. Then the set $\mathcal{C}(\mathbb{X}, \mathcal{SF}(X))$ of continuous semi-Fredholm families is an open subset of the Banach algebra $\mathcal{C}(\mathbb{X}, L(X))$ endowed with the uniform norm $\|\mathcal{T}\| = \sup_{x \in \mathbb{X}} \|\mathcal{T}_x\|$.*

Proof. Let $\mathcal{T} \in \mathcal{C}(\mathbb{X}, \mathbf{s}\Phi(X))$, then $\forall x \in \mathbb{X}, \exists \epsilon_x > 0$, such that $B(\mathcal{T}_x, \epsilon_x) \subset \mathbf{s}\Phi(X)$. We have $\mathbb{X} \subset \bigcup_{x \in \mathbb{X}} \mathcal{T}^{-1}(B(\mathcal{T}_x, \frac{\epsilon_x}{2}))$. Since \mathbb{X} is compact, there exists x_1, \dots, x_n in \mathbb{X} such that $\mathbb{X} \subset \bigcup_{i=1}^n \mathcal{T}^{-1}(B(\mathcal{T}_{x_i}, \frac{\epsilon_{x_i}}{2}))$. Let $\epsilon = \min\{\frac{\epsilon_{x_i}}{2} | 1 \leq i \leq n\}$ the minimum of the $\frac{\epsilon_{x_i}}{2}, 1 \leq i \leq n$. Let $\mathcal{S} \in \mathcal{C}(\mathbb{X}, L(X))$ such that $\|\mathcal{T} - \mathcal{S}\| < \frac{\epsilon}{2}$. If $x \in \mathbb{X}$, then $\|\mathcal{T}_x - \mathcal{S}_x\| < \frac{\epsilon}{2}$ and there exists $i, 1 \leq i \leq n$, such that $x \in \mathcal{T}^{-1}(B(\mathcal{T}_{x_i}, \epsilon_{x_i}))$. Then $\|\mathcal{S}_x - \mathcal{T}_{x_i}\| \leq \|\mathcal{S}_x - \mathcal{T}_x\| + \|\mathcal{T}_x - \mathcal{T}_{x_i}\| < \epsilon/2 + \epsilon_{x_i}/2 \leq \epsilon_{x_i}$ and so \mathcal{S}_x is a semi-Fredholm operator. Therefore $\mathcal{S} \in \mathcal{C}(\mathbb{X}, \mathbf{s}\Phi(X))$ and $\mathcal{C}(\mathbb{X}, \mathbf{s}\Phi(X))$ is open in $\mathcal{C}(\mathbb{X}, L(X))$.

THEOREM 2.8. *Let \mathbb{X} be a compact topological space. Then the index is a continuous locally constant function from $\mathcal{C}(\mathbb{X}, \mathbf{s}\Phi(X))$ into the group \mathbb{Z}^{nc} .*

Proof. As the set $\mathbf{s}\Phi(X)$ is an open subset of $L(X)$, if $\mathcal{T} \in \mathcal{C}(\mathbb{X}, \mathbf{s}\Phi(X))$, then $\forall x \in \mathbb{X}, \exists \epsilon_x > 0$, such that $B(\mathcal{T}_x, \epsilon_x) \subset \mathbf{s}\Phi(X)$. Then the index is constant on $B(\mathcal{T}_x, \epsilon_x)$, because $B(\mathcal{T}_x, \epsilon_x)$ is connected. We have $\mathbb{X} \subset \bigcup_{x \in \mathbb{X}} \mathcal{T}^{-1}(B(\mathcal{T}_x, \frac{\epsilon_x}{2}))$. Since \mathbb{X} is compact, there exists x_1, \dots, x_n in \mathbb{X} such that $\mathbb{X} \subset \bigcup_{i=1}^n \mathcal{T}^{-1}(B(\mathcal{T}_{x_i}, \frac{\epsilon_{x_i}}{2}))$. Let $\epsilon = \min\{\frac{\epsilon_{x_i}}{2} | 1 \leq i \leq n\}$ the minimum of the $\frac{\epsilon_{x_i}}{2}, 1 \leq i \leq n$, and let $\mathcal{S} \in \mathcal{C}(\mathbb{X}, \mathbf{s}\Phi(X))$ such that $\|\mathcal{T} - \mathcal{S}\| < \frac{\epsilon}{2}$. If $x \in \mathbb{X}$, then $\|\mathcal{T}_x - \mathcal{S}_x\| < \frac{\epsilon}{2}$ and there exists $i, 1 \leq i \leq n$, such that $x \in \mathcal{T}^{-1}(B(\mathcal{T}_{x_i}, \frac{\epsilon_{x_i}}{2}))$. Then $\|\mathcal{S}_x - \mathcal{T}_{x_i}\| \leq \|\mathcal{S}_x - \mathcal{T}_x\| + \|\mathcal{T}_x - \mathcal{T}_{x_i}\| < \epsilon/2 + \epsilon_{x_i}/2 \leq \epsilon_{x_i}$. So $\text{ind}(\mathcal{S}_x) = \text{ind}(\mathcal{T}_{x_i}) = \text{ind}(\mathcal{T}_x)$. Hence the index is a locally constant function, in particular it is a continuous function.

3 upper semiregularities of continuous families of bounded linear operators

In this part we consider various regularities of continuous families of bounded linear operators, parametrized by compact topological space \mathbb{X} . These regularities had been considered in [7] and [1], in the case when \mathbb{X} is reduced to a single element.

DEFINITION 3.1. Let \mathbb{X} be a compact topological space. We define the sets $\mathcal{R}_i, 1 \leq i \leq 16$, as follows:

$$\mathcal{R}_1 = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X)) \mid \forall x \in \mathbb{X}, \mathcal{T}_x \text{ is onto}\}$$

$$\mathcal{R}_2 = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X)) \mid \forall x \in \mathbb{X}, \mathcal{T}_x \in \Phi_-(X) \text{ and } \delta(\mathcal{T}_x) < \infty\}.$$

$$\mathcal{R}_3 = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X)) \mid \forall x \in \mathbb{X}, \delta(\mathcal{T}_x) < \infty \text{ and } R(\mathcal{T}_x^{\delta(\mathcal{T}_x)}) \text{ is closed}\}.$$

$$\mathcal{R}_4 = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X)) \mid \forall x \in \mathbb{X}, \mathcal{T}_x \in \Phi_-(X)\}.$$

$$\mathcal{R}_5 = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X)) \mid \forall x \in \mathbb{X}, \delta_e(\mathcal{T}_x) < \infty \text{ and } R(\mathcal{T}_x^{\delta_e(\mathcal{T}_x)}) \text{ is closed}\}.$$

$$\mathcal{R}_6 = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X)) \mid \forall x \in \mathbb{X}, \mathcal{T}_x \text{ is bounded below}\}.$$

$$\mathcal{R}_7 = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X)) \mid \forall x \in \mathbb{X}, \mathcal{T}_x \in \Phi_+(X) \text{ and } a(\mathcal{T}_x) < \infty\}.$$

$$\mathcal{R}_8 = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X)) \mid \forall x \in \mathbb{X}, a(\mathcal{T}_x) < \infty \text{ and } R(\mathcal{T}_x^{a(\mathcal{T}_x)+1}) \text{ is closed}\}.$$

$$\mathcal{R}_9 = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X)) \mid \forall x \in \mathbb{X}, \mathcal{T}_x \in \Phi_+(X)\}.$$

$$\mathcal{R}_{10} = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X)) \mid \forall x \in \mathbb{X}, a_e(\mathcal{T}_x) < \infty \text{ and } R(\mathcal{T}_x^{a_e(\mathcal{T}_x)+1}) \text{ is closed}\}.$$

$$\mathcal{R}_{11} = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X)) \mid \forall x \in \mathbb{X}, N(\mathcal{T}_x) \subset R^\infty(\mathcal{T}_x) \text{ and } R(\mathcal{T}_x) \text{ is closed}\}, \text{ where } R^\infty(\mathcal{T}_x) = \bigcap_{n \in \mathbb{N}} R(\mathcal{T}_x^n).$$

$$\mathcal{R}_{12} = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X)) \mid \forall x \in \mathbb{X}, N(\mathcal{T}_x) \subset_e R^\infty(\mathcal{T}_x) \text{ and } R(\mathcal{T}_x) \text{ is closed}\}.$$

$$\mathcal{R}_{13} = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X)) \mid \forall x \in \mathbb{X}, \exists d_x \in \mathbb{N} : R(\mathcal{T}_x) + N(\mathcal{T}_x^{d_x}) = R(\mathcal{T}_x) + N^\infty(\mathcal{T}_x) \text{ and } R(\mathcal{T}_x^{d_x+1}) \text{ is closed}\}, \text{ where } N^\infty(\mathcal{T}_x) = \bigcup_{n \in \mathbb{N}} N(\mathcal{T}_x^n).$$

$$\mathcal{R}_{14} = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X)) \mid \forall x \in \mathbb{X}, \forall n \in \mathbb{N}, k_n(\mathcal{T}_x) < \infty \text{ and } R(\mathcal{T}_x) \text{ is closed}\}.$$

$$\mathcal{R}_{15} = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X)) \mid \forall x \in \mathbb{X}, \exists d_x \in \mathbb{N} : k_n(\mathcal{T}_x) < \infty \text{ if } n \geq d_x \text{ and } R(\mathcal{T}_x^{d_x+1}) \text{ is closed}\}.$$

$$\mathcal{R}_{16} = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X)) \mid \forall x \in \mathbb{X}, \exists d_x \in \mathbb{N} : \mathcal{T}_x \text{ is of topological uniform descent for } n \geq d_x\}.$$

We have $\mathcal{R}_1 \subset \mathcal{R}_2 \subset \mathcal{R}_3 \subset \mathcal{R}_3 \cup \mathcal{R}_4 \subset \mathcal{R}_5 \subset \mathcal{R}_{13}$, $\mathcal{R}_6 \subset \mathcal{R}_7 \subset \mathcal{R}_8 \subset \mathcal{R}_8 \cup \mathcal{R}_9 \subset \mathcal{R}_{10} \subset \mathcal{R}_{13}$, $\mathcal{R}_{11} \subset \mathcal{R}_{12} \subset \mathcal{R}_{13} \subset \mathcal{R}_{13} \cup \mathcal{R}_{14} \subset \mathcal{R}_{15} \subset \mathcal{R}_{16}$. The elements of \mathcal{R}_{11} and \mathcal{R}_{12} are called semi-regular and essentially semi-regular continuous families. The elements of \mathcal{R}_{13} are called quasi-Fredholm continuous families and the elements of \mathcal{R}_{16} are called continuous families of topological uniform descent.

The inclusion $\mathcal{R}_5 \cup \mathcal{R}_{10} \subset \mathcal{R}_{13}$ is a consequence of [1, Proposition 3.4].

THEOREM 3.2. *The sets \mathcal{R}_i , for $1 \leq i \leq 16$, are regularities in the Banach algebra $\mathcal{C}(\mathbb{X}, L(X))$.*

Proof. From [7] and [1, Theorem 4.3], it follows that for each single element x of \mathbb{X} ,

the set $\mathcal{R}_{i,x} = \{\mathcal{T}_x \mid \mathcal{T} \in \mathcal{R}_i\}$, $1 \leq i \leq 16$ is a regularity. Thus \mathcal{R}_i , $1 \leq i \leq 16$, are regularities.

COROLLARY 3.3. *Let $\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X))$ and f an analytic function in a neighborhood of the usual spectrum $\sigma(\mathcal{T})$ of \mathcal{T} which is non-constant on any connected component of $\sigma(\mathcal{T})$. Then $f(\sigma_{\mathcal{R}_i}(\mathcal{T})) = \sigma_{\mathcal{R}_i}(f(\mathcal{T}))$, $1 \leq i \leq 16$.*

Proof. From [7] and [1, Theorem 4.3], it follows that for each single element x of \mathbb{X} , the set $\mathcal{R}_{i,x} = \{\mathcal{T}_x \mid \mathcal{T} \in \mathcal{R}_i\}$ for $1 \leq i \leq 16$, is a regularity. As $\sigma(\mathcal{T}) = \bigcup_{x \in \mathbb{X}} \sigma(\mathcal{T}_x)$, $1 \leq i \leq 16$, then for all $x \in \mathbb{X}$, f is an analytic function in a neighborhood of the usual spectrum $\sigma(\mathcal{T}_x)$ of x .

From [7], it follows that $f(\sigma_{\mathcal{R}_i}(\mathcal{T}_x)) = \sigma_{\mathcal{R}_i}(f(\mathcal{T}_x))$, $1 \leq i \leq 15$ and from [1, Theorem 4.3], it follows that $f(\sigma_{\mathcal{R}_{16}}(\mathcal{T}_x)) = \sigma_{\mathcal{R}_{16}}(f(\mathcal{T}_x))$, for all $x \in \mathbb{X}$. As $\sigma_{\mathcal{R}_i}(\mathcal{T}) = \bigcup_{x \in \mathbb{X}} \sigma_{\mathcal{R}_{i,x}}(\mathcal{T}_x)$, then $f(\sigma_{\mathcal{R}_i}(\mathcal{T})) = \sigma_{\mathcal{R}_i}(f(\mathcal{T}))$, $1 \leq i \leq 16$.

DEFINITION 3.4. A continuous family \mathcal{F} from \mathbb{X} to $L(X)$ is said to be of finite rank if \mathcal{F}_x is of finite rank for all $x \in \mathbb{X}$.

COROLLARY 3.5. *Let $\mathcal{T} \in \mathcal{R}_i$, for $i \in \{4, 5, 9, 10, 12, 13, 14, 15\}$. If \mathcal{F} is a continuous family of finite rank, then $\mathcal{T} + \mathcal{F} \in \mathcal{R}_i$.*

Proof. From [7], it follows that for each single element x of \mathbb{X} , the regularity $\mathcal{R}_{i,x}$, $i \in \{4, 5, 9, 10, 11, 12, 13, 14, 15\}$, is stable under the finite rank perturbation \mathcal{F}_x . Thus $\mathcal{T} + \mathcal{F} \in \mathcal{R}_i$.

REMARK 3.6. A continuous family \mathcal{T} in $\mathcal{C}(\mathbb{X}, L(X))$ is said to be a Fredholm family if for all $x \in \mathbb{X}$, \mathcal{T}_x is a Fredholm operator. We observe that the set $\Phi(X)$ of continuous Fredholm families is equal to $\mathcal{R}_4 \cap \mathcal{R}_9$, where \mathcal{R}_4 is the regularity of continuous families of upper semi-Fredholm operators and \mathcal{R}_9 is the regularity of continuous families of lower semi-Fredholm operators. Thus $\Phi(X)$ is a regularity and $\sigma_{\Phi(X)}(\mathcal{T}) = \sigma_{\mathcal{R}_4}(\mathcal{T}) \cup \sigma_{\mathcal{R}_9}(\mathcal{T})$.

Now, we extend some known results [4, Theorem 4.3.2, Theorem 4.3.3, Corollary 1.3.4, Corollary 1.3.5, Corollary 1.3.6] in the case of a single semi-Fredholm operators to the case of continuous families of semi-Fredholm operators. As their proofs are straightforwardly obtained from the case of a single operator, we give them without proof.

THEOREM 3.7. *1. Let $\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X))$. Then \mathcal{T} is in \mathcal{R}_4 if and only if there exists a family of bounded linear operators $(U_x)_{x \in \mathbb{X}}$, a family of bounded linear compact operators $(K_x)_{x \in \mathbb{X}}$, a family of bounded projections $(P_x)_{x \in \mathbb{X}}$, such that for all $x \in \mathbb{X}$, P_x is defined from X onto $N(\mathcal{T}_x)$ and $\mathcal{T}_x U_x = I + K_x$.*

2. Let $\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X))$. Then \mathcal{T} is in \mathcal{R}_9 if and only if there exists a family of bounded linear operators $(V_x)_{x \in \mathbb{X}}$, a family of bounded linear compact operators $(K_x)_{x \in \mathbb{X}}$, a family of bounded projections $(P_x)_{x \in \mathbb{X}}$, such that for all $x \in \mathbb{X}$, P_x is defined from X onto $R(\mathcal{T}_x)$ and $V_x \mathcal{T}_x = I + K_x$.
3. Let $\mathcal{T}_1, \mathcal{T}_2$ in $\mathcal{C}(\mathbb{X}, L(X))$ such that $\mathcal{T}_1 \mathcal{T}_2 \in \mathcal{R}_9$, then $\mathcal{T}_2 \in \mathcal{R}_9$.
4. Let $\mathcal{T}_1, \mathcal{T}_2$ in $\mathcal{C}(\mathbb{X}, L(X))$ such that $\mathcal{T}_1 \mathcal{T}_2 \in \mathcal{R}_4$, then $\mathcal{T}_1 \in \mathcal{R}_4$.
5. Let $\mathcal{T}_1, \mathcal{T}_2$ in $\mathcal{C}(\mathbb{X}, L(X))$ such that $\mathcal{T}_1 \mathcal{T}_2 \in \Phi(X)$, then $\mathcal{T}_1 \in \mathcal{R}_4$ and $\mathcal{T}_2 \in \mathcal{R}_9$.

4 Upper and lower semiregularities of continuous families of bounded linear operators

Following [8], the axioms i) and ii) of regularities in Definition 1.1, are splitted in two halves, each of them implying a one-way spectral mapping theorem and giving raises to the notions of upper and lower semiregularities. We first consider lower semiregularities.

In all what follows, \mathbb{X} will be a compact topological space. If \mathcal{R} is a nonempty subset of $\mathcal{C}(\mathbb{X}, L(X))$, then for all $x \in \mathbb{X}$, we set $\mathcal{R}_x = \{\mathcal{T}_x \mid \mathcal{T} \in \mathcal{R}\}, x \in \mathbb{X}$.

A- Upper semiregularities

DEFINITION 4.1. Let A be a unital Banach algebra, with unit \mathbf{e} . A non-empty subset \mathcal{R} of A is called a lower semiregularity if it satisfies the following conditions:

- i) If $a \in A$ and $a^n \in \mathbf{R}$, $n \geq 1$, then $a \in \mathcal{R}$.
- ii) If $a, b, c, d \in A$ are mutually commuting elements of A satisfying $ac + bd = \mathbf{e}$, such that $ab \in \mathcal{R}$, then $a, b \in \mathcal{R}$.

If \mathcal{R} is a lower semiregularity of A and $a \in A$, we define the \mathbf{R} -spectrum of a by $\sigma_{\mathbf{R}}(a) = \{\lambda \in \mathbf{C} / a - \lambda \mathbf{e} \notin \mathcal{R}\}$.

Recall the following result from [8, Lemma 2]

Lemma 4.1. *Let A be a unital Banach algebra with unit \mathbf{e} and let \mathcal{R} a lower semiregularity of A . then:*

1. $\mathbf{e} \in \mathcal{R}$.
2. $\text{Inv}(A) \subset \mathcal{R}$, where $\text{Inv}(A)$ is the set of invertible elements of A .
3. If $a \in \mathcal{R}, b \in \text{Inv}(A)$ and $ab = ba$, then $ab \in \mathcal{R}$.
4. $\sigma_{\mathcal{R}}(a) \subset \sigma(a)$.

5. (Translation property) $\sigma_{\mathcal{R}}(a + \lambda \mathbf{e}) = \lambda \mathbf{e} + \sigma_{\mathcal{R}}(a)$, for all $\lambda \in \mathbb{C}$.

Let \mathbb{X} be a compact topological space, let \mathcal{R} be an empty subset of $\mathcal{C}(\mathbb{X}, L(X))$. For all $x \in \mathbb{X}$, let $\mathcal{R}_x = \{\mathcal{T}_x \mid \mathcal{T} \in \mathcal{R}\}$, $x \in \mathbb{X}$. We assume that \mathcal{R} satisfies the following two conditions, summarized as condions (*):

- i) $\forall x \in \mathbb{X}, \forall \mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X)), \mathcal{T}_x \in \mathcal{R}_x \Rightarrow \mathcal{T} \in \mathcal{R}$.
- ii) $\forall x \in \mathbb{X}, \mathcal{R}_x$ is a lower semiregularity of $L(X)$.

Using the properties of lower semiregularities, we obtain a one way spectral mapping theorem.

THEOREM 4.2. *Let \mathbb{X} be a compact topological space and let \mathcal{R} be a nonempty subset of $\mathcal{C}(\mathbb{X}, L(X))$, satisfying the conditions (*). Then:*

1. \mathcal{R} is a lower semiregularity.
2. If f an analytic function in a neighborhood of the usual spectrum $\sigma(\mathcal{T})$ of an element $\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X))$, which is non-constant on any connected component of $\sigma(\mathcal{T})$, then $f(\sigma_{\mathcal{R}}(\mathcal{T})) \subset \sigma_{\mathcal{R}}(f(\mathcal{T}))$.

Proof. 1. Assume for all $x \in \mathbb{X}$, the set \mathcal{R}_x is a lower semiregularity of $L(X)$. Let us show that \mathcal{R} is a regularity.

For if $\mathcal{T} \in \mathcal{R}$ and $n \geq 1$ such that $\mathcal{T}^n \in \mathcal{R}$, then for all $x \in \mathbb{X}$, we have $(\mathcal{T}^n)_x = (\mathcal{T}_x)^n \in \mathcal{R}_x$. As \mathcal{R}_x is a lower semiregularity, then for all $x \in \mathbb{X}$, we have $\mathcal{T}_x \in \mathcal{R}_x$. Since \mathcal{R} satisfies the conditions (*), then $\mathcal{T} \in \mathcal{R}$.

Now if $\mathcal{S}, \mathcal{T}, \mathcal{U}, \mathcal{V}$, are commuting elements of \mathcal{R} satisfying $\mathcal{U}\mathcal{S} + \mathcal{V}\mathcal{T} = \mathcal{I}$, such that $\mathcal{S}\mathcal{T} \in \mathcal{R}$. Then for all $x \in \mathbb{X}$, we have $\mathcal{U}_x\mathcal{S}_x + \mathcal{V}_x\mathcal{T}_x = \mathcal{I}$ and $\mathcal{S}_x, \mathcal{T}_x, \mathcal{U}_x, \mathcal{V}_x$, are commuting elements of \mathcal{R}_x . As we have for all $x \in \mathbb{X}$, $\mathcal{S}_x\mathcal{T}_x = (\mathcal{S}\mathcal{T})_x \in \mathcal{R}_x$, and since \mathcal{R}_x is a lower semiregularity, then \mathcal{S}_x and $\mathcal{T}_x \in \mathcal{R}_x$. As \mathcal{R} satisfies conditions (*), then \mathcal{S} and \mathcal{T} are in \mathcal{R} . Consequently \mathcal{R} is a lower semiregularity in $\mathcal{C}(\mathbb{X}, L(X))$.

2. Let us show that $\sigma_{\mathcal{R}}(\mathcal{T}) = \bigcup_{x \in \mathbb{X}} \sigma_{\mathcal{R}_x}(\mathcal{T}_x)$. If $\lambda \notin \sigma_{\mathcal{R}}(\mathcal{T})$, then $\mathcal{T} - \lambda \mathcal{I} \in \mathcal{R}$. Therefore for all $x \in \mathbb{X}$, we have $\mathcal{T}_x - \lambda \mathcal{I} \in \mathcal{R}_x$. Hence $\lambda \notin \sigma_{\mathcal{R}_x}(\mathcal{T}_x)$. Conversely if for all $x \in \mathbb{X}, \lambda \notin \sigma_{\mathcal{R}_x}(\mathcal{T}_x)$, then for all $x \in \mathbb{X}, \mathcal{T}_x - \lambda \mathcal{I} \in \mathcal{R}_x$. As \mathcal{R} satisfies the conditions (*), then $\mathcal{T} - \lambda \mathcal{I} \in \mathcal{R}$. So $\sigma_{\mathcal{R}}(\mathcal{T}) = \bigcup_{x \in \mathbb{X}} \sigma_{\mathcal{R}_x}(\mathcal{T}_x)$.

As $\sigma(\mathcal{T}) = \bigcup_{x \in \mathbb{X}} \sigma(\mathcal{T}_x)$, then for all $x \in \mathbb{X}$, f is an analytic function in a neighborhood of the usual spectrum $\sigma(\mathcal{T}_x)$ of \mathcal{T}_x . From [8, Theorem 4], we have $f(\sigma_{\mathcal{R}_x}(\mathcal{T}_x)) \subset \sigma_{\mathcal{R}_x}(f(\mathcal{T}_x))$. Then $f(\sigma_{\mathcal{R}}(\mathcal{T})) = f(\bigcup_{x \in \mathbb{X}} \sigma_{\mathcal{R}_x}(\mathcal{T}_x)) = \bigcup_{x \in \mathbb{X}} f(\sigma_{\mathcal{R}_x}(\mathcal{T}_x)) \subset \bigcup_{x \in \mathbb{X}} \sigma_{\mathcal{R}_x}(f(\mathcal{T}_x)) = \sigma_{\mathcal{R}}(f(\mathcal{T}))$, because by the properties of the functional calculus, we have $f(\mathcal{T}_x) = f(\mathcal{T})_x$.

EXAMPLE 4.3. Let $(m_x)_{x \in \mathbb{X}}$ be a sequence in $\mathbb{N}^{\mathbb{X}}$. From [8] and Proposition 4.2, we deduce that the following sets are lower semiregularities of $\mathcal{C}(\mathbb{X}, L(X))$.

1. $\mathcal{LSR}_1 = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X)) : \forall x \in \mathbb{X}, \alpha(\mathcal{T}_x) \leq m_x\}$
2. $\mathcal{LSR}_2 = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X)) : \forall x \in \mathbb{X}, \beta(\mathcal{T}_x) \leq m_x\}$
3. $\mathcal{LSR}_3 = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X)) : \forall x \in \mathbb{X}, \alpha(\mathcal{T}_x) \leq m_x \text{ and } R(\mathcal{T}_x) \text{ is closed } \}$

B- Upper semiregularities

We assume that \mathcal{R} satisfies the following two conditions, summarized as conditions (**):

- i) $\forall x \in \mathbb{X}, \forall \mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X)), \mathcal{T}_x \in \mathcal{R}_x \Rightarrow \mathcal{T} \in \mathcal{R}$.
- ii) $\forall x \in \mathbb{X}, \mathcal{R}_x$ is an upper semiregularity of $L(X)$.

DEFINITION 4.4. Let A be a unital Banach algebra, with unit \mathbf{e} . A non-empty subset \mathcal{R} of A is called an upper semiregularity if it satisfies the following conditions:

- i) If $a \in A$ and $a \in \mathcal{R}$, then for all $n \geq 1$, then $a^n \in \mathcal{R}$.
- ii) If $a, b, c, d \in L(X)$ are mutually commuting elements of A satisfying $ac + bd = \mathbf{e}$, then if $a, b \in \mathcal{R}$, then $ab \in \mathcal{R}$.
- iii) \mathcal{R} contains a neighbourhood of the unit element \mathbf{e} .

If \mathcal{R} is a lower semiregularity of A and $a \in A$, we define the \mathcal{R} -spectrum of a by $\sigma_{\mathcal{R}}(a) = \{\lambda \in \mathbf{C}/a - \lambda \mathbf{e} \notin \mathcal{R}\}$.

Using the properties of upper semiregularities, we obtain a one way spectral mapping theorem. Here $Inv(\mathcal{C}(\mathbb{X}, L(X)))$ and $Inv(L(X))$, denote respectively the set of invertible elements of $\mathcal{C}(\mathbb{X}, L(X))$ and $L(X)$.

THEOREM 4.5. *Let \mathbb{X} be a compact topological space and let \mathcal{R} be a nonempty subset of $\mathcal{C}(\mathbb{X}, L(X))$, containing an open neighborhood of the identity \mathcal{I} and satisfying the conditions (**). Then:*

1. \mathcal{R} is an upper semiregularity of $\mathcal{C}(\mathbb{X}, L(X))$.
2. Assume that:

$$\forall x \in \mathbb{X}, \text{ if } \mathcal{T}_x \in \mathcal{R}_x \cap Inv(L(X)), \text{ then } (\mathcal{T}_x)^{-1} \in \mathcal{R}. (***)$$

Then if f an analytic function in a neighborhood of the usual spectrum $\sigma(\mathcal{T})$ of \mathcal{T} which is non-constant on any connected component of $\sigma(\mathcal{T})$, then $\sigma_{\mathcal{R}}(f(\mathcal{T})) \subset f(\sigma_{\mathcal{R}}(\mathcal{T}))$.

Proof.

1. Assume that for all $x \in \mathbb{X}$, the set \mathcal{R}_x is an upper semiregularity of $L(X)$. Let us show that \mathcal{R} is an upper regularity.

For if $\mathcal{T} \in \mathcal{R}$, then for all $x \in \mathbb{X}$, we have $\mathcal{T}_x \in \mathcal{R}_x$. As \mathcal{R}_x is an upper semiregularity, then for all $x \in \mathbb{X}$, we have $(\mathcal{T}_x)^n = (\mathcal{T}^n)_x \in \mathcal{R}_x$. Since \mathcal{R} satisfies the conditions (**), then $\mathcal{T}^n \in \mathcal{R}$.

Now if $\mathcal{S}, \mathcal{T}, \mathcal{U}, \mathcal{V}$, are commuting elements of \mathcal{R} satisfying $\mathcal{U}\mathcal{S} + \mathcal{V}\mathcal{T} = \mathcal{I}$, then for all $x \in \mathbb{X}$, we have $\mathcal{U}_x\mathcal{S}_x + \mathcal{V}_x\mathcal{T}_x = I$ and $\mathcal{S}_x, \mathcal{T}_x, \mathcal{U}_x, \mathcal{V}_x$, are commuting elements of \mathcal{R}_x . As \mathcal{R}_x is an upper semiregularity, then for all $x \in \mathbb{X}$, we have $\mathcal{S}_x\mathcal{T}_x = (\mathcal{S}\mathcal{T})_x \in \mathcal{R}_x$. Since \mathcal{R} satisfies the conditions (**), then $\mathcal{S}\mathcal{T} \in \mathcal{R}$.

As we know by hypothesis that \mathcal{R} contains an open neighborhood of the identity \mathcal{I} , then \mathcal{R} is an upper semiregularity in $\mathcal{C}(\mathbb{X}, L(X))$.

2. As for all $x \in \mathbb{X}$, \mathcal{R}_x is an upper semiregularity satisfying (***) , then from [8, Theorem 20], we have $\sigma_{\mathcal{R}_x}(f(\mathcal{T}_x)) \subset f(\sigma_{\mathcal{R}_x}(\mathcal{T}_x))$. Then $\sigma_{\mathcal{R}}(f(\mathcal{T})) = \bigcup_{x \in \mathbb{X}} \sigma_{\mathcal{R}_x}(f(\mathcal{T})_x) = \bigcup_{x \in \mathbb{X}} \sigma_{\mathcal{R}_x}(f(\mathcal{T}_x))$, because by the properties of the functional calculus, we have $f(\mathcal{T})_x = f(\mathcal{T}_x)$. So $\sigma_{\mathcal{R}}(f(\mathcal{T})) \subset \bigcup_{x \in \mathbb{X}} f(\sigma_{\mathcal{R}_x}(\mathcal{T}_x))$ and $\sigma_{\mathcal{R}}(f(\mathcal{T})) \subset f(\sigma_{\mathcal{R}}(\mathcal{T}))$, because $\bigcup_{x \in \mathbb{X}} \sigma_{\mathcal{R}_x}(\mathcal{T}_x) = \sigma_{\mathcal{R}}(\mathcal{T})$ (see the proof of Theorem 4.2) and $f(\bigcup_{x \in \mathbb{X}} \sigma_{\mathcal{R}_x}(\mathcal{T}_x)) = \bigcup_{x \in \mathbb{X}} f(\sigma_{\mathcal{R}_x}(\mathcal{T}_x))$.

EXAMPLE 4.6. Let $(m_x)_{x \in \mathbb{X}}$ be a sequence in $\mathbb{N}^{\mathbb{X}}$. From Theorem 4.4, we deduce that the following sets are upper semiregularities of $\mathcal{C}(\mathbb{X}, L(X))$.

1. $USR_1 = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, \Phi(X)) : \forall x \in \mathbb{X}, \text{ind}(\mathcal{T}_x) \geq 0\}$
2. $USR_2 = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, \Phi(X)) : \forall x \in \mathbb{X}, \text{ind}(\mathcal{T}_x) \leq 0\}$
3. $USR_3 = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, \Phi(X)) : \forall x \in \mathbb{X}, \text{ind}(\mathcal{T}_x) \in m_x\mathbb{Z}\}$
4. $USR_4 = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, \Phi(X)) : \forall x \in \mathbb{X}, \text{ind}(\mathcal{T}_x) = 0\}$
5. $USR_5 = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, \Phi_+(X)) : \forall x \in \mathbb{X}, \text{ind}(\mathcal{T}_x) \leq 0\}$
6. $USR_6 = \{\mathcal{T} \in \mathcal{C}(\mathbb{X}, \Phi_-(X)) : \forall x \in \mathbb{X}, \text{ind}(\mathcal{T}_x) \geq 0\}$

In fact $USR_4 = USR_1 \cap USR_2$, is the upper semiregularity of Weyl families (see [3] for more details about Weyl families).

In the case of a single operator, the spectra corresponding to the upper semiregularities USR_5 and USR_6 , were called respectively the essential approximate point spectrum and the essential defect spectrum. See [9] and the references cited there.

Open question: In [6, Theorem 6.5.2], it is proved that every Weyl operator T can be written as $T = V + F$, where V is an invertible operator and F a finite rank one.

Now let $\mathcal{T} \in \mathcal{C}(\mathbb{X}, L(X))$ be a Weyl family. Does there exists an invertible family $\mathcal{V} \in \mathcal{C}(\mathbb{X}, L(X))$ and a finite rank family $\mathcal{F} \in \mathcal{C}(\mathbb{X}, L(X))$ such that $\mathcal{T} = \mathcal{V} + \mathcal{F}$?

References

- [1] M. Berkani, *Restriction of an operator to the range of its powers*, Studia Mathematica, 140 (2) (2000), 163-174.
- [2] M. Berkani, *A new approach in index theory* , To appear in Mathematische Zeitschrift. <https://arxiv.org/abs/1912.07267>
- [3] M. Berkani, *Index of continuous families of bounded linear operators in Banach spaces and applications*. Submitted.
- [4] S.R. Caradus, Pfaffenberger W.E., B. Yood , *Calkin algebras and algebras of operators on banach spaces*, Marcel Dekker, Inc. New York, 1974.
- [5] S. Grabiner, *Uniform ascent and descent of bounded operators* ; J. Math. Soc. Japan 34 , no 2 , (1982), 317-337.
- [6] R. Harte, *Invertibility and singularity for bounded linear operators*. Marcel Dekker. New York, Basel, 1988.
- [7] M. Mbekhta ; V. Muller *On the axiomatic theory of the spectrum, II*; Studia - Math. 119 (1996), no . 2, 129 - 147.
- [8] V. Muller, *On the axiomatic theory of the spectrum III: Semiregularities*, Studia - Math. 142 (2000), no . 2, 159 - 169.
- [9] V. Rakočević, *Apostol spectrum and generalizations: A brief survey*, Facta Universatis, (Niš), Ser. Math. Inform. 14 (1999), 79-108.

Mohammed Berkani,
 Science faculty of Oujda,
 University Mohammed I,
 Laboratory LAGA,
 Morocco
 berkanimo@aim.com