

SPECTRALITY OF GENERALIZED SIERPINSKI-TYPE SELF-AFFINE MEASURES

JING-CHENG LIU, YING ZHANG, ZHI-YONG WANG AND MING-LIANG CHEN*

ABSTRACT. For an expanding integer matrix $M \in M_2(\mathbb{Z})$ and an integer digit set $D = \{(0, 0)^t, (\alpha_1, \alpha_2)^t, (\beta_1, \beta_2)^t\}$ with $\alpha_1\beta_2 - \alpha_2\beta_1 \neq 0$, let $\mu_{M,D}$ be the Sierpinski-type self-affine measure defined by $\mu_{M,D}(\cdot) = \frac{1}{3} \sum_{d \in D} \mu_{M,D}(M(\cdot) - d)$. In [5, 36], the authors separately investigated the spectral property of the measure $\mu_{M,D}$ in the case of $\det(M) \notin 3\mathbb{Z}$ or $\alpha_1\beta_2 - \alpha_2\beta_1 \notin 3\mathbb{Z}$. In this paper, we consider the remaining case where $\det(M) \in 3\mathbb{Z}$ and $\alpha_1\beta_2 - \alpha_2\beta_1 \in 3\mathbb{Z}$, and give the necessary and sufficient conditions for $\mu_{M,D}$ to be a spectral measure. This completely settles the spectrality of the Sierpinski-type self-affine measure $\mu_{M,D}$.

1. Introduction

Let μ be a Borel probability measure with compact support on \mathbb{R}^n . We call it a *spectral measure* if there exists a countable subset $\Lambda \subset \mathbb{R}^n$ such that the family of exponential functions $E(\Lambda) := \{e^{2\pi i(\lambda, x)} : \lambda \in \Lambda\}$ forms an orthonormal basis for $L^2(\mu)$. In this case, the set Λ is called a *spectrum* of μ , and we also say that (μ, Λ) is a *spectral pair*. In particular, if μ is the normalized Lebesgue measure supported on a Borel set Ω , then Ω is called a *spectral set*. The classical example of a spectral set is the unit cube $\Omega = [-\frac{1}{2}, \frac{1}{2}]^n$, for which the set $\Lambda = \mathbb{Z}^n$ serves as a spectrum.

Spectral theory for the Lebesgue measures on sets has been studied extensively since it initiated by Fuglede [19], whose famous conjecture asserted that Ω is a spectral set on \mathbb{R}^n if and only if Ω is a translational tile on \mathbb{R}^n . By *translational tile* we mean that there exists a set \mathcal{J} (called a *tiling set*) such that

$$\mathbb{R}^n = \bigcup_{t \in \mathcal{J}} (\Omega + t) \quad \text{and} \quad |(\Omega + t) \cap (\Omega + t')| = 0 \quad \text{for all } t \neq t' \in \mathcal{J},$$

Date: January 12, 2022.

2010 Mathematics Subject Classification. Primary 28A78, 28A80; Secondary 42C05, 46C05.

Key words and phrases. Sierpinski-type self-affine measure, Spectral measure, Spectrum, Hadamard triple.

The research is supported in part by the NNSF of China (Nos. 12071125, 12001183, 11831007 and 11971500), the Hunan Provincial NSF (Nos. 2019JJ20012 and 2020JJ5097), the SRF of Hunan Provincial Education Department (Nos.17B158 and 19B117).

*Corresponding author.

where $|\cdot|$ denotes the Lebesgue measure. Fuglede proved that this conjecture holds if the spectrum or the tiling set is assumed to be a lattice. The conjecture remains open for 30 years until Tao [39] gave the first counterexample: there exists a spectral subset of \mathbb{R}^n with $n \geq 5$ which is not a tile. After that, the counterexample was modified to show that the conjecture is false in both directions for $n \geq 3$, see [25, 26]. However, the conjecture is still open in low dimensions $n = 1, 2$. As of today there has been an effort to prove or disprove Fuglede's conjecture, one can refer to [4, 18, 23, 27] and the references therein.

After the original work of Fuglede, the study of spectral measures is also blooming in the fractal community. Recently, He, Lai and Lau [20] proved that a spectral measure μ must be of pure type, that is, μ is discrete with finite support, singularly continuous or absolutely continuous with respect to Lebesgue measure. For the absolutely continuous case, a general result was proved by Dutkay and Lai [11] that an absolutely continuous measure is a spectral one only if it is a normalized Lebesgue restricting on a domain $\Omega \subset \mathbb{R}^n$, which indeed involves the Fuglede's conjecture. For the singular continuous case, the first spectral measure was constructed by Jorgensen and Pedersen in 1998 [24]. They proved that the set $\Lambda = \{\sum_{k=0}^n 4^k d_k : d_k \in \{0, 1\}, n \in \mathbb{N}\}$ is a spectrum of the standard middle-fourth Cantor measure. In the same paper, they also showed that the usual middle-third Cantor measure is not spectral. Since then, many interesting spectral measures have been found, at the same time some singular phenomena different from the spectral theory of Lebesgue measures have been discovered, please see [1–3, 6–10, 12, 14–16, 22, 24, 28, 35, 36] and the references therein. Furthermore, the theory of spectral measures has been connected with many popular mathematical fields, such as the Fourier frame [20, 37], wavelet and Riesz basis [13] and so on.

Consider the iterated function system (IFS) on \mathbb{R}^n ,

$$\phi_d(x) = M^{-1}(x + d), \quad x \in \mathbb{R}^n, \quad d \in D,$$

where $M \in M_n(\mathbb{R})$ is an expanding real matrix (i.e., all eigenvalues of M have modulus strictly greater than one) and $D \subset \mathbb{R}^n$ is a finite digit set with cardinality $\#D$. It follows that there exists a unique nonempty compact subset $T(M, D) \subset \mathbb{R}^n$ satisfying $T(M, D) = \bigcup_{d \in D} \phi_d(T(M, D))$ (see [21] for details). Also there exists a unique probability measure $\mu_{M,D}$ supported on $T(M, D)$ such that

$$\mu_{M,D} = \frac{1}{\#D} \sum_{d \in D} \mu_{M,D} \circ \phi_d^{-1}. \quad (1.1)$$

We call $T(M, D)$ a *self-affine set* (or *attractor*) and $\mu_{M,D}$ a *Sierpinski-type self-affine measure*, respectively. In particular, if $M \in M_n(\mathbb{Z})$ and $D \subset \mathbb{Z}^n$, we call the $\mu_{M,D}$ an integral Sierpinski-type self-affine measure.

For a given measure $\mu_{M,D}$, the most interesting problem in this respect is to determine the spectrality or non-spectrality of $\mu_{M,D}$. And in the case when it is a spectral measure, one needs to determine the Fourier bases in the Hilbert space $L^2(\mu_{M,D})$. All these

are directly connected with the Fourier transform $\hat{\mu}_{M,D}$ (see (2.1)) of the measure $\mu_{M,D}$ simply due to the fact that $\mu_{M,D}$ is a spectral measure with a spectrum Λ if and only if $\sum_{\lambda \in \Lambda} |\hat{\mu}(\xi + \lambda)|^2 = 1$ for all $\xi \in \mathbb{R}^n$ (see [24, Lemma 3.3]). Moreover, the appearance of compatible pair (known also as Hadamard triple), following the terminology of [38], leads the research to an approachable way.

Definition 1.1. *Let $M \in M_n(\mathbb{Z})$ be an $n \times n$ expanding integer matrix, and let $D, S \subset \mathbb{Z}^n$ be two finite digit sets with the same cardinality. We say that (M, D) is admissible (or $(M^{-1}D, S)$ forms a compatible pair or (M, D, S) forms a Hadamard triple) if the matrix*

$$H = \frac{1}{\sqrt{\#D}} \left[e^{2\pi i \langle M^{-1}d, s \rangle} \right]_{d \in D, s \in S}$$

*is unitary, i.e., $H^*H = I$, where H^* denotes the conjugate transposed matrix of H .*

Given a discrete set $A \subset \mathbb{R}^n$, we define the discrete measure on A by $\delta_A = \frac{1}{\#A} \sum_{a \in A} \delta_a$, where δ_a is the Dirac measure at the point a . It is easy to see that the discrete measure $\delta_{M^{-1}D}$ is a spectral measure if and only if (M, D) is admissible. The well-known result of Jorgensen and Pedersen [24] shows that if $(M^{-1}D, S)$ is a compatible pair, then $E(\Lambda(M, S))$ is an infinite orthogonal system in $L^2(\mu_{M,D})$, where

$$\Lambda(M, S) = \left\{ \sum_{j=0}^{k-1} M^{*j} s_j : k \geq 1, s_j \in S \right\}.$$

Moreover, Dutkay and Jorgensen [16, Conjecture 2.5] [17, Conjecture 1.1] formulated the following well-known conjecture:

Conjecture. *The integral Sierpinski-type self-affine measure $\mu_{M,D}$ is a spectral measure if (M, D) is admissible.*

The conjecture implies that the existence of a set S such that (M, D, S) is a Hadamard triple is sufficient to obtain orthonormal bases of exponential functions in $L^2(\mu_{M,D})$. It was first proved on \mathbb{R} by Łaba and Wang [28] and later refined in [14]. The situation becomes more complicated when $n > 1$. In the paper [16], the authors showed that the conjecture is true if (M, D) satisfies a technical condition called the *reducibility condition*. The conjecture is true under some additional assumptions, introduced by Strichartz [38]. Some high dimensional special cases were also considered by Li [32, 33]. Eventually, Dutkay, Haussermann and Lai [12] proved that this conjecture is true, Hadamard triples always generate self-affine spectral measures.

Theorem 1.2. *[12, Theorem 1.3] The integral Sierpinski-type self-affine measure $\mu_{M,D}$ is a spectral measure if (M, D) is admissible.*

In particular, for the integral self-affine measure $\mu_{M,D}$ generated by an expanding matrix $M \in M_2(\mathbb{Z})$ and the digit set $D = \{(0, 0)^t, (1, 0)^t, (0, 1)^t\}$, there are many papers studied the spectrality and non-spectrality of it [15, 29, 30, 34]. Finally, the spectrality of $\mu_{M,D}$ has

been completely characterized by An, He and Tao [3], who showed that $\mu_{M,D}$ is a spectral measure if and only if (M, D) is admissible. For the general integer digit set

$$D = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \alpha_1 \\ \alpha_2 \end{pmatrix}, \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix} \right\} \quad (1.2)$$

with $\alpha_1\beta_2 - \alpha_2\beta_1 \neq 0$, to the best of our knowledge, Chen and Liu [5] first proved that if $\det(M) \notin 3\mathbb{Z}$, then the mutually orthogonal exponential functions in $L^2(\mu_{M,D})$ is finite. Recently, Liu and Wang [36] further considered the case $\alpha_1\beta_2 - \alpha_2\beta_1 \notin 3\mathbb{Z}$, and gave the necessary and sufficient conditions for $\mu_{M,D}$ to be a spectral measure. In order to characterize the spectral property of $\mu_{M,D}$, they introduced the general linear group. For a prime p , let $\mathbb{F}_p = \mathbb{Z}/p\mathbb{Z}$ be the residue class fields. All nonsingular $n \times n$ matrices over \mathbb{F}_p form a finite group under matrix multiplication, which is called the *general linear group* $GL_n(p)$. The following is their main result.

Theorem 1.3. [36, Theorem 1.10] *Let $\mu_{M,D}$ be defined by (1.1), where $M \in M_2(\mathbb{Z})$ is an expanding matrix and the digit set D is given by (1.2) with $\alpha_1\beta_2 - \alpha_2\beta_1 \notin 3\mathbb{Z}$. Then the following statements are equivalent.*

- (i) $\mu_{M,D}$ is a spectral measure.
- (ii) (M, D) is admissible.
- (iii) $(AMB)^*(1, -1)^t \in 3\mathbb{Z}^2$, where $B = \begin{bmatrix} \alpha_1 & \beta_1 \\ \alpha_2 & \beta_2 \end{bmatrix}$ and $A \in GL_2(3)$ satisfies $AB = I \pmod{M_2(3\mathbb{Z})}$.

As stated above, for an expanding matrix $M \in M_2(\mathbb{Z})$ and the digit set D given by (1.2), the spectral properties of the corresponding measure $\mu_{M,D}$ in the case of $\det(M) \notin 3\mathbb{Z}$ or $\alpha_1\beta_2 - \alpha_2\beta_1 \notin 3\mathbb{Z}$ have been completely characterized. A natural subsequent question is that

(Qu 1): For an expanding matrix $M \in M_2(\mathbb{Z})$ with $\det(M) \in 3\mathbb{Z}$ and the digit set D given by (1.2) with $\alpha_1\beta_2 - \alpha_2\beta_1 \in 3\mathbb{Z}$, what is the sufficient and necessary condition for $\mu_{M,D}$ to be a spectral measure? Is it (M, D) admissible?

In this paper, our main purpose is to initiate a study on the question **(Qu 1)**. Without loss of generality, we can assume that $\gcd(\alpha_1, \alpha_2, \beta_1, \beta_2) = 1$ by Lemma 2.1. The main result in the paper is the following:

Theorem 1.4. *Let $\mu_{M,D}$ be defined by (1.1), where $M \in M_2(\mathbb{Z})$ is an expanding matrix with $\det(M) \in 3\mathbb{Z}$ and the digit set D is given by (1.2) with $\alpha_1\beta_2 - \alpha_2\beta_1 \in 3\mathbb{Z}$. Then $\mu_{M,D}$ is a spectral measure if and only if there exists a matrix $Q \in M_2(\mathbb{Z})$ such that (M', D') is admissible, where $M' = QMQ^{-1}$ and $D' = QD$. In particular, if $2\alpha_1 - \beta_1, 2\alpha_2 - \beta_2 \in 3\mathbb{Z}$, then $\mu_{M,D}$ is a spectral measure if and only if $L^2(\mu_{M,D})$ contains an infinite orthogonal set of exponential functions.*

We remark that Theorem 1.4 settles completely the spectrality question **(Qu 1)** for the planar self-affine measure $\mu_{M,D}$. In Theorem 1.4, it is worth noting that Q is determined

by the matrix M . That is, for different M , the corresponding Q may be different (see Theorem 3.4). We also notice that Q is nontrivial, i.e., there exist M and D such that (M, D) is not admissible, but (QMQ^{-1}, QD) is admissible (see Remark 4.9).

We now outline the strategy of the proof of Theorem 1.4. First, we give an equivalent form of Theorem 1.4 under a similarity transformation (see Theorem 2.4), which implies that the spectrality of $\mu_{M,D}$ is the same as that of $\mu_{\tilde{M},\tilde{D}}$, where \tilde{D} and \tilde{M} are given by (2.6) and (2.7) respectively. For the first result of Theorem 1.4, the sufficiency will follow directly from Theorem 1.2 and Lemma 2.1, the difficult part of the proof is the necessity. A key point in the proof is to analyze the characteristics of the zero set of the mask polynomial of \tilde{D} (see Proposition 2.6). Moreover, our approach to the result is based on how to express the matrix \tilde{M} . We will divide the proof of Theorem 1.4 into the following two cases:

Case I: $2\alpha_1 - \beta_1, 2\alpha_2 - \beta_2 \in 3\mathbb{Z}$.

Case II: $2\alpha_1 - \beta_1 \notin 3\mathbb{Z}$ or $2\alpha_2 - \beta_2 \notin 3\mathbb{Z}$.

In each case, we will use different methods to characterize the spectral properties of the measure $\mu_{\tilde{M},\tilde{D}}$ corresponding to different matrices \tilde{M} (see Theorems 3.4 and 4.6).

For the second result of Theorem 1.4, the necessity is essentially trivial from the definition of spectral measure. To complete the proof the sufficiency, we first use the properties of \tilde{M} and \tilde{D} to show that $\mu_{\tilde{M},\tilde{D}}$ is a non-spectral measure for $\tilde{M} \in \mathfrak{B}$ (see Lemma 3.2), and then prove the completeness of the infinite orthogonal set of exponential functions (see Theorem 3.3), where the completeness is established by checking the existence of Hadamard triple.

The paper is organized as follows. In Section 2, we recall a few basic concepts and notations, establish several lemmas that will be needed in the proof of our main results. In Section 3, we focus on proving Theorem 1.4 in Case I. Finally, we settle Case II, and give some remarks and an open problem in Section 4.

2. Preliminaries

The purpose of this section is to collect necessary facts that we need in the following sections. Let $M \in M_n(\mathbb{R})$ be an expanding real matrix, $D \subset \mathbb{R}^n$ be a finite digit set with cardinality $\#D$, and let $\mu_{M,D}$ be defined by (1.1). The Fourier transform of $\mu_{M,D}$ is defined as usual,

$$\hat{\mu}_{M,D}(\xi) = \int e^{2\pi i \langle x, \xi \rangle} d\mu_{M,D}(x) = \prod_{j=1}^{\infty} m_D(M^{*-j}\xi), \quad \xi \in \mathbb{R}^n, \quad (2.1)$$

where M^* denotes the conjugate transposed matrix of M , and $m_D(x)$ is the mask polynomial of D , which is defined by

$$m_D(x) = \frac{1}{\#D} \sum_{d \in D} e^{2\pi i \langle d, x \rangle}, \quad x \in \mathbb{R}^n.$$

Let $\mathcal{Z}(f)$ denote the zero set of the function f , i.e., $\mathcal{Z}(f) = \{x \in \mathbb{R}^n : f(x) = 0\}$, and define

$$\mathcal{Z}_D^n = \mathcal{Z}(m_D) \cap [0, 1)^n. \quad (2.2)$$

It is easy to see that m_D is a \mathbb{Z}^n -periodic function for $D \subset \mathbb{Z}^n$. In this case,

$$\mathcal{Z}(m_D) = \mathcal{Z}_D^n + \mathbb{Z}^n. \quad (2.3)$$

It follows from (2.1) that

$$\mathcal{Z}(\hat{\mu}_{M,D}) = \bigcup_{j=1}^{\infty} M^{*j}(\mathcal{Z}(m_D)). \quad (2.4)$$

For any $\lambda_1 \neq \lambda_2 \in \mathbb{R}^n$, the orthogonality condition

$$0 = \langle e^{2\pi i \langle \lambda_1, x \rangle}, e^{2\pi i \langle \lambda_2, x \rangle} \rangle_{L^2(\mu_{M,D})} = \int e^{2\pi i \langle \lambda_1 - \lambda_2, x \rangle} d\mu_{M,D}(x) = \hat{\mu}_{M,D}(\lambda_1 - \lambda_2)$$

relates to the zero set $\mathcal{Z}(\hat{\mu}_{M,D})$ directly. For a countable subset $\Lambda \subset \mathbb{R}^n$, it is easy to see that $E(\Lambda) = \{e^{2\pi i \langle \lambda, x \rangle} : \lambda \in \Lambda\}$ is an orthonormal family of $L^2(\mu_{M,D})$ if and only if

$$(\Lambda - \Lambda) \setminus \{0\} \subset \mathcal{Z}(\hat{\mu}_{M,D}). \quad (2.5)$$

We call Λ satisfying (2.5) a bi-zero set of $\mu_{M,D}$. Since the properties of bi-zero sets (or spectra) are invariant under a translation, it will be convenient to assume that $0 \in \Lambda$ in this paper, and hence $\Lambda \subset (\Lambda - \Lambda)$.

The following lemma indicates that the spectral properties of $\mu_{M,D}$ are invariant under a similarity transformation. The proof is the same as that of Lemma 4.1 in [15].

Lemma 2.1. *Let $D, D_1 \subset \mathbb{R}^n$ be two finite digit sets with the same cardinality, and let $M, M_1 \in M_n(\mathbb{R})$ be two expanding matrices. If there exists a matrix $A \in M_n(\mathbb{R})$ such that $M_1 = AMA^{-1}$ and $D_1 = AD$, then Λ is a bi-zero set of $\mu_{M,D}$ if and only if $A^{*-1}\Lambda$ is a bi-zero set of μ_{M_1,D_1} . Moreover, $\mu_{M,D}$ is a spectral measure with spectrum Λ if and only if μ_{M_1,D_1} is a spectral measure with spectrum $A^{*-1}\Lambda$.*

In [31], Li gave a necessary and sufficient condition for the finite $\mu_{M,D}$ -orthogonality, which will be used to prove our main results.

Theorem 2.2. *[31, Theorem 2.1] Let $M \in M_n(\mathbb{Z})$ be an expanding matrix and $D \subset \mathbb{Z}^n$ be a finite digit set with $0 \in D$, and let \mathcal{Z}_D^n be defined by (2.2). Suppose that $\mathcal{Z}_D^n \subset \mathbb{Q}^n$ is a finite set, then there exist at most finite mutually orthogonal exponential functions in $L^2(\mu_{M,D})$ if and only if $M^{*j}\mathcal{Z}_D^n \cap \mathbb{Z}^n = \emptyset$ for all $j \in \mathbb{N}$.*

The following lemma is an effective method to illustrate that a countable set Λ cannot be a spectrum of a measure μ , which will be used in Proposition 4.1.

Lemma 2.3. [8, Lemma 2.2] *Let $\mu = \mu_1 * \mu_2$ be the convolution of two probability measures μ_i , $i = 1, 2$, and they are not Dirac measures. Suppose that Λ is a bi-zero set of μ_1 , then Λ is also a bi-zero set of μ , but Λ cannot be a spectrum of μ .*

In order to prove Theorem 1.4, we will give an equivalent form of Theorem 1.4 under a similarity transformation. Before stating the form, some technical work needs to be done.

For the matrix M and the digit set D given in Theorem 1.4, we can let $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in M_2(\mathbb{Z})$ and $\alpha_1\beta_2 - \alpha_2\beta_1 = 3^n\gamma$ for some integers $\eta \geq 1$ and $3 \nmid \gamma$. Without loss of generality, we assume $\gcd(\alpha_1, \alpha_2) = \sigma$ with $3 \nmid \sigma$ (Otherwise, we can choose $\sigma = \gcd(\beta_1, \beta_2)$ with $3 \nmid \sigma$, since $\gcd(\alpha_1, \alpha_2, \beta_1, \beta_2) = 1$). Let $\alpha_1 = \sigma t_1$ and $\alpha_2 = \sigma t_2$ with $\gcd(t_1, t_2) = 1$, then there exist two integers p and q such that $pt_1 + qt_2 = 1$. Clearly, $\sigma = p\alpha_1 + q\alpha_2$ and $\sigma \mid \gamma$. For convenience, we denote $\omega = p\beta_1 + q\beta_2$ and $\vartheta = \gamma/\sigma \notin 3\mathbb{Z}$. Let $P = \begin{bmatrix} p & q \\ -t_2 & t_1 \end{bmatrix}$. By noting that $t_2\alpha_1 = t_1\alpha_2$ and $t_1\beta_2 - t_2\beta_1 = 3^n\vartheta$, we have

$$\tilde{D} = PD = \begin{bmatrix} p & q \\ -t_2 & t_1 \end{bmatrix} \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \alpha_1 \\ \alpha_2 \end{pmatrix}, \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix} \right\} = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \sigma \\ 0 \end{pmatrix}, \begin{pmatrix} \omega \\ 3^n\vartheta \end{pmatrix} \right\} \subset \mathbb{Z}^2 \quad (2.6)$$

and

$$\tilde{M} = PMP^{-1} = \begin{bmatrix} (pa + qc)t_1 + (pb + qd)t_2 & (pb + qd)p - (pa + qc)q \\ (ct_1 - at_2)t_1 + (dt_1 - bt_2)t_2 & (dt_1 - bt_2)p - (ct_1 - at_2)q \end{bmatrix} \quad (2.7)$$

It is easy to verify that \tilde{M} is an expanding integer matrix with $\det(\tilde{M}) = \det(M) \in 3\mathbb{Z}$. We remark that $2\sigma - \omega \in 3\mathbb{Z}$ if $2\alpha_1 - \beta_1, 2\alpha_2 - \beta_2 \in 3\mathbb{Z}$, and $2\sigma - \omega \notin 3\mathbb{Z}$ if $2\alpha_1 - \beta_1 \notin 3\mathbb{Z}$ or $2\alpha_2 - \beta_2 \notin 3\mathbb{Z}$ (see Proposition 2.5).

It is known that the spectrality of $\mu_{\tilde{M}, \tilde{D}}$ is the same as that of $\mu_{M, D}$ by Lemma 2.1. Hence the proof of Theorem 1.4 is equivalent to proving the following

Theorem 2.4. *Let $\mu_{\tilde{M}, \tilde{D}}$ be defined by (1.1), where \tilde{D} and \tilde{M} are given by (2.6) and (2.7) respectively. Then $\mu_{\tilde{M}, \tilde{D}}$ is a spectral measure if and only if there exists a matrix $Q \in M_2(\mathbb{Z})$ such that (\tilde{M}, \tilde{D}) is admissible, where $\tilde{M} = Q\tilde{M}Q^{-1}$ and $\tilde{D} = Q\tilde{D}$. In particular, if $2\sigma - \omega \in 3\mathbb{Z}$, then $\mu_{\tilde{M}, \tilde{D}}$ is a spectral measure if and only if $L^2(\mu_{\tilde{M}, \tilde{D}})$ contains an infinite orthogonal set of exponential functions.*

Now we introduce some properties of the digit set \tilde{D} given in Theorem 2.4. It is known that $1 + e^{2\pi i x_1} + e^{2\pi i x_2} = 0$ if and only if

$$\begin{cases} x_1 = \frac{1}{3} + k_1, \\ x_2 = \frac{2}{3} + k_2, \end{cases} \quad \text{or} \quad \begin{cases} x_1 = \frac{2}{3} + k_3, \\ x_2 = \frac{1}{3} + k_4, \end{cases}$$

where $k_1, k_2, k_3, k_4 \in \mathbb{Z}$. By a direct calculation, we have that

$$\mathcal{Z}(m_{\tilde{D}}) = Z_0 \cup \tilde{Z}_0, \quad (2.8)$$

where

$$Z_0 = \left\{ \left(\frac{1}{3^{\eta+1}\gamma} (2\sigma - \omega - \frac{1+3k_1}{3\sigma} + 3\omega k_1 + 3\sigma k_2) \right) : k_1, k_2 \in \mathbb{Z} \right\},$$

and

$$\tilde{Z}_0 = \left\{ \left(\frac{1}{3^{\eta+1}\gamma} (\sigma - 2\omega - \frac{2+3k_3}{3\sigma} + 3\omega k_3 + 3\sigma k_4) \right) : k_3, k_4 \in \mathbb{Z} \right\}.$$

The following result was proved in [5, Proposition 3.8] and illustrates some inner relationships between two digit sets D and \tilde{D} . Since the proof is simple, we give it here.

Proposition 2.5. *Let D and \tilde{D} be defined by (1.2) and (2.6), respectively. Then*

- (i) $2\sigma - \omega \in 3\mathbb{Z}$ if $2\alpha_1 - \beta_1, 2\alpha_2 - \beta_2 \in 3\mathbb{Z}$.
- (ii) $2\sigma - \omega \notin 3\mathbb{Z}$ if $2\alpha_1 - \beta_1 \notin 3\mathbb{Z}$ or $2\alpha_2 - \beta_2 \notin 3\mathbb{Z}$.

Proof. It is easy to verify that $2\sigma - \omega = p(2\alpha_1 - \beta_1) + q(2\alpha_2 - \beta_2)$, thus (i) holds. For (ii), without loss of generality, we assume $2\alpha_1 - \beta_1 \notin 3\mathbb{Z}$. Since $\alpha_1 = \sigma t_1$, $\alpha_2 = \sigma t_2$ and $t_1\beta_2 - t_2\beta_1 = 3^n\vartheta$, we have $t_2\alpha_1 = t_1\alpha_2$ and

$$q3^n\vartheta = q(t_1\beta_2 - t_2\beta_1) = qt_1(\beta_2 - 2\alpha_2) + qt_2(2\alpha_1 - \beta_1). \quad (2.9)$$

Multiplying $2\alpha_1 - \beta_1$ on both sides of $pt_1 + qt_2 = 1$, we get $pt_1(2\alpha_1 - \beta_1) + qt_2(2\alpha_1 - \beta_1) = 2\alpha_1 - \beta_1$. This together with (2.9) yields that $t_1(p(2\alpha_1 - \beta_1) + q(2\alpha_2 - \beta_2)) = -q3^n\vartheta + 2\alpha_1 - \beta_1$. Hence $2\sigma - \omega = p(2\alpha_1 - \beta_1) + q(2\alpha_2 - \beta_2) \notin 3\mathbb{Z}$ by $2\alpha_1 - \beta_1 \notin 3\mathbb{Z}$. \square

In order to characterize the properties of $\mathcal{Z}(m_{\tilde{D}})$, we first define the following four sets. Let

$$\mathcal{H} = \left\{ \left(\frac{\ell_1}{\frac{3\gamma}{3^{\eta}\gamma}} \right) : \ell_1 \in \mathbb{Z} \setminus 3\mathbb{Z}, \ell_2 \in \mathbb{Z} \right\}, \quad \mathcal{G} = \left\{ \left(\frac{\ell_1}{\frac{3\gamma}{3^{\eta+1}\gamma}} \right) : \ell_1, \ell_2 \in \mathbb{Z} \setminus 3\mathbb{Z} \right\},$$

$$\mathcal{G}_1 = \left\{ \left(\frac{\ell_1}{\frac{3}{3^{\eta+1}}} \right) : \ell_1, \ell_2 \in \mathbb{Z} \setminus 3\mathbb{Z}, \ell_1 = \ell_2 \pmod{3} \right\}$$

and

$$\mathcal{G}_2 = \left\{ \left(\frac{\ell_1}{\frac{3}{3^{\eta+1}}} \right) : \ell_1, \ell_2 \in \mathbb{Z} \setminus 3\mathbb{Z}, \ell_1 \neq \ell_2 \pmod{3} \right\}.$$

The following is an elementary but useful fact in our investigation.

Proposition 2.6. *With the above notations, we have the following assertions:*

- (i) If $2\sigma - \omega \in 3\mathbb{Z}$, then $\gamma\mathcal{H} \subset \mathcal{Z}(m_{\tilde{D}}) \subset \mathcal{H}$.
- (ii) If $2\sigma - \omega \notin 3\mathbb{Z}$, then $\mathcal{G}_i \subset \mathcal{Z}(m_{\tilde{D}}) \subset \mathcal{G}$ for $i = 1$ or 2 .

Proof. (i) Since $2\sigma - \omega \in 3\mathbb{Z}$ and $\gamma = \sigma\vartheta$, it follows from (2.8) that $\mathcal{Z}(m_{\bar{D}}) \subset \mathcal{H}$. Now we prove $\gamma\mathcal{H} \subset \mathcal{Z}(m_{\bar{D}})$. It is equivalent to proving that for any $(\frac{\ell_1}{3}, \frac{\ell_2}{3\eta})^t \in \gamma\mathcal{H}$ with $\ell_1 \notin 3\mathbb{Z}$ and $\ell_2 \in \mathbb{Z}$, there exist $k_1, k_2 \in \mathbb{Z}$ such that

$$\begin{cases} 1 + 3k_1 = \sigma\ell_1, \\ 2\sigma - \omega - 3\omega k_1 + 3\sigma k_2 = 3\gamma\ell_2, \end{cases} \quad \text{or} \quad \begin{cases} 2 + 3k_1 = \sigma\ell_1, \\ \sigma - 2\omega - 3\omega k_1 + 3\sigma k_2 = 3\gamma\ell_2. \end{cases} \quad (2.10)$$

Notice that $\sigma, \ell_1 \notin 3\mathbb{Z}$, then there must exist $k_1 \in \mathbb{Z}$ such that $\sigma\ell_1 = 3k_1 + 1$ or $\sigma\ell_1 = 3k_1 + 2$. Without loss of generality, we assume that $\sigma\ell_1 = 3k_1 + 1$, which yields $\sigma = \ell_1 \pmod{3}$. In view of $\gamma = \sigma\vartheta$ and (2.10), we only need to prove

$$k_2 = \vartheta\ell_2 + \frac{\omega\ell_1 - 2}{3} \in \mathbb{Z}.$$

According to $\sigma \notin 3\mathbb{Z}$, $2\sigma - \omega \in 3\mathbb{Z}$ and $\sigma = \ell_1 \pmod{3}$, we conclude $\omega \notin 3\mathbb{Z}$ and $\omega \neq \ell_1 \pmod{3}$. This implies that $\omega\ell_1 = 3a + 2$ for some $a \in \mathbb{Z}$, thus $k_2 = \vartheta\ell_2 + a \in \mathbb{Z}$. Hence (2.10) holds. This proves $\gamma\mathcal{H} \subset \mathcal{Z}(m_{\bar{D}})$.

(ii) As $2\sigma - \omega \notin 3\mathbb{Z}$ and $\gamma = \sigma\vartheta$, it follows from (2.8) that $\mathcal{Z}(m_{\bar{D}}) \subset \mathcal{G}$. Now we prove $\mathcal{G}_i \subset \mathcal{Z}(m_{\bar{D}})$ for $i = 1$ or 2 . It suffices to prove that for any $(\frac{\ell_1}{3}, \frac{\ell_2}{3\eta+1})^t \in \mathcal{G}_i$ with $\ell_1, \ell_2 \notin 3\mathbb{Z}$, $\ell_1 = \ell_2 \pmod{3}$ or $\ell_1 \neq \ell_2 \pmod{3}$, there exist $k_1, k_2 \in \mathbb{Z}$ such that

$$\begin{cases} 1 + 3k_1 = \sigma\ell_1, \\ 2\sigma - \omega - 3\omega k_1 + 3\sigma k_2 = \gamma\ell_2, \end{cases} \quad \text{or} \quad \begin{cases} 2 + 3k_1 = \sigma\ell_1, \\ \sigma - 2\omega - 3\omega k_1 + 3\sigma k_2 = \gamma\ell_2. \end{cases} \quad (2.11)$$

It follows from $\sigma, \ell_1 \notin 3\mathbb{Z}$ that $\sigma\ell_1 = 3k_1 + 1$ or $\sigma\ell_1 = 3k_1 + 2$ for some $k_1 \in \mathbb{Z}$. Without loss of generality, we assume that $\sigma\ell_1 = 3k_1 + 1$, which means that $\ell_1 = \sigma \pmod{3}$. Using $\gamma = \sigma\vartheta$ and (2.11), we only need to prove

$$k_2 = \frac{\vartheta\ell_2 + \omega\ell_1 - 2}{3} \in \mathbb{Z} \quad \text{for } \ell_2 = \ell_1 \pmod{3} \text{ or } \ell_2 \neq \ell_1 \pmod{3}.$$

We divide the proof into the following two cases.

If $\omega \in 3\mathbb{Z}$, one can write $\omega = 3\omega_1$ with $\omega_1 \in \mathbb{Z}$. Since $\vartheta \notin 3\mathbb{Z}$ and $\ell_1 = \sigma \pmod{3}$, there exists $b \in \mathbb{Z}$ so that $\vartheta\ell_2 = 3b + 2$ for $\ell_2 = \ell_1 \pmod{3}$ or $\ell_2 \neq \ell_1 \pmod{3}$. Thus $k_2 = \omega_1\ell_1 + b \in \mathbb{Z}$.

If $\omega \notin 3\mathbb{Z}$, then $\omega = \sigma \pmod{3}$ by $2\sigma - \omega \notin 3\mathbb{Z}$. This together with $\sigma = \ell_1 \pmod{3}$ yields that $\omega\ell_1 = 3c + 1$ for some $c \in \mathbb{Z}$. Note that $\vartheta \notin 3\mathbb{Z}$, then there must exist $d \in \mathbb{Z}$ so that $\vartheta\ell_2 = 3d + 1$ for $\ell_2 = \ell_1 \pmod{3}$ or $\ell_2 \neq \ell_1 \pmod{3}$. Therefore, $k_2 = c + d \in \mathbb{Z}$.

This proves (2.11), and hence $\mathcal{G}_i \subset \mathcal{Z}(m_{\bar{D}})$ for $i = 1$ or 2 . \square

The following lemma is an effective method to judge whether (\tilde{M}, \tilde{D}) is admissible for some special forms of \tilde{M} and \tilde{D} . It is needed in the proof of Theorem 2.4.

Lemma 2.7. *Let $0 \in \mathcal{J}$ be a set with $\#\mathcal{J} = 3$. If $(\mathcal{J} - \mathcal{J}) \setminus \{0\} \subset \mathcal{Z}(m_{\bar{D}})$ and $\tilde{M}^*\mathcal{J} \subset \mathbb{Z}^2$, then (\tilde{M}, \tilde{D}) is admissible.*

Proof. Let $S = \tilde{M}^* \mathcal{J}$. Since $0 \in \mathcal{J}$ and $\#\mathcal{J} = \#\tilde{D} = 3$, by Definition 1.1, it is clear that $(\tilde{M}^{-1}\tilde{D}, S)$ is a compatible pair if and only if $(\mathcal{J} - \mathcal{J}) \setminus \{0\} \subset \mathcal{Z}(m_{\tilde{D}})$. Hence the lemma follows. \square

At the end of this section, we give two remarks that will be used to prove our main results. Throughout the paper, we denote

$$\mathcal{J}_0 = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 3 \end{pmatrix}, \begin{pmatrix} 2 \\ 3 \end{pmatrix} \right\}, \quad \mathcal{J}_1 = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 3 \\ 2 \end{pmatrix}, \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix} \right\} \quad \text{and} \quad \mathcal{J}_2 = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 3 \\ 1 \end{pmatrix}, \begin{pmatrix} 2 \\ 3 \\ 2 \end{pmatrix} \right\}. \quad (2.12)$$

Remark 2.8. By (2.2) and Proposition 2.6, it is easy to verify that $\mathcal{Z}_{\tilde{D}}^2 \subset \mathbb{Q}^2$ is a finite set. Moreover, if $2\sigma - \omega \in 3\mathbb{Z}$ and $\eta > 0$, then $\mathcal{J}_i \setminus \{0\}, (\mathcal{J}_i - \mathcal{J}_i) \setminus \{0\} \subset \mathcal{Z}(m_{\tilde{D}})$ for $i \in \{0, 1, 2\}$.

Remark 2.9. If $\eta = 0$ in \tilde{D} , using the same proof as Proposition 2.6, the following statements hold.

- (i) If $2\sigma - \omega \in 3\mathbb{Z}$, then $\mathcal{J}_0 \setminus \{0\}, (\mathcal{J}_0 - \mathcal{J}_0) \setminus \{0\} \subset \mathcal{Z}(m_{\tilde{D}})$.
- (ii) If $2\sigma - \omega \notin 3\mathbb{Z}$, then $\mathcal{J}_i \setminus \{0\}, (\mathcal{J}_i - \mathcal{J}_i) \setminus \{0\} \subset \mathcal{Z}(m_{\tilde{D}})$ for $i = 1$ or 2 .

3. Proof of Theorem 1.4 in Case I

In the present section, we focus on proving Theorem 1.4 in Case I, which is equivalent to proving Theorem 2.4 in the case of $2\sigma - \omega \in 3\mathbb{Z}$. To this end, we use the residue system of modulo 3 and rewrite the matrix \tilde{M} given by (2.7) in the following form:

$$\tilde{M} = 3 \begin{bmatrix} a & b \\ 3^{s-1}c & d \end{bmatrix} + M_k := M' + M_k, \quad (3.1)$$

where $s \geq 1$, $a, b, d \in \mathbb{Z}$ and $c \in (\mathbb{Z} \setminus 3\mathbb{Z}) \cup \{0\}$, and the entries of the matrix M_k are from the set $\{0, 1, 2\}$. It is obvious that s can be any positive integer if $c = 0$. Without loss of generality, in the rest of this paper, we always assume that $s \geq \eta$ in this case. As $\det(\tilde{M}) \in 3\mathbb{Z}$, there are 10 different matrices M_k as following:

$$\begin{aligned} M_1 &= \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \quad M_2 = \begin{bmatrix} p_1 & 0 \\ 0 & 0 \end{bmatrix}, \quad M_3 = \begin{bmatrix} 0 & p_2 \\ 0 & 0 \end{bmatrix}, \quad M_4 = \begin{bmatrix} 0 & 0 \\ p_3 & 0 \end{bmatrix}, \quad M_5 = \begin{bmatrix} 0 & 0 \\ 0 & p_4 \end{bmatrix}, \\ M_6 &= \begin{bmatrix} p_1 & 0 \\ p_3 & 0 \end{bmatrix}, \quad M_7 = \begin{bmatrix} p_1 & p_2 \\ 0 & 0 \end{bmatrix}, \quad M_8 = \begin{bmatrix} 0 & 0 \\ p_3 & p_4 \end{bmatrix}, \quad M_9 = \begin{bmatrix} 0 & p_2 \\ 0 & p_4 \end{bmatrix}, \quad M_{10} = \begin{bmatrix} p_1 & p_2 \\ p_3 & p_4 \end{bmatrix}, \end{aligned}$$

where $p_1, p_2, p_3, p_4 \in \{1, 2\}$ and $p_1p_4 - p_2p_3 \in 3\mathbb{Z}$. Fix $k \in \{1, 2, \dots, 10\}$, we denote

$$\mathfrak{M}_k = \left\{ \tilde{M} : \tilde{M} = M' + M_k \right\}. \quad (3.2)$$

The following lemma illustrates a basic property of the matrix $\tilde{M} \in \mathfrak{M}_2$, which will be needed in proving Lemma 3.2.

Lemma 3.1. *If $\tilde{M} \in \mathfrak{M}_2$, then for any $\ell \in \mathbb{N}$, there exist $a_\ell, b_\ell, d_\ell \in \mathbb{Z}$ and $c_\ell \in (\mathbb{Z} \setminus 3\mathbb{Z}) \cup \{0\}$ such that*

$$\tilde{M}^{*\ell} = \begin{bmatrix} 3a_\ell + p_1^\ell & 3^s c_\ell \\ 3b_\ell & 3d_\ell \end{bmatrix}.$$

In particular, $c_\ell = 0$ if $c = 0$.

Proof. We prove the lemma by induction. It is obvious that the lemma holds for $\ell = 1$. Inductively, we assume that it holds for $\ell = k$. That is,

$$\tilde{M}^{*k} = \begin{bmatrix} 3a_k + p_1^k & 3^s c_k \\ 3b_k & 3d_k \end{bmatrix},$$

where $a_k, b_k, d_k \in \mathbb{Z}$ and $c_k \in (\mathbb{Z} \setminus 3\mathbb{Z}) \cup \{0\}$. In particular, $c_k = 0$ if $c = 0$.

We then consider $\ell = k + 1$. By inductive hypothesis, we have

$$\begin{aligned} \tilde{M}^{*k+1} &= \begin{bmatrix} 3a_k + p_1^k & 3^s c_k \\ 3b_k & 3d_k \end{bmatrix} \begin{bmatrix} 3a + p_1 & 3^s c \\ 3b & 3d \end{bmatrix} \\ &= \begin{bmatrix} 3(3^s b c_k + 3a a_k + a p_1^k + a_k p_1) + p_1^{k+1} & 3^s(3c a_k + 3d c_k + c p_1^k) \\ 3(3ab_k + 3bd_k + b_k p_1) & 3(3^s c b_k + 3d d_k) \end{bmatrix} \\ &:= \begin{bmatrix} 3a_{k+1} + p_1^{k+1} & 3^s c_{k+1} \\ 3b_{k+1} & 3d_{k+1} \end{bmatrix}. \end{aligned}$$

It is easy to see that $a_{k+1}, b_{k+1}, d_{k+1} \in \mathbb{Z}$. Since $c, c_k \in (\mathbb{Z} \setminus 3\mathbb{Z}) \cup \{0\}$, $p_1 \in \{1, 2\}$ and $c_k = 0$ if $c = 0$, we conclude that $c_{k+1} \in (\mathbb{Z} \setminus 3\mathbb{Z}) \cup \{0\}$. In particular, if $c = 0$, we have $c_{k+1} = 0$ by $c_k = 0$. This means that the lemma holds for $\ell = k + 1$. Hence the proof is completed. \square

For the set \mathfrak{M}_k given by (3.2), we denote

$$\mathfrak{B} = \left\{ \tilde{M} : \tilde{M} \in \mathfrak{M}_k, k \in \{2, 7\}, s \geq \eta \right\}. \quad (3.3)$$

It should be noted that the set \mathfrak{B} contains the case $c = 0$ in \tilde{M} . Under the assumption of $2\sigma - \omega \in 3\mathbb{Z}$, we will prove that $\mu_{\tilde{M}, \tilde{D}}$ is a non-spectral measure for any $\tilde{M} \in \mathfrak{B}$.

Lemma 3.2. *Let \mathfrak{B} be given by (3.3). If $\tilde{M} \in \mathfrak{B}$ and $2\sigma - \omega \in 3\mathbb{Z}$ in \tilde{D} , then there exist at most finite mutually orthogonal exponential functions in $L^2(\mu_{\tilde{M}, \tilde{D}})$.*

Proof. First, we consider $\tilde{M} \in \mathfrak{B} \cap \mathfrak{M}_2$. We claim that $\tilde{M}^{*\ell} \mathcal{Z}_{\tilde{D}}^2 \cap \mathbb{Z}^2 = \emptyset$ for any $\ell \in \mathbb{N}$. If not, there exists $m \in \mathbb{N}$ such that $\tilde{M}^{*m} \mathcal{Z}_{\tilde{D}}^2 \cap \mathbb{Z}^2 \neq \emptyset$. Since $2\sigma - \omega \in 3\mathbb{Z}$, it follows from Proposition 2.6 that

$$\mathcal{Z}_{\tilde{D}}^2 \subset \mathcal{H} \cap [0, 1)^2 = \left\{ \left(\begin{array}{c} \frac{\ell_1}{3\gamma} \\ \frac{\ell_2}{3^{\eta}\gamma} \end{array} \right) : 1 \leq \ell_1 \leq 3\gamma - 1, 0 \leq \ell_2 \leq 3^{\eta}\gamma - 1, \ell_1 \in \mathbb{Z} \setminus 3\mathbb{Z} \right\}.$$

Thus there exist $(\frac{\ell_1}{3\gamma}, \frac{\ell_2}{3^{\eta}\gamma})^t \in \mathcal{H} \cap [0, 1)^2$ and $(s_1, s_2)^t \in \mathbb{Z}^2$ such that

$$\tilde{M}^{*m} \begin{pmatrix} \frac{\ell_1}{3\gamma} \\ \frac{\ell_2}{3^{\eta}\gamma} \end{pmatrix} = \begin{pmatrix} s_1 \\ s_2 \end{pmatrix}. \quad (3.4)$$

By Lemma 3.1, there exist $a_m, b_m, d_m \in \mathbb{Z}$ and $c_m \in (\mathbb{Z} \setminus 3\mathbb{Z}) \cup \{0\}$ such that

$$\tilde{M}^{*m} = \begin{bmatrix} 3a_m + p_1^m & 3^s c_m \\ 3b_m & 3d_m \end{bmatrix}.$$

Then multiplying both sides of equation (3.4) by $3^\eta \gamma$, we get

$$\ell_1 p_1^m = 3(\gamma s_1 - a_m \ell_1 - 3^{s-\eta} c_m \ell_2) \quad \text{and} \quad d_m \ell_2 = 3^{\eta-1}(\gamma s_2 - b_m \ell_1). \quad (3.5)$$

Since $\ell_1 \notin 3\mathbb{Z}$ and $p_1 \in \{1, 2\}$, on can get $\ell_1 p_1^m \notin 3\mathbb{Z}$. However, it follows from $s \geq \eta$ that $3(\gamma s_1 - a_m \ell_1 - 3^{s-\eta} c_m \ell_2) \in 3\mathbb{Z}$. Thus (3.5) does not hold, and hence the claim follows. According to the claim and Theorem 2.2, there exist at most finite mutually orthogonal exponential functions in $L^2(\mu_{\tilde{M}, \tilde{D}})$.

Second, we consider $\tilde{M} \in \mathfrak{B} \cap \mathfrak{M}_7$. It follows from (3.2) and (3.3) that

$$\tilde{M} = \begin{bmatrix} 3a + p_1 & 3b + p_2 \\ 3^s c & 3d \end{bmatrix},$$

where $s \geq \eta$, $a, b, d \in \mathbb{Z}$, $c \in (\mathbb{Z} \setminus 3\mathbb{Z}) \cup \{0\}$ and $p_1, p_2 \in \{1, 2\}$. It is clear that there exists $\tau \in \{1, 2\}$ such that $p_2 - \tau p_1 \in 3\mathbb{Z}$. Let $Q = \begin{bmatrix} 1 & \tau \\ 0 & 1 \end{bmatrix}$, then we have

$$\tilde{M}' = Q\tilde{M}Q^{-1} = \begin{bmatrix} 3(3^{s-1}\tau c + a) + p_1 & 3(-3^{s-1}c\tau^2 - (a-d)\tau + b) + p_2 - \tau p_1 \\ 3^s c & 3(d - 3^{s-1}\tau c) \end{bmatrix}$$

and

$$\tilde{D}' = Q\tilde{D} = \begin{bmatrix} 1 & \tau \\ 0 & 1 \end{bmatrix} \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \sigma \\ 0 \end{pmatrix}, \begin{pmatrix} \omega \\ 3^\eta \vartheta \end{pmatrix} \right\} = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \sigma \\ 0 \end{pmatrix}, \begin{pmatrix} \omega + 3^\eta \tau \vartheta \\ 3^\eta \vartheta \end{pmatrix} \right\} \subset \mathbb{Z}^2.$$

Combining $p_1 \in \{1, 2\}$ with $p_2 - \tau p_1 \in 3\mathbb{Z}$, we can easily know that $\tilde{M}' \in \mathfrak{M}_2 \cap \mathfrak{B}$. Moreover, it follows from $2\sigma - \omega \in 3\mathbb{Z}$ and $\eta \geq 1$ that $2\sigma - (\omega + 3^\eta \tau \vartheta) \in 3\mathbb{Z}$, thus \tilde{D}' has the same properties as \tilde{D} . Therefore, using the result of $\tilde{M} \in \mathfrak{B} \cap \mathfrak{M}_2$ and Lemma 2.1, one can conclude that there exist at most finite mutually orthogonal exponential functions in $L^2(\mu_{\tilde{M}, \tilde{D}})$ for all $\tilde{M} \in \mathfrak{B} \cap \mathfrak{M}_7$.

Hence we complete the proof of Lemma 3.2. \square

Recall that \mathcal{J}_i ($i \in \{0, 1, 2\}$) are given by (2.12), and let

$$Q_n = \begin{bmatrix} 1 & 0 \\ 0 & \frac{1}{3^n} \end{bmatrix} \quad \text{for any } n \in \mathbb{N}. \quad (3.6)$$

They will be used many times in the rest of this paper.

Now we begin to prove Theorem 2.4 in the case of $2\sigma - \omega \in 3\mathbb{Z}$, which is equivalent to proving Theorems 3.3 and 3.4.

Theorem 3.3. *If $2\sigma - \omega \in 3\mathbb{Z}$ in \tilde{D} , then $\mu_{\tilde{M}, \tilde{D}}$ is a spectral measure if and only if $L^2(\mu_{\tilde{M}, \tilde{D}})$ contains an infinite orthogonal set of exponential functions.*

Proof. According to the definition of spectral measure, the necessity is obvious. We now prove the sufficiency. Suppose that $L^2(\mu_{\tilde{M}, \tilde{D}})$ contains an infinite orthogonal set of exponential functions, then Lemma 3.2 and (3.2) imply that $\tilde{M} \in \mathfrak{B}_1 \cup \mathfrak{B}_2$, where

$$\mathfrak{B}_1 = \left\{ \tilde{M} : \tilde{M} \in \mathfrak{M}_k, k \in \{1, 3, 4, 5, 6, 8, 9, 10\} \right\}, \quad (3.7)$$

and

$$\mathfrak{B}_2 = \left\{ \tilde{M} : \tilde{M} \in \mathfrak{M}_k, k \in \{2, 7\}, s < \eta \right\}. \quad (3.8)$$

The proof will be divided into the following three cases.

Case 1: $\tilde{M} \in \mathfrak{B}_1 \setminus \mathfrak{M}_3$. We first claim that $\tilde{M}^* \mathcal{J}_\kappa \subset \mathbb{Z}^2$ for some $\kappa \in \{0, 1, 2\}$. If $\tilde{M} \in \{\tilde{M} : \tilde{M} \in \mathfrak{M}_k, k \in \{1, 4, 5, 8\}\}$, it follows from (3.2) that \tilde{M} can be written as

$$\tilde{M} = \begin{bmatrix} 3a & 3b \\ c' & d' \end{bmatrix},$$

where $a, b, c', d' \in \mathbb{Z}$. By a simple calculation, we obtain that

$$\tilde{M}^* \mathcal{J}_0 = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} 2a \\ 2b \end{pmatrix} \right\} \subset \mathbb{Z}^2. \quad (3.9)$$

If $\tilde{M} \in \mathfrak{M}_6$, we denote

$$\tilde{M} = \tilde{M}_{6,1} \quad \text{if } p_1 = p_3 \quad \text{and} \quad \tilde{M} = \tilde{M}_{6,2} \quad \text{if } p_1 \neq p_3.$$

If $\tilde{M} \in \mathfrak{M}_9$, we denote

$$\tilde{M} = \tilde{M}_{9,1} \quad \text{if } p_2 = p_4 \quad \text{and} \quad \tilde{M} = \tilde{M}_{9,2} \quad \text{if } p_2 \neq p_4.$$

Then by using (2.12) and (3.2), it is easy to verify that

$$\tilde{M}_{k,i}^* \mathcal{J}_i \subset \mathbb{Z}^2 \quad \text{for } k \in \{6, 9\} \text{ and } i \in \{1, 2\}. \quad (3.10)$$

If $\tilde{M} \in \mathfrak{M}_{10}$, we denote

$$\tilde{M} = \tilde{M}_{10,1} \quad \text{if } p_1 = p_3 \quad \text{and} \quad \tilde{M} = \tilde{M}_{10,2} \quad \text{if } p_1 \neq p_3.$$

Note that $p_1, p_2, p_3, p_4 \in \{1, 2\}$ and $p_1 p_4 - p_2 p_3 \in 3\mathbb{Z}$, then $p_2 = p_4$ if $p_1 = p_3$, and $p_2 \neq p_4$ if $p_1 \neq p_3$. Hence, it follows from (2.12) and (3.2) that $\tilde{M}_{10,i}^* \mathcal{J}_i \subset \mathbb{Z}^2$ for $i \in \{1, 2\}$. Combining this with (3.9) and (3.10), the claim follows.

Since $2\sigma - \omega \in 3\mathbb{Z}$, it follows from Remark 2.8 that $\mathcal{J}_i \setminus \{0\}, (\mathcal{J}_i - \mathcal{J}_i) \setminus \{0\} \subset \mathcal{Z}(m_{\tilde{D}})$ for $i \in \{0, 1, 2\}$. Together with the claim and Lemma 2.7, it shows that (\tilde{M}, \tilde{D}) is admissible. Therefore, $\mu_{\tilde{M}, \tilde{D}}$ is a spectral measure by Theorem 1.2.

Case 2: $\tilde{M} \in \mathfrak{M}_3$. Applying (3.2) and (3.6), we have

$$A_1 = Q_1 \tilde{M} Q_1^{-1} = \begin{bmatrix} 3a & 9b + 3p_2 \\ 3^{s-1}c & 3d \end{bmatrix} \quad \text{and} \quad D_1 = Q_1 \tilde{D} = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \sigma \\ 0 \end{pmatrix}, \begin{pmatrix} \omega \\ 3^{\eta-1}\vartheta \end{pmatrix} \right\},$$

where $s \geq 1, a, b, d \in \mathbb{Z}, c \in (\mathbb{Z} \setminus 3\mathbb{Z}) \cup \{0\}$ and $p_2 \in \{1, 2\}$. It follows from $\eta \geq 1$ that $\eta - 1 \geq 0$. Note that $2\sigma - \omega \in 3\mathbb{Z}$, then Remarks 2.8 and 2.9 give that $\mathcal{J}_0 \setminus \{0\}, (\mathcal{J}_0 - \mathcal{J}_0) \setminus$

$\{0\} \subset \mathcal{Z}(m_{D_1})$. Moreover, it is a direct check to see $A_1^* \mathcal{J}_0 \subset \mathbb{Z}^2$. In view of Lemma 2.7, (A_1, D_1) is admissible. Hence $\mu_{\tilde{M}, \tilde{D}}$ is a spectral measure by Theorem 1.2 and Lemma 2.1.

Case 3: $\tilde{M} \in \mathfrak{B}_2$. Form (3.2), we can write $\tilde{M} \in \mathfrak{B}_2$ as

$$\tilde{M} = \begin{bmatrix} 3a + p_1 & b' \\ 3^s c & 3d \end{bmatrix},$$

where $1 \leq s < \eta$, $a, b', d \in \mathbb{Z}$, $c \in \mathbb{Z} \setminus 3\mathbb{Z}$ and $p_1 \in \{1, 2\}$. Applying (3.6), we have

$$A_2 = Q_s \tilde{M} Q_s^{-1} = \begin{bmatrix} 3a + p_1 & 3^s b' \\ c & 3d \end{bmatrix} \quad \text{and} \quad D_2 = Q_s \tilde{D} = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \sigma \\ 0 \end{pmatrix}, \begin{pmatrix} \omega \\ 3^{\eta-s} \vartheta \end{pmatrix} \right\},$$

As $c \notin 3\mathbb{Z}$ and $\eta - s \geq 1$, one can easily know that $A_2 \in \mathfrak{M}_6$ and D_2 has the same form as \tilde{D} . By Case 1, we conclude that (A_2, D_2) is admissible and μ_{A_2, D_2} is a spectral measure. Therefore, $\mu_{\tilde{M}, \tilde{D}}$ is a spectral measure by Lemma 2.1.

Finally, according to the results of the above three cases, we know that $\mu_{\tilde{M}, \tilde{D}}$ is a spectral measure. Hence the sufficiency follows. This ends the proof of Theorem 3.3. \square

Theorem 3.4. *If $2\sigma - \omega \in 3\mathbb{Z}$ in \tilde{D} , then $\mu_{\tilde{M}, \tilde{D}}$ is a spectral measure if and only if there exists a matrix $Q \in M_2(\mathbb{Z})$ such that (\bar{M}, \bar{D}) is admissible, where $\bar{M} = Q\tilde{M}Q^{-1}$ and $\bar{D} = Q\tilde{D}$.*

Proof. The sufficiency follows directly from Theorem 1.2 and Lemma 2.1. Now we are devoted to proving the necessity. Suppose that $\mu_{\tilde{M}, \tilde{D}}$ is a spectral measure, applying Lemma 3.2, we obtain $\tilde{M} \in \mathfrak{B}_1 \cup \mathfrak{B}_2$, where \mathfrak{B}_1 and \mathfrak{B}_2 are given by (3.7) and (3.8) respectively. From the proof of Theorem 3.3, we have that (\bar{M}, \bar{D}) is admissible, where

$$\begin{aligned} \bar{M} &= \tilde{M} \quad \text{and} \quad \bar{D} = \tilde{D} \quad \text{if } \tilde{M} \in \mathfrak{B}_1 \setminus \mathfrak{M}_3, \\ \bar{M} &= Q_1 \tilde{M} Q_1^{-1} \quad \text{and} \quad \bar{D} = Q_1 \tilde{D} \quad \text{if } \tilde{M} \in \mathfrak{M}_3, \\ \bar{M} &= Q_s \tilde{M} Q_s^{-1} \quad \text{and} \quad \bar{D} = Q_s \tilde{D} \quad \text{if } \tilde{M} \in \mathfrak{B}_2. \end{aligned}$$

This proves the necessity, and hence the proof is completed. \square

4. Proof of Theorem 1.4 in Case II

In this section, we are devoted to proving Theorem 1.4 in Case II, which is equivalent to proving Theorem 2.4 in the case of $2\sigma - \omega \notin 3\mathbb{Z}$. Before starting to prove the theorem, we first characterize the spectral properties of some special measures $\mu_{\tilde{M}, \tilde{D}}$. Recall that \mathfrak{M}_k is defined by (3.2), where $k \in \{1, 2, \dots, 10\}$.

Proposition 4.1. *If $\tilde{M} \in \mathfrak{M}_4$ and $2\sigma - \omega \notin 3\mathbb{Z}$ in \tilde{D} , then $\mu_{\tilde{M}, \tilde{D}}$ is a non-spectral measure.*

Proof. For simplicity, we denote $\mathcal{Z}_j := \tilde{M}^{*j} \mathcal{Z}(m_{\tilde{D}})$. Let

$$\mu_1 = \delta_{\tilde{M}^{-1}\tilde{D}} * \delta_{\tilde{M}^{-3}\tilde{D}} * \cdots \quad \text{and} \quad \mu_2 = \delta_{\tilde{M}^{-2}\tilde{D}} * \delta_{\tilde{M}^{-3}\tilde{D}} * \cdots. \quad (4.1)$$

Then $\mu_{\tilde{M}, \tilde{D}} = \mu_1 * \delta_{\tilde{M}^{-2}\tilde{D}} = \mu_2 * \delta_{\tilde{M}^{-1}\tilde{D}}$. It follows from (2.4) that

$$\mathcal{Z}(\hat{\mu}_1) = \bigcup_{j \geq 1, j \neq 2} \mathcal{Z}_j \quad \text{and} \quad \mathcal{Z}(\hat{\mu}_2) = \bigcup_{j \geq 2} \mathcal{Z}_j. \quad (4.2)$$

Let Λ be a bi-zero set of $\mu_{\tilde{M}, \tilde{D}}$ with $0 \in \Lambda$.

We first claim that $\Lambda \setminus \{0\} \subset \mathcal{Z}(\hat{\mu}_1)$ or $\Lambda \setminus \{0\} \subset \mathcal{Z}(\hat{\mu}_2)$. Otherwise, there exist $\lambda_1, \lambda_2 \in \Lambda \setminus \{0\}$ such that $\lambda_1 \in \mathcal{Z}_2$ and $\lambda_2 \in \mathcal{Z}_1$. According to the orthogonality of Λ , we have

$$\lambda_1 - \lambda_2 = \tilde{M}^{*2} \xi_1 - \tilde{M}^* \xi_2 = \tilde{M}^{*\kappa} \xi_3 \quad (4.3)$$

for some $\xi_1, \xi_2, \xi_3 \in \mathcal{Z}(m_{\tilde{D}})$ and $\kappa \geq 1$. Multiplying both sides of (4.3) by $3^n \gamma$, we get

$$\tilde{M}^{*2} 3^n \gamma \xi_1 - \tilde{M}^* 3^n \gamma \xi_2 = \tilde{M}^{*\kappa} 3^n \gamma \xi_3. \quad (4.4)$$

Since $\tilde{M} \in \mathfrak{M}_4$, it follows from (3.2) that

$$\tilde{M}^* = \begin{bmatrix} 3a & 3^s c + p_3 \\ 3b & 3d \end{bmatrix} \quad \text{and} \quad \tilde{M}^{*2} = 3 \begin{bmatrix} 3a^2 + 3^s bc + bp_3 & (a+d)(3^s c + p_3) \\ 3b(a+d) & 3^s bc + 3d^2 + bp_3 \end{bmatrix} \in M_2(3\mathbb{Z}),$$

where $s \geq 1$, $a, b, d \in \mathbb{Z}$, $c \in (\mathbb{Z} \setminus 3\mathbb{Z}) \cup \{0\}$ and $p_3 \in \{1, 2\}$. Applying $2\sigma - \omega \notin 3\mathbb{Z}$ and Proposition 2.6, we conclude that

$$\tilde{M}^* 3^n \gamma \mathcal{Z}(m_{\tilde{D}}) \cap \mathbb{Z}^2 = \emptyset \quad \text{and} \quad \tilde{M}^{*j} 3^n \gamma \mathcal{Z}(m_{\tilde{D}}) \subset \mathbb{Z}^2 \quad \text{for } j \geq 2. \quad (4.5)$$

This and (4.4) give $\kappa = 1$. Hence (4.4) becomes

$$\tilde{M}^* 3^n \gamma (\xi_2 + \xi_3) = \tilde{M}^{*2} 3^n \gamma \xi_1. \quad (4.6)$$

If $3^n \gamma (\xi_2 + \xi_3) \in \mathbb{Z}^2$, then (4.6) implies $\tilde{M}^* 3^n \gamma \xi_1 \in \mathbb{Z}^2$, which contradicts to (4.5). If $3^n \gamma (\xi_2 + \xi_3) \notin \mathbb{Z}^2$, it follows from $2\sigma - \omega \notin 3\mathbb{Z}$ and Proposition 2.6 that there exist $\ell_1, \ell_2, \ell_3, \ell_4 \notin 3\mathbb{Z}$ and $\ell_2 + \ell_4 \notin 3\mathbb{Z}$ such that $\xi_2 = (\frac{\ell_1}{3^\gamma}, \frac{\ell_2}{3^{n+1}\gamma})^t$ and $\xi_3 = (\frac{\ell_3}{3^\gamma}, \frac{\ell_4}{3^{n+1}\gamma})^t$. Consequently,

$$\tilde{M}^* 3^n \gamma (\xi_2 + \xi_3) = \begin{pmatrix} 3^n a(\ell_1 + \ell_3) + 3^{s-1} c(\ell_2 + \ell_4) + \frac{p_3(\ell_2 + \ell_4)}{3} \\ 3^n b(\ell_1 + \ell_3) + d(\ell_2 + \ell_4) \end{pmatrix} \notin \mathbb{Z}^2$$

since $s \geq 1$ and $p_3, \ell_2 + \ell_4 \notin 3\mathbb{Z}$. Together with (4.5), it implies that the left-hand side of (4.6) is not in \mathbb{Z}^2 , while the right-hand side is in \mathbb{Z}^2 . This is a contradiction, and hence the claim follows.

Next, we prove that $(\Lambda - \Lambda) \setminus \{0\} \subset \mathcal{Z}(\hat{\mu}_1)$ or $(\Lambda - \Lambda) \setminus \{0\} \subset \mathcal{Z}(\hat{\mu}_2)$. According to the claim, we decompose the proof into the following two cases.

If $\Lambda \setminus \{0\} \subset \mathcal{Z}(\hat{\mu}_1)$, we have $(\Lambda - \Lambda) \setminus \{0\} \subset \mathcal{Z}(\hat{\mu}_1)$. Suppose, on the contrary, that there exist $\lambda_1, \lambda_2 \in \Lambda \setminus \{0\}$ such that

$$\lambda_1 - \lambda_2 = \tilde{M}^{*j_1} \xi_1 - \tilde{M}^{*j_2} \xi_2 = \tilde{M}^{*2} \xi_3, \quad (4.7)$$

where $\xi_1, \xi_2, \xi_3 \in \mathcal{Z}(m_{\tilde{D}})$ and $j_1, j_2 \in \mathbb{N} \setminus \{2\}$. Similar to (4.6), it is easy to verify that $j_1 \neq j_2$. Without loss of generality, we assume $j_1 > j_2$. If $j_2 = 1$, using (4.5) and $j_1 > 2$,

we have $\tilde{M}^{*j_1} 3^n \gamma \xi_1, \tilde{M}^{*2} 3^n \gamma \xi_3 \in \mathbb{Z}^2$ and $\tilde{M}^* 3^n \gamma \xi_2 \notin \mathbb{Z}^2$. Thus (4.7) does not hold. If $j_2 > 1$, then $j_1 > j_2 > 2$. Multiplying both sides of (4.7) by $\tilde{M}^{*-1} 3^n \gamma$, we get

$$\tilde{M}^{*j_1-1} 3^n \gamma \xi_1 - \tilde{M}^{*j_2-1} 3^n \gamma \xi_2 = \tilde{M}^* 3^n \gamma \xi_3.$$

In view of (4.5), the above equation does not hold. Hence $(\Lambda - \Lambda) \setminus \{0\} \subset \mathcal{Z}(\hat{\mu}_1)$.

If $\Lambda \setminus \{0\} \subset \mathcal{Z}(\hat{\mu}_2)$, we have $(\Lambda - \Lambda) \setminus \{0\} \subset \mathcal{Z}(\hat{\mu}_2)$. The proof is similar to that of $\Lambda \setminus \{0\} \subset \mathcal{Z}(\hat{\mu}_1)$, so we omit it here.

Finally, we show that Λ cannot be a spectrum of $\mu_{\tilde{M}, \tilde{D}}$. Since $(\Lambda - \Lambda) \setminus \{0\} \subset \mathcal{Z}(\hat{\mu}_1)$ or $(\Lambda - \Lambda) \setminus \{0\} \subset \mathcal{Z}(\hat{\mu}_2)$, it follows from (2.5) that Λ is a bi-zero set of μ_1 or μ_2 . Therefore, Lemma 2.3 means that Λ cannot be a spectrum of $\mu_{\tilde{M}, \tilde{D}}$.

This completes the proof of Proposition 4.1. \square

The following lemma is a basic property of the matrix $\tilde{M} \in \mathfrak{M}_6$.

Lemma 4.2. *If $\tilde{M} \in \mathfrak{M}_6$, then for any $\ell \in \mathbb{N}$, there exist $a_\ell, b_\ell, c_\ell, d_\ell \in \mathbb{Z}$ such that*

$$\tilde{M}^{*\ell} = \begin{bmatrix} 3a_\ell + p_1^\ell & 3c_\ell + p_1^{\ell-1} p_3 \\ 3b_\ell & 3d_\ell \end{bmatrix},$$

where $p_1, p_3 \in \{1, 2\}$.

Proof. The proof is similar to that of Lemma 3.1, so we omit it here. \square

Based on Lemma 4.2, we can further characterize the spectral property of $\mu_{\tilde{M}, \tilde{D}}$ for $\tilde{M} \in \mathfrak{M}_6$.

Proposition 4.3. *If $\tilde{M} \in \mathfrak{M}_6$ and $2\sigma - \omega \notin 3\mathbb{Z}$ in \tilde{D} , then $\mu_{\tilde{M}, \tilde{D}}$ is a non-spectral measure.*

Proof. For any $\tilde{M} \in \mathfrak{M}_6$, we first prove that $\tilde{M}^{*j} \mathcal{Z}_{\tilde{D}}^2 \cap \mathbb{Z}^2 = \emptyset$ for $j \in \mathbb{N}$. Suppose, on the contrary, that $\tilde{M}^{*n} \mathcal{Z}_{\tilde{D}}^2 \cap \mathbb{Z}^2 \neq \emptyset$ for some $n \in \mathbb{N}$. By $2\sigma - \omega \notin 3\mathbb{Z}$ and Proposition 2.6, we have

$$\mathcal{Z}_{\tilde{D}}^2 \subset \mathcal{G} \cap [0, 1)^2 = \left\{ \left(\begin{array}{c} \ell_1 \\ \frac{3\gamma}{3^{n+1}\gamma} \ell_2 \end{array} \right) : 1 \leq \ell_1 \leq 3\gamma - 1, 1 \leq \ell_2 \leq 3^{n+1}\gamma - 1, \ell_1, \ell_2 \in \mathbb{Z} \setminus 3\mathbb{Z} \right\}.$$

Thus there exist $(\frac{\ell_1}{3\gamma}, \frac{\ell_2}{3^{n+1}\gamma})^t \in \mathcal{G} \cap [0, 1)^2$ and $(s_1, s_2)^t \in \mathbb{Z}^2$ such that

$$\tilde{M}^{*n} \begin{pmatrix} \frac{\ell_1}{3\gamma} \\ \frac{\ell_2}{3^{n+1}\gamma} \end{pmatrix} = \begin{pmatrix} s_1 \\ s_2 \end{pmatrix}. \quad (4.8)$$

According to Lemma 4.2, there exist $a_n, b_n, c_n, d_n \in \mathbb{Z}$ such that

$$\tilde{M}^{*n} = \begin{bmatrix} 3a_n + p_1^n & 3c_n + p_1^{n-1} p_3 \\ 3b_n & 3d_n \end{bmatrix},$$

where $p_1, p_3 \in \{1, 2\}$. Now multiplying both sides of (4.8) by $3^{\eta+1}\gamma$, one may get that

$$\begin{pmatrix} \ell_2 p_1^{n-1} p_3 \\ d_n \ell_2 \end{pmatrix} = \begin{pmatrix} 3(3^\eta(\gamma s_1 - a_n \ell_1) - 3^{\eta-1} \ell_1 p_1^n + c_n \ell_2) \\ 3^\eta(\gamma s_2 - b_n \ell_1) \end{pmatrix} \in 3\mathbb{Z}^2.$$

This is impossible since $\ell_2 \notin 3\mathbb{Z}$ and $p_1, p_3 \in \{1, 2\}$. Thus $\tilde{M}^{*j} \mathcal{Z}_{\tilde{D}}^2 \cap \mathbb{Z}^2 = \emptyset$ for all $j \in \mathbb{N}$. By Theorem 2.2, $\mu_{\tilde{M}, \tilde{D}}$ is a non-spectral measure. \square

In Proposition 4.3, if $\eta = 0$ in \tilde{D} , it is interesting that for different matrices $\tilde{M}_1, \tilde{M}_2 \in \mathfrak{M}_6$, the spectrality of $\mu_{\tilde{M}_1, \tilde{D}}$ and $\mu_{\tilde{M}_2, \tilde{D}}$ may be different. The following example is devoted to display the fact.

Example 4.4. Let

$$\tilde{M}_1 = \begin{bmatrix} 4 & 0 \\ 1 & 3 \end{bmatrix}, \quad \tilde{M}_2 = \begin{bmatrix} 4 & 0 \\ 2 & 3 \end{bmatrix} \quad \text{and} \quad \tilde{D} = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right\}.$$

Then $\mu_{\tilde{M}_1, \tilde{D}}$ is a spectral measure, while $\mu_{\tilde{M}_2, \tilde{D}}$ is a non-spectral measure.

Proof. By (3.2), one can easily see that $\tilde{M}_1, \tilde{M}_2 \in \mathfrak{M}_6$. Let

$$S = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 2 \\ 2 \end{pmatrix}, \begin{pmatrix} 3 \\ 1 \end{pmatrix} \right\} \quad \text{and} \quad H = \frac{1}{\sqrt{3}} \left[e^{2\pi i \langle \tilde{M}_1^{-1} d, s \rangle} \right]_{d \in \tilde{D}, s \in S}.$$

By a simple calculation, we obtain $H^*H = I$, which implies that (\tilde{M}_1, \tilde{D}) is admissible. Then with Theorem 1.2, $\mu_{\tilde{M}_1, \tilde{D}}$ is a spectral measure.

Next we prove that $\mu_{\tilde{M}_2, \tilde{D}}$ is a non-spectral measure. Let $A = B = I$, it is evident that $A \in GL_2(3)$ and $AB = I \pmod{M_2(3\mathbb{Z})}$. Then we have $(A\tilde{M}_2B)^*(1, -1)^t = (2, -3)^t \notin 3\mathbb{Z}^2$. Together with $\det(\tilde{D}) = 1 \notin 3\mathbb{Z}$ and Theorem 1.3, it means that $\mu_{\tilde{M}_2, \tilde{D}}$ is a non-spectral measure. \square

Recall that \mathfrak{M}_k is defined by (3.2). To simplify the notation, in what follows we denote

$$\begin{aligned} \mathfrak{R}_1 &= \left\{ \tilde{M} : \tilde{M} \in \mathfrak{M}_k, k \in \{4, 6, 8, 10\} \right\}, \\ \mathfrak{R}_2 &= \left\{ \tilde{M} : \tilde{M} \in \mathfrak{M}_k, k \in \{1, 2, 3, 5, 7, 9\}, s < \eta \right\}, \\ \mathfrak{R}_3 &= \left\{ \tilde{M} : \tilde{M} \in \mathfrak{M}_k, k \in \{1, 2, 3, 5, 7, 9\}, s \geq \eta \right\}. \end{aligned}$$

To complete the proof of Theorem 2.4 in the case of $2\sigma - \omega \notin 3\mathbb{Z}$, we need the following proposition, which indicates the non-spectrality of the measure $\mu_{\tilde{M}, \tilde{D}}$ for $\tilde{M} \in \mathfrak{R}_1 \cup \mathfrak{R}_2$.

Proposition 4.5. *If $\tilde{M} \in \mathfrak{R}_1 \cup \mathfrak{R}_2$ and $2\sigma - \omega \notin 3\mathbb{Z}$ in \tilde{D} , then $\mu_{\tilde{M}, \tilde{D}}$ is a non-spectral measure.*

Proof. We first prove that $\mu_{\tilde{M}, \tilde{D}}$ is a non-spectral measure for $\tilde{M} \in \mathfrak{R}_1$.

(a) : $\tilde{M} \in \mathfrak{M}_4 \cup \mathfrak{M}_6$. Since $2\sigma - \omega \notin 3\mathbb{Z}$, it follows from Propositions 4.1 and 4.3 that $\mu_{\tilde{M}, \tilde{D}}$ is a non-spectral measure.

(b) : $\tilde{M} \in \mathfrak{M}_8 \cup \mathfrak{M}_{10}$. By (3.2), one can write $\tilde{M} \in \mathfrak{M}_8 \cup \mathfrak{M}_{10}$ as

$$\tilde{M} = \begin{bmatrix} a' & b' \\ 3^s c + p_3 & 3d + p_4 \end{bmatrix},$$

where $s \geq 1$, $a', b', d \in \mathbb{Z}$, $c \in (\mathbb{Z} \setminus 3\mathbb{Z}) \cup \{0\}$, $p_3, p_4 \in \{1, 2\}$ and $a'p_4 - b'p_3 \in 3\mathbb{Z}$. Then we can choose $\tau \in \{1, 2\}$ such that $\tau p_3 + p_4 \in 3\mathbb{Z}$. Combining this with $a'p_4 - b'p_3 \in 3\mathbb{Z}$, we obtain that

$$(\tau a' + b')p_3 = (\tau p_3 + p_4)a' - (a'p_4 - b'p_3) \in 3\mathbb{Z}. \quad (4.9)$$

Notice that $p_3 \in \{1, 2\}$, so (4.9) implies $\tau a' + b' \in 3\mathbb{Z}$. Let $\tilde{Q} = \begin{bmatrix} 1 & -\tau \\ 0 & 1 \end{bmatrix}$. Then

$$\bar{M}_1 = \tilde{Q}\tilde{M}\tilde{Q}^{-1} = \begin{bmatrix} 3^s c\tau + a' - \tau p_3 & -3^s c\tau^2 - 3d\tau + \tau a' + b' - \tau(\tau p_3 + p_4) \\ 3^s c + p_3 & 3^s c\tau + 3d + \tau p_3 + p_4 \end{bmatrix},$$

and

$$\bar{D}_1 = \tilde{Q}\tilde{D} = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \sigma \\ 0 \end{pmatrix}, \begin{pmatrix} \omega - 3^\eta \tau \vartheta \\ 3^\eta \vartheta \end{pmatrix} \right\}.$$

As $\tau_2, p_3 \in \{1, 2\}$ and $\tau a' + b', \tau_2 p_3 + p_4 \in 3\mathbb{Z}$, it can be easily seen that $\bar{M}_1 \in \mathfrak{M}_4$ if $a' - \tau p_3 \in 3\mathbb{Z}$, and $\bar{M}_1 \in \mathfrak{M}_6$ if $a' - \tau p_3 \notin 3\mathbb{Z}$. Moreover, it follows from $2\sigma - \omega \notin 3\mathbb{Z}$ and $\eta \geq 1$ that $2\sigma - (\omega - 3^\eta \tau \vartheta) \notin 3\mathbb{Z}$. Thus \bar{D}_1 has the similar properties as \tilde{D} . Hence Proposition 4.1 or Proposition 4.3 still holds for $\mu_{\bar{M}_1, \bar{D}_1}$, which implies that $\mu_{\bar{M}_1, \bar{D}_1}$ is a non-spectral measure. Therefore, $\mu_{\bar{M}, \bar{D}}$ is a non-spectral measure by Lemma 2.1.

According to (a) and (b), $\mu_{\bar{M}, \bar{D}}$ is a non-spectral measure for any $\tilde{M} \in \mathfrak{R}_1$.

Now we prove that $\mu_{\bar{M}, \bar{D}}$ is a non-spectral measure for $\tilde{M} \in \mathfrak{R}_2$. It follows from (3.2) that $\tilde{M} \in \mathfrak{R}_2$ can be expressed as

$$\tilde{M} = \begin{bmatrix} a' & b' \\ 3^s c & d' \end{bmatrix},$$

where $1 \leq s < \eta$, $a', b', d' \in \mathbb{Z}$, $c \in \mathbb{Z} \setminus 3\mathbb{Z}$ and $a'd' \in 3\mathbb{Z}$. By (3.6), we obtain that

$$\bar{M}_2 = Q_s \tilde{M} Q_s^{-1} = \begin{bmatrix} a' & 3^s b' \\ c & d' \end{bmatrix} \quad \text{and} \quad \bar{D}_2 = Q_s \tilde{D} = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \sigma \\ 0 \end{pmatrix}, \begin{pmatrix} \omega \\ 3^{\eta-s} \vartheta \end{pmatrix} \right\}.$$

In view of $a'd' \in 3\mathbb{Z}$ and $\eta - s \geq 1$, using (3.2), one can easily see that $\bar{M}_2 \in \mathfrak{R}_1$, and \bar{D}_2 has the similar form as \tilde{D} . It has been proved that $\mu_{\bar{M}, \bar{D}}$ is a non-spectral measure for $\tilde{M} \in \mathfrak{R}_1$, which implies that $\mu_{\bar{M}_2, \bar{D}_2}$ is a non-spectral measure. Then by Lemma 2.1, $\mu_{\bar{M}, \bar{D}}$ is a non-spectral measure.

The proof of Proposition 4.5 is now completed. \square

We have all ingredients for the proof of Theorem 2.4 in the case of $2\sigma - \omega \notin 3\mathbb{Z}$. That is to prove the following theorem:

Theorem 4.6. *Let \tilde{D} and \tilde{M} be defined by (2.6) and (2.7), respectively. If $2\sigma - \omega \notin 3\mathbb{Z}$ in \tilde{D} , then $\mu_{\tilde{M},\tilde{D}}$ is a spectral measure if and only if there exists a matrix $Q \in M_2(\mathbb{Z})$ such that (\bar{M}, \bar{D}) is admissible, where $\bar{M} = Q\tilde{M}Q^{-1}$ and $\bar{D} = Q\tilde{D}$.*

Proof. We first prove the sufficiency. Suppose that (\bar{M}, \bar{D}) is admissible, where $\bar{M} = Q\tilde{M}Q^{-1}$ and $\bar{D} = Q\tilde{D}$, by Theorem 1.2, $\mu_{\bar{M},\bar{D}}$ is a spectral measure. This together with Lemma 2.1 shows that $\mu_{\tilde{M},\tilde{D}}$ is a spectral measure. Hence the sufficiency follows.

Now we are devoted to proving the necessity. Suppose that $\mu_{\tilde{M},\tilde{D}}$ is a spectral measure, then Proposition 4.5 implies $\tilde{M} \in \mathfrak{R}_3$. According to the definition of \mathfrak{R}_3 and (3.2), we can write the matrix $\tilde{M} \in \mathfrak{R}_3$ as

$$\tilde{M} = \begin{bmatrix} a' & b' \\ 3^s c & d' \end{bmatrix},$$

where $s \geq \eta$, $a', b', d' \in \mathbb{Z}$, $c \in (\mathbb{Z} \setminus 3\mathbb{Z}) \cup \{0\}$ and $a'd' \in 3\mathbb{Z}$. Applying (3.6), we get

$$\bar{M}_3 = Q_\eta \tilde{M} Q_\eta^{-1} = \begin{bmatrix} a' & 3^\eta b' \\ 3^{s-\eta} c & d' \end{bmatrix} \quad \text{and} \quad \bar{D}_3 = Q_\eta \tilde{D} = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \sigma \\ 0 \end{pmatrix}, \begin{pmatrix} \omega \\ \vartheta \end{pmatrix} \right\}.$$

Since $\mu_{\bar{M},\bar{D}}$ is a spectral measure, it follows from Lemma 2.1 that $\mu_{\bar{M}_3,\bar{D}_3}$ is also a spectral measure. Together with $\sigma\vartheta \notin 3\mathbb{Z}$ and Theorem 1.3, it gives that (\bar{M}_3, \bar{D}_3) is admissible.

This completes the proof of Theorem 4.6. \square

Remark 4.7. Under the assumption of $2\sigma - \omega \notin 3\mathbb{Z}$ in \tilde{D} , from the proof of Theorem 2.4 and Theorem 1.3, we can easily see that $\mu_{\tilde{M},\tilde{D}}$ is a spectral measure if and only if $s \geq \eta$ and $(AQ_\eta \tilde{M} Q_\eta^{-1} B)^*(1, -1)^t \in 3\mathbb{Z}^2$, where $B = \begin{bmatrix} \sigma & \omega \\ 0 & \vartheta \end{bmatrix}$ and $A \in GL_2(3)$ satisfies

$AB = I \pmod{M_2(3\mathbb{Z})}$. By noting that $\sigma, \vartheta \notin 3\mathbb{Z}$, it is clear that $A = \sigma\vartheta \begin{bmatrix} \vartheta & -\omega \\ 0 & \sigma \end{bmatrix}$ satisfies $AB = (\sigma\vartheta)^2 I = I \pmod{M_2(3\mathbb{Z})}$.

Now we give an example to illustrate the fact in the above remark.

Example 4.8. Let

$$M_1 = \begin{bmatrix} 8 & -5 \\ 4 & -1 \end{bmatrix}, \quad M_2 = \begin{bmatrix} 5 & -1 \\ 2 & 2 \end{bmatrix} \quad \text{and} \quad D = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 2 \\ 4 \end{pmatrix} \right\}.$$

Then $\mu_{M_1,D}$ is a spectral measure, while $\mu_{M_2,D}$ is a non-spectral measure.

Proof. Let $P = \begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix}$. Then we obtain that

$$\tilde{M}_1 = PM_1P^{-1} = \begin{bmatrix} 4 & 0 \\ 3 & 3 \end{bmatrix}, \quad \tilde{M}_2 = PM_2P^{-1} = \begin{bmatrix} 3 & 0 \\ 3 & 4 \end{bmatrix},$$

and

$$\tilde{D} = PD = \left\{ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} -2 \\ 6 \end{pmatrix} \right\}.$$

According to (2.6), it is easy to see that $\sigma = \eta = 1$, $\omega = -2$ and $\vartheta = 2$. Therefore, A, B and Q_η in Remark 4.7 can be written as

$$A = \begin{bmatrix} 4 & 4 \\ 0 & 2 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & -2 \\ 0 & 2 \end{bmatrix} \quad \text{and} \quad Q_\eta = \begin{bmatrix} 1 & 0 \\ 0 & \frac{1}{3} \end{bmatrix}.$$

By calculations, we get

$$(AQ_\eta \tilde{M}_1 Q_\eta^{-1} B)^* \begin{pmatrix} 1 \\ -1 \end{pmatrix} = \begin{bmatrix} 20 & 2 \\ -16 & 8 \end{bmatrix} \begin{pmatrix} 1 \\ -1 \end{pmatrix} = \begin{pmatrix} 18 \\ -24 \end{pmatrix} \in 3\mathbb{Z}^2,$$

and

$$(AQ_\eta \tilde{M}_2 Q_\eta^{-1} B)^* \begin{pmatrix} 1 \\ -1 \end{pmatrix} = \begin{bmatrix} 16 & 2 \\ 0 & 12 \end{bmatrix} \begin{pmatrix} 1 \\ -1 \end{pmatrix} = \begin{pmatrix} 14 \\ -12 \end{pmatrix} \notin 3\mathbb{Z}^2.$$

Then by using Theorem 1.3, one can conclude that $\mu_{Q_\eta \tilde{M}_1 Q_\eta^{-1}, Q_\eta \tilde{D}}$ is a spectral measure and $\mu_{Q_\eta \tilde{M}_2 Q_\eta^{-1}, Q_\eta \tilde{D}}$ is a non-spectral measure. Hence the assertion follows by Lemma 2.1. \square

Remark 4.9. Form the proof of Example 4.8, we know that $(Q_\eta \tilde{M}_1 Q_\eta^{-1}, Q_\eta \tilde{D})$ is admissible. However, (\tilde{M}_1, \tilde{D}) is not admissible since $\mathcal{Z}(m_{\tilde{D}}) \subset \mathfrak{G} := \{(\ell_1/3, \ell_2/18)^t : \ell_1, \ell_2 \notin 3\mathbb{Z}\}$ and $\tilde{M}_1^* \mathfrak{G} \cap \mathbb{Z}^2 = \emptyset$. This means that the matrix Q is nontrivial in Theorem 1.4.

At the end of this paper, we give some remarks and an open problem which related to our main results. In the paper, the integer digit set $D = \{(0, 0)^t, (\alpha_1, \alpha_2)^t, (\beta_1, \beta_2)^t\}$ satisfies $\alpha_1\beta_2 - \alpha_2\beta_1 \neq 0$, it is of interest to consider the following question:

(Qu 2): For an expanding matrix $M \in M_2(\mathbb{Z})$ and the digit set D given by (1.2) with $\alpha_1\beta_2 - \alpha_2\beta_1 = 0$, what is the sufficient and necessary condition for $\mu_{M,D}$ to be a spectral measure?

It is obvious that $\alpha_1\beta_2 - \alpha_2\beta_1 = 0$ implies the digit set D is collinear, i.e., $D = \{0, a, b\}v$ for some $a, b \in \mathbb{Z}$ and $v \in \mathbb{Z}^2$. At this time, it is easy to show that $\mathcal{Z}(m_D) \neq \emptyset$ if and only if $\{a, b\} = \{1, 2\} \pmod{3}$. If $\{a, b\} = \{1, 2\} \pmod{3}$, by using the similar proof as Theorem 1.2 in [35], we can prove that $\mu_{M,D}$ is a spectral measure if $\det(M) \in 3\mathbb{Z}$. However, we know nothing about the necessary condition. We conjecture that $\det(M) \in 3\mathbb{Z}$ is also the necessary condition for $\mu_{M,D}$ to be a spectral measure.

References

- [1] L.X. An, X.G. He, A class of spectral Moran measures, *J. Funct. Anal.*, **266**(1) (2014), 343–354.
- [2] L.X. An, X.G. He, K.S. Lau, Spectrality of a class of infinite convolutions, *Adv. Math.*, **283** (2015), 362–376.
- [3] L.X. An, X.G. He, L. Tao, Spectrality of the planar Sierpinski family, *J. Math. Anal. Appl.*, **432**(2) (2015), 725–732.

- [4] D. Bose, S. Madan, “Spectral implies tiling” for three intervals revisited, *Forum Math.*, **26**(4) (2014), 1247–1260.
- [5] M.L. Chen, J.C. Liu, The cardinality of orthogonal exponentials of planar self-affine measures with three-element digit sets, *J. Funct. Anal.*, **277**(1) (2019), 135–156.
- [6] X.R. Dai, When does a Bernoulli convolution admit a spectrum? *Adv. Math.*, **231** (2012), 1681–1693.
- [7] X.R. Dai, X.G. He, C.K. Lai, Spectral property of Cantor measures with consecutive digits, *Adv. Math.*, **242** (2013), 187–208.
- [8] X.R. Dai, X.G. He, K.S. Lau, On spectral N-Bernoulli measures, *Adv. Math.*, **259** (2014), 511–531.
- [9] Q.R. Deng, On the spectra of Sierpinski-type self-affine measures, *J. Funct. Anal.*, **270**(12) (2016), 4426–4442.
- [10] Q.R. Deng, K.S. Lau, Sierpinski-type spectral self-similar measures, *J. Funct. Anal.*, **269**(5) (2015), 1310–1326.
- [11] D. Dutkay, C.K. Lai, Uniformity of measures with Fourier frames, *Adv. Math.*, **252** (2014), 684–707.
- [12] D. Dutkay, J. Haussermann, C.K. Lai, Hadamard triples generate self-affine spectral measures, *Trans. Amer. Math. Soc.*, **371**(2) (2019), 1439–1481.
- [13] D. Dutkay, P. Jorgensen, Wavelets on fractals, *Rev. Mat. Iberoam.*, **22**(1) (2006), 131–180.
- [14] D. Dutkay, P. Jorgensen, Iterated function systems, Ruelle operators, and invariant projective measures, *Math. Comp.*, **75** (2006), 1931–1970.
- [15] D. Dutkay, P. Jorgensen, Analysis of orthogonality and of orbits in affine iterated function systems, *Math. Z.*, **256**(4) (2007), 801–823.
- [16] D. Dutkay, P. Jorgensen, Fourier frequencies in affine iterated function systems, *J. Funct. Anal.*, **247**(1) (2007), 110–137.
- [17] D. Dutkay, P. Jorgensen, Probability and Fourier duality for affine iterated function systems, *Acta Appl. Math.*, **107** (2009), 293–311.
- [18] A.H. Fan, S.L. Fan, L.M. Liao, R.X. Shi, Fuglede’s conjecture holds in \mathbb{Q}_p , *Math. Ann.*, **375** (2019), 315–341.
- [19] B. Fuglede, Commuting self-adjoint partial differential operators and a group theoretic problem, *J. Funct. Anal.*, **16** (1974), 101–121.
- [20] X.G. He, C.K. Lai, K.S. Lau, Exponential spectra in $L^2(\mu)$, *Appl. Comput. Harmon. Anal.*, **34**(3) (2013), 327–338.
- [21] J. Hutchinson, Fractals and self-similarity, *Indiana Univ. Math. J.*, **30** (1981), 713–747.
- [22] T.Y. Hu, K.S. Lau, Spectral property of the Bernoulli convolutions, *Adv. Math.*, **219** (2008), 554–567.
- [23] A. Iosevich, N. Katz, T. Tao, The Fuglede spectral conjecture holds for convex planar domains, *Math. Res. Lett.*, **10** (2003), 559–569.
- [24] P. Jorgensen, S. Pedersen, Dense analytic subspaces in fractal L^2 -spaces, *J. Anal. Math.*, **75** (1998), 185–228.
- [25] M. Kolountzakis, M. Matolcsi, Complex Hadamard matrices and the spectral set conjecture, *Collect. Math.*, **57** (2006), 281–291.
- [26] M. Kolountzakis, M. Matolcsi, Tiles with no spectra, *Forum Math.*, **18** (2006), 519–528.
- [27] I. Łaba, Fuglede’s conjecture for a union of two intervals, *Proc. Am. Math. Soc.*, **129**(10) (2001), 2965–2972.
- [28] I. Łaba, Y. Wang, On spectral Cantor measures, *J. Funct. Anal.*, **193** (2002), 409–420.
- [29] J.L. Li, Non-spectral problem for a class of planar self-affine measures, *J. Funct. Anal.*, **255**(11) (2008), 3125–3148.
- [30] J.L. Li, On the $\mu_{M,D}$ -orthogonal exponentials, *Nonlinear Anal.*, **73**(4) (2010), 940–951.
- [31] J.L. Li, A necessary and sufficient condition for the finite $\mu_{M,D}$ -orthogonality, *Sci. China Math.*, **58**(12) (2015), 2541–2548.

- [32] J.L. Li, Spectrality of self-affine measures on the three-dimensional Sierpinski gasket, *Proc. Edinb. Math. Soc.*, **55**(2) (2012), 477–496.
- [33] J.L. Li, Spectral self-affine measures on the planar Sierpinski family, *Sci. China Math.*, **56**(8) (2013), 1619–1628.
- [34] J.C. Liu, X.H. Dong, J.L. Li, Non-spectral problem for the self-affine measures, *J. Funct. Anal.*, **273**(2) (2017), 705–720.
- [35] J.C. Liu, J.J. Luo, Spectral property of self-affine measures on \mathbb{R}^n , *J. Funct. Anal.*, **272**(2) (2017), 599–612.
- [36] J.C. Liu, Z.Y. Wang, The spectrality of self-affine measure under the similarity transformation of $GL_n(p)$, arXiv:2008.07047.
- [37] S. Nitzan, A. Olevskii, A. Ulanovskii, Exponential frames on unbounded sets, *Proc. Amer. Math. Soc.*, **144**(1) (2016), 109–118.
- [38] R. Strichartz, Mock Fourier series and transforms associated with certain Cantor measures, *J. Anal. Math.*, **81** (2000), 209–238.
- [39] T. Tao, Fuglede’s conjecture is false in 5 and higher dimensions, *Math. Res. Lett.*, **11** (2004), 251–258.

KEY LABORATORY OF HIGH PERFORMANCE COMPUTING AND STOCHASTIC INFORMATION PROCESSING (MINISTRY OF EDUCATION OF CHINA), COLLEGE OF MATHEMATICS AND STATISTICS, HUNAN NORMAL UNIVERSITY, CHANGSHA, HUNAN 410081, CHINA

Email address: jcliu@hunnu.edu.cn

Email address: zhangying19952020@163.com

COLLEGE OF MATHEMATICS AND COMPUTATIONAL SCIENCE, HUNAN FIRST NORMAL UNIVERSITY, CHANGSHA, HUNAN 410205, P. R. CHINA

Email address: wzyzzq1@163.com

SCHOOL OF MATHEMATICS, SUN YAT-SEN UNIVERSITY, GUANGZHOU 510275, P. R. CHINA

Email address: mathcm1@163.com