

LOCAL AND 2-LOCAL DERIVATIONS OF THE OCTONION ALGEBRA

F.N. ARZIKULOV, I.A. KARIMJANOV, AND S. UGUZ

ABSTRACT. In the present paper, we prove that a local derivation on the octonion (Cayley) algebra \mathbb{O} over an arbitrary field, satisfying some conditions is a derivation, and every 2-local derivation on \mathbb{O} is a Jordan derivation.

1. INTRODUCTION

In many classes of algebras, local and 2-local derivations are characterized by the algebra derivations. At present, many specialists in this field are dealing with the problem of describing local and 2-local derivations of various type of algebras. Currently, this problem is more attracting and more interesting.

Local derivations were first introduced and investigated in 1990 in [4] by Kadison and in [5] by Larson and Sourour. Kadison proved that every continuous local derivation of a von Neumann algebra M into a dual Banach M -bimodule is a derivation. Larson and Sourour proved that every linear local derivation on $B(X)$, the algebra of all bounded linear operators on a Banach space X , is a derivation. The result of Kadison gave way to a cascade of results and studies on derivations on C^* -algebras, culminating with a definitive contribution due to Johnson, which asserts that every local derivation (no continuity is assumed) of a C^* -algebra A into a Banach A -bimodule is a derivation [3].

In 1997, P. Šemrl, in his paper [6], introduced and studied 2-local derivations on the algebra of bounded linear operators on a Hilbert space. P. Šemrl proved that each 2-local derivation on the algebra of bounded linear operators on a separable Hilbert space is a derivation. The theory of 2-local derivations received the most rapid development after the paper [1] by Sh. Ayupov and K. Kudaybergenov, who proved that every 2-local derivation on the algebra of bounded linear operators on an arbitrary Hilbert space (no separability is assumed) is a derivation.

In the present paper we prove that every local and 2-local derivation on the octonion algebra over an arbitrary field are derivations. The octonion algebra constructed by Arthur Cayley in 1945. The octonion algebra satisfies the next weak-associative law: $(aa)b = a(ab)$, $(ab)b = a(bb)$. Algebras which satisfies two previous identities called alternative algebras. It is well known that the octonion algebra is unique non-associative simple alternative algebra. Therefore,

2010 *Mathematics Subject Classification.* 17A36, 17D10.

Key words and phrases. Local derivation, 2-local derivation, derivation, octonion algebra.

This work was supported by Agencia Estatal de Investigacion on (Spain), grant MTM2016-79661-P and by Xunta de Galicia, Grant ED431C 2019/10 (European FEDER support included, UE10.13039/501100002924).

we have the description of local and 2-local derivations of finite-dimensional semi-simple alternative algebras under some conditions. We proved that every 2-local derivation of a finite-dimensional semi-simple alternative algebra is a Jordan derivation.

The outline of paper is given as follows: we set up the basic definitions of octonion algebra and the notions of local and 2-local derivations of algebra in Section 2. The local and 2-local derivations of octonion algebra are presented in Section 3. The conclusions of the present study are presented in Section 4.

2. PRELIMINARIES

An octonion algebra or Cayley number algebra \mathbb{O} over an arbitrary field \mathbb{F} is 8-dimensional algebra generated by $\mathcal{B} = \{e_0, e_1, \dots, e_7\}$ with the next product table

		e_j							
		e_0	e_1	e_2	e_3	e_4	e_5	e_6	e_7
e_i	e_0	e_0	e_1	e_2	e_3	e_4	e_5	e_6	e_7
	e_1	e_1	$-e_0$	e_3	$-e_2$	e_5	$-e_4$	$-e_7$	e_6
	e_2	e_2	$-e_3$	$-e_0$	e_1	e_6	e_7	$-e_4$	$-e_5$
	e_3	e_3	e_2	$-e_1$	$-e_0$	e_7	$-e_6$	e_5	$-e_4$
	e_4	e_4	$-e_5$	$-e_6$	$-e_7$	$-e_0$	e_1	e_2	e_3
	e_5	e_5	e_4	$-e_7$	e_6	$-e_1$	$-e_0$	$-e_3$	e_2
	e_6	e_6	e_7	e_4	$-e_5$	$-e_2$	e_3	$-e_0$	$-e_1$
	e_7	e_7	$-e_6$	e_5	e_4	$-e_3$	$-e_2$	e_1	$-e_0$

The table can be summarized as follows:

$$e_i e_j = \begin{cases} e_j, & \text{if } i = 0; \\ e_i, & \text{if } j = 0; \\ -\delta_{ij} e_0 + \varepsilon_{ijk} e_k, & \text{otherwise.} \end{cases}$$

where δ_{ij} is the Kronecker delta and ε_{ijk} is a completely antisymmetric tensor with value +1 when $ijk = 123, 145, 176, 246, 257, 347, 365$ (cf. [7], [2, Chapter I, §5]).

Definition 2.1. Let \mathcal{A} be an algebra. A linear map $D : \mathcal{A} \rightarrow \mathcal{A}$ is called a derivation of \mathcal{A} , if

$$d(xy) = d(x)y + xd(y), \quad \text{for any } x, y \in \mathcal{A}.$$

The algebra of all derivations is denoted by $\mathcal{D}(\mathcal{A})$. Now we define the notions of local and 2-local derivations of algebra.

Definition 2.2. Let \mathcal{A} be an algebra. A linear map $\nabla : \mathcal{A} \rightarrow \mathcal{A}$ is called a local derivation, if for any element $x \in \mathcal{A}$ there exists a derivation $D_x : \mathcal{A} \rightarrow \mathcal{A}$ such that $\nabla(x) = D_x(x)$.

Definition 2.3. Let \mathcal{A} be an algebra. A (not necessary linear) map $\Delta : \mathcal{A} \rightarrow \mathcal{A}$ is called a 2-local derivation, if for any two elements $x, y \in \mathcal{A}$ there exists a derivation $D_{x,y} : \mathcal{A} \rightarrow \mathcal{A}$ such that $\Delta(x) = D_{x,y}(x)$, $\Delta(y) = D_{x,y}(y)$.

Definition 2.4. Let \mathcal{A} be an algebra. A linear map $D : \mathcal{A} \rightarrow \mathcal{A}$ is called a Jordan derivation, if, for any two elements $x, y \in \mathcal{A}$,

$$D(xy + yx) = D(x)y + yD(x) + xD(y) + D(y)x.$$

3. LOCAL AND 2-LOCAL DERIVATIONS OF THE ALGEBRA OF OCTONIONS

Proposition 3.1. A linear map $D : \mathbb{O} \rightarrow \mathbb{O}$ is a derivation if and only if the matrix of D has the following matrix form:

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & a_{12} & a_{13} & a_{14} & a_{15} & a_{16} & a_{17} \\ 0 & -a_{12} & 0 & a_{23} & a_{24} & a_{25} & a_{26} & a_{27} \\ 0 & -a_{13} & -a_{23} & 0 & a_{16} - a_{25} & a_{17} + a_{24} & -a_{14} + a_{27} & -a_{15} - a_{26} \\ 0 & -a_{14} & -a_{24} & -a_{16} + a_{25} & 0 & a_{45} & a_{46} & a_{47} \\ 0 & -a_{15} & -a_{25} & -a_{17} - a_{24} & -a_{45} & 0 & a_{12} + a_{47} & a_{13} - a_{46} \\ 0 & -a_{16} & -a_{26} & a_{14} - a_{27} & -a_{46} & -a_{12} - a_{47} & 0 & a_{23} + a_{45} \\ 0 & -a_{17} & -a_{27} & a_{15} + a_{26} & -a_{47} & -a_{13} + a_{46} & -a_{23} - a_{45} & 0 \end{pmatrix}.$$

Proof. Let D be a derivation of the algebra \mathbb{O} . It is easy to see that element e_0 is identity element and from the equalities

$$D(e_0) = D(e_0e_0) = D(e_0)e_0 + e_0D(e_0) = 2D(e_0)$$

we obtain $D(e_0) = 0$.

We set

$$D(e_i) = \sum_{j=0}^7 a_{ji}e_j, \quad 1 \leq j \leq 7.$$

Note that, the multiplications of basis elements $\{e_1, e_2, \dots, e_7\}$ are anti-commutative. So, from

$$0 = -D(e_0) = D(e_i e_i) = D(e_i)e_i + e_i D(e_i) = 2a_{0i}e_i - 2a_{ii}e_0$$

we have $a_{0i} = a_{ii} = 0$ where $1 \leq i \leq 7$.

Applying again the identity of derivation for all possible cases we obtain the matrix of derivation. \square

Theorem 3.2. Let ∇ be a local derivation of \mathbb{O} with a matrix

$$A = \begin{pmatrix} b_{00} & b_{01} & b_{02} & b_{03} & b_{04} & b_{05} & b_{06} & b_{07} \\ b_{10} & b_{11} & b_{12} & b_{13} & b_{14} & b_{15} & b_{16} & b_{17} \\ b_{20} & b_{21} & b_{22} & b_{23} & b_{24} & b_{25} & b_{26} & b_{27} \\ b_{30} & b_{31} & b_{32} & b_{33} & b_{34} & b_{35} & b_{36} & b_{37} \\ b_{40} & b_{41} & b_{42} & b_{43} & b_{44} & b_{45} & b_{46} & b_{47} \\ b_{50} & b_{51} & b_{52} & b_{53} & b_{54} & b_{55} & b_{56} & b_{57} \\ b_{60} & b_{61} & b_{62} & b_{63} & b_{64} & b_{65} & b_{66} & b_{67} \\ b_{70} & b_{71} & b_{72} & b_{73} & b_{74} & b_{75} & b_{76} & b_{77} \end{pmatrix}.$$

Then the following conditions are equivalent

- (1) ∇ is a derivation,

(2) the matrix A satisfies the following conditions

$$\begin{aligned} b_{16} - b_{25} &= b_{34} = -b_{43}, b_{17} + b_{24} = b_{35} = -b_{53}, \\ -b_{14} + b_{27} &= b_{36} = -b_{63}, -b_{15} - b_{26} = b_{37} = -b_{73}, \\ b_{12} + b_{47} &= b_{56} = -b_{65}, b_{13} - b_{46} = b_{57} = -b_{75}, b_{23} + b_{45} = b_{67} = -b_{76}. \end{aligned}$$

Proof. The implication (1) \implies (2) is obvious. We prove the implication (2) \implies (1). Let ∇ be an arbitrary local derivation of \mathbb{O} . By the definition for all $x \in \mathbb{O}$ there exists a derivation D_x on \mathbb{O} such that

$$\nabla(x) = D_x(x).$$

By proposition 3.1, the derivation D_x has the following matrix form:

$$A^x = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & a_{12}^x & a_{13}^x & a_{14}^x & a_{15}^x & a_{16}^x & a_{17}^x \\ 0 & -a_{12}^x & 0 & a_{23}^x & a_{24}^x & a_{25}^x & a_{26}^x & a_{27}^x \\ 0 & -a_{13}^x & -a_{23}^x & 0 & a_{16}^x - a_{25}^x & a_{17}^x + a_{24}^x & -a_{14}^x + a_{27}^x & -a_{15}^x - a_{26}^x \\ 0 & -a_{14}^x & -a_{24}^x & -a_{16}^x + a_{25}^x & 0 & a_{45}^x & a_{46}^x & a_{47}^x \\ 0 & -a_{15}^x & -a_{25}^x & -a_{17}^x - a_{24}^x & -a_{45}^x & 0 & a_{12}^x + a_{47}^x & a_{13}^x - a_{46}^x \\ 0 & -a_{16}^x & -a_{26}^x & a_{14}^x - a_{27}^x & -a_{46}^x & -a_{12}^x - a_{47}^x & 0 & a_{23}^x + a_{45}^x \\ 0 & -a_{17}^x & -a_{27}^x & a_{15}^x + a_{26}^x & -a_{47}^x & -a_{13}^x + a_{46}^x & -a_{23}^x - a_{45}^x & 0 \end{pmatrix}$$

Let A be the matrix of ∇ , then by choosing subsequently $x = e_0, x = e_1, \dots, x = e_7$, and using $\nabla(x) = D_x(x)$, i.e. $A\bar{x} = A^x\bar{x}$, where \bar{x} is the vector corresponding to x , it is easy to see that

$$A = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & a_{12}^{e_2} & a_{13}^{e_3} & a_{14}^{e_4} & a_{15}^{e_5} & a_{16}^{e_6} & a_{17}^{e_7} \\ 0 & -a_{12}^{e_1} & 0 & -a_{23}^{e_3} & a_{24}^{e_4} & a_{25}^{e_5} & a_{26}^{e_6} & a_{27}^{e_7} \\ 0 & -a_{13}^{e_1} & a_{23}^{e_2} & 0 & a_{16}^{e_4} - a_{25}^{e_4} & a_{17}^{e_5} + a_{24}^{e_5} & -a_{14}^{e_6} + a_{27}^{e_6} & -a_{15}^{e_7} - a_{26}^{e_7} \\ 0 & -a_{14}^{e_1} & a_{24}^{e_2} & -a_{16}^{e_3} + a_{25}^{e_3} & 0 & a_{45}^{e_5} & a_{46}^{e_6} & a_{47}^{e_7} \\ 0 & -a_{15}^{e_1} & a_{25}^{e_2} & -a_{17}^{e_3} - a_{24}^{e_3} & -a_{45}^{e_4} & 0 & a_{12}^{e_6} + a_{47}^{e_6} & a_{13}^{e_7} - a_{46}^{e_7} \\ 0 & -a_{16}^{e_1} & a_{26}^{e_2} & a_{14}^{e_3} - a_{27}^{e_3} & -a_{46}^{e_4} & -a_{12}^{e_5} - a_{47}^{e_5} & 0 & a_{23}^{e_7} + a_{45}^{e_7} \\ 0 & -a_{17}^{e_1} & -a_{27}^{e_2} & a_{15}^{e_3} + a_{26}^{e_3} & -a_{47}^{e_4} & -a_{13}^{e_5} + a_{46}^{e_5} & -a_{23}^{e_6} - a_{45}^{e_6} & 0 \end{pmatrix}.$$

From the following equalities $\nabla(e_1 + e_i) = \nabla(e_1) + \nabla(e_i)$ for $2 \leq i \leq 7$, we derive that

$$a_{1i}^{e_1+e_i} = a_{1i}^{e_1}, \quad a_{1i}^{e_1+e_i} = a_{1i}^{e_i}, \quad \text{i.e. } a_{1i}^{e_1} = a_{1i}^{e_i}, \quad 2 \leq i \leq 7.$$

Then the equalities $\nabla(e_2 + e_i) = \nabla(e_2) + \nabla(e_i)$ for $3 \leq i \leq 7$, imply

$$a_{2i}^{e_2+e_i} = a_{2i}^{e_2}, \quad a_{2i}^{e_1+e_i} = a_{2i}^{e_i}, \quad \text{i.e. } a_{2i}^{e_2} = a_{2i}^{e_i}, \quad 3 \leq i \leq 7.$$

From the equalities $\nabla(e_3 + e_i) = \nabla(e_3) + \nabla(e_i)$ for $4 \leq i \leq 7$, we conclude that

$$\begin{aligned} a_{16}^{e_3} - a_{25}^{e_3} &= a_{16}^{e_4} - a_{25}^{e_4}, & a_{17}^{e_3} + a_{24}^{e_3} &= a_{17}^{e_5} + a_{24}^{e_5}, \\ a_{14}^{e_3} - a_{27}^{e_3} &= a_{14}^{e_6} - a_{27}^{e_6}, & a_{15}^{e_3} + a_{26}^{e_3} &= a_{15}^{e_7} + a_{26}^{e_7}. \end{aligned}$$

From $\nabla(e_4 + e_i) = \nabla(e_4) + \nabla(e_i)$ for $5 \leq i \leq 7$, we deduce that

$$a_{4i}^{e_4} = a_{4i}^{e_i}, \quad 5 \leq i \leq 7.$$

Similarly as above, from the equalities: $\nabla(e_5 + e_i) = \nabla(e_3) + \nabla(e_i)$ for $i = 5, 6$ we obtain

$$a_{12}^{e_5} + a_{47}^{e_5} = a_{12}^{e_6} + a_{47}^{e_6}, \quad a_{13}^{e_5} - a_{46}^{e_5} = a_{13}^{e_7} - a_{46}^{e_7}.$$

Finally, from $\nabla(e_6 + e_7) = \nabla(e_6) + \nabla(e_7)$ we get

$$a_{23}^{e_6} + a_{45}^{e_6} = a_{23}^{e_7} + a_{45}^{e_7}.$$

By re-denoting parameters $a_{ij}^{e_k}$ to α_{ij} we have the next matrix

$$A = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \alpha_{12} & \alpha_{13} & \alpha_{14} & \alpha_{15} & \alpha_{16} & \alpha_{17} \\ 0 & -\alpha_{12} & 0 & \alpha_{23} & \alpha_{24} & \alpha_{25} & \alpha_{26} & \alpha_{27} \\ 0 & -\alpha_{13} & -\alpha_{23} & 0 & \alpha'_{16} - \alpha'_{25} & \alpha'_{17} + \alpha'_{24} & -\alpha'_{14} + \alpha'_{27} & -\alpha'_{15} - \alpha'_{26} \\ 0 & -\alpha_{14} & -\alpha_{24} & -\alpha'_{16} + \alpha'_{25} & 0 & \alpha_{45} & \alpha_{46} & \alpha_{47} \\ 0 & -\alpha_{15} & -\alpha_{25} & -\alpha'_{17} - \alpha'_{24} & -\alpha_{45} & 0 & \alpha'_{12} + \alpha'_{47} & \alpha'_{13} - \alpha'_{46} \\ 0 & -\alpha_{16} & -\alpha_{26} & \alpha'_{14} - \alpha'_{27} & -\alpha_{46} & -\alpha'_{12} - \alpha'_{47} & 0 & \alpha'_{23} + \alpha'_{45} \\ 0 & -\alpha_{17} & -\alpha_{27} & \alpha'_{15} + \alpha'_{26} & -\alpha_{47} & -\alpha'_{13} + \alpha'_{46} & -\alpha'_{23} - \alpha'_{45} & 0 \end{pmatrix}$$

Then we can represent the matrix A as the sum of the following two matrices:

$$A = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \alpha_{12} & \alpha_{13} & \alpha_{14} & \alpha_{15} & \alpha_{16} & \alpha_{17} \\ 0 & -\alpha_{12} & 0 & \alpha_{23} & \alpha_{24} & \alpha_{25} & \alpha_{26} & \alpha_{27} \\ 0 & -\alpha_{13} & -\alpha_{23} & 0 & \alpha_{16} - \alpha_{25} & \alpha_{17} + \alpha_{24} & -\alpha_{14} + \alpha_{27} & -\alpha_{15} - \alpha_{26} \\ 0 & -\alpha_{14} & -\alpha_{24} & -\alpha_{16} + \alpha_{25} & 0 & \alpha_{45} & \alpha_{46} & \alpha_{47} \\ 0 & -\alpha_{15} & -\alpha_{25} & -\alpha_{17} - \alpha_{24} & -\alpha_{45} & 0 & \alpha_{12} + \alpha_{47} & \alpha_{13} - \alpha_{46} \\ 0 & -\alpha_{16} & -\alpha_{26} & \alpha_{14} - \alpha_{27} & -\alpha_{46} & -\alpha_{12} - \alpha_{47} & 0 & \alpha_{23} + \alpha_{45} \\ 0 & -\alpha_{17} & -\alpha_{27} & \alpha_{15} + \alpha_{26} & -\alpha_{47} & -\alpha_{13} + \alpha_{46} & -\alpha_{23} - \alpha_{45} & 0 \end{pmatrix}$$

$$+ \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \lambda_{16}^{25} & \lambda_{17}^{24} & \lambda_{14}^{27} & \lambda_{15}^{26} \\ 0 & 0 & 0 & -\lambda_{16}^{25} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -\lambda_{17}^{24} & 0 & 0 & \lambda_{12}^{47} & \lambda_{13}^{46} \\ 0 & 0 & 0 & -\lambda_{14}^{27} & 0 & -\lambda_{12}^{47} & 0 & \lambda_{23}^{45} \\ 0 & 0 & 0 & -\lambda_{15}^{26} & 0 & -\lambda_{13}^{46} & -\lambda_{23}^{45} & 0 \end{pmatrix}$$

$$\lambda_{16}^{25} = \alpha'_{16} - \alpha'_{25} - (\alpha_{16} - \alpha_{25}), \lambda_{17}^{24} = \alpha'_{17} + \alpha'_{24} - (\alpha_{17} + \alpha_{24}), \lambda_{14}^{27} = -\alpha'_{14} + \alpha'_{27} - (-\alpha_{14} + \alpha_{27}),$$

$$\lambda_{15}^{26} = -\alpha'_{15} - \alpha'_{26} - (-\alpha_{15} - \alpha_{26}), \lambda_{12}^{47} = \alpha'_{12} + \alpha'_{47} - (\alpha_{12} + \alpha_{47}), \lambda_{13}^{46} = \alpha'_{13} - \alpha'_{46} - (\alpha'_{13} - \alpha_{46}),$$

$$\lambda_{23}^{45} = \alpha'_{23} + \alpha'_{45} - (\alpha_{23} + \alpha_{45}).$$

By the assumption

$$\lambda_{16}^{25} = 0, \lambda_{17}^{24} = 0, \lambda_{14}^{27} = 0, \lambda_{15}^{26} = 0, \lambda_{12}^{47} = 0, \lambda_{13}^{46} = 0, \lambda_{23}^{45} = 0.$$

By Proposition 3.1 this ends the proof. \square

Conjecture 3.3. Let ∇ be a linear operator on \mathbb{O} with a nonzero matrix

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \lambda_{16}^{25} & \lambda_{17}^{24} & \lambda_{14}^{27} & \lambda_{15}^{26} \\ 0 & 0 & 0 & -\lambda_{16}^{25} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -\lambda_{17}^{24} & 0 & 0 & \lambda_{12}^{47} & \lambda_{13}^{46} \\ 0 & 0 & 0 & -\lambda_{14}^{27} & 0 & -\lambda_{12}^{47} & 0 & \lambda_{23}^{45} \\ 0 & 0 & 0 & -\lambda_{15}^{26} & 0 & -\lambda_{13}^{46} & -\lambda_{23}^{45} & 0 \end{pmatrix},$$

where $\lambda_{16}^{25}, \lambda_{17}^{24}, \lambda_{14}^{27}, \lambda_{15}^{26}, \lambda_{12}^{47}, \lambda_{13}^{46}, \lambda_{23}^{45}$ are elements in the field \mathbb{F} . Then ∇ is a local derivation which is not a derivation.

Theorem 3.4. *Every 2-local derivation of the algebra of octonions \mathbb{O} is linear, i.e., a local derivation.*

Proof. Let Δ be an arbitrary 2-local derivation of \mathbb{O} . By the definition, for all $x, y \in \mathbb{O}$ there exists a derivation $D_{x,y}$ of \mathbb{O} such that

$$\Delta(x) = D_{x,y}(x), \quad \Delta(y) = D_{x,y}(y).$$

By proposition 3.1, the derivation $D_{x,y}$ has the following matrix form:

$$A^{x,y} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & a_{12}^{xy} & a_{13}^{xy} & a_{14}^{xy} & a_{15}^{xy} & a_{16}^{xy} & a_{17}^{xy} \\ 0 & -a_{12}^{xy} & 0 & a_{23}^{xy} & a_{24}^{xy} & a_{25}^{xy} & a_{26}^{xy} & a_{27}^{xy} \\ 0 & -a_{13}^{xy} & -a_{23}^{xy} & 0 & a_{16}^{xy} - a_{25}^{xy} & a_{17}^{xy} + a_{24}^{xy} & -a_{14}^{xy} + a_{27}^{xy} & -a_{15}^{xy} - a_{26}^{xy} \\ 0 & -a_{14}^{xy} & -a_{24}^{xy} & -a_{16}^{xy} + a_{25}^{xy} & 0 & a_{45}^{xy} & a_{46}^{xy} & a_{47}^{xy} \\ 0 & -a_{15}^{xy} & -a_{25}^{xy} & -a_{17}^{xy} - a_{24}^{xy} & -a_{45}^{xy} & 0 & a_{12}^{xy} + a_{47}^{xy} & a_{13}^{xy} - a_{46}^{xy} \\ 0 & -a_{16}^{xy} & -a_{26}^{xy} & a_{14}^{xy} - a_{27}^{xy} & -a_{46}^{xy} & -a_{12}^{xy} - a_{47}^{xy} & 0 & a_{23}^{xy} + a_{45}^{xy} \\ 0 & -a_{17}^{xy} & -a_{27}^{xy} & a_{15}^{xy} + a_{26}^{xy} & -a_{47}^{xy} & -a_{13}^{xy} + a_{46}^{xy} & -a_{23}^{xy} - a_{45}^{xy} & 0 \end{pmatrix}$$

Let $x = \sum_{i=0}^7 \lambda_i e_i$ be an arbitrary element from \mathbb{O} . For every $z \in \mathbb{O}$ there exists a derivation $D_{z,x}$ such that

$$\Delta(z) = D_{z,x}(z), \quad \Delta(x) = D_{z,x}(x).$$

In accordance with the equality

$$D_{e_1,x}(e_1) = D_{e_1,z}(e_1)$$

we obtain

$$\begin{aligned} a_{12}^{e_1x} e_2 + a_{13}^{e_1x} e_3 + a_{14}^{e_1x} e_4 + a_{15}^{e_1x} e_5 + a_{16}^{e_1x} e_6 + a_{17}^{e_1x} e_7 = \\ a_{12}^{e_1z} e_2 + a_{13}^{e_1z} e_3 + a_{14}^{e_1z} e_4 + a_{15}^{e_1z} e_5 + a_{16}^{e_1z} e_6 + a_{17}^{e_1z} e_7, \end{aligned}$$

which implies

$$a_{1i}^{e_1x} = a_{1i}^{e_1z}, \quad 2 \leq i \leq 7.$$

Then the matrix $A^{e_1, x}$ has the following form

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & a_{12}^{e_1 z} & a_{13}^{e_1 z} & a_{14}^{e_1 z} & a_{15}^{e_1 z} & a_{16}^{e_1 z} & a_{17}^{e_1 z} & a_{17}^{e_1 z} \\ 0 & -a_{12}^{e_1 z} & 0 & a_{23}^{e_1 x} & a_{24}^{e_1 x} & a_{25}^{e_1 x} & a_{26}^{e_1 x} & a_{27}^{e_1 x} & a_{27}^{e_1 x} \\ 0 & -a_{13}^{e_1 z} & -a_{23}^{e_1 x} & 0 & a_{16}^{e_1 z} - a_{25}^{e_1 x} & a_{17}^{e_1 z} + a_{24}^{e_1 x} & -a_{14}^{e_1 z} + a_{27}^{e_1 x} & -a_{15}^{e_1 z} - a_{26}^{e_1 x} & -a_{15}^{e_1 z} - a_{26}^{e_1 x} \\ 0 & -a_{14}^{e_1 z} & -a_{24}^{e_1 x} & -a_{16}^{e_1 z} + a_{25}^{e_1 x} & 0 & a_{45}^{e_1 x} & a_{46}^{e_1 x} & a_{47}^{e_1 x} & a_{47}^{e_1 x} \\ 0 & -a_{15}^{e_1 z} & -a_{25}^{e_1 x} & -a_{17}^{e_1 z} - a_{24}^{e_1 x} & -a_{45}^{e_1 x} & 0 & a_{12}^{e_1 z} + a_{47}^{e_1 x} & a_{13}^{e_1 z} - a_{46}^{e_1 x} & a_{13}^{e_1 z} - a_{46}^{e_1 x} \\ 0 & -a_{16}^{e_1 z} & -a_{26}^{e_1 x} & a_{14}^{e_1 z} - a_{27}^{e_1 x} & -a_{46}^{e_1 x} & -a_{12}^{e_1 z} - a_{47}^{e_1 x} & 0 & a_{23}^{e_1 x} + a_{45}^{e_1 x} & a_{23}^{e_1 x} + a_{45}^{e_1 x} \\ 0 & -a_{17}^{e_1 z} & -a_{27}^{e_1 x} & a_{15}^{e_1 z} + a_{26}^{e_1 x} & -a_{47}^{e_1 x} & -a_{13}^{e_1 z} + a_{46}^{e_1 x} & -a_{23}^{e_1 x} - a_{45}^{e_1 x} & 0 & 0 \end{pmatrix}$$

Hence,

$$\begin{aligned} \Delta(x) = D_{e_1, x}(x) &= (a_{12}^{e_1 z} \lambda_1 + a_{13}^{e_1 z} \lambda_1 + a_{14}^{e_1 z} \lambda_1 + a_{15}^{e_1 z} \lambda_1 + a_{16}^{e_1 z} \lambda_1 + a_{17}^{e_1 z} \lambda_1) e_1 + \\ &\quad \mu_2^{e_1 x} e_2 + \mu_3^{e_1 x} e_3 + \mu_4^{e_1 x} e_4 + \mu_5^{e_1 x} e_5 + \mu_6^{e_1 x} e_6 + \mu_7^{e_1 x} e_7 \end{aligned}$$

for some $\mu_2^{e_1 x}, \mu_3^{e_1 x}, \mu_4^{e_1 x}, \mu_5^{e_1 x}, \mu_6^{e_1 x}, \mu_7^{e_1 x} \in \mathbb{F}$.

Considering the equality

$$D_{e_2, x} = D_{e_2, z}$$

we find that

$$a_{12}^{e_2 x} = a_{12}^{e_2 z}, \quad a_{2i}^{e_2 x} = a_{2i}^{e_2 z}, \quad 3 \leq i \leq 7.$$

Then we can write

$$\begin{aligned} \Delta(x) = D_{e_2, x}(x) &= \mu_1^{e_2 x} e_1 + \\ &\quad (-a_{12}^{e_2 z} \lambda_2 + a_{23}^{e_2 z} \lambda_2 + a_{24}^{e_2 z} \lambda_2 + a_{25}^{e_2 z} \lambda_2 + a_{26}^{e_2 z} \lambda_2 + a_{27}^{e_2 z} \lambda_2) e_2 + \\ &\quad \mu_3^{e_2 x} e_3 + \mu_4^{e_2 x} e_4 + \mu_5^{e_2 x} e_5 + \mu_6^{e_2 x} e_6 + \mu_7^{e_2 x} e_7 \end{aligned}$$

for some $\mu_1^{e_2 x}, \mu_3^{e_2 x}, \mu_4^{e_2 x}, \mu_5^{e_2 x}, \mu_6^{e_2 x}, \mu_7^{e_2 x} \in \mathbb{F}$.

From the equality

$$D_{e_3, x} = D_{e_3, z}$$

we deduce that

$$\begin{aligned} a_{13}^{e_3 x} = a_{13}^{e_3 z}, \quad a_{23}^{e_3 x} = a_{23}^{e_3 z}, \quad -a_{16}^{e_3 x} + a_{25}^{e_3 x} = -a_{16}^{e_3 z} + a_{25}^{e_3 z}, \quad -a_{17}^{e_3 x} - a_{24}^{e_3 x} = -a_{17}^{e_3 z} - a_{24}^{e_3 z}, \\ a_{14}^{e_3 x} - a_{27}^{e_3 x} = a_{14}^{e_3 z} - a_{27}^{e_3 z}, \quad a_{15}^{e_3 x} + a_{26}^{e_3 x} = a_{15}^{e_3 z} + a_{26}^{e_3 z}. \end{aligned}$$

Follows, we have

$$\begin{aligned} \Delta(x) = D_{e_3, x}(x) &= \mu_1^{e_3 x} e_1 + \mu_2^{e_3 x} e_2 + \\ &\quad (-a_{13}^{e_3 z} \lambda_3 - a_{23}^{e_3 z} \lambda_3 + (a_{16}^{e_3 z} - a_{25}^{e_3 z}) \lambda_3 + (a_{17}^{e_3 z} + a_{24}^{e_3 z}) \lambda_3 + (-a_{14}^{e_3 z} + a_{27}^{e_3 z}) \lambda_3 - (a_{15}^{e_3 z} + a_{26}^{e_3 z}) \lambda_3) e_3 + \\ &\quad \mu_4^{e_3 x} e_4 + \mu_5^{e_3 x} e_5 + \mu_6^{e_3 x} e_6 + \mu_7^{e_3 x} e_7 \end{aligned}$$

for some $\mu_1^{e_3 x}, \mu_2^{e_3 x}, \mu_4^{e_3 x}, \mu_5^{e_3 x}, \mu_6^{e_3 x}, \mu_7^{e_3 x} \in \mathbb{F}$.

The equality

$$D_{e_4, x} = D_{e_4, z}$$

imply that

$$a_{14}^{e_4 x} = a_{14}^{e_4 z}, \quad a_{24}^{e_4 x} = a_{24}^{e_4 z}, \quad a_{16}^{e_4 x} - a_{25}^{e_4 x} = a_{16}^{e_4 z} - a_{25}^{e_4 z}, \quad a_{4i}^{e_4 x} = a_{4i}^{e_4 z}, \quad 5 \leq i \leq 7.$$

Then we can write

$$\begin{aligned} \Delta(x) = D_{e_4, x}(x) &= \mu_1^{e_4 x} e_1 + \mu_2^{e_4 x} e_2 + \mu_3^{e_4 x} e_3 + \\ &\quad (-a_{14}^{e_4 z} \lambda_4 - a_{24}^{e_4 z} \lambda_4 + (-a_{16}^{e_4 z} + a_{25}^{e_4 z}) \lambda_4 + a_{45}^{e_4 z} \lambda_4 + a_{46}^{e_4 z} \lambda_4 + a_{47}^{e_4 z} \lambda_4) e_4 + \\ &\quad \mu_5^{e_4 x} e_5 + \mu_6^{e_4 x} e_6 + \mu_7^{e_4 x} e_7 \end{aligned}$$

for some $\mu_1^{e_4x}, \mu_2^{e_4x}, \mu_3^{e_4x}, \mu_5^{e_4x}, \mu_6^{e_4x}, \mu_7^{e_4x} \in \mathbb{F}$.

Similarly, considering the equality

$$D_{e_5,x} = D_{e_5,z}$$

we have

$$\begin{aligned} a_{15}^{e_5x} &= a_{15}^{e_5z}, & a_{25}^{e_5x} &= a_{25}^{e_5z}, & a_{17}^{e_5x} + a_{24}^{e_5x} &= a_{17}^{e_5z} + a_{24}^{e_5z}, \\ a_{45}^{e_5x} &= a_{45}^{e_5z}, & a_{12}^{e_5x} + a_{47}^{e_5x} &= a_{12}^{e_5z} + a_{47}^{e_5z}, & -a_{13}^{e_5x} + a_{46}^{e_5x} &= -a_{13}^{e_5z} + a_{46}^{e_5z} \end{aligned}$$

which implies

$$\begin{aligned} \Delta(x) &= D_{e_5,x}(x) = \mu_1^{e_5x} e_1 + \mu_2^{e_5x} e_2 + \mu_3^{e_5x} e_3 + \mu_4^{e_5x} e_4 + \\ &(-a_{15}^{e_5z} \lambda_5 - a_{25}^{e_5z} \lambda_5 - (a_{17}^{e_5z} + a_{24}^{e_5z}) \lambda_5 - a_{45}^{e_5z} \lambda_5 + (a_{12}^{e_5z} + a_{47}^{e_5z}) \lambda_5 + (a_{13}^{e_5z} - a_{46}^{e_5z}) \lambda_5) e_5 + \\ &\mu_6^{e_5x} e_6 + \mu_7^{e_5x} e_7 \end{aligned}$$

for some $\mu_1^{e_5x}, \mu_2^{e_5x}, \mu_3^{e_5x}, \mu_4^{e_5x}, \mu_6^{e_5x}, \mu_7^{e_5x} \in \mathbb{F}$.

Further from

$$D_{e_6,x} = D_{e_6,z}$$

we obtain that

$$\begin{aligned} a_{16}^{e_6x} &= a_{16}^{e_6z}, & a_{26}^{e_6x} &= a_{26}^{e_6z}, & a_{14}^{e_6x} - a_{27}^{e_6x} &= a_{14}^{e_6z} - a_{27}^{e_6z}, \\ a_{46}^{e_6x} &= a_{46}^{e_6z}, & a_{12}^{e_6x} + a_{47}^{e_6x} &= a_{12}^{e_6z} + a_{47}^{e_6z}, & a_{23}^{e_6x} + a_{45}^{e_6x} &= a_{23}^{e_6z} + a_{45}^{e_6z} \end{aligned}$$

Hence we may assume

$$\begin{aligned} \Delta(x) &= D_{e_6,x}(x) = \mu_1^{e_6x} e_1 + \mu_2^{e_6x} e_2 + \mu_3^{e_6x} e_3 + \mu_4^{e_6x} e_4 + \mu_5^{e_6x} e_5 + \\ &(-a_{16}^{e_6z} \lambda_6 - a_{26}^{e_6z} \lambda_6 + (a_{14}^{e_6z} - a_{27}^{e_6z}) \lambda_6 - a_{46}^{e_6z} \lambda_6 - (a_{12}^{e_6z} + a_{47}^{e_6z}) \lambda_6 + (a_{23}^{e_6z} + a_{45}^{e_6z}) \lambda_6) e_6 + \\ &+ \mu_7^{e_6x} e_7 \end{aligned}$$

for some $\mu_1^{e_6x}, \mu_2^{e_6x}, \mu_3^{e_6x}, \mu_4^{e_6x}, \mu_5^{e_6x}, \mu_7^{e_6x} \in \mathbb{F}$.

Similarly, from

$$D_{e_7,x} = D_{e_7,z}$$

it follows that

$$\begin{aligned} a_{17}^{e_7x} &= a_{17}^{e_7z}, & a_{27}^{e_7x} &= a_{27}^{e_7z}, & a_{15}^{e_7x} + a_{26}^{e_7x} &= a_{15}^{e_7z} - a_{26}^{e_7z}, \\ a_{47}^{e_7x} &= a_{47}^{e_7z}, & a_{13}^{e_7x} - a_{46}^{e_7x} &= a_{13}^{e_7z} - a_{46}^{e_7z}, & a_{23}^{e_7x} + a_{45}^{e_7x} &= a_{23}^{e_7z} + a_{45}^{e_7z} \end{aligned}$$

Then we conclude

$$\begin{aligned} \Delta(x) &= D_{e_7,x}(x) = \mu_1^{e_7x} e_1 + \mu_2^{e_7x} e_2 + \mu_3^{e_7x} e_3 + \mu_4^{e_7x} e_4 + \mu_5^{e_7x} e_5 + \mu_6^{e_7x} e_6 + \\ &(-a_{17}^{e_7z} \lambda_7 - a_{27}^{e_7z} \lambda_7 + (a_{15}^{e_7z} + a_{26}^{e_7z}) \lambda_7 - a_{47}^{e_7z} \lambda_7 + (-a_{13}^{e_7z} + a_{46}^{e_7z}) \lambda_7 - (a_{23}^{e_7z} + a_{45}^{e_7z}) \lambda_7) e_7 \end{aligned}$$

for some $\mu_1^{e_7x}, \mu_2^{e_7x}, \mu_3^{e_7x}, \mu_4^{e_7x}, \mu_5^{e_7x}, \mu_6^{e_7x} \in \mathbb{F}$.

Summarizing, we obtain the following

$$\begin{aligned} \Delta(x) &= D_{e_1,x}(x) = D_{e_2,x}(x) = D_{e_3,x}(x) = D_{e_4,x}(x) = D_{e_5,x}(x) = D_{e_6,x}(x) = D_{e_7,x}(x) = \\ &(a_{12}^{e_1z_1} \lambda_1 + a_{13}^{e_1z_1} \lambda_1 + a_{14}^{e_1z_1} \lambda_1 + a_{15}^{e_1z_1} \lambda_1 + a_{16}^{e_1z_1} \lambda_1 + a_{17}^{e_1z_1} \lambda_1) e_1 + \\ &(-a_{12}^{e_2z_2} \lambda_2 + a_{23}^{e_2z_2} \lambda_2 + a_{24}^{e_2z_2} \lambda_2 + a_{25}^{e_2z_2} \lambda_2 + a_{26}^{e_2z_2} \lambda_2 + a_{27}^{e_2z_2} \lambda_2) e_2 + \\ &(-a_{13}^{e_3z_3} \lambda_3 - a_{23}^{e_3z_3} \lambda_3 + (a_{16}^{e_3z_3} - a_{25}^{e_3z_3}) \lambda_3 + (a_{17}^{e_3z_3} + a_{24}^{e_3z_3}) \lambda_3 + (-a_{14}^{e_3z_3} + a_{27}^{e_3z_3}) \lambda_3 - (a_{15}^{e_3z_3} + a_{26}^{e_3z_3}) \lambda_3) e_3 + \\ &(-a_{14}^{e_4z_4} \lambda_4 - a_{24}^{e_4z_4} \lambda_4 + (-a_{16}^{e_4z_4} + a_{25}^{e_4z_4}) \lambda_4 + a_{45}^{e_4z_4} \lambda_4 + a_{46}^{e_4z_4} \lambda_4 + a_{47}^{e_4z_4} \lambda_4) e_4 + \\ &(-a_{15}^{e_5z_5} \lambda_5 - a_{25}^{e_5z_5} \lambda_5 - (a_{17}^{e_5z_5} + a_{24}^{e_5z_5}) \lambda_5 - a_{45}^{e_5z_5} \lambda_5 + (a_{12}^{e_5z_5} + a_{47}^{e_5z_5}) \lambda_5 + (a_{13}^{e_5z_5} - a_{46}^{e_5z_5}) \lambda_5) e_5 + \\ &(-a_{16}^{e_6z_6} \lambda_6 - a_{26}^{e_6z_6} \lambda_6 + (a_{14}^{e_6z_6} - a_{27}^{e_6z_6}) \lambda_6 - a_{46}^{e_6z_6} \lambda_6 - (a_{12}^{e_6z_6} + a_{47}^{e_6z_6}) \lambda_6 + (a_{23}^{e_6z_6} + a_{45}^{e_6z_6}) \lambda_6) e_6 + \end{aligned}$$

$$(-a_{17}^{e_7 z_7} \lambda_7 - a_{27}^{e_7 z_7} \lambda_7 + (a_{15}^{e_7 z_7} + a_{26}^{e_7 z_7}) \lambda_7 - a_{47}^{e_7 z_7} \lambda_7 + (-a_{13}^{e_7 z_7} + a_{46}^{e_7 z_7}) \lambda_7 - (a_{23}^{e_7 z_7} + a_{45}^{e_7 z_7}) \lambda_7) e_7$$

for any $z_1, z_2, z_3, z_4, z_5, z_6, z_7 \in \mathbb{O}$. Note that the components in this last sum does not depend on the element x . Therefore the map Δ is linear, and it is a local derivation. This completes the proof. \square

Theorem 3.5. *Every 2-local derivation of the algebra of octonions \mathbb{O} is a Jordan derivation.*

Proof. Let Δ be a 2-local derivation on \mathbb{O} . Since Δ is a 2-local derivation, for any element x in \mathbb{O} , there exists a derivation D_{x,x^2} on \mathbb{O} such that

$$\Delta(x) = D_{x,x^2}(x), \Delta(x^2) = D_{x,x^2}(x^2).$$

Hence,

$$\Delta(x^2) = D_{x,x^2}(x^2) = D_{x,x^2}(x)x + xD_{x,x^2}(x) = \Delta(x)x + x\Delta(x).$$

Therefore, for any two elements x, y in \mathbb{O} , we get

$$\Delta((x+y)^2) = \Delta(x+y)(x+y) + (x+y)\Delta(x+y).$$

At the same time,

$$\Delta((x+y)^2) = \Delta(x^2 + xy + yx + y^2) = \Delta(x^2) + \Delta(xy + yx) + \Delta(y^2)$$

and

$$\begin{aligned} \Delta(x+y)(x+y) + (x+y)\Delta(x+y) &= \Delta(x)x + x\Delta(x) + \Delta(x)y + y\Delta(x) \\ &\quad + x\Delta(y) + \Delta(y)x + \Delta(y)y + y\Delta(y) \\ &= \Delta(x^2) + \Delta(x)y + y\Delta(x) + x\Delta(y) + \Delta(y)x + \Delta(y^2). \end{aligned}$$

Hence,

$$\Delta(xy + yx) = \Delta(x)y + y\Delta(x) + x\Delta(y) + \Delta(y)x.$$

By theorem 3.4 Δ is linear. Therefore, Δ is a Jordan derivation. This ends the proof. \square

4. CONCLUSIONS

In the present study, we prove that a local derivation on the octonion (Cayley) algebra \mathbb{O} over an arbitrary field, satisfying some conditions is a derivation, and every 2-local derivation on \mathbb{O} is a Jordan derivation. The special algebra \mathbb{O} over an arbitrary field \mathbb{F} is 8-dimensional algebra. The algebra \mathbb{O} satisfies the next weak-associative law: $(aa)b = a(ab)$, $(ab)b = a(bb)$. Algebras which satisfies two previous identities called alternative algebras. It is well known that the octonion algebra \mathbb{O} is unique non-associative simple alternative algebra. Therefore, we have the description of local and 2-local derivations of finite-dimensional semi-simple alternative algebras under some conditions. We proved that every 2-local derivation of a finite-dimensional semi-simple alternative algebra is a Jordan derivation.

REFERENCES

- [1] Sh. Ayupov, K. Kudaybergenov, 2-local derivations and automorphisms on $B(H)$, J. Math. Anal. Appl. 395 (2012) 15–18.
- [2] N. Jacobson, Structure and Representations of Jordan Algebras, Amer. Math. Soc. Colloq. Publ. v. 39, Providence, (1968) pp. 453.
- [3] B. Johnson, “Local derivations on C*-algebras are derivations,” Trans. Amer. Math. Soc. **353**, 313–325 (2001).
- [4] Kadison R., Local derivations, Journal of Algebra, 130 (1990), 2, 494–509.
- [5] D.R. Larson, A.R. Sourour, Local derivations and local automorphisms of $B(X)$, Proc. Sympos. Pure Math., 51, Part 2, Providence, Rhode Island 1990, 187–194.
- [6] Šemrl P., Local automorphisms and derivations on $B(H)$, Proceedings of the American Mathematical Society, 125 (1997) 2677–2680.
- [7] Zorn M., Alternativekörper und quadratische Systeme, Abhandlungen aus dem mathematischen seminar der Universität Hamburg, 9(3/4) (1931) 395–402.

(F.N. Arzikulov) V.I.ROMANOVSKIY INSTITUTE OF MATHEMATICS UZBEKISTAN ACADEMY OF SCIENCES 81, MIRZO ULUGHBEK STR., TASHKENT 100170, UZBEKISTAN

AND

DEPARTMENT OF MATHEMATICS, ANDIJAN STATE UNIVERSITY, 129, UNIVERSITET STREET, ANDIJAN, 170100, UZBEKISTAN

Email address: arzikulovfn@rambler.ru

(I.A. Karimjanov) DEPARTMENT OF MATHEMATICS, ANDIJAN STATE UNIVERSITY, 129, UNIVERSITET STREET, ANDIJAN, 170100, UZBEKISTAN

AND

V.I.ROMANOVSKIY INSTITUTE OF MATHEMATICS UZBEKISTAN ACADEMY OF SCIENCES 81, MIRZO ULUGHBEK STR., TASHKENT 100170, UZBEKISTAN

Email address: iqboli@gmail.com

(S. Uguz) UT DALLAS, DEPT. MATH. SCIENCES, RICHARDSON, TX 75080

Email address: selman.uguz@utdallas.edu