

On minimal subspace Z_p -null designs*

Denis S. Krotov

Abstract

Let q be a power of a prime p , and let V be an n -dimensional space over the field $\text{GF}(q)$. A Z_p -valued function C on the set of k -dimensional subspaces of V is called a k -uniform Z_p -null design of strength t if for every t -dimensional subspace y of V the sum of C over the k -dimensional superspaces of y equals 0. For $q = p = 2$ and $0 \leq t < k < n$, we prove that the minimum number of non-zeros of a non-void k -uniform Z_p -null design of strength t equals 2^{t+1} . For $q > 2$, we give lower and upper bounds for that number.

1. Introduction

Let p be a prime number, and let $q = p^s$ where s is a positive integer. We denote by \mathbb{F}_q the finite field $\text{GF}(q)$ of order q and by \mathbb{Z}_s the factor ring $\mathbb{Z}/s\mathbb{Z}$. By $J_q(n, k)$, we denote the set of k -dimensional subspaces of the n -dimensional space \mathbb{F}_q^n over \mathbb{F}_q . We will also use the notation

$$J_q(n, \geq t) := \bigcup_{k=t}^n J_q(n, k).$$

Let \mathbb{G} be an abelian group. A function $\mathcal{C} : J_q(n, \geq t) \rightarrow \mathbb{G}$ is called a *subspace \mathbb{G} -null design of strength t* (depending on the context, we will omit “subspace”, “ \mathbb{G} ”, and/or “of strength t ”) if for every \mathbf{y} from $J_q(n, t)$ we have $\mathcal{C}(\geq \mathbf{y}) = 0$, where

$$\mathcal{C}(\geq \mathbf{y}) := \sum_{\mathbf{x} \in J_q(n, \geq t) : \mathbf{y} \subset \mathbf{x}} \mathcal{C}(\mathbf{x}).$$

The identity-zero null design is called *void*. A null design is called *k-uniform* or simply *uniform* if all its nonzeros lie in $J_q(n, k)$. Our main goal is to

*The work was funded by the Russian Science Foundation under grant 18-11-00136

study the minimum number of non-zeros of uniform null designs; however, it is convenient for some statements to be proved in a general form, for unrestricted null designs.

The following treatment of the strength is well known for many kinds of designs; however, usually it holds only for uniform designs but not in the unrestricted case in general. So, it is notable that we do not need the uniformity in the following proposition.

Proposition 1. *Let q and r be powers of a prime p such that $p \leq r \leq q$. If $\mathcal{C} : J_q(n, \geq t) \rightarrow \mathbb{Z}_r$ satisfies $\mathcal{C}(\geq \mathbf{z}) = \lambda$ for all \mathbf{z} from $J_q(n, t)$ and some constant λ from \mathbb{Z}_r , then \mathcal{C} (continued by zeros to the extended domain) satisfies $\mathcal{C}(\geq \mathbf{y}) = \lambda$ for all \mathbf{z} in $J_q(n, \tau)$, $\tau \in \{0, \dots, t\}$. In particular, if \mathcal{C} is a \mathbb{Z}_r -null design of strength t , then it is a \mathbb{Z}_r -null design of strength τ for any $\tau \in \{0, \dots, t\}$.*

Proof. For \mathbf{y} from $J_q(n, t-1)$ and \mathbf{x} from $J_q(n, \geq t)$, we note that

$$\begin{aligned} |\{\mathbf{z} \in J_q(n, t) : \mathbf{y} \subset \mathbf{z} \subset \mathbf{x}\}| &= \frac{q^{\dim(\mathbf{x})-t+1} - 1}{q-1} \\ &= q^{\dim(\mathbf{x})-t} + q^{\dim(\mathbf{x})-t-1} + \dots + q^0 \equiv 1 \pmod{r}. \end{aligned} \quad (1)$$

So, for every \mathbf{y} from $J_q(n, t-1)$ we have

$$\begin{aligned} \sum_{\mathbf{x} \in J_q(n, \geq t) : \mathbf{y} \subset \mathbf{x}} \mathcal{C}(\mathbf{x}) &\stackrel{(1)}{=} \sum_{\mathbf{x} \in J_q(n, \geq t) : \mathbf{y} \subset \mathbf{x}} \sum_{\mathbf{z} \in J_q(n, t) : \mathbf{y} \subset \mathbf{z} \subset \mathbf{x}} \mathcal{C}(\mathbf{x}) \\ &= \sum_{\mathbf{z} \in J_q(n, t) : \mathbf{y} \subset \mathbf{z}} \sum_{\mathbf{x} \in J_q(n, \geq t) : \mathbf{z} \subset \mathbf{x}} \mathcal{C}(\mathbf{x}) = \sum_{\mathbf{z} \in J_q(n, t) : \mathbf{y} \subset \mathbf{z}} \lambda \stackrel{(1)}{=} \lambda, \end{aligned}$$

and the claim is true for $\tau = t-1$. By induction, it holds for $\tau \in \{0, \dots, t\}$. \square

2. Main results

Theorem 2. *For any abelian group \mathbb{G} , the minimum number of non-zeros in a non-void \mathbb{G} -null design $\mathcal{C} : J_q(n, \geq t) \rightarrow \mathbb{G}$ of strength t , $0 \leq t < n$, is $1 + \frac{q^{t+1}-1}{q-1}$.*

Proof. Let e be any non-identity element of \mathbb{G} , and let $-e$ be its inverse. If $\mathcal{C}(\mathbf{v}) = e$ for some \mathbf{v} from $J_q(n, t+1)$, $\mathcal{C}(\mathbf{u}) = -e$ for every t -dimensional subset \mathbf{u} of \mathbf{v} , and $\mathcal{C}(\mathbf{w}) = 0$ for any other \mathbf{w} from $J_q(n, \geq t)$, then \mathcal{C} is a null design of strength t with $1 + \frac{q^{t+1}-1}{q-1}$ non-zeros.

By backward induction on t , we will prove that *the number of non-zeros of non-void null design of strength t is not less than $1 + \frac{q^{t+1}-1}{q-1}$* . For $t = n$, the claim is trivial because there are no non-void null designs. Assume that the claim is true for $t = \tau$, where $0 < \tau \leq n$; let us show it for $t = \tau - 1$. Consider a non-void null design \mathcal{C} of strength t . Denote $\mathcal{C}' = \mathcal{C}|_{J_q(n, \geq t+1)}$. Clearly, \mathcal{C}' is not constantly zero (otherwise, \mathcal{C} is void as well). If \mathcal{C}' is a null design of strength $t + 1$, then the claim is true by the induction hypothesis. Otherwise, there is \mathbf{v} in $J_q(n, t + 1)$ such that $\mathcal{C}(\geq \mathbf{v}) \neq 0$. By the definition of null design, for every t -dimensional subspace \mathbf{u} of \mathbf{v} , we have $\mathcal{C}(\geq \mathbf{u}) = 0$. So, the sum of \mathcal{C} over $U_{\mathbf{u}}$, where

$$U_{\mathbf{u}} := \{\mathbf{w} \in J_q(n, \geq t+1) : \mathbf{u} \subset \mathbf{w}, \mathbf{v} \not\subset \mathbf{w}\},$$

equals $-\mathcal{C}(\geq \mathbf{v})$. Hence, $U_{\mathbf{u}}$ contains a non-zero of \mathcal{C} . Since $U_{\mathbf{u}}$ and $U_{\mathbf{u}'}$ are disjoint for different t -dimensional subspaces \mathbf{u} and \mathbf{u}' of \mathbf{v} , we have at least $1 + \frac{q^{t+1}-1}{q-1}$ non-zeros. \square

The null design constructed in the proof is not uniform. So, for the number of non-zeros in a non-void uniform null design, Theorem 2 only gives a lower bound.

Theorem 3. *For every $k, t < k < n$, there is a k -uniform \mathbb{Z}_q -null design $\mathcal{C} : J_q(n, k) \rightarrow \{0, 1\}$ of strength t with q^{t+1} non-zeros.*

Note that a $\{0, 1\}$ -valued \mathbb{Z}_q -null design is a \mathbb{Z}_r -null design for every $r, r|q$.

Proof. Take any \mathbf{w} from $J_q(n, k + 1)$, \mathbf{u} from $J_q(n, k - t - 1)$, and \mathbf{v} from $J_q(n, k - t)$ such that $\mathbf{u} \subset \mathbf{v} \subset \mathbf{w}$. Denote $U := \{\mathbf{x} \in J_q(n, k) : \mathbf{u} \subset \mathbf{x} \subset \mathbf{w}\}$ and $V := \{\mathbf{x} \in J_q(n, k) : \mathbf{v} \subset \mathbf{x} \subset \mathbf{w}\}$. We state that the characteristic $\{0, 1\}$ -function of $U \setminus V$ is a required null design.

At first, we note that $|U \setminus V| = |U| - |V| = q^{t+1}$ because

$$|U| = \frac{q^{(k+1)-(k-t-1)} - 1}{q-1} = \frac{q^{t+2} - 1}{q-1} \quad \text{and} \quad |V| = \frac{q^{(k+1)-(k-t)} - 1}{q-1} = \frac{q^{t+1} - 1}{q-1}.$$

Next, consider \mathbf{y} from $J_q(n, t)$ and count the numbers

$$\alpha = |\{\mathbf{x} \in U : \mathbf{y} \subset \mathbf{x}\}| \quad \text{and} \quad \beta = |\{\mathbf{x} \in V : \mathbf{y} \subset \mathbf{x}\}|.$$

If $\mathbf{y} \not\subset \mathbf{w}$, then obviously $\alpha = \beta = 0$. If $\mathbf{y} \subset \mathbf{w}$, then we have

$$\alpha = \frac{q^{(k+1)-\dim(\mathbf{u}+\mathbf{y})} - 1}{q-1} \equiv 1 \pmod{q} \quad \text{and} \quad \beta = \frac{q^{(k+1)-\dim(\mathbf{v}+\mathbf{y})} - 1}{q-1} \equiv 1 \pmod{q}.$$

Anyway, $\alpha - \beta \equiv 0 \pmod{q}$. Hence, $U \setminus V$ is a \mathbb{Z}_q -null design of strength t . \square

For $q = 2$, the lower and upper bounds in Theorems 2 and 3 coincide.

Corollary 4. *For $0 \leq t < k < n$, the minimum number of non-zeros in a non-void k -uniform \mathbb{Z}_2 -null design $\mathcal{C} : J_2(n, \geq t) \rightarrow \mathbb{Z}_2$ of strength t is 2^{t+1} .*

3. On Wilson matrices

The Wilson matrix $W_{t,k}$ is a $\{0, 1\}$ -matrix whose rows are indexed by t -subsets of $\{1, \dots, n\}$ and columns are indexed by k -subsets of $\{1, \dots, n\}$; the element of $W_{t,k}$ in the \mathbf{y} th row and \mathbf{x} th column equals 1 if and only if $\mathbf{y} \subset \mathbf{x}$. By analogy, the subspace Wilson matrix $W_{q;t,k}$ is a $\{0, 1\}$ -matrix whose rows and columns are indexed by $J_q(n, t)$ and $J_q(n, k)$, respectively; $W_{q;t,k}(\mathbf{y}, \mathbf{x}) = 1$ if and only if $\mathbf{y} \subset \mathbf{x}$. For the case $q = 2$, we have established (Corollary 4) that the binary linear code with check matrix $W_{q;t,k}$ has minimum distance 2^{t+1} . (A similar result for $W_{k-1,k}$ was obtained in [5, Prop. 9].) However, the dimension of this code (equivalently, the rank of $W_{q;t,k}$) remains unknown for $t > 1$. In the case $t = 1$, $W_{q;1,k}^T$ is a generator matrix of the punctured Reed–Muller code $\mathcal{R}^*(n - k, n)$, and $\text{rank}(W_{q;1,k}) = \sum_{i=0}^k \binom{n}{i}$, see e.g. [1, Theorem 3.14]. Much less is known for $q > 2$. It is also known that $W_{q;t,k}$ is full-rank over \mathbb{R} , see [3]; the minimum number of non-zeros in a non-void \mathbb{R} -null design $J_q(n, k) \rightarrow \mathbb{R}$ is $\prod_{i=0}^t (1 + q^i)$ if $k = t + 1$ and is conjectured to be $\prod_{i=0}^t (1 + q^i)$ if $t + 2 \leq k < n - t$, see [2] (with the additional restriction on the null design to be $\{0, 1, -1\}$ -valued, the conjecture was proved in [4]).

References

- [1] Jr. Assmus, E. F. and J. D. Key. Polynomial codes and finite geometries. In V.S. Pless and W. C. Huffman, editors, *Handbook of Coding Theory*, chapter 16, page 1269–1343. Elsevier, New York, 1998.
- [2] S. Cho. Minimal null designs of subspace lattice over finite fields. *Linear Algebra Appl.*, 282(1-3):199–220, 1998. DOI: 10.1016/S0024-3795(98)10062-9.
- [3] W. M. Kantor. On incidence matrices of finite projective and affine spaces. *Mathematische Zeitschrift*, 124(4):315–318, Dec. 1972. DOI: 10.1007/BF01113923.
- [4] D. S. Krotov. The minimum volume of subspace trades. *Discrete Math.*, 340(12):2723–2731, 2017. DOI: 10.1016/j.disc.2017.08.012.

- [5] V. N. Potapov. Splitting of hypercube into k -faces and DP-colorings of hypergraphs. E-print 1905.04461v4, arXiv.org, 2020. Available at <http://arxiv.org/abs/1905.04461v4>.