

DFT quantization of the Gibbs Free Energy of a quantum body

S. Selenu

(Dated: February 15, 2022)

In this article it is introduced a theoretical model made in order to perform calculations of the quantum heat of a body that could be acquired or delivered during a thermal transformation of its quantum states. Here the model is mainly targeted to the electronic structure of matter[1] at the nano and micro scale where DFT models have been frequently developed of the total energy of quantum systems. De

fining an Entropy functional $S[\rho]$ makes us able optimizing a free energy $G[\rho]$ of the quantum system at finite temperatures. Due to the generality of the model, the latter can be also applied to several first principles computational codes where ab initio modelling of quantum matter is asked.

INTRODUCTION

This article will be focused on a study based on the derivation of the quantum electrical heat absorbed or either released by a physical system made of atoms and electrons either at the nano or at the microscopic scale, extending to it the concept of thermal heat within a DFT model. Also, the latter being fundamental in physics, can be employed for a full understanding of the thermal phase transformation of the electronic subsystem involved in the physics of crystals, where the thermal effects may directly affects their thermal stability while indirectly enhancing their functionality, to it associated a lowering of the classical thermal gibbs free energy of electrons. Within this classical model shown here having a quantum counterpart we shall introduce in the next part of the article a comprehensive Density Functional Theory (DFT)[2–5] modelling taking into account the nature of thermal quantum phenomena that has been avoided since the advent of DFT. A new DFT model based on either a direct statistical interpretation of the electronic heat, as a thermodynamical degree of freedom, allows to characterize the energetics of the quantum body on a base where a variational principle still can be applied on the searching of the quantum electronic wave functions, either via computational simulations of quantum matter[1]. In the following section it will be reported and developed a new DFT model of the quantum electronic heat while in the last section are reported conclusions of the article.

DFT MODEL OF THE ELECTRONIC QUANTUM HEAT

Since 1960's the work on DFT modelling of quantum matter has been growing more until it becomes one of the most diffused theoretical model employed for the understanding of the interactions of electrons in matter. Effort is mainly put onto the development of this matter modelling in order showing for the first time how to perform calculations of the quantum electronic heat, either absorbed or released by a quantum body during its

thermal transformations at a fi

nite temperature T . This work could then be considered as the beginning part of a more general theory can include an ab initio modelling of the first law and second law of thermodynamics[13], where heat and free energy variations of a quantum body are taken into account. In fact it is actually considered a quantum system of electrons at a given temperature T in thermal equilibrium with its surrounding. Electrons also are considered either interacting with themselves by a potential V_{ee} [3], an ionic external potential V_{ie} [3], having also an exchange correlation potential[3] V_{xc} ruling their correlation, while an associated electronic charge density of the system is given by the following relation $\rho = \sum_n f_n \Psi_n^* \Psi_n$ times the electronic charge, i.e. a product of electronic elementary charge e and of the quantum probability distribution ρ of an event happening. The latter is calculated as the weighted average of the products of wave functions of the quantum system at a state n , with f_n the number of electrons per state. The probability density can be directly used in order to perform routinely calculations of the potential energies usually encountered in quantum DFT, considered then an eligible candidate for a further study of the quantum Gibbs free energy and its related Entropy[9, 10, 12, 14, 15]. In order to base our statistics on a generalisation of the Shannon model of the Entropy, by optimizing at a variational level a quantum Gibbs free energy functional, a free energy operator is derived, and related to a new set of differential eigenvalue equations. The latter, built as a functional of the electronic probability density allows then deriving a set of quantum differential equations by considering the very quantum mechanical interpretation of the electronic density as being the not normalized probability of delocalized electrons being found in a position in space[7]. It is possible then recognizing to it being associated to a quantum Entropy whose expression is given by:

$$S = -k_B \int \rho \ln \rho \quad (1)$$

$$S = -k_b \sum_n f_n \langle \Psi_n | \ln \rho | \Psi_n \rangle \quad (2)$$

in agreement with the representation of the entropy of continuous distributions of probabilities as given by Shannon, C. and Weaver, W.[15]. Here a Lagrange multiplier $S = N \ln N$ appears on variations of the energy functional, with $N = \int \rho$, being the latter the total number of electrons in the volume of the body. Latter result implies the writing of a new functional of the probability density $\rho' = \rho N$, that shall be reported in the next part of this section. By firstly considering Gibbs free energy of the thermodynamical system begin given by the following classical formula $G = H - TS$, where $Q = TS$ is the electronic heat at a given entropy S for a fi

xed temperature T of the quantum system, an Enthalpy functional $H[\rho] = E[\rho] + PV$ is calculated, being $E[\rho]$ the internal energy of the body. The latter energy functional, is customarily calculated in quantum DFT and kept the volume V of the body constant in order to neglect dependence of the energy of the system on the pressure P . The Enthalpy is then given by the sum of the internal energy plus a trivial constant PV not varying with wave functions. The Gibbs free energy functional is then finally written making use of eq.(1) as it follows:

$$G[N\rho'] = H[N\rho'] - NTS[\rho'] + k_B T N \ln N \quad (3)$$

made by the sum of the Helmholtz free energy[11, 13] and the electronic heat of the body. In what follows it is not varied only the Helmholtz free energy, where the product PV is kept constant in order calculating the Gibbs free energy variation, but also the Entropy functional. A stationary differential equation is reached and directly interpreted as a quantum mechanical eigenvalue equation whose eigen values ϵ_G are the eigen free energies of the quantum body. The functional derivatives of G , H and S are calculated by referring to eq.(3), obtaining the following result:

$$\begin{aligned} \hat{G}\Psi_n &= \epsilon_{n,G}\Psi_n \quad (4) \\ \hat{G} &= \left[\frac{-\hbar^2 \nabla^2}{2m} + V_{ee} + V_{ie} + V_{XC} + k_B T \ln \rho' + k_B T \right] \end{aligned}$$

An Hamiltonian eigenvalue set of equations can also be reached by the shifting of energies $\epsilon_n = \epsilon_{n,G} + k_B T$, due to the increase of the latter of a thermal energy amount equal to $k_B T$, related to the thermal motion of electrons, allowing to reach the following stationary equation:

$$\begin{aligned} \hat{H}\Psi_n &= \epsilon_n \Psi_n \quad (5) \\ \hat{H} &= \left[\frac{-\hbar^2 \nabla^2}{2m} + V_{ee} + V_{ie} + V_{XC} + TV_S \right] \end{aligned}$$

It appears a new thermodynamical potential on right hand side of equation (5) directly dependent on the normalized probability density, $V_S = k_B \ln \rho'$, whose expectation values with respect the n th state of the electronic

wave Ψ_n is the Entropy per electron S' of the system so as to call it the entropic potential. It becomes evident we shall add the latter to the new DFT operator, reported in eq.(4), in order calculating the eigenstates of quantum matter in its thermal states. Due to the de

inition of the quantum electronic heat, being $Q = TS$, it is straightforward showing by eq.(1) the expression of the electronic heat to be written as follows:

$$Q = -Nk_B T \int \rho' \ln \rho' \quad (6)$$

with respect to the electronic probability density ρ' , making the free energy a minimum as it is the natural tendency of a physical system due to the second law of thermodynamics[11, 13]. This has been the first attempt to study the quantum thermodynamics of matter in its steady states in a Hamiltonian formalism, brought us introducing the quantum free energy operator, then calculate eigenstates along any thermal changings of the quantum state of matter. Having reached for the first time a quantization of the Gibbs free energy the article will be concluded in the next section.

CONCLUSIONS

This article is concluded having shown how to calculate the quantum electronic heat of a body, at a DFT variational level, also defining the electronic Entropy functional $S[\rho]$, it interpreted as the Entropy of the system. Latter result makes us able optimizing the quantum free energy $G[\rho]$ by variations of the eigenstates of the quantum system at finite temperature T . Due to the generality of our model we have reached a set of differential equations being expressed in (4), showing that it can be applied to several first principle codes of an ab initio modelling of matter, either at the nano or microscopic scale. A future computational modelling of the electronic heat delivered or either absorbed by a quantum system during its thermal phase transformations can then be targeted.

-
- [1] R.M. Martin, Electronic structure (Cambridge University-Press, 2004)
 - [2] Hohenberg, P.; Kohn, W. (1964). "Inhomogeneous Electron Gas". Physical Review. 136 (3B): B864. Bibcode:1964PhRv..136..864H. doi:10.1103/PhysRev.136.B864.
 - [3] Kohn, W.; Sham, L. J. (1965). "Self-Consistent Equations Including Exchange and Correlation Effects". Physical Review. 140 (4A): A1133. Bibcode:1965PhRv..140.1133K. doi:10.1103/PhysRev.140.A1133.

- [4] March, N. H. (1992). *Electron Density Theory of Atoms and Molecules*. Academic Press. p. 24. ISBN 978-0-12-470525-8.
- [5] Parr, R. G.; Yang, W. (1989). *Density-Functional Theory of Atoms and Molecules*. New York: Oxford University Press. ISBN 978-0-19-504279-5.
- [6] Thomas, L. H. (1927). "The calculation of atomic elds". *Proc. Camb. Phil. Soc.* 23 (5): 542548. Bib- code:1927PCPS...23..542T. doi:10.1017/S0305004100011683.
- [7] Neil W. Ashcroft and N. David Mermin (1976). *Solid State Physics*. Orlando: Harcourt. ISBN 0-03-049346-3.
- [8] Kittel, Charles (1996). *Introduction to Solid State Physics*. New York: Wiley. ISBN 0-471-14286-7.
- [9] Licker, Mark D. (2004). *McGraw-Hill concise encyclopedia of chemistry*. New York: McGraw-Hill Professional. ISBN 978-0-07-143953-4.
- [10] Sethna, James P. (2006). *Statistical mechanics : entropy, order parameters, and complexity* ([Online-Ausg.]. ed.). Oxford: Oxford University Press. p. 78. ISBN 978-0-19-856677-9.
- [11] Entropy, Helmholtz Free Energy and the Partition Function". theory.physics.manchester.ac.uk. Retrieved 2016-12-06.
- [12] Clark, John O.E. (2004). *The essential dictionary of science*. New York: Barnes and Noble. ISBN 978-0-7607-4616-5.
- [13] Enrico Fermi, *thermodynamics*, Dover Publication Inc. New York, (1957).
- [14] Frigg, R. and Werndl, C. "Entropy A Guide for the Perplexed". In *Probabilities in Physics*; Beisbart C. and Hartmann, S. Eds; Oxford University Press, Oxford, 2010
- [15] Shannon, C. and Weaver, W. (1949), *The Mathematical Theory of Communication*, University of Illinois Press, Urbana, Chicago, IL & London.