

The role of the Axiom of Choice in proper and distinguishing colourings

Marcin Stawiski
stawiski@agh.edu.pl

AGH University, Department of Discrete Mathematics,
al. Mickiewicza 30, 30-059 Krakow, Poland

June 29, 2021

Abstract

Call a colouring of a graph *distinguishing* if the only automorphism of a graph which preserves it is the identity. We investigate the role of the Axiom of Choice in the existence of certain proper or distinguishing colourings in both vertex and edge variants with special emphasis on locally finite connected graphs. In particular, we show that every locally finite connected graph has a distinguishing colouring or every locally finite connected graph has a proper colouring with at most countable number of colours if and only if König's Lemma holds. Furthermore, we show that it is improvable in ZF that such colourings exists even for every connected graph with maximum degree three. We also formulate few conditions about distinguishing and proper colourings which are equivalent to the Axiom of Choice.

AMS Subject Classification: 05C15, 05C25, 05C63, 03E25

Keywords: Proper Colourings, Distinguishing Colourings, Asymmetric Colourings, Infinite Graphs, Graph Automorphisms, Axiom of Choice

1 Introduction

Through the paper we use standard terminology of graph theory and set theory. For notions which are not defined here see [5] or [8]. Colourings in this paper are not necessarily proper unless stated otherwise. Let c be a vertex or an edge colouring of a graph G . We say that an automorphism φ of G *preserves* c if each vertex of G is mapped to a vertex of the same colour or each edge of G is mapped to an edge of the same colour. We say that c *breaks* an automorphism φ of G if c is not preserved by φ . Call a colouring c *distinguishing* if the only automorphism which preserves c is the identity. *Distinguishing number* $D(G)$ of a graph G is the least number of colours in a distinguishing vertex colouring of G . Similarly, *distinguishing index* $D'(G)$ of a graph

G is the least number of colours in a distinguishing edge colouring of G . Distinguishing vertex colourings were introduced by Babai [2] in 1977 under the name *asymmetric* colourings during his study of the complexity of the graph isomorphism problem [1]. Distinguishing edge colourings were introduced by Kalinowski and Pilśniak [9].

In this paper we study proper and distinguishing colourings in ZF so without assuming the Axiom of Choice. Proper colourings without the assumption of the Axiom of Choice were investigated by Galvin and Komjáth [6], who proved that the existence of the chromatic number of every graph is equivalent to the Axiom of Choice. Distinguishing colourings and proper edge colourings in ZF were not previously investigated.

In most of the papers about infinite graphs some version of the Axiom of Choice is used though not always explicitly. The most popular methods often involve Zorn's Lemma or König's Lemma. In particular, proofs for general bounds for chromatic number [4], chromatic index [10], distinguishing number [11] and distinguishing index [13] of connected infinite graphs in terms of their maximum degree use König's Lemma in the case of locally finite graphs and the Axiom of Choice for general bounds. We show that in all these cases the use of the Axiom of Choice or respectively König's Lemma is necessary. Note that similar problems for not connected graphs often lead to different results (see for example [7] p. 109–116).

Arguably most of the results related to proper or distinguishing colourings in graph theory concern only locally finite connected graphs. From the results mentioned in the previous paragraph, it follows that one cannot prove in ZF that every locally finite connected graph has a distinguishing or a proper colouring with at most countable number of colours. We show that one cannot prove the existence of such colourings in ZF even in the simplest case of connected graphs with maximum degree three.

Definitions of chromatic number, chromatic index, distinguishing number and distinguishing index involve the notion of cardinal number. By a *cardinal number* we mean an ordinal which is not equinumerous with any smaller ordinal. If for every set there exists a cardinal number equinumerous with it, then the Axiom of Choice holds. In Section 3 we investigate the existence of the minimal (in a sense of irreducibility) proper or distinguishing colourings which do not involve the notion of cardinal numbers.

Let G be an arbitrary graph. Denote by Ω the set of vertices of G or the set of its edges. Let c be a colouring of Ω and let $a, b \in c(\Omega)$ be different colours. Define colouring d of Ω by $d(x) = a$ if $c(x) = b$ and $d(x) = x$ otherwise. We call d a *reduction* of c . In other words, a reduction of a colouring is obtained by recolouring every vertex of a chosen colour to another chosen colour. We say that a proper colouring c of Ω is *irreducible* if there is no reduction of c which is a proper colouring. Similarly, we say that a distinguishing colouring c of Ω is *irreducible* if there is no reduction of c which is a distinguishing colouring. One can define irreducible colourings in a similar fashion for other types of colourings.

We show that the existence of an irreducible distinguishing vertex colouring, irreducible distinguishing edge colouring or an irreducible proper edge colouring for every graph is equivalent to the Axiom of Choice. The equivalence between the existence of irreducible proper vertex colourings and the Axiom of Choice was proved by Galvin and Komjáth [6]. However, they did not show the connection in ZF between the existence

of chromatic number of a given graph and the existence of an irreducible proper colouring of said graph. We show that if a graph has the chromatic number, then it also has an irreducible proper colouring. Moreover, if the Axiom of Choice fails, then this implication cannot be reversed. Therefore, the existence of an irreducible proper colouring of a given graph is a strictly weaker condition than the existence of its chromatic number. The similar connection will be established for distinguishing vertex colourings, distinguishing edge colourings and proper edge colourings.

2 The Axiom of Choice in proper and distinguishing colourings

Let Γ be a group acting on a set Ω . We say that a set $A \subset \Omega$ is *fixed* if every $\varphi \in \Gamma$ acts trivially on A . We say that a set $A \subset \Omega$ is *stabilized* if for every $\varphi \in \Gamma$, we have $\varphi(A) = A$. The *orbit* of a set $A \in \Omega$ is the set $\{\varphi(a) : a \in A, \varphi \in \Gamma\}$. In the definitions in this paragraph, if $A = \{a\}$ is a singleton, then we often refer to a instead of set $\{a\}$. The meaning of Γ and Ω will follow from the context.

Let κ be an arbitrary non-zero cardinal. Call a family \mathcal{A} *acceptable* if \mathcal{A} is a countable family of pairwise disjoint non-empty sets. We say that \mathcal{A} *almost κ -acceptable* if \mathcal{A} is acceptable and every set in \mathcal{A} has cardinality less than κ . We say that \mathcal{A} is *κ -acceptable* if it is almost κ^+ -acceptable. In other words, \mathcal{A} is *κ -acceptable* if \mathcal{A} is acceptable and every set in \mathcal{A} has cardinality at most κ .

Let $\mathcal{A} = \{A_i : i \in \omega\}$ be an acceptable family and put $Y = \bigcup \mathcal{A}$. Let $Z = \{z_i : i \in \omega\}$ and $Z' = \{z'_i : i \in \omega\}$ be disjoint sets which are also disjoint from Y . We now define graphs $G_{\mathcal{A}}$ and $H_{\mathcal{A}}$. Let $V(G_{\mathcal{A}}) = Y \cup Z \cup Z'$ be a set of vertices of $G_{\mathcal{A}}$ and define set of edges of $G_{\mathcal{A}}$ by

$$E(G_{\mathcal{A}}) = \{z_i z_{i+1} : i \in \omega\} \cup \{z_i z'_i : i \in \omega\} \cup \{a_i z'_i : i \in \omega, a_i \in A_i\}.$$

Define $V(H_{\mathcal{A}}) = V(G_{\mathcal{A}})$ and $E(H_{\mathcal{A}}) = E(G_{\mathcal{A}}) \cup \{ab : a \neq b, a, b \in A_i, i \in \omega\}$.

From the definitions of $G_{\mathcal{A}}$ and $H_{\mathcal{A}}$ it follows that every vertex in $Z \setminus \{z_0\}$ has degree three in both graphs, and vertex z_0 has degree two. For the rest of the paragraph, assume that for every $i \in \omega$ the cardinality of A_i is well-defined. Vertex z'_i has degree $|A_i| + 1$ for every $i \in \omega$ in both $G_{\mathcal{A}}$ and $H_{\mathcal{A}}$. Every vertex in Y has degree one in $G_{\mathcal{A}}$. However, every vertex $a \in A_i$ has degree $|A_i|$ in graph $H_{\mathcal{A}}$ because it has edges to every other vertex in A_i and to z'_i . Summarizing, we obtain $\Delta(G_{\mathcal{A}}) = \Delta(H_{\mathcal{A}}) = \max\{3, \sup\{|A_i| + 1 : i \in \omega\}\}$.

Claim 2.1. *Let $\mathcal{A} = \{A_i : i \in \omega\}$ be an acceptable family. Then for every $i \in \omega$ vertices from A_i form an orbit with respect to the groups of automorphisms of $G_{\mathcal{A}}$ and $H_{\mathcal{A}}$. The rest of the vertices in both graphs are fixed with respect to these groups.*

Proof. First we show that in both graphs the ray R induced by Z is fixed. First we show that z_0 is fixed. Let P be a maximal induced path with endvertex z_0 , which is edge-disjoint from R . Assume that there exists a path P' which does not contain z_0 as endvertex, and which can be mapped into P by some automorphism of $G_{\mathcal{A}}$ or $H_{\mathcal{A}}$. It follows that there exists an induced ray $R' \neq R$ whose endvertex $z' \neq z_0$ is an endvertex of P' and z' has degree two in $G_{\mathcal{A}}$ ($H_{\mathcal{A}}$ respectively). As $z' \neq z_0$, we have that z' is not

in Z because every vertex in $Z \setminus \{z_0\}$ has degree three. If $z' \in Z'$, then P' has length one, and if $z' \in Y$, then P' has length zero or one. This leads to contradiction as path P has length two. Therefore, P can be mapped to path P' only if P' contains z_0 as endvertex. Only one of endvertices of P is also an endvertex of some maximal induced ray in $G_{\mathcal{A}}$ ($H_{\mathcal{A}}$ respectively). Hence, z_0 is fixed. It follows that the ray R is fixed.

As every vertex of a fixed set Z has exactly one neighbour outside Z , then all of these neighbours are fixed. Hence, Z' is fixed, and A_i is stabilized for every $i \in \omega$. Vertices from A_i form an independent set in $G_{\mathcal{A}}$ (or clique in $H_{\mathcal{A}}$). Hence, they form an orbit with respect to the group of automorphisms of $\mathcal{G}_{\mathcal{A}}$, and also with respect to the group of automorphisms of $\mathcal{H}_{\mathcal{A}}$. \square

Notice that from the above lemma we obtained that the groups of automorphisms of $G_{\mathcal{A}}$ and $H_{\mathcal{A}}$ acts the same way on the set of vertices of these graphs. Now, we prove a lemma which allow us to restrict part of the later considerations to the problem of the existence of the distinguishing number of graph $G_{\mathcal{A}}$. With the above lemma we are able to obtain results about distinguishing colourings and proper colourings in both vertex and edge versions at the same time.

Lemma 2.2. *Let \mathcal{A} be an acceptable family. Then the following conditions are equivalent.*

- a) *There exists the distinguishing number of $G_{\mathcal{A}}$.*
- b) *There exists the distinguishing index of $G_{\mathcal{A}}$.*
- c) *There exists the chromatic index of $G_{\mathcal{A}}$.*
- d) *There exists the chromatic number of $H_{\mathcal{A}}$.*

Proof. By Claim 2.1 if c is a distinguishing vertex colouring of $G_{\mathcal{A}}$, then for every $i \in \omega$ vertices in A_i have distinct colours. If for every $i \in \omega$ and each vertex $v \in A_i$ we colour edge vz'_i with colour $c(v)$, and we colour the rest of the edges of $G_{\mathcal{A}}$ arbitrarily, then we obtain a distinguishing edge colouring of $G_{\mathcal{A}}$. Hence, condition a) implies b).

Now, let c be a distinguishing edge colouring of $G_{\mathcal{A}}$. Let c' be a colouring in which edges incident to vertices in Y have the same colour as in c , the edges between Z and Z' are coloured with the same new colour, and the edges between the vertices in Z are coloured alternately with two new colours. The colouring c' is a proper edge colouring. Therefore, condition b) implies condition c).

Let c be a proper edge colouring of $G_{\mathcal{A}}$. We now define a proper vertex colouring c' of $H_{\mathcal{A}}$. First, for every $i \in \omega$, we colour each vertex $v \in A_i$ with colour $c'(v) = c(vz'_i)$. Next, we colour the vertices in Z' with the same new colour, and we colour vertices in Z alternately with two new colours. Colouring c' is a proper vertex colouring of $H_{\mathcal{A}}$. Hence, condition c) implies condition d).

Implication between d) and a) follows directly from Claim 2.1 because every proper vertex colouring of $H_{\mathcal{A}}$ is a distinguishing vertex colouring of $G_{\mathcal{A}}$. \square

We can now proceed to the study of relations between the existence of certain colourings and the Axiom of Choice. The first step is Lemma 2.3, which shows that the existence of the distinguishing number of $G_{\mathcal{A}}$ implies the existence of a choice function for \mathcal{A} .

Lemma 2.3. *Let \mathcal{A} be an acceptable family and assume that there exists the distinguishing number of $G_{\mathcal{A}}$. Then there exists a choice function for \mathcal{A} .*

Proof. Let c be a vertex colouring of $G_{\mathcal{A}}$ with elements of some cardinal number. Then $f(A) = \min\{c(a) : a \in A\}$ is a choice function for \mathcal{A} . \square

We now prove the next lemma which in the case of non-zero natural number k and k -acceptable family \mathcal{A} allows us to construct a distinguishing colouring of $G_{\mathcal{A}}$ using a choice function for \mathcal{A} . One can verify that the assumption that k is a natural number (and not an arbitrary cardinal) is necessary. However, we will not show it here.

Lemma 2.4. *Let k be an arbitrary non-zero natural number and assume that for every k -acceptable family there exists its choice function. Then for every k -acceptable family \mathcal{A} graph $G_{\mathcal{A}}$ has distinguishing number at most k .*

Proof. The proof is by induction on k . Let \mathcal{A} be a k -acceptable family. If $k = 1$, then G has no non-trivial automorphism so its distinguishing number is equal to one. Assume that $k \geq 2$, and that the statement of the lemma holds for every $l < k$. Let f be a choice function for \mathcal{A} . From the inductive assumption $G_{\mathcal{A}} - f(\mathcal{A})$ has a distinguishing vertex colouring c' using at most $k - 1$ colours. Colouring c which agrees with c' on $V(G_{\mathcal{A}}) \setminus f(\mathcal{A})$ and which assigns the rest of vertices of $G_{\mathcal{A}}$ the same, new colour is a distinguishing colouring using at most k colours. \square

Lemmas 2.2–2.4 allows us to formulate the following corollary about the existence of certain parameters for k -acceptable families in the case of natural k .

Theorem 2.5. *Let $k \geq 2$ be an arbitrary natural number. Then the following conditions are equivalent.*

- a) *For every k -acceptable family \mathcal{A} there exists its choice function.*
- b) *For every k -acceptable family \mathcal{A} graph $G_{\mathcal{A}}$ has the distinguishing number.*
- c) *For every k -acceptable family \mathcal{A} graph $G_{\mathcal{A}}$ has the distinguishing index.*
- d) *For every k -acceptable family \mathcal{A} graph $G_{\mathcal{A}}$ has the chromatic index.*
- e) *For every k -acceptable family \mathcal{A} graph $H_{\mathcal{A}}$ has the chromatic number.*

In particular for $k = 2$ condition a) is the Axiom of Countable Choice for Pairs which is independent of ZF. It follows that in ZF one cannot prove the existence of said parameters even for every connected subcubic graph. The proof that the Axiom of Countable Choice for Pairs cannot be proven in ZF is due to Cohen (see [3]). Note that

the Axiom of Countable Choice for Pairs does not imply the Axiom of Countable Choice for Finite Sets. The proof is basically due to Mostowski [12] although it was established not for ZF but for the related system ZFU.

Theorem 2.5 tells us about the existence of certain parameters for connected graphs with finite maximal degree. Now, we establish the relations between König's Lemma and the existence of proper colourings and distinguishing colourings using at most countable number of colours in the case of locally finite connected graphs.

Theorem 2.6. *The following conditions are equivalent.*

- (KL) *König's Lemma.*
- (KL1) *Every infinite locally finite connected graph has the distinguishing number.*
- (KL2) *Every infinite locally finite connected graph has the distinguishing index.*
- (KL3) *Every infinite locally finite connected graph has the chromatic index.*
- (KL4) *Every infinite locally finite connected graph has the chromatic number.*
- (KL5) *For every almost \aleph_0 -acceptable family \mathcal{A} graph $G_{\mathcal{A}}$ has the distinguishing number.*
- (KL6) *For every almost \aleph_0 -acceptable family \mathcal{A} graph $G_{\mathcal{A}}$ has the distinguishing index.*
- (KL7) *For every almost \aleph_0 -acceptable family \mathcal{A} graph $G_{\mathcal{A}}$ has the distinguishing index.*
- (KL8) *For every almost \aleph_0 -acceptable family \mathcal{A} graph $H_{\mathcal{A}}$ has the chromatic number.*

Proof. First, we show that König's Lemma implies conditions (KL1)–(KL4). Let G be an infinite locally finite connected graph. König's Lemma is equivalent to the statement that the sum of every countable family of finite sets is countable. Therefore, the set $V(G) = \bigcup\{B(v, d) : d < \omega\}$ is at most countable for every $v \in V(G)$, where $B(v, d)$ denotes the ball of radius d and centre v . As $E(G) \subseteq V(G) \times V(G)$, then $E(G)$ is also at most countable. As both of the sets $V(G)$ and $E(G)$ are countable, we can obtain the desired colourings by assigning every vertex (edge respectively) a unique natural number.

Implications (KL1) \Rightarrow (KL5), (KL2) \Rightarrow (KL6), (KL3) \Rightarrow (KL7) and (KL4) \Rightarrow (KL8) are trivial. Equivalence of conditions (KL5)–(KL8) follows from Lemma 2.2.

It remains to show that the condition (KL5) implies König's Lemma. From (KL5) and Lemma 2.3, we have that for every countable family of finite sets there exists its choice function. This is the Axiom of Countable Choice for Finite Sets, which is known to be equivalent to König's Lemma. \square

The existence of the distinguishing number of $G_{\mathcal{A}}$ for every almost \aleph_0 -acceptable family \mathcal{A} is equivalent to König's Lemma and therefore to the Axiom of Countable Choice for Finite Sets. One may think that the existence of the distinguishing number of every graph of the form $G_{\mathcal{A}}$ for some acceptable family \mathcal{A} is equivalent to the Axiom

of Countable Choice. It turns out that this condition is much stronger and it implies the full Axiom of Choice.

Theorem 2.7. *If for every acceptable family \mathcal{A} graph $G_{\mathcal{A}}$ has the distinguishing number, then the Axiom of Choice holds.*

Proof. Let X be a non-empty set and let \mathcal{A} be an acceptable family such that $X \in \mathcal{A}$. By the assumption there exists a distinguishing vertex colouring c of graph $G_{\mathcal{A}}$ using colours from some cardinal κ . As colouring c is distinguishing, the elements of X have distinct colours in c . It follows that $c|_X$ is injection from X to cardinal number κ . Hence, Zermelo's Theorem holds and so does the Axiom of Choice. \square

The above theorem let us formulate the list of conditions equivalent to the Axiom of Choice. Conditions (AC1)–(AC4) in the theorem below are equivalent to their restrictions to connected graphs. Recall that the equivalence of the Axiom of Choice, the existence of chromatic number of every graph and the existence of chromatic number of every connected graph was proved by Galvin and Komjáth [6].

Theorem 2.8. *The following conditions are equivalent.*

- (AC) *Axiom of Choice.*
- (AC1) *Every graph has the distinguishing number.*
- (AC2) *Every graph without component isomorphic to K_1 or K_2 has the distinguishing index.*
- (AC3) *Every graph has chromatic index.*
- (AC4) *Every graph has chromatic number.*
- (AC5) *For every acceptable family \mathcal{A} graph $G_{\mathcal{A}}$ has the distinguishing number.*
- (AC6) *For every acceptable family \mathcal{A} graph $G_{\mathcal{A}}$ has the distinguishing index.*
- (AC7) *For every acceptable family \mathcal{A} graph $G_{\mathcal{A}}$ has the chromatic index.*
- (AC8) *For every acceptable family \mathcal{A} graph $G_{\mathcal{A}}$ has the chromatic number.*

Proof. From the Zermelo's theorem we can well-order sets of vertices and edges of a given graph and then colour each vertex (edge respectively) of said graph with a unique colour, which means that Axiom of Choice implies conditions (AC1)–(AC4). Each of the condition (AC1)–(AC4) imply its restriction to connected graph and also the corresponding condition (AC5)–(AC8). By Lemma 2.2 conditions (AC5)–(AC8) are equivalent. Implication between condition (AC5) and the Axiom of Choice is Theorem 2.7. \square

3 Irreducible colourings and the Axiom of Choice

Theorem 2.8 shows the equivalence between the existence of certain colour-related parameters and the Axiom of Choice. However, all these parameters are by definition linked to cardinal numbers. For every set there exists a cardinal number equinumerous with it if and only if the Axiom of Choice holds. Hence, Theorem 2.8 is not surprising. In this section we investigate irreducible colourings in the same fashion as it was done in [6] for proper vertex colourings. We show that the existence of an irreducible proper or distinguishing colouring for every graph is equivalent to the Axiom of Choice. Moreover, we show that if the Axiom of Choice does not hold, then the existence of an irreducible colouring of a given graph is a strictly stronger condition than the existence of the corresponding parameter of said graph.

If graph G has the distinguishing number, then it also has an irreducible colouring. The analogous implication holds for the distinguishing index, the chromatic number and the chromatic index. We prove this fact in a more general version.

Let c be a colouring of Ω into an ordinal α . We define

$$\begin{aligned} C_c^0 &= \{c\}, \\ C_c^{\beta+1} &= C_c^\beta \cup \{s : s \text{ is a reduction of some } d \in C_c^\beta\}, \text{ for every ordinal } \beta, \\ C_c^\beta &= \bigcup \{C_c^\gamma : \gamma < \beta\}, \text{ for every non-zero limit ordinal } \beta. \end{aligned}$$

Denote $C_c = \bigcup \{C_c^\gamma : \gamma \text{ is an ordinal}\}$. We define a well-ordering \leq of C_c . We denote by $f^{-1}(X)$ the inverse image of X by f . Let $c_1, c_2 \in C_c$. We write $c_1 < c_2$ if one of the above conditions is satisfied:

1. There exists an ordinal $\beta < \alpha$ such that $\beta \in c_2(\Omega)$ and $\beta \notin c_1(\Omega)$ and for every $\gamma < \beta$ we have either $\gamma \in c_2(\Omega) \cap c_1(\Omega)$ or $\gamma \notin c_1(\Omega) \cup c_2(\Omega)$.
2. $c_1(\Omega) = c_2(\Omega)$ and let $\beta < \alpha$ be the least ordinal such that $c_1^{-1}(\beta) \neq c_2^{-1}(\beta)$. Denote by δ_1 the least ordinal such that $c^{-1}(\delta_1) \subseteq c_1^{-1}(\beta)$ and $c^{-1}(\delta_1) \not\subseteq c_2^{-1}(\beta)$. Similarly, denote by δ_2 the least ordinal such that $c^{-1}(\delta_2) \not\subseteq c_1^{-1}(\beta)$ and $c^{-1}(\delta_2) \subseteq c_2^{-1}(\beta)$. Then $\delta_1 < \delta_2$.

The following theorem is easy to check.

Theorem 3.1. *Let c be a colouring of Ω into an ordinal number α . Then \leq is a well-ordering of C_c . \square*

Note that if a colouring c' is the least element of a set $C \subseteq C_c$ with respect to \leq , then c' has no reduction in C . From the above theorem it follows that if a graph G has the chromatic number, the chromatic index, the distinguishing number or the distinguishing index, then it has a corresponding irreducible colouring. This also holds for similarly defined parameters of different colourings. To see that, let Φ be some property of colourings of Ω . Let $c : \Omega \mapsto \kappa$ be a colouring satisfying Φ and let $C' \subseteq C_c$ be the set of colourings in C_c satisfying Φ . The least element of (C', \leq) satisfies Φ and it has no reduction satisfying Φ .

As the definitions of irreducible colourings do not refer to cardinal numbers, one can expect that the existence of irreducible colouring of some graph is a stronger condition in ZF than the existence of corresponding parameter of said graph. We show that this is always the case if the Axiom of Choice fails. Namely, we show that if the Axiom of Choice does not hold, then there exists a graph with irreducible distinguishing vertex colouring and without the distinguishing number. The same holds for distinguishing edge colourings and for proper colourings. For this purpose we define graphs $DS(X, Y)$ and $DC(X, Y)$.

Let $X, Y, \{x'\}$ and $\{y'\}$ be pairwise disjoint non-empty sets. Graph $DS(X, Y)$ is a graph with vertex set $V(DS(X, Y)) = X \cup Y \cup \{x', y'\}$ such that e is an edge in $DS(X, Y)$ if and only if $e = x'y'$, $e = x'x$ or $e = y'y$ for some $x \in X$ or some $y \in Y$. Graph $DC(X, Y)$ is a graph obtained from $DS(X, Y)$ by adding edges between each pair of different vertices in X and each pair of different vertices in Y .

Lemma 3.2. *Let X and Y be disjoint non-empty sets. Then the following conditions are equivalent.*

- a) $DS(X, Y)$ has an irreducible distinguishing vertex colouring.
- b) $DS(X, Y)$ has an irreducible distinguishing edge colouring.
- c) $DC(X, Y)$ has an irreducible proper vertex colouring.
- d) $DS(X, Y)$ has an irreducible proper edge colouring.
- e) $|X| \leq |Y|$ or $|Y| \leq |X|$.

Proof. Denote by x' the neighbour in $DS(X, Y)$ ($DC(X, Y)$ respectively) of every vertex in X which is not in X and denote by y' the neighbour in $DS(X, Y)$ ($DC(X, Y)$ respectively) of every vertex in Y which is not in Y . Note that in both graphs vertices x' and y' are uniquely defined.

First, we show the equivalence of conditions a) and e). Assume that e) does not hold and there exists an irreducible distinguishing vertex colouring c of $DS(X, Y)$. In particular, it means that every vertex in X has different colour and every vertex in Y has different colour. Hence, $|c(X)| = |X|$ and $|c(Y)| = |Y|$. As condition e) does not hold, there exists a vertex $x \in X$ such that $c(x) \notin c(Y)$ and a vertex $y \in Y$ such that $c(y) \notin c(X)$. Let c' be a colouring obtained from c by recolouring every vertex of colour $c(x)$ with colour $c(y)$. Colouring c' is a distinguishing reduction of c which contradicts the assumption of irreducibility of c .

Assume now that e) holds. Without loss of generality we can assume that $|X| \leq |Y|$ and that Y has at least two elements. Let f be an injection from X to Y . Let $c(x) = f(x)$ for $x \in X$ and let $c(y) = y$ for $y \in Y$. If f is bijection, then there exists an automorphism φ of $DS(X, Y)$ which maps X into Y . In that case we colour x' and y' in c with arbitrary but different colours from set Y . Note that φ is not preserved by c . If f is not a bijection, then we colour x and y with the same colour from $c(Y)$. The defined colouring c is irreducible and distinguishing which finishes the proof of the equivalence of conditions a) and e).

Now we prove the equivalence between a) and c). Recall that $DS(X, Y)$ and $DC(X, Y)$ have the same set of vertices. Let c be an irreducible distinguishing vertex colouring of $DS(X, Y)$. We first recolour x' and y' with new, distinct colours to obtain a proper vertex colouring c' of $DC(X, Y)$. Either c' is irreducible or we can find an irreducible proper vertex colouring by applying a reduction operation to c' at most three times.

Now, let c be an irreducible proper vertex colouring of $DC(X, Y)$. As $c(x') \neq c(y')$, colouring c is a distinguishing colouring of $DS(X, Y)$. Either c is an irreducible distinguishing colouring or we can find an irreducible distinguishing vertex colouring by applying a reduction operation to c at most two times. This finishes the proof of the equivalence of conditions a) and c).

The proof of the equivalence between b) and d) is trivial. We obtain the desired colourings from each other by recolouring the edge $x'y'$.

It remains to show the equivalence between conditions c) and d). Let c be a proper irreducible vertex colouring of $DC(X, Y)$. We construct an edge colouring c' in the following way. We assign colour $c(x)$ to the edge $x'x$ for every $x \in X$, colour $c(y)$ to the edge $y'y$ for every $y \in Y$. Furthermore, we assign a new colour to the edge $x'y'$. We can obtain an irreducible proper edge colouring of $DS(X, Y)$ from c' using reduction operation at most three times.

Similarly, if c' is an irreducible proper edge colouring of $DS(X, Y)$, then we assign to each vertex $x \in X$ colour $c(x) = c'(xx')$, and we assign to each vertex $y \in Y$ colour $c(y) = c'(yy')$. We colour x' in c with an arbitrary colour from the set $c(Y) \setminus c(X)$ or a new colour if this set is empty, and we colour y' in c with an arbitrary colour from the set $c(X) \setminus c(Y)$ or a new colour different from $c(x)$ if this set is empty. Colouring c is an irreducible proper vertex colouring of $DC(X, Y)$. This finishes the proof of equivalence between conditions c) and d) and thus the proof of the lemma. \square

Let X and Y be non-empty sets and assume that there exists an irreducible proper or distinguishing colouring of $DS(X, Y)$ ($DC(X, Y)$ respectively). From Lemma 3.2 we obtain that $|X| \leq |Y|$ or $|X| \leq |Y|$. It follows that if for every graph of the form $DS(X, Y)$ ($DC(X, Y)$ respectively) we have irreducible proper or distinguishing colouring, then the law of trichotomy holds and so does the Axiom of Choice.

Moreover, if the Axiom of Choice does not hold, then, by Zermelo's Theorem, there exists a set Y which is not well-orderable. By Lemma 3.2 and the fact that $|Y| \leq |Y|$, it follows that graph $DS(Y, Y)$ ($DC(Y, Y)$ respectively) has an irreducible colouring but none of the parameters $D(DS(Y, Y))$, $D'(DS(Y, Y))$, $\chi'(DS(Y, Y))$, $\chi'(DC(Y, Y))$ exists. Hence, if the Axiom of Choice fails, then the existence of an irreducible colouring of a graph G is not equivalent to the existence of the corresponding parameter for G , and it is indeed a strictly weaker condition for each of the considered types of colourings.

Summarizing the above paragraph and Lemma 3.2, we can formulate a list of conditions equivalent to the Axiom of Choice. Note that conditions (AC9)–(AC16) can be replaced with their restrictions to connected graphs.

Theorem 3.3. *The following conditions are equivalent.*

(AC1) *The Axiom of Choice.*

- (AC9) *Every graph has an irreducible distinguishing vertex colouring.*
- (AC10) *Every graph without component isomorphic to K_1 or K_2 has an irreducible distinguishing edge colouring.*
- (AC11) *Every graph has an irreducible proper vertex colouring.*
- (AC12) *Every graph has an irreducible proper edge colouring.*
- (AC13) *Every graph with an irreducible proper vertex colouring has the chromatic number.*
- (AC14) *Every graph with an irreducible proper edge colouring has the chromatic index.*
- (AC15) *Every graph with an irreducible distinguishing vertex colouring has the distinguishing number.*
- (AC16) *Every graph with an irreducible distinguishing edge colouring has the distinguishing index.*

References

- [1] L. Babai. Graph Isomorphism in Quasipolynomial Time. arXiv:1512.03547v2.
- [2] L. Babai. Asymmetric trees with two prescribed degrees. *Acta Math. Acad. Sci. Hung.*, 29:193–200, 1977.
- [3] P. J. Cohen. *Set Theory and the Continuum Hypothesis*. Benjamin, 1966.
- [4] N. G. de Bruijn and P. Erdős. A Colour Problem for Infinite Graphs and a Problem in the Theory of Relations. *Period. Math. Hung.*, 54:371–373, 1951.
- [5] R. Diestel. *Graph Theory*. Springer-Verlag Berlin Heidelberg, 2017.
- [6] F. Galvin and P. Komjáth. Graph colorings and the axiom of choice. *Period. Math. Hung.*, 22:71–75, 1991.
- [7] H. Herrlich. *Axiom of Choice*. Springer-Verlag, 2006.
- [8] T. Jech. *Set Theory*. Springer-Verlag Berlin Heidelberg New York, 2003.
- [9] R. Kalinowski and M. Pilśniak. Distinguishing graphs by edge-colourings. *European J. Combin.*, 45:124–131, 2015.
- [10] J. Lake. A Generalization of a Theorem of de Bruijn and Erdős on the Chromatic Number of Infinite Graphs. *J. Combin. Theory Ser. B*, 18(2):170–174, 1975.
- [11] F. Lehner, M. Pilśniak, and M. Stawiski. Distinguishing infinite graphs with bounded degrees. *preprint*, arXiv:1810.03932.

- [12] A. Mostowski. Axiom of choice for finite sets. *Fund. Math.*, 33:137–168, 1945.
- [13] M. Pilśniak and M. Stawiski. The Optimal General Upper Bound for the Distinguishing Index of Infinite Graphs. *J. Graph Theory*, 93:463–469, 2020.