

DELTA-POINTS IN BANACH SPACES GENERATED BY ADEQUATE FAMILIES

TROND A. ABRAHAMSEN, VEGARD LIMA, AND ANDRÉ MARTINY

ABSTRACT. We study delta-points in Banach spaces $h_{\mathcal{A},p}$ generated by adequate families \mathcal{A} where $1 \leq p < \infty$. When the family \mathcal{A} is regular and $p = 1$, these spaces are known as combinatorial Banach spaces. When $p > 1$ we prove that neither $h_{\mathcal{A},p}$ nor its dual contain delta-points. Under the extra assumption that \mathcal{A} is regular, we prove that the same is true when $p = 1$. In particular the Schreier spaces and their duals fail to have delta-points. If \mathcal{A} consists of finite sets only we are able to rule out the existence of delta-points in $h_{\mathcal{A},1}$ and Daugavet-points in its dual.

We also show that if $h_{\mathcal{A},1}$ is polyhedral, then it is either (I)-polyhedral or (V)-polyhedral (in the sense of Fonf and Veselý).

1. INTRODUCTION

According to Talagrand [Tal79, Tal84], a family \mathcal{A} of subsets of \mathbb{N} is *adequate* if \mathcal{A} contains the singletons, is hereditary and is compact with respect to the topology of pointwise convergence. We study the spaces $h_{\mathcal{A},p}$, $1 \leq p < \infty$, generated by an adequate family \mathcal{A} by completing the space c_{00} of all finitely supported sequences with the norm

$$\left\| \sum_{i=1}^n a_i e_i \right\| = \sup_{A \in \mathcal{A}} \left(\sum_{i \in A} |a_i|^p \right)^{1/p}.$$

Examples of $h_{\mathcal{A},p}$ spaces are c_0 , ℓ_p , $\ell_1(c_0)$ and the (higher order) Schreier space(s). We study the presence or absence of *Daugavet*- and *delta*-points in $h_{\mathcal{A},p}$ spaces and their duals.

For a Banach space X we denote by B_X , S_X and X^* the closed unit ball, the unit sphere and the topological dual of X respectively. Recall that a *slice* of B_X is a set

$$S(x^*, \delta) = \{x \in B_X : x^*(x) > 1 - \delta\},$$

where $x^* \in S_{X^*}$ and $\delta > 0$. Following [AHL20] we say that $x \in S_X$ is a *Daugavet*-point (resp. *delta*-point) if every element in the unit ball (resp. x itself) is in the closed convex hull of unit ball elements that

2010 *Mathematics Subject Classification*. Primary 46B20, 46B22, 46B04.

Key words and phrases. delta-point, Daugavet-point, 1-unconditional basis, adequate families.

V. Lima acknowledges the support by the Estonian Research Council grant PRG877.

are almost at distance 2 from x . In [AHL20, Lemmas 2.2 and 2.3], we find the following characterization of Daugavet- and delta-points which will serve as our definitions.

Lemma 1.1. *Let X be a Banach space. Then $x \in S_X$ is a*

- (i) *delta-point if and only if for every slice $S(x^*, \delta)$ of B_X with $x \in S(x^*, \delta)$ and for every $\varepsilon > 0$ there exists $y \in S(x^*, \delta)$ such that $\|x - y\| \geq 2 - \varepsilon$.*
- (ii) *Daugavet-point if and only if for every slice $S(x^*, \delta)$ of B_X and for every $\varepsilon > 0$ there exists $y \in S(x^*, \delta)$ such that $\|x - y\| \geq 2 - \varepsilon$.*

A Banach space has the Daugavet property if (and only if) every $x \in S_X$ is a Daugavet-point. For more on Daugavet- and delta-points see [AHL20, ALMT20, HPV21, JR20, RZ20].

It is known that if a Banach space X has the Daugavet property, then X does not have an unconditional basis [Kad96, Corollary 2.3]. In [ALMT20, Theorem 4.7] it was shown that there exists an $h_{\mathcal{A},1}$ space (with 1-unconditional basis) with “lots” of Daugavet-points, in the sense that its unit ball is the weak closure of its Daugavet-points. After some preliminaries in Section 2, we show in Section 3 that if $1 < p < \infty$ then neither $h_{\mathcal{A},p}$ nor $h_{\mathcal{A},p}^*$ have delta-points.

The case $p = 1$ studied in Section 4 is not so clear cut. As mentioned above there exist $h_{\mathcal{A},1}$ spaces with delta-points, but if \mathcal{A} consists of all subsets of \mathbb{N} , then $h_{\mathcal{A},1} = \ell_1$ and by [ALMT20, Theorem 2.17] ℓ_1 does not have delta-points. We show that if the adequate family \mathcal{A} contains an infinite set then $h_{\mathcal{A},1}^*$ always has delta-points. Our main focus in Section 4 is on adequate families of finite sets. If \mathcal{A} is an adequate family of finite subsets it follows from the results of [ALMT20] that $h_{\mathcal{A},1}$ does not have delta-points. If \mathcal{A} is in addition spreading (see Section 2) then the $h_{\mathcal{A},1}$ spaces we get are the combinatorial Banach spaces studied by Antunes, Beanland and Chu in [ABC21]. Using results from [ABC21] we show that if $h_{\mathcal{A},1}$ is a combinatorial Banach space, then also $h_{\mathcal{A},1}^*$ fail to have delta-points.

In Section 4 we also show that all extreme points of $h_{\mathcal{A},1}^*$ are w^* -exposed for all families \mathcal{A} . If \mathcal{A} consists of finite sets only, then the extreme points in $h_{\mathcal{A},1}^*$ are in fact w^* -strongly exposed. We use this observation to show that such $h_{\mathcal{A},1}^*$ spaces fail to have Daugavet-points.

Finally, in Section 5 we study polyhedrality in $h_{\mathcal{A},1}$ spaces. A Banach space is *polyhedral* if the unit ball of every finite-dimensional subspace of X is a polytope [Kle60]. In [ABC21] it was shown that combinatorial Banach spaces are all (V)-polyhedral (in the sense of Fonf and Vesely [FV04]). In fact, their proofs also work without the spreading property and show that all $h_{\mathcal{A},1}$ spaces where the adequate family consists of finite subsets of \mathbb{N} are (V)-polyhedral. We observe that $h_{\mathcal{A},1}$ is polyhedral if and only if $h_{\mathcal{A},1}$ is (V)-polyhedral if and only if \mathcal{A} is an adequate

family of finite sets. Furthermore, we show that $h_{\mathcal{A},1}$ is (I)-polyhedral if and only if $h_{\mathcal{A},1}$ is (IV)-polyhedral if and only if $\{A \in \mathcal{A} : i \in A\}$ is finite for all $i \in \mathbb{N}$.

We conclude the paper with some questions arising from the present work.

2. PRELIMINARIES

Recall that a Schauder basis $(e_i)_{i \in \mathbb{N}}$ of a Banach space X is a *1-unconditional basis* if for all $N \in \mathbb{N}$ and all scalars $a_1, \dots, a_N, b_1, \dots, b_N$ such that $|a_i| \leq |b_i|$ for $i = 1, \dots, N$ then the following inequality holds:

$$\left\| \sum_{i=1}^N a_i e_i \right\| \leq \left\| \sum_{i=1}^N b_i e_i \right\|.$$

A basis $(e_i)_{i \in \mathbb{N}}$ is *normalized* if $\|e_i\| = 1$ for all i and it is *shrinking* if the biorthogonal functionals $(e_i^*)_{i \in \mathbb{N}}$ is a basis for X^* . For $x \in X$ the *support* of x is defined by $\text{supp}(x) = \{i \in \mathbb{N} : e_i^*(x) \neq 0\}$.

If $(e_i)_{i \in \mathbb{N}}$ is 1-unconditional, then by the classic result of James it is shrinking if and only if X does not contain an isomorphic copy of ℓ_1 . If $(e_i)_{i \in \mathbb{N}}$ is a 1-unconditional basis then for any $A \subset \mathbb{N}$ the projection P_A defined by

$$P_A\left(\sum_{i \in \mathbb{N}} x_i e_i\right) = \sum_{i \in A} x_i e_i$$

satisfies $\|P_A\| \leq 1$.

As mentioned earlier we will study the existence of delta- and Daugavet-points in sequence spaces with 1-unconditional bases generated by adequate families of subsets of \mathbb{N} .

Definition 2.1. A family \mathcal{A} of subsets of \mathbb{N} is *adequate* if

- (i) \mathcal{A} contains the empty set and the singletons: $\{i\} \in \mathcal{A}$ for all $i \in \mathbb{N}$.
- (ii) \mathcal{A} is hereditary: If $A \in \mathcal{A}$ and $B \subseteq A$, then $B \in \mathcal{A}$.
- (iii) \mathcal{A} is compact with respect to the topology of pointwise convergence: Given $A \subset \mathbb{N}$, if every finite subset of A is in \mathcal{A} , then $A \in \mathcal{A}$.

We denote by \mathcal{A}^{MAX} the maximal elements of \mathcal{A} , that is $A \in \mathcal{A}^{\text{MAX}}$ if $A \in \mathcal{A}$ and $B \in \mathcal{A}$ with $A \subseteq B$ implies that $A = B$.

Let c_{00} be the vector space of all finitely supported sequences with standard basis $(e_i)_{i \in \mathbb{N}}$. If \mathcal{A} is adequate and $1 \leq p < \infty$, then $h_{\mathcal{A},p}$ is the completion of c_{00} with respect to the norm

$$\left\| \sum_{i=1}^n a_i e_i \right\| = \sup_{A \in \mathcal{A}} \left(\sum_{i \in A} |a_i|^p \right)^{1/p}.$$

It is clear that $(e_i)_{i \in \mathbb{N}}$ is a normalized 1-unconditional basis for $h_{\mathcal{A},p}$.

If \mathcal{A} is an adequate family of finite sets of \mathbb{N} which is *spreading*, that is, if $\{k_1, \dots, k_n\} \in \mathcal{A}$ and $k_i \leq l_i$, then $\{l_1, \dots, l_n\} \in \mathcal{A}$, then \mathcal{A} is often called a *regular* family of subsets of \mathbb{N} and the space $h_{\mathcal{A},1}$ is called a combinatorial Banach space and $h_{\mathcal{A},p}$ is its p -convexification (see e.g. [ABC21]).

From [ALMT20] we need the notion of *minimal norming subsets* for vectors in a Banach space with 1-unconditional basis (see also the notion of *1-sets* in [BDHQ19]).

Definition 2.2. For any Banach space X with 1-unconditional basis $(e_i)_{i \in \mathbb{N}}$ and for $x \in X$, define

$$M(x) := \{A \subseteq \mathbb{N} : \|P_A x\| = \|x\|, \|P_A x - x_i e_i\| < \|x\|, \text{ for all } i \in A\},$$

and

$$M^\infty(x) := \{A \in M(x) : |A| = \infty\}.$$

For any $B = (b_i)_{i \in \mathbb{N}} \subseteq \mathbb{N}$, assume that the elements are ordered, that is, $b_i < b_{i+1}$ for all i . For $n \in \mathbb{N}$ we define $B(n) := (b_i)_{i=1}^n$. Let X be a Banach space with a 1-unconditional basis $(e_i)_{i \in \mathbb{N}}$, if $n \in \mathbb{N}$, then by [ALMT20, Lemma 2.8] the set $\bigcup_{D \in M^\infty(x)} \{D(n)\}$ is finite whenever $x \in S_X$. This means that there exists $s = s(n) \in \mathbb{N}$ and sets D_1, \dots, D_s in $M^\infty(x)$ such that

$$\bigcup_{D \in M^\infty(x)} \{D(n)\} = \{D_1(n), D_2(n), \dots, D_s(n)\}.$$

Our next goal is to use [ALMT20, Lemma 2.14] to show that if the number of elements s in this set does not grow too fast as n increases, then x is not a delta-point.

Theorem 2.3. *Let X be a Banach space with 1-unconditional basis. If for $x \in S_X$ there exists $n \in \mathbb{N}$ such that for $s = |\bigcup_{D \in M^\infty(x)} \{D(n)\}|$*

$$\|P_E x\| > 1 - \frac{1}{2s} \quad \text{for all } E \in \bigcup_{D \in M^\infty(x)} \{D(n)\},$$

then x is not a delta-point.

In particular, if $|M^\infty(x)| < \infty$, then x is not a delta-point.

Proof. Let X have 1-unconditional basis $(e_i)_{i \in \mathbb{N}}$ and let $x \in S_X$. Since changing signs of the coordinates of vectors is an isometry on spaces with 1-unconditional basis and delta-points are preserved by isometries we may assume that $x = (x_i)_{i \in \mathbb{N}} \in S_X$ with $x_i \geq 0$ for all $i \in \mathbb{N}$. By assumption there exists $n \in \mathbb{N}$, $s \in \mathbb{N}$ and $D_1, \dots, D_s \in M^\infty(x)$ such that $\|P_{D_k(n)} x\| > 1 - \frac{1}{2s}$ for all $k = 1, \dots, s$.

For each $k \leq s$ let $x_k^* \in S_{X^*}$ be such that $x_k^*(P_{D_k(n)} x) = \|P_{D_k(n)} x\|$ and $x_k^*(e_j) = 0$ for all $j \notin D_k(n)$.

Let $x^* = \frac{1}{s} \sum_{k=1}^s x_k^*$. Then $x^*(x) > 1 - \frac{1}{2s}$, so $x \in S(x^*, \frac{1}{2s})$. If $y \in S(x^*, \frac{1}{2s})$ it follows that $x_k^*(y) > 1 - \frac{1}{2}$ for all $k = 1, \dots, s$. Fix k .

If $y_i \leq x_i/2$ for all $i \in D_k(n)$, then $x_k^*(y) \leq \frac{1}{2}x_k^*(x) \leq \frac{1}{2}$. Thus for each $1 \leq k \leq s$, there exists $i \in D_k(n)$ such that $y_i > \frac{1}{2}x_i$ and therefore we can apply [ALMT20, Lemma 2.14] with $S(x^*, \frac{1}{2s})$, n and $\eta = \frac{1}{2}$ and conclude that x is not a delta-point. \square

Example 2.4. Recall that a norm in a Banach lattice X is called *strictly monotone* if $\|x + y\| > \|x\|$ whenever $x, y \in X$ with $x, y \geq 0$ and $y \neq 0$. From Theorem 2.3 we immediately get that a Banach space X with a 1-unconditional basis and such a norm has no delta-points since in this case $M(x) = \{\text{supp}(x)\}$ for any $x \in S_X$.

Example 2.5. The space $X = \ell_1 \oplus_\infty \ell_1$ has a 1-unconditional basis. Moreover, it is easily seen that $|M^\infty(x)| \leq 2$ for any $x \in S_X$, so X has no delta-points.

3. THE CASE $1 < p < \infty$

In this section we prove the following theorem.

Theorem 3.1. *Let \mathcal{A} be an adequate family of subsets of \mathbb{N} and let $1 < p < \infty$. Then*

- (i) $h_{\mathcal{A},p}$ does not have delta-points.
- (ii) $h_{\mathcal{A},p}^*$ does not have delta-points.

Proof of Theorem 3.1 (i). Let $x \in S_{h_{\mathcal{A},p}}$. As in the proof of Theorem 2.3 we assume without loss of generality that $x_i \geq 0$ for all i .

As noted above Theorem 2.3 we can find $k \in \mathbb{N}$ and $D_j \in M^\infty(x)$ for $j = 1, \dots, k$ such that

$$\bigcup_{D \in M^\infty(x)} \{D(1)\} = \{D_1(1), \dots, D_k(1)\}.$$

For each $j = 1, \dots, k$ define

$$x_j^* = \sum_{i \in D_j} x_i^{p-1} e_i.$$

Let $q > 1$ such that $\frac{1}{p} + \frac{1}{q} = 1$. Then by Hölders inequality we have

$$\begin{aligned} |x_j^*(y)| &\leq \sum_{i \in D_j} |x_i^{p-1}| |y_i| \leq \left(\sum_{i \in D_j} (|x_i^{p-1}|)^q \right)^{1/q} \left(\sum_{i \in D_j} |y_i|^p \right)^{1/p} \\ &\leq \|x\|^{p/q} \|y\| \leq 1 \end{aligned}$$

As $x_j^*(x) = 1$ we conclude that $x_j^* \in S_{X^*}$.

For each j write $D_j = (d_i^j)_{i \in \mathbb{N}}$ with $d_i^j < d_{i+1}^j$. Define

$$\xi_j := (x_{d_1^j}^{p-1}, x_{d_2^j}^{p-1}, \dots) \in S_{\ell_q}$$

and $T_j : h_{\mathcal{A},p} \rightarrow \ell_p$ by

$$T_j\left(\sum_{i=1}^{\infty} a_i e_i\right) = (a_{d_1^j}, a_{d_2^j}, \dots).$$

Note that $\|T_j(x)\|_p = 1$ since $D_j \in M^\infty(x)$. Let $\varepsilon := \frac{1}{2} \min_j x_{d_1^j} > 0$. Clarkson [Cla36] showed that ℓ_p is uniformly convex so there exists $\delta_j > 0$ such that if

$$y \in S(\xi_j, \delta_j) \subseteq B_{\ell_p}$$

then $\|T_j(x) - y\|_p < \varepsilon$.

Define $\delta := \min_j \delta_j$ and $x^* := \frac{1}{k} \sum_{j=1}^k x_j^*$. We have $x^*(x) = 1$ and if $z \in S(x^*, \frac{\delta}{k})$, then

$$T_j(z) \in S(\xi_j, \delta) \subseteq S(\xi_j, \delta_j),$$

hence $|x_{d_1^j} - z_{d_1^j}| \leq \|T_j(x) - T_j(z)\| < \varepsilon$. By definition of ε we get $z_{d_1^j} \geq \frac{1}{2} x_{d_1^j} > 0$. Applying [ALMT20, Lemma 2.14] with x^* , δ/k , $\eta = \frac{1}{2}$ and $n = 1$ the result follows. \square

For the proof that $h_{\mathcal{A},p}^*$ does not have delta-points we first need to show that the standard basis is shrinking.

Lemma 3.2. *Let \mathcal{A} be an adequate family and let $1 < p < \infty$. Then the standard basis $(e_i)_{i \in \mathbb{N}}$ of $h_{\mathcal{A},p}$ is shrinking. In particular, $h_{\mathcal{A},p}^*$ is separable and has an unconditional basis.*

Proof. Let $(x_n)_{n \in \mathbb{N}}$ be a normalized block basis of $(e_i)_{i \in \mathbb{N}}$. This means that there is a sequence $1 \leq p_1 < p_2 < \dots$ and coefficients (a_i^n) such that $x_n = \sum_{i=p_n}^{p_{n+1}-1} a_i^n e_i$ satisfies $\|x_n\| = 1$.

Define an operator $S : \ell_p \rightarrow h_{\mathcal{A},p}$ by

$$S((\lambda_n)) = \sum_{n=1}^{\infty} \lambda_n x_n.$$

We have that S is a bounded linear operator. Indeed, for $(\lambda_n) \in \ell_p$ and $A \in \mathcal{A}$ we define $A_n = A \cap \text{supp}(x_n)$. Then we have

$$\sum_{i \in A} |S((\lambda_n))_i|^p = \sum_{n=1}^{\infty} \sum_{i \in A_n} |\lambda_n a_i^n|^p = \sum_{n=1}^{\infty} |\lambda_n|^p \sum_{i \in A_n} |a_i^n|^p \leq \sum_{n=1}^{\infty} |\lambda_n|^p \|x_n\|^p$$

and hence $\|S\| \leq 1$. If (f_n) denotes the standard basis in ℓ_p , then $S(f_n) = x_n$. Since S is weak-weak continuous we get that (x_n) is weakly null since (f_n) is weakly null in ℓ_p . By [AK06, Proposition 3.2.7] $(e_i)_{i \in \mathbb{N}}$ is shrinking. \square

Proof of Theorem 3.1 (ii). By Lemma 3.2 the standard basis $(e_i)_{i \in \mathbb{N}}$ for $X := h_{\mathcal{A},p}$ is shrinking and hence the biorthogonal functionals $(e_i^*)_{i \in \mathbb{N}}$ is an 1-unconditional basis for X^* . Furthermore, we know that X^{**} is a sequence space and that for $x^{**} = (a_i)_{i \in \mathbb{N}} \in X^{**}$ we have $\|x^{**}\| = \sup_N \|P_N x^{**}\|$ where $P_N(x^{**}) = \sum_{i=1}^N a_i e_i \in X$.

Let $x^* = (x_j^*)_{j \in \mathbb{N}} \in S_{X^*}$. Without loss of generality we may assume that $x_j^* \geq 0$ for each j . By Theorem 2.3 it is enough to show that $M^\infty(x^*)$ is finite. To this end we will show that $M(x^*) = \{\text{supp}(x^*)\}$.

Assume for contradiction that there exists $j \in \text{supp}(x^*)$ such that $\|x^* - x_j^* e_j^*\| = 1$. Find $x^{**} \in S_{X^{**}}$ such that $x^{**}(x^* - x_j^* e_j^*) = 1$ and $x^{**}(e_k^*) = 0$ for all $k \notin \text{supp}(x^* - x_j^* e_j^*)$. Let $y^{**} = x_j^{*q-1} e_j + (1 - x_j^{*q})^{1/p} x^{**}$. For $A \in \mathcal{A}$ and $N \in \mathbb{N}$ denote $A_N = A \cap \{1, \dots, N\}$. We have

$$\begin{aligned} \|P_N y^{**}\|^p &= \sup_{A \in \mathcal{A}} \sum_{i \in A_N} |y_i^{**}|^p \\ &\leq (x_j^{*q-1})^p + (1 - x_j^{*q}) \sup_{A \in \mathcal{A}} \sum_{i \in A_N} |x_i^{**}|^p \\ &\leq x_j^{*q} + (1 - x_j^{*q}) = 1. \end{aligned}$$

This yields $\|y^{**}\| = \sup_N \|P_N y^{**}\| \leq 1$ and since $x^* \in S_{X^*}$ we arrive at the contradiction

$$y^{**}(x^*) = x_j^{*q} + (1 - x_j^{*q})^{1/p} > x_j^{*q} + (1 - x_j^{*q}) = 1.$$

Hence we can leave no index behind and $M(x^*) = \{\text{supp}(x^*)\}$. \square

4. THE CASE $p = 1$

We now turn to $h_{\mathcal{A},1}$ spaces. In this case the situation is not as clear as for $p > 1$. Let us first note that the extreme points of the dual space has a well-known characterization.

Lemma 4.1 (Lemma 2.3 in [AM93]). *Let \mathcal{A} be an adequate family of \mathbb{N} . Then*

$$\text{ext } B_{h_{\mathcal{A},1}^*} = \left\{ \sum_{i \in A} \varepsilon_i e_i^* : A \in \mathcal{A}^{\text{MAX}}, \varepsilon_i \in \{-1, 1\} \right\}.$$

In [ALMT20, Theorem 3.1] it was shown that if we build an adequate family on \mathbb{N} by using the branches of the binary tree, then we get an $h_{\mathcal{A},1}$ space with delta-points. This adequate family contains many infinite sets. We have the following general result about duals of $h_{\mathcal{A},1}$ spaces when the adequate family contains an infinite set.

Proposition 4.2. *Let \mathcal{A} be an adequate family that contains an infinite set. Then there exists $x^* \in h_{\mathcal{A},1}^*$ such that x^* is an extreme point and a delta-point.*

Proof. As \mathcal{A} contains an infinite set there exist $A \in \mathcal{A}^{\text{MAX}}$ with $|A| = \infty$. Write $A = (a_i)_{i=1}^\infty$ with $a_i < a_{i+1}$ for all i . For $x^{**} \in S_{h_{\mathcal{A},1}^{**}}$ we have $y^* = \sum_{i=1}^\infty \text{sgn}(x^{**}(e_{a_i}^*)) e_{a_i}^* \in B_{h_{\mathcal{A},1}^*}$ by Lemma 4.1. Hence

$$\sum_{i=1}^\infty |x^{**}(e_{a_i}^*)| = x^{**}(y^*) \leq 1$$

and this implies that $(e_{a_i}^*)_{i=1}^\infty$ is weakly null.

Now let $x^* = \sum_{i \in A} e_i^* \in \text{ext } B_{h_{\mathcal{A},1}^*}$. Let $x^{**} \in S_{h_{\mathcal{A},1}^{**}}$ and $\varepsilon > 0$ with $x^* \in S(x^{**}, \varepsilon)$. Define $x_i^* = x^* - 2e_{a_i}^* \in \text{ext } B_{h_{\mathcal{A},1}^*}$ so that $\|x^* - x_i^*\| = \|2e_{a_i}^*\| = 2$. For i large enough we have $x_i^* \in S(x^{**}, \varepsilon)$ and thus x is a delta-point. \square

We shall henceforth exclusively consider $h_{\mathcal{A},1}$ spaces where the adequate family consists of finite sets only. Recall that a Banach spaces is said to be *polyhedral* if the unit ball of each of its finite-dimensional subspaces is a polyhedron [Kle60].

The following result is well-known but we include a proof for easy reference.

Proposition 4.3. *Let \mathcal{A} be an adequate family of subsets of \mathbb{N} . The following are equivalent:*

- (i) \mathcal{A} consists of finite sets only;
- (ii) $h_{\mathcal{A},1}$ is polyhedral;
- (iii) The standard basis $(e_i)_{i \in \mathbb{N}}$ is shrinking.

Proof. (i) \implies (ii) Note that the proof of Theorem 4.5 (see also Remark 4.4) [ABC21] also holds for adequate families of finite sets, not just for regular families, so that $h_{\mathcal{A},1}$ is (V)-polyhedral in the sense Fonf and Veselý [FV04]. In particular, $h_{\mathcal{A},1}$ is polyhedral.

(ii) \implies (iii) If $h_{\mathcal{A},1}$ is polyhedral, then it is c_0 saturated by [Fon80] (see the remark following Theorem 5). Hence $h_{\mathcal{A},1}$ cannot contain an isomorphic copy of ℓ_1 and this yields that $(e_i)_{i \in \mathbb{N}}$ is shrinking since it is unconditional.

(iii) \implies (i). Assume \mathcal{A} contains an infinite set A . Then the basis vectors $(e_i)_{i \in A}$ span an isometric copy of ℓ_1 in $h_{\mathcal{A},1}$ and $(e_i)_{i \in \mathbb{N}}$ is not shrinking. \square

We will have more to say about polyhedrality in $h_{\mathcal{A},1}$ spaces below in Section 5.

Next we note that extreme points in $h_{\mathcal{A},1}^*$ are actually w^* -exposed.

Proposition 4.4. *Let \mathcal{A} be an adequate family of \mathbb{N} and $x^* \in \text{ext } B_{h_{\mathcal{A},1}^*}$. Then the following are equivalent:*

- (i) x^* is an extreme point of B_{X^*} ;
- (ii) x^* is a w^* -exposed point of B_{X^*} .

In particular, if \mathcal{A} is an adequate family of finite sets, then x^ is an extreme point if and only if x^* is a w^* -strongly exposed point.*

Remark 4.5. We showed above in Proposition 4.2 that if the adequate family \mathcal{A} contains an infinite set, then there is an extreme point in $h_{\mathcal{A},1}^*$ which is also a delta-point. Proposition 4.4 shows that this extreme point is also exposed, but being a delta-point it is far from being a w^* -strongly exposed point.

Proof. One direction is trivial, so we only need to show (i) \implies (ii). Let $x^* \in \text{ext } B_{h_{\mathcal{A},1}^*}$. By Lemma 4.1 we can write $x^* = \sum_{i \in B} \varepsilon_i e_i^*$ with $B \in \mathcal{A}^{\text{MAX}}$ and $\varepsilon_i \in \{-1, 1\}$. Find some $y \in S_{\ell_1}$ with $\text{supp}(y) = B$ and $y_i \geq 0$ for all $i \in \mathbb{N}$. Define $x = \sum_{i \in B} y_i \varepsilon_i e_i \in S_{h_{\mathcal{A},1}}$ and notice that $x^*(x) = 1$.

Take some $y^* \in S_{h_{\mathcal{A},1}^*}$ with $y^*(x) = 1$. Then $y_i^* = y^*(e_i) = \varepsilon_i$ for all $i \in B$. For $j \in \mathbb{N} \setminus B$, there must exist, by compactness of \mathcal{A} , some finite $A \subset B$ such that $A \cup \{j\} \notin \mathcal{A}$. Define

$$\tilde{x} = x + \text{sgn } y_i^* \min_{i \in A} |x_i| e_j,$$

and observe that $\|\tilde{x}\| = 1$. Indeed, take $C \in \mathcal{A}$ with $j \in C$. Then $A \setminus C \neq \emptyset$, and so

$$\sum_{i \in C} |\tilde{x}_i| = \sum_{i \in C \cap B} |\tilde{x}_i| + |\tilde{x}_j| \leq \sum_{i \in B} |x_i| - \min_{i \in A} |x_i| + \min_{i \in A} |x_i| = 1.$$

We now get

$$1 \geq y^*(\tilde{x}) = y^*(x) + \min_{i \in A} |x_i| |y_j^*| = 1 + \min_{i \in A} |x_i| |y_j^*|,$$

implying that $y_j^* = 0$ for all $j \in \mathbb{N} \setminus B$. That is, $y^* = x^*$.

If \mathcal{A} consists of finite sets, then $h_{\mathcal{A},1}$ is polyhedral and for polyhedral spaces the w^* -exposed points and w^* -strongly exposed points of the dual unit ball coincides ([Fon00, Theorem 1.4]). \square

Let \mathcal{A} be an adequate family and let $x \in S_{h_{\mathcal{A},1}}$. Since \mathcal{A} is compact in the topology of pointwise convergence in \mathbb{N} there exists $A \in \mathcal{A}$ such that $\|P_A x\| = \|x\|$. In particular, $M(x) \subset \mathcal{A}$. The following result is now immediate from Theorem 2.3 (or [ALMT20, Proposition 2.15]).

Proposition 4.6. *If \mathcal{A} is an adequate family of finite subsets of \mathbb{N} , then $h_{\mathcal{A},1}$ does not have delta-points.*

Definition 4.7. A Banach space X has the *convex series representation property* (CSRP) if for each $x \in B_X$, there exists a sequence (f_i) of extreme points of B_X and a sequence of nonnegative real numbers (λ_i) such that $\sum_{i=1}^{\infty} \lambda_i = 1$ and

$$x = \sum_{i=1}^{\infty} \lambda_i f_i.$$

Note that the CSRP is equivalent to the λ -property [ALS91]. The proof of [ABC21, Proposition 4.3] also holds for adequate families of finite sets and thus we have the following.

Proposition 4.8 (Proposition 4.3 in [ABC21]). *Let \mathcal{A} be an adequate family of finite subsets of \mathbb{N} . Then $h_{\mathcal{A},1}^*$ has the CSRP.*

Using this proposition we are able to show that for regular families \mathcal{A} the dual of $h_{\mathcal{A},1}$ does not have delta-points.

Proposition 4.9. *If \mathcal{A} is an adequate family of finite subsets of \mathbb{N} which is spreading, then $h_{\mathcal{A},1}^*$ fails to have any delta-points.*

Proof. By assumption the standard basis $(e_i)_{i \in \mathbb{N}}$ for $h_{\mathcal{A},1}$ is shrinking and hence the biorthogonal functionals $(e_i^*)_{i \in \mathbb{N}}$ is a 1-unconditional basis for $h_{\mathcal{A},1}^*$.

We only need to consider x^* with infinite support by Theorem 2.3. From Lemma 4.1 we have that elements of $\text{ext } h_{\mathcal{A},1}^*$ can be written

$$x_{A,(\varepsilon_i)}^* = \sum_{i \in A} \varepsilon_i e_i^* \quad \text{where } A \in \mathcal{A}^{\text{MAX}} \text{ and } (\varepsilon_i) \in \{-1, 1\}.$$

By Proposition 4.8 we can write $x^* = \sum_{n=1}^{\infty} \lambda_n x_{F_n,(\varepsilon_i^n)}^*$ where for $F_n \in \mathcal{A}$ and $\varepsilon_i^n \in \{-1, 1\}$. There are two possibilities. Either $\|x^* - x_k^* e_k^*\| < 1$, for every $k \in \text{supp}(x^*)$, and then $M^\infty(x^*) = \{\text{supp}(x^*)\}$, and the result follows from Theorem 2.3. The other possibility is that there exists $k \in \text{supp}(x^*)$ such that $\|x^* - x_k^* e_k^*\| = 1$. Write

$$y^* := x^* - x_k^* e_k^* = \sum_{n=1}^{\infty} \lambda_n x_{F'_n,(\varepsilon_i^n)}^*,$$

where $F'_n = F_n \setminus \{k\}$ for all n . Find $y^{**} = (y_i) \in S_{X^{**}}$ such that

$$1 = y^{**}(y^*) = \sum_{n=1}^{\infty} \lambda_n y^{**}(x_{F'_n,(\varepsilon_i^n)}^*).$$

This implies that $y^{**}(x_{F'_n,(\varepsilon_i^n)}^*) = 1$ for every $n \in \mathbb{N}$. Now find $n \in \mathbb{N}$ with $k \in F_n$. Take any integer $j \in (F'_n)^C$ with $j \geq \max F_n$, let $G_n := F'_n \cup \{j\}$, and let $\varepsilon_j = \text{sgn } y_j$. As \mathcal{A} is spreading $G_n \in \mathcal{A}$ and thus $\|x_{G_n,(\varepsilon_i^n)}^*\| = 1$. But then,

$$1 \geq y^{**}(x_{G_n,(\varepsilon_i^n)}^*) = y^{**}(x_{F'_n,(\varepsilon_i^n)}^*) + y^{**}(\varepsilon_j e_j^*) = 1 + |y_j| \geq 1,$$

which forces y_j to be zero. From this it follows that $y^{**} = (y_i)$ is zero on all coordinates $i \geq \max F_n$ (at least). That is, every $A \in M(x^*)$ must be a subset of $\{1, \dots, \max F_n\}$, so $M^\infty(x^*) = \emptyset$ and the result follows from Theorem 2.3. \square

If \mathcal{A} is an adequate family of finite sets which is not spreading, we do not know whether or not $h_{\mathcal{A},1}^*$ can have delta-points. But we are able to say something about Daugavet-points.

Let X be a Banach space. Recall that $x \in S_X$ is a Daugavet-point if and only if for every slice $S(x^*, \delta)$ of B_X and for every $\varepsilon > 0$ there exists $y \in S(x^*, \delta)$ such that $\|x - y\| \geq 2 - \varepsilon$ (see e.g. [AHL20, Lemma 2.3]).

Proposition 4.10. *If $x^* \in S_{X^*}$ can be written as*

$$x^* = \sum_{i=1}^{\infty} \lambda_i f_i$$

where $\lambda_i \geq 0$, $\sum_{i=1}^{\infty} \lambda_i = 1$, f_i are (w^*) -strongly exposed points in B_{X^*} , then x^* is not a Daugavet-point.

In particular, if \mathcal{A} is an adequate family of finite sets, then $h_{\mathcal{A},1}^*$ does not have Daugavet-points.

Proof. Choose j with $\lambda_j > 0$. Find a slice $S(x^{**}, \delta)$ containing f_j with diameter less than 1. Then for any $y^* \in S(x^{**}, \delta)$ we have

$$\begin{aligned} \|x^* - y^*\| &\leq \left\| \sum_{i \neq j} \lambda_i f_i \right\| + \|y^* - \lambda_j f_j\| \\ &\leq \sum_{i \neq j} \lambda_i + (1 - \lambda_j) + \lambda_j \|y^* - f_j\| \\ &< 2(1 - \lambda_j) + \lambda_j = 2 - \lambda_j. \end{aligned}$$

So the distance from x^* to y^* is bounded away from 2. \square

5. POLYHEDRALITY

In this section we study polyhedrality in $h_{\mathcal{A},1}$ spaces. The goal is to describe the polyhedrality of $h_{\mathcal{A},1}$ spaces in terms of the structure of the adequate family \mathcal{A} . Let us begin by recalling some concepts and results.

If X is a Banach space and $A \in X^*$ then A' denotes the set of all w^* -limit points of A , that is

$$A' = \left\{ f \in X^* : f \in \overline{A \setminus \{f\}}^{w^*} \right\}.$$

In [FV04] Fonf and Vesely identified eight known definitions of polyhedrality from the literature and gave examples showing that in general they are different. We will use the following three definitions from their paper.

Definition 5.1. Let X be a Banach space. Then

- (i) X is (I)-polyhedral if $(\text{ext } B_{X^*})' \subseteq \{0\}$;
- (ii) X is (IV)-polyhedral if $f(x) < 1$ whenever $x \in S_X$ and $f \in (\text{ext } B_{X^*})'$;
- (iii) X is (V)-polyhedral if $\sup \{f(x) : f \in \text{ext } B_{X^*} \setminus D(x)\} < 1$ for each $x \in S_X$, where $D(x) = \{f \in S_{X^*} : f(x) = 1\}$.

In the proof of Proposition 4.3 we met the following:

Theorem 5.2 (Theorem 4.5 in [ABC21]). *Let \mathcal{A} be an adequate family of finite subsets of \mathbb{N} . Then $h_{\mathcal{A},1}$ is (V)-polyhedral.*

It is natural to ask whether any (V)-polyhedral $h_{\mathcal{A},1}$ can be (IV)- or even (I)-polyhedral. Our next goal is to show the following: If $h_{\mathcal{A},1}$ is polyhedral, then it is either (I)-polyhedral or (V)-polyhedral. Considering Proposition 4.3 and Theorem 5.2 we only need to show the following.

Theorem 5.3. *Let \mathcal{A} be an adequate family of finite sets. Then the following are equivalent:*

- (i) $\{A \in \mathcal{A} : i \in A\}$ is finite for all $i \in \mathbb{N}$.
- (ii) $h_{\mathcal{A},1}$ is (I)-polyhedral.
- (iii) $h_{\mathcal{A},1}$ is (IV)-polyhedral.

Proof. (ii) \implies (iii) is trivial.

(i) \implies (ii). By assumption and Lemma 4.1, for any $i \in \mathbb{N}$, there are only a finite number of extreme points that have support on i . It follows that if $f \in (\text{ext } B_{X^*})'$, then $f(e_i) = 0$ for all $i \in \mathbb{N}$. Therefore f is 0.

(iii) \implies (i). First observe that if $\{A \in \mathcal{A} : i \in A\}$ is infinite, then the set $\mathcal{C}_i = \{A \in \mathcal{A}^{\text{MAX}} : i \in A\}$ is also infinite.

Assume that for $i \in \mathbb{N}$ the set \mathcal{C}_i is infinite and let $(A_n)_{n=1}^{\infty} \subset \mathcal{C}_i$ be a sequence of distinct sets. As elements of \mathcal{A} are finite and \mathcal{A} is compact in the topology of pointwise convergence on \mathbb{N} , we can by passing to a subsequence if necessary assume that (A_n) converges pointwise to some $A \in \mathcal{A}$ and that $A \subset A_n$ for all $n \in \mathbb{N}$.

If $k \in \mathbb{N}$ then there exists $N \in \mathbb{N}$ such that $\{1, 2, \dots, k\} \cap A_n = \{1, 2, \dots, k\} \cap A$ for all $n > N$, i.e. $\min(A_n \setminus A) \rightarrow \infty$.

With the sequence (A_n) , we wish to show that $h_{\mathcal{A},1}$ is not (IV)-polyhedral. That is, we wish to show that there exist some $x^* \in (\text{ext } B_{h_{\mathcal{A},1}^*})'$ and some $x \in S_{h_{\mathcal{A},1}}$ such that $x^*(x) = 1$. To this end, define $x^* = \sum_{i \in A} e_i^*$ and $x = \frac{1}{|A|} \sum_{i \in A} e_i \in S_{h_{\mathcal{A},1}}$. As $x^*(x) = 1$, it remains to show that x^* is a w^* -limit of elements in $\text{ext } B_{h_{\mathcal{A},1}^*} \setminus \{x^*\}$.

By Lemma 4.1 $x_n^* := \sum_{i \in A_n} e_i^* = x^* + \sum_{i \in A_n \setminus A} e_i^* \in \text{ext } B_{h_{\mathcal{A},1}^*}$. If $y \in h_{\mathcal{A},1}$ we get that

$$\begin{aligned} |x_n^*(y) - x^*(y)| &= |x^*(y) + \sum_{i \in A_n \setminus A} y_i - x^*(y)| \\ &\leq \sum_{i \in A_n \setminus A} |y_i|. \end{aligned}$$

Since $\sum_{i \in A_n \setminus A} |y_i| \rightarrow 0$ because $\min(A_n \setminus A) \rightarrow \infty$ we get $x_n^* \xrightarrow{w^*} x^*$ and thus $h_{\mathcal{A},1}$ is not (IV)-polyhedral. \square

The above theorem shows that the only combinatorial Banach space which is (IV)-polyhedral is c_0 . Antunes, Beanland and Chu [ABC21, Theorem 4.5] (see also their Remark 4.4) proved that combinatorial Banach spaces are (V)-polyhedral. So our next corollary shows that this is in fact best possible.

Corollary 5.4. *If \mathcal{A} is a regular family of sets such that $h_{\mathcal{A},1}$ is (IV)-polyhedral, then $h_{\mathcal{A},1} = c_0$.*

Proof. Assume that $A \in \mathcal{A}$ with $|A| > 1$. Let $i = \min A$. Since there are infinitely many spreads of A we have that $\{A \in \mathcal{A} : i \in A\}$ is infinite. \square

We end the paper with some questions.

Let \mathcal{A} be an adequate family. Proposition 4.3 gives a simple characterization of when $h_{\mathcal{A},1}$ contains a copy of ℓ_1 and $h_{\mathcal{A},p}$ never contains ℓ_1 by Lemma 3.2 (and James' characterization of shrinking unconditional bases).

Note that $h_{\mathcal{A},1}$ and even $h_{\mathcal{A},p}$ contains an isometric copy of c_0 if there exists an infinite subset $E \subseteq \mathbb{N}$ such that $|A \cap E| \leq 1$ for all $A \in \mathcal{A}$. We ask:

Question 5.5. What is a natural condition on \mathcal{A} such that $h_{\mathcal{A},1}$ (or $h_{\mathcal{A},p}$) does not contain c_0 . That is, when does $h_{\mathcal{A},p}$ have a boundedly complete basis?

We have seen that if \mathcal{A} is an adequate family of finite sets which is spreading, then neither $h_{\mathcal{A},1}$ nor $h_{\mathcal{A},1}^*$ have delta-points. We do not know if that is also the case if \mathcal{A} is not spreading.

Question 5.6. If $h_{\mathcal{A},1}$ is (V)-polyhedral, does then $h_{\mathcal{A},1}^*$ fail to have delta-points?

One can even ask: If $h_{\mathcal{A},1}$ is (I)-polyhedral, does then $h_{\mathcal{A},1}^*$ fail to have delta-points? Note that if X is (I)-polyhedral, then X is isometric to a subspace of c_0 [FV04, Theorem 1.2].

ACKNOWLEDGEMENTS

The authors would like to thank Stanimir Troyanski for discussions on the topic of the paper.

REFERENCES

- [ABC21] L. Antunes, K. Beanland, and H. V. Chu, *On the geometry of higher order Schreier spaces*, Illinois Journal of Mathematics (to appear) (2021).
- [AHL20] T. A. Abrahamsen, R. Haller, V. Lima, and K. Pirk, *Delta- and Daugavet points in Banach spaces*, Proc. Edinb. Math. Soc. (2) **63** (2020), no. 2, 475–496. MR 4085036
- [AK06] F. Albiac and N. J. Kalton, *Topics in Banach space theory*, Graduate Texts in Mathematics, vol. 233, Springer, New York, 2006. MR 2192298 (2006h:46005)
- [ALMT20] T. A. Abrahamsen, V. Lima, A. Martiny, and S. Troyanski, *Daugavet- and delta-points in Banach spaces with unconditional bases*, preprint (2020), arXiv:2007.04946.
- [ALS91] R. M. Aron, R. H. Lohman, and A. Suárez, *Rotundity, the C.S.R.P., and the λ -property in Banach spaces*, Proc. Amer. Math. Soc. **111** (1991), no. 1, 151–155. MR 1030732
- [AM93] S. Argyros and S. Mercourakis, *On weakly Lindelöf Banach spaces*, Rocky Mountain J. Math. **23** (1993), no. 2, 395–446. MR 1226181

- [BDHQ19] K. Beanland, N. Duncan, M. Holt, and J. Quigley, *Extreme points for combinatorial Banach spaces*, *Glasg. Math. J.* **61** (2019), no. 2, 487–500. MR 3928650
- [Cla36] J. A. Clarkson, *Uniformly convex spaces*, *Trans. Amer. Math. Soc.* **40** (1936), no. 3, 396–414. MR 1501880
- [Fon80] V. P. Fonf, *Some properties of polyhedral Banach spaces*, *Funktsional. Anal. i Prilozhen.* **14** (1980), no. 4, 89–90. MR 595744
- [Fon00] ———, *On the boundary of a polyhedral Banach space*, *Extracta Math.* **15** (2000), no. 1, 145–154. MR 1792984
- [FV04] V. P. Fonf and L. Veselý, *Infinite-dimensional polyhedrality*, *Canad. J. Math.* **56** (2004), no. 3, 472–494. MR 2057283
- [HPV21] R. Haller, K. Pirk, and T. Veeorg, *Daugavet- and delta-points in absolute sums of Banach spaces*, *J. Convex Anal.* **28** (2021), no. 1, 41–54. MR 4199633
- [JR20] M. Jung and A. Rueda Zoca, *Daugavet points and Δ -points in Lipschitz-free spaces*, preprint (2020), arXiv:2010.09357.
- [Kad96] V. M. Kadets, *Some remarks concerning the Daugavet equation*, *Quaestiones Math.* **19** (1996), 225–235.
- [Kle60] V. Klee, *Polyhedral sections of convex bodies*, *Acta Math.* **103** (1960), 243–267. MR 139073
- [RZ20] A. Rueda Zoca, *L-orthogonality in Daugavet centers and narrow operators*, preprint (2020), arXiv:2010.16270.
- [Tal79] M. Talagrand, *Espaces de Banach faiblement \mathcal{K} -analytiques*, *Ann. of Math. (2)* **110** (1979), no. 3, 407–438. MR 554378
- [Tal84] ———, *A new countably determined Banach space*, *Israel J. Math.* **47** (1984), no. 1, 75–80. MR 736065

(T. A. Abrahamsen) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF AGDER, POSTBOKS 422, 4604 KRISTIANSAND, NORWAY.

Email address: `trond.a.abrahamsen@uia.no`

URL: `http://home.uia.no/trondaa/index.php3`

(V. Lima) DEPARTMENT OF ENGINEERING SCIENCES, UNIVERSITY OF AGDER, POSTBOKS 509, 4898 GRIMSTAD, NORWAY.

Email address: `Vegard.Lima@uia.no`

(A. Martiny) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF AGDER, POSTBOKS 422, 4604 KRISTIANSAND, NORWAY.

Email address: `andre.martiny@uia.no`