

FAMILIES OF FINITE SETS IN WHICH NO SET IS COVERED BY THE UNION OF THE OTHERS

GUILLERMO ALESANDRONI

ABSTRACT. Let \mathcal{F} be a finite nonempty family of finite nonempty sets. We prove the following:

- (1) \mathcal{F} satisfies the condition of the title if and only if for every pair of distinct subfamilies $\{A_1, \dots, A_r\}, \{B_1, \dots, B_s\}$ of \mathcal{F} , $\bigcup_{i=1}^r A_i \neq \bigcup_{i=1}^s B_i$.
- (2) If \mathcal{F} satisfies the condition of the title, then the number of subsets of $\bigcup_{A \in \mathcal{F}} A$ containing at least one set of \mathcal{F} is odd.

We give two applications of these results, one to number theory and one to commutative algebra.

1. MOTIVATION AND DEFINITION

This work is not to be confused with [2] or [3], two articles whose titles are almost identical to the title of this paper; but it is to be compared to them. In [2] and [3], Erdős, Frankl, and Füredi study a family \mathcal{F} of sets in which no set is contained in the union of r others, for various values of r . In this article, we address the particular case where r is one less than the cardinality of \mathcal{F} . In other words, we consider a family of sets, such that no set is contained in the union of the others.

In spite of these similarities, our work is different from that of Erdős, Frankl, and Füredi both in its approach and scope. While they are interested in bounds on the cardinality of \mathcal{F} , we investigate intrinsic properties of \mathcal{F} . Once our main results are proven, we give two applications: one to number theory and one to commutative algebra.

In the next definition, we give a special name to the families to be studied in the present article.

Definition 1.1. Let \mathcal{F} be a finite nonempty family of finite nonempty sets. We will say that \mathcal{F} is a **dominant family of sets**, if no set in \mathcal{F} is contained in the union of the others. Equivalently, we will say that \mathcal{F} is dominant, if every set of \mathcal{F} contains some element not shared by any of the other sets of \mathcal{F} .

The term dominant family of sets comes from its algebraic counterpart; namely, a dominant ideal [1]. This is how these concepts are related.

Each monomial ideal can be represented as a squarefree monomial ideal using a technique known as polarization [5]. Suppose that, in its polarized form, a monomial ideal M is minimally generated by squarefree monomials m_1, \dots, m_q . We say that M is a dominant ideal if, for each m_i there is a variable x_{m_i} that appears in the factorization of m_i but not in the factorizations of $m_1, \dots, \widehat{m_i}, \dots, m_q$.

Denote by $set(m_i)$ the set of variables that appear in the factorization of m_i . Then the condition above can be restated as follows: M is dominant if each set $set(m_i)$ contains a variable x_{m_i} that is not shared by any of the sets $set(m_1), \dots, set(\widehat{m_i}), \dots, set(m_q)$.

In other words, suppose that, in its polarized form, M can be represented as $M = (m_1, \dots, m_q)$. Then M is a dominant ideal if and only if $\{\text{set}(m_1), \dots, \text{set}(m_q)\}$ is a dominant family of sets.

Note: Free resolutions of monomial ideals is a core topic of commutative algebra. Perhaps, its most emblematic construction is the Taylor resolution [6], which yields a free resolution for each monomial ideal. The class of dominant ideals gives a complete characterization of when the Taylor resolution is minimal [1].

2. MAIN RESULTS

Theorem 2.1. \mathcal{F} is a dominant family of sets if and only if for every pair of distinct subfamilies $\{A_1, \dots, A_r\}$, $\{B_1, \dots, B_s\}$ of \mathcal{F} , $\bigcup_{i=1}^r A_i \neq \bigcup_{i=1}^s B_i$.

Proof. First, let us consider the case in which \mathcal{F} is dominant. If $\{A_1, \dots, A_r\}$ and $\{B_1, \dots, B_s\}$ are different subfamilies of \mathcal{F} , there must be a set C that belongs to one of the subfamilies but not to the other; say that $C \in \{A_1, \dots, A_r\}$. Since \mathcal{F} is dominant, there is an element $x \in C$, not shared by any other set of \mathcal{F} . Therefore, $x \in \bigcup_{i=1}^r A_i$ and $x \notin \bigcup_{i=1}^s B_i$, which proves the first implication.

Now, let us assume that $\bigcup_{i=1}^r A_i \neq \bigcup_{i=1}^s B_i$, for every pair of distinct subfamilies $\{A_1, \dots, A_r\}$, $\{B_1, \dots, B_s\}$ of \mathcal{F} . Let $A \in \mathcal{F}$. By hypothesis, $\bigcup_{B \in \mathcal{F} \setminus \{A\}} B \neq \bigcup_{B \in \mathcal{F}} B$. That is, $\bigcup_{B \in \mathcal{F} \setminus \{A\}} B \neq \left(\bigcup_{B \in \mathcal{F} \setminus \{A\}} B \right) \cup A$. Thus, $A \not\subseteq \bigcup_{B \in \mathcal{F} \setminus \{A\}} B$, which proves the second implication. \square

Theorem 2.2. Let $\mathcal{A} = \{A_1, \dots, A_n\}$ be a dominant family of sets, and let $A = \bigcup_{i=1}^n A_i$. Then, the number of subsets of A that contain at least one member of \mathcal{A} is odd.

Proof. Since the power set of A has even cardinality, it is enough to show that the number of subsets of A not containing any of the elements of \mathcal{A} is odd. The proof is by induction on n . If $n = 1$, then $\mathcal{A} = \{A_1\}$, $A = A_1$, and the number of subsets of A that do not contain any sets of \mathcal{A} is $2^{|A|} - 1$, which is odd. Suppose that the theorem holds for $n - 1$. Let \mathcal{B} be the family of all subsets of A not containing any elements of \mathcal{A} . Since \mathcal{A} is dominant, there exists an element $x \in A_1 \setminus \bigcup_{i=2}^n A_i$. Define $\mathcal{B}_{x^-} = \{B \in \mathcal{B} : x \notin B\}$, and $\mathcal{B}_{x^+} = \{B \in \mathcal{B} : x \in B\}$. Note that \mathcal{B} is the disjoint union of \mathcal{B}_{x^-} and \mathcal{B}_{x^+} . Consider the function

$$\begin{aligned} f &: \mathcal{B}_{x^+} \rightarrow \mathcal{B}_{x^-} \\ B &\rightarrow B \setminus \{x\}. \end{aligned}$$

It is clear that f is one-to-one. Therefore,

$$\begin{aligned} |\mathcal{B}| &= |\mathcal{B}_{x^+}| + |f(\mathcal{B}_{x^+})| + |\mathcal{B}_{x^-} \setminus f(\mathcal{B}_{x^+})| \\ &= 2|\mathcal{B}_{x^+}| + |\mathcal{B}_{x^-} \setminus f(\mathcal{B}_{x^+})|. \end{aligned}$$

Thus, in order to prove that $|\mathcal{B}|$ is odd, it is enough to prove that $|\mathcal{B}_{x^-} \setminus f(\mathcal{B}_{x^+})|$ is odd. This fact follows from two assertions that we will consider previously.

The first assertion is as follows:

$$\mathcal{B}_{x^-} \setminus f(\mathcal{B}_{x^+}) = \{B \in \mathcal{B}_{x^-} : A_1 \subseteq B \cup \{x\}\}.$$

To show this, choose a set $D \in \mathcal{B}_{x^-}$, such that $A_1 \subseteq D \cup \{x\}$. We want to prove that $D \in \mathcal{B}_{x^-} \setminus f(\mathcal{B}_{x^+})$. Suppose not. Then there is a set $C \in \mathcal{B}_{x^+}$, such that $f(C) = D$. By construction, $D = C \setminus \{x\}$. Hence, $D \cup \{x\} = C$, which implies that $A_1 \subseteq C$, a contradiction. We conclude that $\{B \in \mathcal{B}_{x^-} : A_1 \subseteq B \cup \{x\}\} \subseteq \mathcal{B}_{x^-} \setminus f(\mathcal{B}_{x^+})$. Conversely, if $D \in \mathcal{B}_{x^-} \setminus f(\mathcal{B}_{x^+})$, we must have that $D \cup \{x\} \notin \mathcal{B}_{x^+}$. Therefore, for some j , $A_j \subseteq D \cup \{x\}$, but $A_j \not\subseteq D$. Since $x \in A_1 \setminus \bigcup_{i=2}^n A_n$, it follows that $j = 1$, and $D \in \{B \in \mathcal{B}_{x^-} : A_1 \subseteq B \cup \{x\}\}$, which proves our assertion.

The second assertion is as follows: if $\mathcal{A}' = \{A_2 \setminus A_1, \dots, A_n \setminus A_1\}$, $A' = \bigcup_{i=2}^n (A_i \setminus A_1)$, and $\Lambda = \{B \setminus A_1 : B \in \mathcal{B}_{x^-} \setminus f(\mathcal{B}_{x^+})\}$, then Λ is the class of all subsets of A' not containing any of the elements of \mathcal{A}' . To prove this fact, consider a set $B \setminus A_1 \in \Lambda$. Since $B \in \mathcal{B}_{x^-} \setminus f(\mathcal{B}_{x^+})$, $A_1 \subseteq B \cup \{x\}$. For each $i = 2, \dots, n$, let $y_i \in A_i \setminus B$. Since $x \notin A_i$, $y_i \in A_i \setminus (B \cup \{x\}) \subseteq A_i \setminus A_1$. And since $y_i \notin B$, $A_i \setminus A_1 \not\subseteq B \setminus A_1$. Therefore, $B \setminus A_1$ is a subset of A' not containing any elements of \mathcal{A}' , which proves the first inclusion. To prove the other inclusion, suppose that C is a subset of A' not containing any of $A_2 \setminus A_1, \dots, A_n \setminus A_1$. This implies that $C \cup (A_1 \setminus \{x\})$ is a subset of A not containing any of A_1, \dots, A_n . Hence, $C \cup (A_1 \setminus \{x\}) \in \mathcal{B}$ and, given that $x \notin C \cup (A_1 \setminus \{x\})$, we have that $C \cup (A_1 \setminus \{x\}) \in \mathcal{B}_{x^-}$. Finally, $A_1 \subseteq [C \cup (A_1 \setminus \{x\})] \cup \{x\}$, which implies that $C \cup (A_1 \setminus \{x\}) \in \mathcal{B}_{x^-} \setminus f(\mathcal{B}_{x^+})$. Thus, $C = (C \cup (A_1 \setminus \{x\})) \setminus A_1 \in \Lambda$. This proves the second inclusion and, hence, our assertion.

Now we can complete the proof. Since \mathcal{A}' is a dominant family of sets, of cardinality $n - 1$, it follows that $|\Lambda|$ is odd, by induction hypothesis. But $|\mathcal{B}_{x^-} \setminus f(\mathcal{B}_{x^+})| = |\Lambda|$, which proves the theorem. \square

Below we sketch an alternative proof to Theorem 2.2 (which was ingeniously created by Darij Grinberg). The notation A and \mathcal{A} is as in Theorem 2.2.

Let p be the number of all pairs (X, J) with $X \subseteq A$ and $J \subseteq \mathcal{A}$ satisfying $X \supseteq \bigcup_{B \in J} B$. We shall compute p in two ways:


One way is: $p = \sum_{X \subseteq A} q_X$, where q_X is the number of all $J \subseteq \mathcal{A}$ satisfying $X \supseteq \bigcup_{B \in J} B$. It is easy to see that $q_X = 2^m$, where m is the number of all $B \in \mathcal{A}$ satisfying $X \supseteq B$. Hence, q_X is even if and only if $X \supseteq B$ for some $B \in \mathcal{A}$. Therefore, p is congruent modulo 2 to the number of all subsets $X \subseteq A$ that contain none of the sets in \mathcal{A} .

Another way is: $p = \sum_{J \subseteq \mathcal{A}} r_J$, where r_J is the number of all $X \subseteq A$ satisfying $X \supseteq \bigcup_{B \in J} B$. It is easy to see that $r_J = 2^r$, where r is the size of the set $A \setminus \bigcup_{B \in J} B$. Hence, q_X is even if and only if $A \setminus \bigcup_{B \in J} B$ is nonempty; in other words, q_X is even if and only if $J \neq \mathcal{A}$ (here we use the dominance requirement). Thus, $p \equiv 1 \pmod{2}$.

Comparing these two results, we conclude that the number of subsets of A that contain none of the sets in \mathcal{A} is $\equiv 1 \pmod{2}$. In other words, this number is odd. Since $2^{|A|}$ is even, it follows that the number of subsets of A containing at least one member of \mathcal{A} is also odd.

3. APPLICATION TO NUMBER THEORY

Suppose that a_1, a_2, a_3, a_4 are 4 squarefree numbers (i.e., each of them is the product of distinct primes). Label the vertices of a solid tetrahedron T with a_1, a_2, a_3, a_4 , as shown in the figure below.



Next, label each edge of T with the lcm of its 2 vertices (for example, the edge with vertices a_1, a_2 gets the label $\text{lcm}(a_1, a_2)$). Then, label each face of T with the lcm of its 3 vertices and, finally, label T itself with the lcm of its 4 vertices. That is, T receives the label $a = \text{lcm}(a_1, a_2, a_3, a_4)$. Each vertex, edge, and face of T , as well as T itself, will be called a multiface. That is, the term multiface encompasses all objects that have been given a label.

We will consider the following problems.

(i) State a condition C such that, all multifaces have distinct labels if and only if condition C is satisfied.

(ii) Suppose that condition C is satisfied. Let N be the set of divisors of a that are divisible by one of a_1, a_2, a_3, a_4 . Is $|N|$ even or odd?

(i) For each squarefree integer n , denote by $\text{set}(n)$ the set of prime factors of n . Note that $\text{set}(\text{lcm}(a_i, a_j)) = \text{set}(a_i) \cup \text{set}(a_j)$ and, more generally, $\text{set}(\text{lcm}(a_{i_1}, \dots, a_{i_r})) = \bigcup_{t=1}^r \text{set}(a_{i_t})$.

Let $\text{lcm}(a_{i_1}, \dots, a_{i_r}), \text{lcm}(a_{j_1}, \dots, a_{j_s})$ be the labels of two multifaces. Then

$$\begin{aligned} \text{lcm}(a_{i_1}, \dots, a_{i_r}) &\neq \text{lcm}(a_{j_1}, \dots, a_{j_s}) \\ &\text{if and only if} \\ \text{set}(\text{lcm}(a_{i_1}, \dots, a_{i_r})) &\neq \text{set}(\text{lcm}(a_{j_1}, \dots, a_{j_s})) \\ &\text{if and only if} \\ \bigcup_{t=1}^r \text{set}(a_{i_t}) &\neq \bigcup_{t=1}^s \text{set}(a_{j_t}). \end{aligned}$$

Therefore, by Theorem 2.1,

$$\begin{aligned} &\text{all multifaces have different labels} \\ &\text{if and only if} \\ &\{\text{set}(a_1), \dots, \text{set}(a_4)\} \text{ is a dominant family of sets} \\ &\text{if and only if} \end{aligned}$$

each $\text{set}(a_i)$ contains a prime p_i not shared by any of $\text{set}(a_1), \dots, \widehat{\text{set}(a_i)}, \dots, \text{set}(a_4)$
if and only if

there is a prime p_i that divides a_i , but not any of the integers $a_1, \dots, \widehat{a_i}, \dots, a_4$.

In consequence, condition C can be stated as follows.

C : for each $i = 1, \dots, 4$, there is a prime p_i that is a factor of a_i but not a factor of any of $a_1, \dots, \widehat{a_i}, \dots, a_4$.

(ii) Consider the set $N = \{n \in \mathbb{N} : n \mid a, \text{ and } a_i \mid n, \text{ for some } 1 \leq i \leq 4\}$. Note that $n \in N$ if and only if $\text{set}(a_i) \subseteq \text{set}(n) \subseteq \text{set}(\text{lcm}(a_1, \dots, a_4)) = \bigcup_{t=1}^4 \text{set}(a_t)$, for some

$1 \leq i \leq 4$. Hence, $|N|$ equals the number of subsets of $\bigcup_{t=1}^4 \text{set}(a_t)$ that contain one set of $\mathcal{F} = \{\text{set}(a_1), \dots, \text{set}(a_4)\}$. By Theorem 2.2, $|N|$ is odd.

4. APPLICATION TO COMMUTATIVE ALGEBRA

We close this paper with an application to free resolutions of monomial ideals. Even though we will use simple terms, some background material may be needed, and can be found in [4, 5].

Let $A = \{m_1, \dots, m_q\}$ be a set of monomials, and denote $m = \text{lcm}(A)$. We will say that a subfamily $\{m_{i_1}, \dots, m_{i_j}\}$ of A is a **generator of m in A** (or just a generator of m), if $\text{lcm}(m_{i_1}, \dots, m_{i_j}) = m$. In addition, a generator of m , none of whose proper subsets is a generator of m , will be called a **minimal generator of m in A** .

Consider the class $\mathcal{F} = \{A_1, \dots, A_r\}$ of all minimal generators of m in A , and suppose that $A = \bigcup_{i=1}^r A_i$. Notice that every generator of m must be a subset of A that contains one of A_1, \dots, A_r . By Theorem 2.2, if \mathcal{F} is a dominant family of sets, the number of generators of m is odd.

We will now explain how this result fits in the study of monomial resolutions. One important question in this topic is: if m is a multidegree of the Taylor resolution of a monomial ideal M , does the minimal resolution of M contain basis elements of multidegree m ?

It is known that the minimal resolution of M can be obtained from its Taylor resolution by doing a series of computations known as consecutive cancellations [5], each of which eliminates two basis elements of equal multidegree. This amounts to saying that the number of basis elements of multidegree m in Taylor's resolution, and the number of basis elements of multidegree m in the minimal resolution of M differ by an even number. Therefore, if the number of basis elements of multidegree m in the Taylor resolution is odd, then the minimal resolution of M will contain at least one element of multidegree m .

And how can we tell if the number of basis elements of multidegree m in the Taylor resolution is odd? Informally speaking, these basis elements are precisely the sets that we call generators of m in A , where A is the set of all minimal generators of M that divide m [4]. (Note that the minimal generators of M are monomials, while the minimal generators of m are sets of monomials.) It is here where Theorem 2.2 becomes a useful tool.

Summarizing, suppose that m is a multidegree that occurs in the Taylor resolution of a monomial ideal M . Let A be the class of all minimal generators of M that divide m . Suppose that $A = \bigcup_{i=1}^r A_i$, where $\mathcal{F} = \{A_1, \dots, A_r\}$ is the class of all minimal generators of m . If \mathcal{F} is a dominant family of sets, then the number of basis elements of multidegree m in the Taylor resolution of M is odd, and hence, the minimal resolution of M contains at least one basis element of multidegree m .

For example, consider the ideal $M = (x_1x_2, x_3x_4, x_5x_6, x_1x_3x_5, x_2x_4x_6, x_7)$, and let $m = x_1x_2x_3x_4x_5x_6x_7$, which is a multidegree of the Taylor resolution of M . Then, $A = \{x_1x_2, x_3x_4, x_5x_6, x_1x_3x_5, x_2x_4x_6, x_7\}$ (and $m = \text{lcm}(A)$). It is not difficult to see that $A_1 = \{x_1x_2, x_3x_4, x_5x_6, x_7\}$ and $A_2 = \{x_1x_3x_5, x_2x_4x_6, x_7\}$ are the only minimal generators of m in A . Since $A = A_1 \cup A_2$, and given that $\mathcal{F} = \{A_1, A_2\}$ is a dominant family of sets, we conclude that the minimal resolution of M contains at least one basis element of multidegree m .

Acknowledgements: A few days after I posted the first version of this paper to the arxiv, Darij Grinberg emailed me an elegant alternative proof to Theorem 2.2, which is included in this new version. Thanks, Darij! My wife Danisa always types and proofreads my papers, which I deeply appreciate. Thanks, sweetheart!

REFERENCES

- [1] G. Alesandroni, *Minimal resolutions of dominant and semidominant ideals*, J. Pure Appl. Algebra **221** (2017), no. 4, 780-798.
- [2] P. Erdős, P. Frankl, and Z. Füredi, *Families of finite sets in which no set is covered by the union of two others*, J. Comb. Theory, Ser. A **33** (1982), 158-166.
- [3] P. Erdős, P. Frankl, and Z. Füredi, *Families of finite sets in which no set is covered by the union of r others*, Israel J. Math. **51** (1985), 75-89.
- [4] J. Mermin, *Three simplicial resolutions*, *Progress in Commutative Algebra 1*, Edited by Francisco, Christopher / Klingler, Lee C. / Sather-Wagstaff, Sean / Vassilev, Janet C. De Gruyter (2012).
- [5] I. Peeva, *Graded Syzygies*, Algebra and Applications, vol. 14, Springer, London (2010).
- [6] D. Taylor, *Ideals generated by monomials in an R -sequence*, Ph.D. thesis, University of Chicago (1960).

2000 ROSARIO, SANTA FE, ARGENTINA

Email address: guillea@okstate.edu, alesangc@wfu.edu, alesandronig@yahoo.com