

BOUNDING THE SPECTRAL RADIUS AND POWERS OF NONNEGATIVE MATRICES

VUONG BUI

ABSTRACT. Given a nonnegative square matrix A , the n -th root of the largest entry of A^n is well known to converge to the spectral radius $\rho(A)$ of A when n tends to infinity. We give another proof of the convergence by showing

$$\lim_{n \rightarrow \infty} \max_i \max_j \sqrt[n]{(A^n)_{i,j}} = \max_i \sup_n \sqrt[n]{(A^n)_{i,i}}.$$

As the sequence $\{(A^n)_{i,i}\}_n$ for every i is shown to be both supermultiplicative and submultiplicative in a weak form, an extension of Fekete's lemma concludes that the subsequence of positive $\sqrt[n]{(A^n)_{i,i}}$, if not empty, converges to $\sup_n \sqrt[n]{(A^n)_{i,i}} = \inf\{\sqrt[n]{K_i(A^n)_{i,i}} : (A^n)_{i,i} > 0\}$ for some constant K_i . A corollary is the following bound of $\rho(A)$:

$$\max_i \sqrt[m_i]{(A^{m_i})_{i,i}} \leq \rho(A) \leq \max_i \sqrt[m_i]{K_i(A^{m_i})_{i,i}},$$

where m_i satisfies $(A^{m_i})_{i,i} > 0$ for each i , or we just set $m_i = 1$ if there is no such m_i . An unoptimized setting for the constants is $K_i = D^3(\frac{U}{V})^{6D^2}$ for every i , where $D \times D$ is the dimension of A , and U, V are respectively the largest and smallest entry of A .

The order of $(A^n)_{i,i}$ also gives another proof of the formula $\lambda = \max_i \sup_n \sqrt[n]{(A^n)_{i,i}}$ for the limit using Wimmer's formula. Yet another proof is however also given. Although the third one is much more complex, it gives a more precise growth: There exists some nonnegative integer r such that for every n ,

$$\text{const } n^r \lambda^n \leq \max_i \max_j (A^n)_{i,j} \leq \text{const } n^r \lambda^n.$$

The approach in use is more combinatorial than the traditional algebraic approach.

1. INTRODUCTION

Given a *nonnegative* square matrix A (i.e. all entries are nonnegative), it is well known by Gelfand's formula that the largest entry in A^n grows exponentially as n tends to infinity. The n -th root of this largest value converges to the spectral radius $\rho(A)$, which is the largest absolute value of an eigenvalue of A . In fact, Gelfand's formula is more general as the matrix is not required to be nonnegative and the norm is not necessarily the maximum norm. The popular proof, which is algebraic in its nature, involves eigenvalues and eigenvectors.

In this text, we provide another approach to the convergence of the n -th root of the largest entry of A^n . Note that the maximum norm is chosen throughout the work for convenience without the loss of generality in the asymptotic statements since any two arbitrary norms are within constant factors of each other. Instead of using the machinery of eigenvalue/eigenvector, we show that:

2010 *Mathematics Subject Classification.* 15A18, 15A60.

Key words and phrases. Nonnegative matrix, Matrix norm, Spectral radius.

The author is supported by the Deutsche Forschungsgemeinschaft (DFG) Graduiertenkolleg "Facets of Complexity" (GRK 2434).

Theorem 1. *Given a nonnegative matrix A , the following limit λ exists and can be expressed as:*

$$\lambda := \lim_{n \rightarrow \infty} \max_i \max_j \sqrt[n]{(A^n)_{i,j}} = \max_i \sup_n \sqrt[n]{(A^n)_{i,i}},$$

where $(A^n)_{i,j}$ is the (i, j) -th entry of the matrix power A^n ¹.

Although Gelfand's formula allows A to be any complex matrix, the requirement that A is nonnegative in our formula is essential. Even after relaxing the statement with absolute values $\lim_{n \rightarrow \infty} \max_i \max_j \sqrt[n]{|(A^n)_{i,j}|}$ and $\max_i \sup_n \sqrt[n]{|(A^n)_{i,i}|}$, the formula still does not hold for the following matrix A , where the former value is 1 while the latter value is at least 2:

$$(1) \quad A = \begin{bmatrix} 0 & -1 \\ 1 & 2 \end{bmatrix}, \quad A^n = \begin{bmatrix} -(n-1) & -n \\ n & n+1 \end{bmatrix}.$$

The limit λ here is identical to the spectral radius $\rho(A)$, and this relation allows us to estimate $\rho(A)$ by estimating λ . In fact, we provide an interval bounding λ using the diagonals of some matrices as in Equation (3) (in Page 3). We give below the method and state the supporting lemmas/theorems, which will be verified in Section 4. From now on, we will not use any property of the spectral radius $\rho(A)$ but only the equivalent limit $\lambda = \max_i \sup_n \sqrt[n]{(A^n)_{i,i}}$. However, the two notations $\rho(A)$ and λ will be both used.

The method is to show some weak forms of submultiplicativity and supermultiplicativity for the sequence of the entries on the diagonal, and later apply the approach of Fekete's lemma [1]. However, we first need to give an extended version of the original Fekete's lemma in the way that the extended version does not require the sequence to be positive.

Lemma 1 (An extension of Fekete's lemma for nonnegative supermultiplicative sequences). *Given a nonnegative sequence $\{a_n\}_n$ such that for every ℓ, m , we have $a_{\ell+m} \geq a_\ell a_m$. Then the subsequence $\{\sqrt[n]{a_n} : a_n > 0\}$ is either empty or has a limit² that is equal to $\sup_n \{\sqrt[n]{a_n}\}$.*

The sequence $\{(A^n)_{i,i}\}_n$ for every i satisfies the condition of the lemma since $(A^{\ell+m})_{i,i} \geq (A^\ell)_{i,i} (A^m)_{i,i}$ for every ℓ, m . However, the whole sequence $\sqrt[n]{(A^n)_{i,i}}$ may not converge, e.g. $(A^n)_{i,i} = 1$ for even n and $(A^n)_{i,i} = 0$ for odd n .

We have the following dual form of Lemma 1.

Lemma 2 (An extension of Fekete's lemma for nonnegative *almost* submultiplicative sequences). *Given a nonnegative sequence $\{a_n\}_n$ such that for every ℓ, m , either $a_\ell a_m = 0$ or $0 < a_{\ell+m} \leq a_\ell a_m$. Then the subsequence $\{\sqrt[n]{a_n} : a_n > 0\}$ is either empty or has a limit that is equal to $\inf \{\sqrt[n]{a_n} : a_n > 0\}$.*

It should be noted that given a nonnegative sequence $\{a_n\}_n$, if this sequence is submultiplicative (in contrast to supermultiplicativity in Lemma 1), then the convergence of $\sqrt[n]{a_n}$ is always ensured. This condition is however stronger than the condition in Lemma 2.

Lemma 2 can be proved using the same argument as Lemma 1 (see the proof in Section 4), or one can obtain Lemma 2 by applying Lemma 1 to the sequence $\{b_n\}_n$ where $b_n = 0$ if $a_n = 0$ and $b_n = 1/a_n$ otherwise³.

¹Throughout the work we never consider zeroth power of a matrix and assume it undefined.

²We permit the limit ∞ here.

³If the limit of $\{\sqrt[n]{b_n} : b_n > 0\}$ is ∞ , the limit of $\{\sqrt[n]{a_n} : a_n > 0\}$ is 0.

The sequence $\{(A^n)_{i,i}\}_n$ is always supermultiplicative but it is in general not submultiplicative even in the sense of Lemma 2. However, the following slightly modified form is always guaranteed.

Theorem 2 (Weak submultiplicativity). *Given a nonnegative matrix A . For each index i , there is a constant K_i such that for every ℓ, m , either $(A^\ell)_{i,i}(A^m)_{i,i} = 0$, or $0 < (A^{\ell+m})_{i,i} \leq K_i(A^\ell)_{i,i}(A^m)_{i,i}$. The constant K_i for every i can be set to $D^3(\frac{U}{V})^{6D^2}$ where $D \times D$ is the dimension of A , and U, V are respectively the largest and smallest entry of A .*

By Theorem 2, the sequence $\{K_i(A^n)_{i,i}\}_n$ is submultiplicative in the sense of Lemma 2.

If the sequence $\{(A^n)_{i,i}\}_n$ for an i is not a sequence of zeros, we obtain two expressions for the same following limit:

$$\theta_i := \lim\{\sqrt[n]{(A^n)_{i,i}} : (A^n)_{i,i} > 0\} = \sup_n \sqrt[n]{(A^n)_{i,i}} = \inf\{\sqrt[n]{K_i(A^n)_{i,i}} : (A^n)_{i,i} > 0\}.$$

For convenience, when $\{(A^n)_{i,i}\}_n$ is a sequence of zeros, we set $\theta_i = 0$.

The two expressions imply that for any $(A^n)_{i,i} > 0$, we have

$$(2) \quad \sqrt[n]{(A^n)_{i,i}} \leq \theta_i \leq \sqrt[n]{K_i(A^n)_{i,i}}.$$

Since $\lambda = \max_i \theta_i$, we can use Equation (2) to bound λ as

$$(3) \quad \max_i \sqrt[m_i]{(A^{m_i})_{i,i}} \leq \lambda \leq \max_i \sqrt[m_i]{K_i(A^{m_i})_{i,i}},$$

where the number m_i satisfies $(A^{m_i})_{i,i} > 0$, or $m_i = 1$ if there no such m_i .

The readers may relate this bound to the bound in Gershgorin circle theorem [2] for a complex matrix A . Each eigenvalue in such a setting is bounded by a disc whose center is some element $A_{i,i}$ on the diagonal and the radius is the sum of the absolute values of the nondiagonal entries of the i -th row. Our bound follows somewhat a similar fashion, but we just need to look at the diagonal and it seems that involving higher powers allows us a narrower gap between the upper and lower bounds.

Actually there is a known upper bound of the spectral radius $\rho(A) \leq \|A^n\|^{\frac{1}{n}}$ that is almost as good as our bound. This works for any matrix and any consistent matrix norm. However, it is slightly worse since the order of $\|A^n\|$ can be polynomial times $(\rho(A))^n$ (as in Theorem 8) while the entries on the diagonal are ensured to be of the order at most $(\rho(A))^n$ (by Corollary 1). The more important contribution is that our upper bound has a corresponding lower bound.

Another popular result that gives a lower bound is Collatz-Wielandt's formula for the spectral radius of a nonnegative matrix A :

$$\max_{x: x \geq 0, x \neq 0} \min_{i: x_i \neq 0} \frac{(Ax)_i}{x_i}.$$

Plugging any x into the formula, we obtain a lower bound of the spectral radius. But how to choose a good x is still not obvious.

Another bound in literature that bounds $\rho(A)$ of a nonnegative matrix A from both above and below is:

$$\min_i \sum_j A_{i,j} \leq \rho(A) \leq \max_i \sum_j A_{i,j}.$$

In general, the gap between these bounds can be large. Following the idea of raising to higher powers as in the bound (3) and the fact $\rho(A^n) = (\rho(A))^n$, one may ask how good is the following bound: For any n ,

$$(4) \quad \sqrt[n]{\min_i \sum_j (A^n)_{i,j}} \leq \rho(A) \leq \sqrt[n]{\max_i \sum_j (A^n)_{i,j}}.$$

We will analyse this bound later after we present the necessary tools. In short, this bound is in general worse, but a modification can make it as good as the bound in (3).

In practice, power iteration (Von Mises's iteration [3]) would be used to provide a good estimation of both spectral radius and the corresponding eigenvector. Starting with a vector x_0 , the next vector at each step is given by $x_{n+1} = (Ax_n)/\|x_n\|$. As the vector x_n converges to the eigenvector, the spectral radius can be estimated accordingly. Although the convergence is in general fast, there is no obvious way to tell how close the actual $\rho(A)$ is to the estimation as in our method.

In both Equation (2) and Equation (3), it still remains to show a way to check if such an m_i exists, and if it exists, how to find an m_i greater than some number N . In practice, we need N large enough for a good bound.

In order to do that, we introduce the following notion of *dependency graph*⁴. The dependency graph is a directed graph whose vertices are the dimensions $1, \dots, D$ and there is a directed edge from i to j if and only if $A_{i,j}$ is nonzero. This directed graph can be partitioned into strongly connected components, which we will call *components* for short. In this paper, acyclic components will be called *single components*. Note that an acyclic component necessarily contains only one vertex but the component containing only one vertex may not be acyclic (e.g. a loop around the only vertex). There exists m_i such that $(A^{m_i})_{i,i} > 0$ if and only if i is in a nonsingle component. This is also equivalent to $\theta_i > 0$.

It remains to show how to choose such an $m_i > N$. Let the distance from i to i in the dependency graph be ℓ_i , we then have $(A^m)_{i,i} > 0$ for every multiple m of ℓ_i . Then for every i , we set $m_i = N + (\ell_i - r)$, where $N \equiv r \pmod{\ell_i}$, i.e. m_i is the least integer greater than N divided by ℓ_i . Note that the difference between N and every m_i is bounded.

Suppose in order to have a good bound, we need the ratio of the upper bound and the lower bound at most $1 + \epsilon$ for some $\epsilon > 0$. This can be obtained if for every i , we have $\sqrt[m_i]{K_i} \leq 1 + \epsilon$. At first, N is chosen so that $N \geq \log_{1+\epsilon} K_i$ for every i . If we allow ourselves to stick to the simple method of complexity D^3 for the product of two $D \times D$ matrices, the complexity for computing A^m is $(D^3)^{\log_2 m}$. Therefore, the computation of the matrices A^{m_i} is of order

$$(D^3)^{\log_2(\log_{1+\epsilon}(D^3(\frac{U}{V})^{6D^2}))} = D^{3 \log_2(6D^2 \log_{1+\epsilon}(U/V) + 3 \log_{1+\epsilon} D)}.$$

In Theorem 2, the constant $D^3(\frac{U}{V})^{6D^2}$ is not optimized and there seems to be further room to reduce this constant. However, the factor U/V , which can be arbitrarily large, is essential. The same applies in Equations (2), (3). For example, consider the following matrix for a small $\epsilon > 0$,

$$A = \begin{bmatrix} \epsilon & 1 \\ 1 & \epsilon \end{bmatrix}.$$

⁴This definition, also known as the adjacency matrix, can be related to the dependency graph in [4].

In the context of Theorem 2, we have $(A^2)_{i,i} = 1 + \epsilon^2 > 1 = \frac{1}{\epsilon^2}(A^1)_{i,i}(A^1)_{i,i} = (U/V)^2(A^1)_{i,i}(A^1)_{i,i}$ for $i = 1, 2$. Since $(A^2)_{1,1} = (A^2)_{2,2} > 1$, it follows that $\theta_1 > 1, \theta_2 > 1, \lambda > 1$. In the context of Equations (2), (3), we have $(A^1)_{i,i} < \epsilon\theta_i = \frac{1}{U/V}\theta_i^1$ and $(A^1)_{i,i} < \epsilon\lambda = \frac{1}{U/V}\lambda^1$ for $i = 1, 2$.

We would add that a method of estimating the spectral radius of a nonnegative matrix is followed by an estimation of the largest absolute value of the solutions (which is also a solution) of

$$c_0 + c_1x + c_2x^2 + \dots + c_{n-1}x^{n-1} = x^n,$$

where all c_i are nonnegative.

It is due to a well known fact that the solutions are the eigenvalues of

$$A = \begin{bmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ 0 & \ddots & \ddots & \ddots & \vdots \\ \vdots & \vdots & \ddots & 0 & 1 \\ c_0 & c_1 & \dots & c_{n-2} & c_{n-1} \end{bmatrix}.$$

The readers may check that given a desired precision of the approximation, our method has the same complexity as the traditional bisection method. However, while the latter requires comparisons at every step to decide on which smaller intervals to take, our method has a somewhat more “closed-form” solution.

Throughout the work, a more important quantity than the distant ℓ_i from i to i for an i not in a single component is

$$d_i = \gcd\{n : (A^n)_{i,i} > 0\}.$$

A basic property of d_i is Lemma 4, which states that: For every n large enough divided by d_i , we have $(A^n)_{i,i} > 0$. Note that d_i is undefined for the vertices i in single components.

Let us first use this quantity to relate the previously shown work to a formula by H. K. Wimmer ([5]): Given any *complex* matrix A , we have

$$\rho(A) = \limsup_{n \rightarrow \infty} \sqrt[n]{|\operatorname{tr}(A^n)|}.$$

We show that our results deduce the following more specific version of Wimmer’s result for nonnegative matrices.

Theorem 3 (A more specific version of Wimmer’s result for nonnegative matrices). *Given a nonnegative matrix A , there exists d such that*

$$\rho(A) = \lim_{n \rightarrow \infty} \sqrt[d n]{\operatorname{tr}(A^{dn})}.$$

It is rather easy to verify Theorem 3 since for $d = d_j$ for a maximal $\theta_j = \rho(A)$ and a large enough m divided by d , Equation (2) gives

$$\frac{1}{K_j}(\theta_j)^m \leq (A^m)_{j,j} \leq \sum_i (A^m)_{i,i} \leq \sum_i (\theta_i)^m \leq D(\theta_j)^m.$$

If d is the minimal d_j over all j satisfying $\theta_j = \rho(A)$, then d is the smallest number for Theorem 3 to hold.

Note that the above manipulation also gives the original Wimmer's formula for nonnegative matrices, since for every n and with the same j , we still have

$$\operatorname{tr}(A^n) = \sum_i (A^n)_{i,i} \leq \sum_i (\theta_i)^n \leq D(\theta_j)^n.$$

On the other hand, one can deduce Theorem 1 from the original form of Wimmer's theorem. Indeed, since $\operatorname{const}(\theta_j)^m \leq \max_i (A^m)_{i,i}$ for a maximal θ_j and every large enough m divided by d_j , and $\max_i (A^n)_{i,i} \leq \operatorname{const}(\theta_j)^n$ for the same j and every n , we have $\max_i \theta_i = \limsup_{n \rightarrow \infty} \sqrt[n]{\max_i (A^n)_{i,i}}$, which is followed by

$$\begin{aligned} \max_i \sup_n \sqrt[n]{(A^n)_{i,i}} &= \max_i \theta_i = \limsup_{n \rightarrow \infty} \sqrt[n]{\max_i (A^n)_{i,i}} \leq \limsup_{n \rightarrow \infty} \sqrt[n]{\sum_i (A^n)_{i,i}} \\ &\leq \limsup_{n \rightarrow \infty} \sqrt[n]{D \max_i (A^n)_{i,i}} = \limsup_{n \rightarrow \infty} \sqrt[n]{\max_i (A^n)_{i,i}}, \end{aligned}$$

which implies

$$\max_i \sup_n \sqrt[n]{(A^n)_{i,i}} = \limsup_{n \rightarrow \infty} \sqrt[n]{\sum_i (A^n)_{i,i}} = \rho(A) = \lim_{n \rightarrow \infty} \max_i \max_j \sqrt[n]{(A^n)_{i,j}}.$$

The readers may notice that Wimmer's result and Theorem 1 look somewhat similar, and the above analysis shown that one can be deduced from the other by using the order of $(A^n)_{i,i}$. However, the ideas behind them are different: the proof of Wimmer's result relies on algebraic statements, e.g. $\operatorname{tr}(A)$ is the sum of eigenvalues; the proof of our result is however somewhat more combinatorial as one may see in Section 2. In fact, the author was not aware of Wimmer's formula at first and Theorem 1 in this paper was inspired by Theorem 1 in [4] instead.

Since we have shown the equivalence between the two results, one may say that we have come up with a second proof of Theorem 1. A third proof, which is much more complex, gives a more precise order $n^r \lambda^n$ of the maximal entries of A^n , as in Theorem 8. The development will be described after Theorem 4, where we close the discussion of the theorems presented so far with a characterization result. A compact version of the third proof is also given in Section 6, which only concludes the same as Theorem 1 but is much shorter.

As the limit of the subsequence of positive $\sqrt[n]{(A^n)_{i,i}}$ is $\sup_n \sqrt[n]{(A^n)_{i,i}}$ for an i not in a single component, the following result characterizes the cases where the supremum is attained in terms of the dependency graph.

Theorem 4. *Given a nonnegative matrix A . For any index i , there exists m such that $\sqrt[m]{(A^m)_{i,i}} = \sup_n \sqrt[n]{(A^n)_{i,i}}$ if and only if in the dependency graph either there is no circuit from i to i , or every circuit from i to i not visiting i in between ⁵ has the same length.*

One can see from the proof of Theorem 4 that in such a situation, the sequence $\{(A^n)_{i,i}\}_n$ is both supermultiplicative and almost submultiplicative, i.e. K_i can be set to 1.

We begin the study of the order of the maximal entry of A^n by a fact deduced from Equation (2): If $(A^n)_{i,i} > 0$, then

$$\operatorname{const}(\theta_i)^n \leq (A^n)_{i,i} \leq (\theta_i)^n.$$

⁵Note that this term is not identical to *cycle* but rather a more general definition.

Therefore, we have a better description than a mere exponential growth for the subsequence of positive $(A^n)_{i,i}$. On the way to the investigation of the order $\max_i \max_j (A^n)_{i,j}$, we also cover the order of $(A^n)_{i,j}$ for a fixed pair i, j .

Let us first remind the condensation of a directed graph for later usage, which is the directed graph whose vertices are the strongly connected components, and there is a directed edge from component C_1 to component C_2 for $C_1 \neq C_2$ if there is an edge in the original graph from i_1 to i_2 with i_1 in C_1 and i_2 in C_2 . We can actually define a partial order between the components: For different C_1, C_2 , we say $C_1 > C_2$ if there is a path from C_1 to C_2 in the condensation.

The proof of Theorem 2 relies on a lemma (Lemma 6 in Section 4) which states that: Given i , if m, n are two integers whose difference is bounded, then either $(A^n)_{i,i}$ is zero, or the ratio $(A^m)_{i,i}/(A^n)_{i,i}$ is also bounded. (In Lemma 6, we even give an explicit bound.)

We generalize the lemma in the following form.

Theorem 5. *Given a nonnegative matrix A and i, j, i', j' (not necessarily different) of the same component. If m, n are two integers whose difference is bounded, then either $(A^n)_{i',j'}$ is zero, or $(A^m)_{i,j}/(A^n)_{i',j'}$ is also bounded.*

The above theorem is immediately followed by the following corollary.

Corollary 1. *For any two i, j of the same component, we have $\theta_i = \theta_j$, and if $(A^n)_{i,j} > 0$ then*

$$\text{const}(\theta_i)^n \leq (A^n)_{i,j} \leq \text{const}(\theta_i)^n.$$

Since all θ_k are equal for the vertices k in the same component C , we allow ourselves to denote $\theta_C := \theta_k$ for any k in C .

We can now analyze the bound in Equation (4). Suppose $\theta_{C_0} = \min_C \theta_C$ and C_0 is a minimal component, then $\min_i \sum_j (A^n)_{i,j}$ is of order $(\theta_{C_0})^n$. If $\theta_{C^*} = \max_C \theta_C > \theta_{C_0}$, then $\max_i \sum_j (A^n)_{i,j}$ is of order at least $(\theta_{C^*})^n$. That is the gap between the lower bound and the upper bound does not decrease when we take higher powers of A .

However, as $\theta_C = \lim_{n \rightarrow \infty} \max_{i,j \in C} (A^n)_{i,j}$, considering the matrix A_C induced from A by the dimensions in C , we see that $\min_i \sum_j (A_C^n)_{i,j}$ and $\max_i \sum_j (A_C^n)_{i,j}$ are of the same order $(\theta_C)^n$ due to Corollary 1 (note that every row of A_C^n has a positive entry). Hence the bound of θ_C for each A_C is effective, and we combine them to obtain

$$\max_C \sqrt[n]{\min_{i \in C} \sum_{j \in C} (A_C^n)_{i,j}} \leq \rho(A) \leq \max_C \sqrt[n]{\max_{i \in C} \sum_{j \in C} (A_C^n)_{i,j}}.$$

The ratio of the lower bound and the upper bound converges to 1 when n tends to ∞ but not as obviously as the bound in Equation (3).

Note that the result of Corollary 1 does not always apply for the matrices that are not nonnegative. For example, the matrix A in Equation (1) has $\|A^n\|$ of order n while the dependency graph is connected.

Moreover, Corollary 1 is not necessarily true when the scope is not limited to the same component. In fact, there exists a matrix A and a pair of indices i, j such that there are infinity many pairs m, n such that their difference $\Delta = m - n$ is fixed, but $(A^m)_{i,j}/(A^n)_{i,j}$ grows exponentially. Consider the graph in Figure 1. Let A be the matrix with $A_{i,j} = 1$ if there is an edge from i to j and $A_{i,j} = 0$ otherwise for every i, j . Then $(A^n)_{i,j}$ is the number




FIGURE 1. A counterexample for the intercomponent case

of walks from i to j of length n . Consider the pair of vertices s, t as in the figure. Let $m = 4\ell$ and $n = 4\ell + 2$, one can see that $(A^m)_{s,t} = 1$ while $(A^n)_{s,t} = 2^\ell + 1$, since the walk of length m must go through the component on the left while a walk of length n can go through the component on the right with more choices in branching there.

However, under the following condition, we obtain an analogue to Corollary 1.

Theorem 6. *Given a nonnegative matrix A and two indices i, j . Suppose there is precisely one path from the component of i to the component of j in the condensation. If $(A^n)_{i,j} > 0$, then*

$$\text{const } n^r \theta^n \leq (A^n)_{i,j} \leq \text{const } n^r \theta^n,$$

where $\theta = \max_C \theta_C$ over all the components C on the path and the degree r is one less than the number of components C on the path such that $\theta_C = \theta$.

In the general setting, we have the following situation.

Theorem 7. *Given a nonnegative matrix A and two indices i, j . For every n we have*

$$(A^n)_{i,j} \leq \text{const } n^r \theta^n,$$

where $n^r \theta^n$ is the maximal order corresponding to a path as in Theorem 6 over all the paths from the component of i to the component of j (if there is no such path, we set $\theta = 0$). Moreover, there is an infinite sequence $\{n_t\}_t$ such that $(A^{n_t})_{i,j} \geq \text{const } n_t^r \theta^{n_t}$ for every t and the difference between two consecutive n_t is the same.

When $(A^n)_{i,j}$ is considered not only for a specific pair i, j , Theorem 7 gives the following two-sided inequality.

Theorem 8. *Given a nonnegative matrix A , there exists θ and r such that for every n ,*

$$\text{const } n^r \theta^n \leq \max_i \max_j (A^n)_{i,j} \leq \text{const } n^r \theta^n.$$

The order $n^r \theta^n$ is the maximal order in Theorem 7 over all pairs i, j (or in other words, the maximal order in Theorem 6 over all the paths in the condensation). That is to say θ is the maximal θ_C over all the components C and r is one less than the maximal number of components C in any path with $\theta_C = \theta$. An obvious equivalence is $\theta = \lambda = \rho(A)$, each one is just defined in a different way.

Note that in the latter results, we do not give an explicit constant as in Theorem 2. The readers may apply the same method as in the proof of Theorem 2 to obtain some constants.

However, the expressions may be complex, and they are not as important as the constant used for bounding the spectral radius.

Since Theorem 1 is actually a corollary of this result, we have given a third proof of Theorem 1. This third proof is much more complex than the first two proofs, but it describes more exactly the growth of the maximum entry of A^n . For example, when $\lambda = 1$, it ensures that it is of order of n^r while the other proofs do not. Since the third proof is significantly longer than the first proof, we give in Section 6 a compact version of the third proof that just proves the convergence to λ but not shows the polynomial factor.

Note that our bounds are best possible in the sense that a result of order $Kn^r\lambda^n$ for a fixed constant K is not guaranteed, e.g. consider the following periodic matrix A :

$$A = \begin{bmatrix} 0 & 1 \\ 2 & 0 \end{bmatrix}, A^{2k} = \begin{bmatrix} 2^k & 0 \\ 0 & 2^k \end{bmatrix} = \begin{bmatrix} \sqrt{2}^{2k} & 0 \\ 0 & \sqrt{2}^{2k} \end{bmatrix}, A^{2k+1} = \begin{bmatrix} 0 & 2^k \\ 2^{k+1} & 0 \end{bmatrix} = \begin{bmatrix} 0 & \frac{1}{\sqrt{2}}\sqrt{2}^{2k+1} \\ \sqrt{2}\sqrt{2}^{2k+1} & 0 \end{bmatrix}.$$

Before proving the lemmas/theorems in the following sections, let us address some remaining questions for complex matrices, which are not necessarily nonnegative:

- Is there a formula like the one in Theorem 1 for a complex matrix?
- Does the bound of the spectral radius in Equation (3) with some different constants still hold for a complex matrix?
- Does the specific form as in Theorem 3 also apply for a complex matrix?
- What is the order of $\|A^n\|$ for a complex matrix?

What we have known so far is only that $\|A^n\| \geq \text{const } \lambda^n$ due to the bound $\rho(A) \leq \sqrt[n]{\|A^n\|}$ for any n and a consistent matrix norm. An upper bound $\|A^n\| \leq \text{const } n^r \lambda^n$ for some degree r would be expected but we do not have any idea how to proceed. We would note that the behaviour of complex matrices that are not nonnegative is rather different. The example in Equation (1) has $\|A^n\|$ of order $n\lambda^n$ even with a connected dependency graph. On the other hands, a chain of components of the same maximal θ_C does not necessarily mean we have a higher degree than $\text{const } \lambda^n$. For example, consider the following case with a chain of two components

$$A = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix}, A^{2k} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, A^{2k+1} = A.$$

That being said the order of $\|A^n\|$ for a general matrix, if any, must be described in a quite different way.

2. PROOF OF THEOREM 1

In order to prove

$$\lambda := \lim_{n \rightarrow \infty} \max_i \max_j \sqrt[n]{(A^n)_{i,j}} = \max_i \sup_n \sqrt[n]{(A^n)_{i,i}},$$

we will prove ⁶ a stronger statement that for each i , there exists k such that

$$\lambda_i := \lim_{n \rightarrow \infty} \max_j \sqrt[n]{(A^n)_{i,j}} = \sup_n \sqrt[n]{(A^n)_{k,k}}.$$

If the latter limits λ_i exist, the former limit λ also exists with $\lambda = \max_i \lambda_i$. It remains to prove the existence of the latter limits λ_i .

⁶The approach we take is very similar to the one in the proof of Theorem 1 in [4].

We assume that every row i has a positive entry, since if there is any row i of all zeros, we can safely remove row i and column i from the matrix as $(A^n)_{i,j} = 0$ for every n and j . The assumption means for every i, n , there exists j such that $(A^n)_{i,j} > 0$, since every vertex in the dependency graph has then an outward edge.

Denote $\underline{\lambda}_i = \liminf_{n \rightarrow \infty} \max_j \sqrt[n]{(A^n)_{i,j}}$ and $\bar{\lambda}_i = \limsup_{n \rightarrow \infty} \max_j \sqrt[n]{(A^n)_{i,j}}$.

We give the following lemma, which is a mere observation whose proof can be easily verified and left to the readers ⁷.

Lemma 3. *If i and j are both in the same component, then*

$$\underline{\lambda}_i = \underline{\lambda}_j, \bar{\lambda}_i = \bar{\lambda}_j.$$

If $i \in C_1, j \in C_2$ and $C_1 > C_2$, then

$$\underline{\lambda}_i \geq \underline{\lambda}_j, \bar{\lambda}_i \geq \bar{\lambda}_j.$$

For a dimension i , we consider two cases of the component containing i . (The second case is quite a corollary of the first case, whose treatment is rather long.)

Case 1: The vertex i is in a component C_0 such that for every k in a component smaller than C_0 , we have $\bar{\lambda}_k < \bar{\lambda}_i$.

At first, we show that

$$\liminf_{n \rightarrow \infty} \max_j \sqrt[n]{(A^n)_{i,j}} \geq \sup_n \sqrt[n]{(A^n)_{i,i}}.$$

Indeed, fix n , for $m = np + q$ ($1 \leq q \leq n$), we have $(A^{np+q})_{i,i} \geq [(A^n)_{i,i}]^p$. There exists j such that $(A^q)_{i,j}$ is positive, which is also bounded as q is bounded. The bound of the limit inferior follows since $(A^{np+q})_{i,j} \geq (A^{np})_{i,i}(A^q)_{i,j}$.

It remains to prove

$$\limsup_{n \rightarrow \infty} \max_j \sqrt[n]{(A^n)_{i,j}} \leq \sup_n \sqrt[n]{(A^n)_{i,i}}.$$

Denote ⁸ $\theta_i = \sup_n \sqrt[n]{(A^n)_{i,i}}$. We prove the above inequality by contradiction. Assume the contrary $\bar{\lambda}_i > \theta_i$.

By the definition of $\bar{\lambda}_i$,

$$\forall \epsilon \exists n(i, \epsilon) \forall n > n(i, \epsilon), \max_j \sqrt[n]{(A^n)_{i,j}} < \bar{\lambda}_i + \epsilon,$$

and

$$\forall \epsilon \forall N \exists n > N, \max_j \sqrt[n]{(A^n)_{i,j}} > \bar{\lambda}_i - \epsilon.$$

Fix some ϵ , let $N = \max_k 2(n(k, \epsilon) + 1)$. We choose n such that $n > N$ and $\max_j \sqrt[n]{(A^n)_{i,j}} > \bar{\lambda}_i - \epsilon$. It means for some j ,

$$(A^n)_{i,j} > (\bar{\lambda}_i - \epsilon)^n.$$

Let $\ell = \lfloor n/2 \rfloor, m = \lceil n/2 \rceil$.

It follows from $(A^n)_{i,j} = \sum_k (A^\ell)_{i,k} (A^m)_{k,j}$ that for some k ,

$$(A^\ell)_{i,k} (A^m)_{k,j} > \text{const} (\bar{\lambda}_i - \epsilon)^n.$$

⁷Hint: If Δ is the distance from i to j , then $\max_k (A^{n+\Delta})_{i,k} \geq \text{const} \max_k (A^n)_{j,k}$.

⁸The value θ_i here is the same as the one defined in the introduction, but we do not use any insight from there.

Since $m > n(k, \epsilon)$ for every k , we have $(A^m)_{k,j} < (\bar{\lambda}_k + \epsilon)^m$, which implies

$$(A^\ell)_{i,k} > \text{const} \frac{(\bar{\lambda}_i - \epsilon)^n}{(\bar{\lambda}_k + \epsilon)^m}.$$

We can see that the component of k is not smaller than the component of i when n is large enough and ϵ is small enough, since otherwise, $\bar{\lambda}_i > \bar{\lambda}_k$ implies $(A^\ell)_{i,k} \leq \max_j (A^\ell)_{i,j}$ is not bounded by $(\bar{\lambda}_i + \epsilon)^\ell$ (note that $\ell > n(i, \epsilon)$).

Therefore, i and k are in the same component in such a situation, that is $\bar{\lambda}_i = \bar{\lambda}_k$ by Lemma 3. We obtain

$$(A^\ell)_{i,k} > \text{const} \frac{(\bar{\lambda}_i - \epsilon)^n}{(\bar{\lambda}_i + \epsilon)^m} > \text{const}(\bar{\lambda}_i - \epsilon')^\ell,$$

where ϵ' depends on ϵ and for every ϵ' there exists ϵ such that the inequality holds.

Let t be the distance from k to i in the dependency graph, we have

$$(A^{\ell+t})_{i,i} \geq (A^\ell)_{i,k} (A^t)_{k,i} \geq \text{const}(\bar{\lambda}_i - \epsilon')^\ell \geq \text{const}(\bar{\lambda}_i - \epsilon')^{\ell+t}.$$

When ϵ' is small enough and ℓ is large enough, we come up with a contradiction by $\sqrt[\ell+t]{(A^{\ell+t})_{i,i}} > \theta_i$ since $\theta_i < \bar{\lambda}_i$ by assumption.

Therefore, $\theta_i \geq \bar{\lambda}_i$. Together with $\theta_i \leq \underline{\lambda}_i$, the limit $\lambda_i = \underline{\lambda}_i = \bar{\lambda}_i = \theta_i$ exists in this considered case.

Case 2: The vertex i is in a component C that does not satisfy the requirement of C_0 .

By Lemma 3, there exists a satisfying component $C_0 < C$ and $\bar{\lambda}_i = \bar{\lambda}_k$ for $k \in C_0$. Note that in the component C_0 , we already have $\bar{\lambda}_k = \underline{\lambda}_k = \theta_k$. All together with Lemma 3, we obtain

$$\bar{\lambda}_i = \bar{\lambda}_k = \underline{\lambda}_k \leq \underline{\lambda}_i \leq \bar{\lambda}_i.$$

It means the limit $\lambda_i = \underline{\lambda}_i = \bar{\lambda}_i = \theta_k$ also exists in this remaining case, that finishes the proof.

3. PROOF OF THEOREM 4

We assume that there is a circuit from i to i , otherwise the theorem becomes trivial.

The \leftarrow direction is rather obvious: Every $(A^m)_{i,i}$ is zero except when m is multiple of the unique length p of a cycle from i to i , since otherwise, a circuit from i to i of length m not divided by p can be decomposed into a sequence of circuits from i to i , each not visiting i in between, and their lengths cannot be all divided by p .

Also, $(A^p)_{i,j} = 0$ (and $(A^p)_{j,i} = 0$) for j in the same component as i and $j \neq i$, since otherwise, there is a circuit from i to i without visiting i in between but longer than p by extending the path from the endpoint j to a new endpoint i .

For $m = pq$, we have

$$(A^m)_{i,i} = (A^{pq})_{i,i} = \sum_j (A^p)_{i,j} (A^{(q-1)p})_{j,i} = (A^p)_{i,i} (A^{(q-1)p})_{i,i} = (A^p)_{i,i} [(A^p)_{i,i}]^{q-1} = [(A^p)_{i,i}]^q,$$

where the next to last equality is obtained by induction with the trivial base case $q = 1$.

The conclusion that $\sqrt[p]{(A^p)_{i,i}} = \sup_n \sqrt[n]{(A^n)_{i,i}}$ follows.

It remains to prove the \rightarrow direction: Assume there is m so that $\sqrt[m]{(A^m)_{i,i}} = \sup_n \sqrt[n]{(A^n)_{i,i}}$.

First, let p be the length of a circuit from i to i not visiting i in between, we show that p divides m . Assume otherwise, let n be the least common multiple of m and p . Due to the assumption, we have $n = mq$ for some $q > 1$. Let $q = q_1 + q_2$ for some $q_1 > 0, q_2 > 0$.

Since n is also a multiple of p , there is a circuit from i to i such that it does not visit i except at the positions being multiples of p (positions are counted from 0). In this walk, the vertex at position q_1m is not i since p does not divide q_1m . Therefore, we have the following strict inequality

$$(A^n)_{i,i} = \sum_j (A^{q_1m})_{i,j} (A^{q_2m})_{j,i} > (A^{q_1m})_{i,i} (A^{q_2m})_{i,i}.$$

Note that

$$(A^{q_1m})_{i,i} (A^{q_2m})_{i,i} \geq [(A^m)_{i,i}]^{q_1} [(A^m)_{i,i}]^{q_2} = [(A^m)_{i,i}]^q.$$

In total, $(A^n)_{i,i} > [(A^m)_{i,i}]^q$, which contradicts with the maximality of $\sqrt[m]{(A^m)_{i,i}}$.

Now we are ready to show that there is only one length p for all circuits from i to i not visiting i in between. Assume there are two different lengths p_1, p_2 with $p_1 < p_2$. As previously shown, m is a multiple of them, say $m = p_1q_1 = p_2q_2$. Since $2m = p_1q_1 + p_2q_2 = (q_1 - 1)p_1 + p_2 + p_1 + (q_2 - 1)p_2$, there exists a circuit of length $2m$ such that it visits i only at positions $0, p_1, 2p_1, \dots, (q_1 - 1)p_1, (q_1 - 1)p_1 + p_2, (q_1 - 1)p_1 + p_2 + p_1, m + 2p_2, m + 3p_2, \dots, 2m$. Note that $(q_1 - 1)p_1 < m < (q_1 - 1)p_1 + p_2$, that is the circuit visits a vertex other than i at position m . Therefore,

$$(A^{2m})_{i,i} = \sum_j (A^m)_{i,j} (A^m)_{j,i} > (A^m)_{i,i} (A^m)_{i,i} = [(A^m)_{i,i}]^2.$$

This contradicts with the maximality of $\sqrt[m]{(A^m)_{i,i}}$. The conclusion follows.

4. PROOF OF THE FACTS SUPPORTING THE BOUND OF $\lambda = \rho(A)$

The proof of Lemma 1 belows can be seen as a modification of the usual proof of Fekete's lemma.

Proof of Lemma 1. If there is any m such that $a_m > 0$, then the subsequence $\{a_n : a_n > 0\}$ is infinite ($a_{mt} > 0$ for every $t \geq 1$). We suppose the subsequence $\{a_n : a_n > 0\}$ is not empty, thus infinite.

It is obvious by definition that $\limsup\{\sqrt[n]{a_n} : a_n > 0\} \leq \sup_n \sqrt[n]{a_n}$. To finish the proof, it remains to show $\liminf\{\sqrt[n]{a_n} : a_n > 0\} \geq \sup_n \sqrt[n]{a_n}$.

Consider any positive integer q . Let R be the set of integer r ($0 \leq r < q$) such that there exists some m_r with $m_r \equiv r \pmod{q}$ and $a_{m_r} > 0$.

For every n such that $a_n > 0$, if $n \equiv r \pmod{q}$, then $r \in R$. Consider the representation $n = pq + m_r$, we obtain

$$\sqrt[n]{a_n} \geq \sqrt[q]{(a_q)^p a_{m_r}}.$$

The lower bound converges to $\sqrt[q]{a_q}$ when $n \rightarrow \infty$ since m_r is bounded.

As the lower bound holds for every q , we have shown the lower bound $\sup_n \sqrt[n]{a_n}$ for the limit inferior and finished the proof. \square

In order to prove Theorem 2, we first prove the following lemmas.

Denote $d_i = \gcd\{n : (A^n)_{i,i} > 0\}$ for an i not in a single component.

Lemma 4. *For every i in a nonsingle component, there exists N such that for every $n \geq N$, if $d_i | n$ then $(A^n)_{i,i} > 0$. In particular, one can set $N = (D - 1)(2D - 1)$ where $D \times D$ is the dimension of A .*

The readers may relate this lemma to a result of Wielandt: A matrix A is primitive if and only if $A^{D^2 - 2D + 2} > 0$. If a matrix is irreducible but imprimitive, then all d_i are equal to some $d > 0$. Hence A^d is primitive. It follows that A^{dt} for every $t \geq D^2 - 2D + 2$ is positive. However, we do not use this approach.

We say 2 subwalks in a walk are disjoint if the number of vertices they share is at most 1 (only at one of the endpoints if there is one). We give first the following observation.

Observation 1. *If a walk does not contain two disjoint circuits, then its length is less than $2D$.*

Proof. We suppose the walk contains at least a circuit as it is trivial otherwise. Assume the walk does not contain two disjoint circuits.

Let the walk be v_0, \dots, v_k . Let i, j be the indices with the minimal i such that the subwalk v_i, \dots, v_j is a circuit that does not contain a subcircuit (i.e. a cycle).

The vertices v_0, \dots, v_{i-1} are distinct, otherwise we will have a circuit to the left of v_i .

The vertices v_i, \dots, v_{j-1} are distinct due to the absence of a subcircuit.

The sets $\{v_0, \dots, v_{i-1}\}$ and $\{v_i, \dots, v_{j-1}\}$ are disjoint, otherwise for $v_{i'} = v_{j'}$ ($0 \leq i' \leq i - 1, i \leq j' \leq j - 1$) with maximal i' , we will have a circuit $v_{i'}, \dots, v_{j'}$ that does not contain a subcircuit but with $i' < i$.

Therefore, the vertices v_0, \dots, v_{j-1} are all distinct.

The vertices v_j, \dots, v_n are also distinct as otherwise, we will have a second circuit, which is disjoint from the first circuit.

In total, the walk contains at most $2D$ vertices, which is followed by the conclusion. \square

The walk $1, \dots, D, D, \dots, 1$ indicates the sharpness of the bound.

Before proving Lemma 4, we state I. Schur's lemma that gives a bound on Frobenius' number.

Lemma 5 (I. Schur 1935). *Let $2 \leq p_1 < p_2 < \dots < p_k$ be k integers such that $\gcd(p_1, \dots, p_k) = 1$, then every integer $n \geq (p_1 - 1)(p_k - 1)$ can be expressed as a linear combination of p_1, \dots, p_k with nonnegative coefficients.*

We are now ready to prove Lemma 4.

Proof. Let $S = \{n : (A^n)_{i,i} > 0\}$.

Let m be the minimal element of S such that the set $S^* = \{n \in S : n \leq m\}$ satisfies $\gcd S^* = d_i$. We prove that $m < 2D$.

Due to the minimality of m , we have $\gcd(S^* \setminus \{m\}) = d^* > d_i$.

Suppose $m \geq 2D$. As $(A^m)_{i,i} > 0$, there is a circuit from i to i of length m , which contains 2 disjoint subcircuits (by Observation 1). Let a and b be the lengths of the two subcircuits. Removing any of these subcircuits results in a circuit of length less than m , which is divided by d^* . That is $d^* | m - a$ and $d^* | m - b$. However, the circuit obtained by removing both subcircuits would have length $m - a - b$, which is not divided by d^* , a contradiction. Therefore, $m < 2D$.

Let $T = \{n/d_i : n \in S^*\}$, we have $\gcd T = 1$. As the smallest element of T is at most D/d_i and the biggest element of T is at most $2D/d_i$, we have $(A^n)_{i,i} > 0$ for every

$n \geq d_i(D/d_i - 1)(2D/d_i - 1)$ and n divided by d_i , by the bound in I. Schur's lemma. The conclusion follows as $d_i(D/d_i - 1)(2D/d_i - 1) \leq (D - 1)(2D - 1)$. \square

The readers would notice that the bound given in Lemma 4 is not optimal, but sometimes it is asymptotically optimal. Suppose the graph has two disjoint cycles around i of length ℓ and m with $\ell + m = D + 1$ and $\gcd(\ell, m) = 1$, then I. Schur's bound $(\ell - 1)(m - 1)$ is optimal. If the two values ℓ, m are chosen to be around $D/2$, then our bound $(D - 1)(2D - 1)$ differs by a constant times only.

Lemma 6. *For any index i , if m, n are two integers whose difference is bounded, then either $(A^n)_{i,i} = 0$ or the ratio $(A^m)_{i,i}/(A^n)_{i,i}$ is bounded. In particular, if $m \geq n$, then the bound can be set to*

$$(UD)^{m-n} \left(\frac{U}{V}\right)^{3D^2-D},$$

where $D \times D$ is the dimension of A and U, V are respectively the largest and smallest entry of A .

We are interested in an explicit constant only for the case $m \geq n$ since it will be applied in the proof of Theorem 2.

Proof. Let $d = \gcd\{t : (A^t)_{i,i} > 0\}$.

Denote $N = (D - 1)(2D - 1)$. By Lemma 4, for every $n \geq N$, if $d|n$ then $(A^n)_{i,i} > 0$.

Let $S_t = \{j : (A^t)_{j,i} > 0\}$ for each $t \geq 1$, that is the set of vertices from which we can reach i by a walk of length t .

Let $S = \bigcup_{t:d|t} S_t$, that is the set of vertices from which we can reach i by a walk of some length divided by d .

Let $M = N + D^2$, we have: If $t \geq M$ and $d|t$, then $S_t = S$. Indeed, for each $j \in S$, let ud be the least multiple of d such that $(A^{ud})_{j,i} > 0$. We have $u \leq D$. Consider any $t = vd$ such that $t \geq M \geq N + ud$. We then have $vd - ud \geq N$, which implies $(A^{vd-ud})_{i,i} > 0$. That is $(A^t)_{j,i} \geq (A^{ud})_{j,i}(A^{vd-ud})_{i,i} > 0$, i.e. the vertex j is also in S_t .

Note that $M = N + D^2 = (D - 1)(2D - 1) + D^2 = 3D^2 - 3D + 1 \leq 3D^2 - 2D$.

Suppose n is large enough and $m > d$ such that there exists a greatest positive integer $t < \min\{m, n\}$ such that $n - t \geq M$ and $d|(n - t)$. Since d also divides $m - t$, we have

$$(A^m)_{i,i} = \sum_{j \in S} (A^t)_{i,j} (A^{m-t})_{j,i} \leq \sum_{j \in S} (A^t)_{i,j} (A^{n-t})_{j,i} \frac{\max_{j \in S} (A^{m-t})_{j,i}}{\min_{j \in S} (A^{n-t})_{j,i}} = \frac{\max_{j \in S} (A^{m-t})_{j,i}}{\min_{j \in S} (A^{n-t})_{j,i}} (A^n)_{i,i}.$$

Note that denominator is positive as $n - t \geq M$.

Since $m - t$ and $n - t$ are bounded, the ratio $\max_{j \in S} (A^{m-t})_{j,i} / \min_{j \in S} (A^{n-t})_{j,i}$ is bounded, and so is the ratio $(A^m)_{i,i} / (A^n)_{i,i}$.

Note that for every $\Delta > 0$, every entry of A^Δ is bounded from above by $D^{\Delta-1}U^\Delta$ and from below by $D^{\Delta-1}V^\Delta$, where U, V are respectively the largest and smallest entry of A .

Although the statement of the lemma asks for an explicit constant only when $m \geq n$, we will give explicit constants for any pair m, n .

Suppose $n - m \leq N$ (the situation $m \geq n$ is included in this case), we have $n - t$ is not too far from M . In particular, $M \leq n - t \leq M + d \leq 3D^2 - D$.

Hence,

$$\frac{(A^m)_{i,i}}{(A^n)_{i,i}} \leq \frac{D^{m-t-1}U^{m-t}}{D^{n-t-1}V^{n-t}} = D^{m-n} \left(\frac{U}{V}\right)^{n-t} U^{m-n} \leq D^{m-n} \left(\frac{U}{V}\right)^{3D^2-D} U^{m-n} = (UD)^{m-n} \left(\frac{U}{V}\right)^{3D^2-D}.$$

If $n - m > N$, then there is a circuit of length $n - m$ from i to i . Therefore,

$$(A^n)_{i,i} \geq (A^m)_{i,i} (A^{n-m})_{i,i} \geq (A^m)_{i,i} D^{n-m-1} V^{n-m},$$

which means

$$\frac{(A^m)_{i,i}}{(A^n)_{i,i}} \leq D^{m-n+1} V^{m-n}.$$

It remains to consider the case when n is not large enough or $m = d$ for which we may not have such t . If $m = d$ and $n > 3D^2 - D$ then $n - m > N$ and the above bound holds. If $n \leq 3D^2 - D$, then we may not have satisfying t , but regardless of m , we still have

$$\frac{(A^m)_{i,i}}{(A^n)_{i,i}} \leq \frac{D^{m-1} U^m}{D^{n-1} V^n} \leq D^{m-n} \left(\frac{U}{V}\right)^n U^{m-n} \leq D^{m-n} \left(\frac{U}{V}\right)^{3D^2-D} U^{m-n} = (UD)^{m-n} \left(\frac{U}{V}\right)^{3D^2-D}.$$

□

Note that U and V can be set to the largest entry and the smallest entry of the matrix induced by the vertices in the component of i . Also, the term D can be replaced by d in some places. However, they are inessential improvements, and we did not intend to optimize the constant very much in the first place.

We are now ready to prove Theorem 2.

Proof of Theorem 2. Let C be the component containing i . We assume $(A^\ell)_{i,i}$ and $(A^m)_{i,i}$ are both positive as it is trivial otherwise.

For every $j \in C$, let Δ be the distance from j to i , we have

$$\begin{aligned} (A^\ell)_{i,j} &\leq \frac{1}{(A^\Delta)_{j,i}} (A^{\ell+\Delta})_{i,i} \leq \frac{1}{D^{\Delta-1} V^\Delta} (A^{\ell+\Delta})_{i,i} \leq \frac{1}{D^{\Delta-1} V^\Delta} (UD)^\Delta \left(\frac{U}{V}\right)^{3D^2-D} (A^\ell)_{i,i} \\ &= D \left(\frac{U}{V}\right)^{3D^2-D+\Delta} (A^\ell)_{i,i} \leq D \left(\frac{U}{V}\right)^{3D^2} (A^\ell)_{i,i}, \end{aligned}$$

where the second inequality is due to the lower bound $D^{\Delta-1} V^\Delta$ for any entry in A^Δ , and the third inequality is due to Lemma 6.

Likewise, for every $j \in C$, we have

$$(A^m)_{j,i} \leq D \left(\frac{U}{V}\right)^{3D^2} (A^m)_{i,i}.$$

We obtain the conclusion by

$$\begin{aligned} (A^{\ell+m})_{i,i} &= \sum_{j \in C} (A^\ell)_{i,j} (A^m)_{j,i} \leq \sum_{j \in C} D \left(\frac{U}{V}\right)^{3D^2} (A^\ell)_{i,i} D \left(\frac{U}{V}\right)^{3D^2} (A^m)_{i,i} \\ &\leq D \left(D \left(\frac{U}{V}\right)^{3D^2}\right)^2 (A^\ell)_{i,i} (A^m)_{i,i} \\ &= D^3 \left(\frac{U}{V}\right)^{6D^2} (A^\ell)_{i,i} (A^m)_{i,i}. \end{aligned}$$

□

Although Theorem 2 is deduced from Lemma 6, if the actual constants are not concerned, we can do the same for the other direction. Indeed, suppose we have Theorem 2 in the first

place, we then have the situation of Equation (2), which can be rewritten as

$$\frac{1}{K_i}(\theta_i)^n \leq (A^n)_{i,i} \leq (\theta_i)^n$$

for every positive $(A^n)_{i,i}$. This is obviously followed by Lemma 6 (with perhaps another constant, e.g. $(\theta_i)^{m-n} D^3 (\frac{U}{V})^{6D^2}$ for every pair m, n not necessarily $m \geq n$).

5. PROOF OF THE BOUNDS FOR THE POWERS OF MATRICES

Denote $d_i = \gcd\{n : (A^n)_{i,i} > 0\}$ for an i in a nonsingle component. We begin with the following lemmas.

Lemma 7. *If i, j are two different vertices in the same component, then $d_i = d_j$.*

Proof. Consider a circuit from j to j that has length td_j such that $\gcd(t, d_i) = 1$ (we can just choose any prime t large enough, which is available by Lemma 4). Then we have a circuit from i to i of length $\delta(i, j) + td_j + \delta(j, i)$, which is divided by d_i , where $\delta(i, j)$ is the distance from i to j .

Since there is also a circuit from i to i of length $\delta(i, j) + \delta(j, i)$, we have $d_i | \delta(i, j) + \delta(j, i)$. Therefore, $d_i | td_j$, which implies $d_i | d_j$.

Exchanging the roles of i and j , one also obtains $d_j | d_i$.

The conclusion $d_i = d_j$ follows. □

Lemma 7 is used in the following lemma.

Lemma 8. *For i, j of the same component, if there is a walk of length n from i to j for some n large enough, then there is also a walk of length n from i to j that begins with a circuit from i to i of length m such that $n - m$ is bounded.*

Proof. Let the walk of length n from i to j be

$$v_0, v_1, \dots, v_n,$$

where $v_0 = i$ and $v_n = j$.

Starting with v_0 , we take the latest instance v_t of v_0 in the walk, that is $v_t = v_0$ and $v_{t'} \neq v_0$ for every $t' > t$. We then consider the subwalk,

$$v_{t+1}, \dots, v_n,$$

which has no instance of v_0 . (Note that if $v_{t'} \neq v_0$ for every $t' \neq 0$, then we just set $t = 0$.)

Repeating the process with the subwalk until reaching an empty walk, we will obtain a decomposition of the original walk into consecutive circuits represented by

$$(\ell_1, r_1), \dots, (\ell_s, r_s),$$

where s is bounded by the number of vertices in the component of i, j ; $\ell_1 = 0, r_s = n$; $\ell_t = r_{t-1} + 1$ for $t = 2, \dots, s$; and the subwalk from v_{ℓ_t} to v_{r_t} is a circuit (of possibly empty length) for $t = 1, \dots, s$.

Denote $d = d_k$ for every k in the component of i, j (by Lemma 7), then $d | (r_t - \ell_t)$ for every t , which means

$$d | \sum_{t=1}^s (r_t - \ell_t).$$

As we only consider n large enough, the above sum will be large and there is a circuit from i to i of that length (by Lemma 4). (If n is small enough, the original walk itself satisfies the requirement.)

Removing the circuit from v_{ℓ_t} to v_{r_t} for every $t = 1, \dots, s$ and inserting the big circuit from i to i of length $m = \sum_{t=1}^s (r_t - \ell_t)$ into the beginning of the original walk, we obtain a new walk from i to j of the same length n satisfying the requirement. \square

We have an immediate corollary.

Corollary 2. *If i, j are in the same component and $(A^n)_{i,j} > 0$ for some n large enough, then there exists some bounded $\Delta \geq 0$ such that $(A^{n-\Delta})_{i,i} > 0$ and*

$$(A^n)_{i,j} \geq \text{const}(A^{n-\Delta})_{i,i}.$$

Proof of Theorem 5. We assume that $(A^n)_{i',j'}$ is positive and prove that the ratio is bounded. Also, we assume m, n are large enough.

By Corollary 2, there exists some bounded Δ such that $(A^{n-\Delta})_{i',i'} > 0$ and

$$(A^n)_{i',j'} \geq \text{const}(A^{n-\Delta})_{i',i'}.$$

Let $\Delta' = \delta(j, i)$ be the distance from j to i , we have

$$(A^m)_{i,j} \leq \frac{1}{(A^{\Delta'})_{j,i}} (A^{m+\Delta'})_{i,i} = \text{const}(A^{m+\Delta'})_{i,i}.$$

On the other hand,

$$(A^{\Delta_1})_{i',i} (A^{m+\Delta'})_{i,i} (A^{\Delta_2})_{i,i'} \leq (A^{m+\Delta'+\Delta_1+\Delta_2})_{i',i'} \leq \text{const}(A^{n-\Delta})_{i',i'},$$

where $\Delta_1 = \delta(i', i)$, $\Delta_2 = \delta(i, i')$, and the latter inequality is due to Lemma 6.

It follows that

$$(A^{m+\Delta'})_{i,i} \leq \text{const}(A^{n-\Delta})_{i',i'}.$$

Combining the inequalities, we obtain the conclusion since

$$(A^m)_{i,j} \leq \text{const}(A^{m+\Delta'})_{i,i} \leq \text{const}(A^{n-\Delta})_{i',i'} \leq \text{const}(A^n)_{i',j'}.$$

\square

Proof of Theorem 6. Let the path from the component of i to the component of j in the condensation be C_1, \dots, C_s .

For a walk of length n from i to j , we decompose it into s disjoint consecutive subwalks, the t -th subwalk is from the position ℓ_t to the position r_t for $t = 1, \dots, s$ and $\ell_1 = 0$, $r_t = n$; $\ell_t = r_{t-1} + 1$ for $t = 2, \dots, s$ such that the whole t -th subwalk for every t contains only the vertices in C_t . Denote $n_t = r_t - \ell_t$, the difference between n and the sum of all n_t is bounded since

$$\sum_{t=1}^s n_t = n - s + 1.$$

Fix a sequence of n_1, \dots, n_s , let $(\underline{A}^{n_1, \dots, n_s})_{i,j}$ be the value with restriction to all walks satisfying the decomposition n_1, \dots, n_s , that is:

$$(\underline{A}^{n_1, \dots, n_s})_{i,j} = \sum_{\substack{i_1, j_1, \dots, i_s, j_s \\ i_t, j_t \in C_t \\ i_1 = i, j_s = j}} (A^{n_1})_{i_1, j_1} A_{j_1, i_2} (A^{n_2})_{i_2, j_2} \dots (A^{n_{s-1}})_{i_{s-1}, j_{s-1}} A_{j_{s-1}, i_s} (A^{n_s})_{i_s, j_s},$$

where i_t, j_t are the two endpoints in the t -th subwalk for $t = 1, \dots, s$.

Since there are finitely many (may be even zero) options for the two endpoints in each subwalk and $(A^{n_t})_{i_t, j_t} \leq \text{const}(\theta_{C_t})^{n_t}$ for every component t (by Corollary 1), we have

$$(\underline{A}^{n_1, \dots, n_s})_{i, j} \leq \text{const} \prod_{1 \leq t \leq s} (\theta_{C_t})^{n_t}.$$

An upper bound of $(\underline{A}^{n_1, \dots, n_s})_{i, j}$ for any n_1, \dots, n_s is therefore of order θ^n where $\theta = \max_t \theta_{C_t}$. The bound is attained when the sum of n_t over all t satisfying $\theta_{C_t} = \theta$ is at a bounded distance from n . Since there are at most polynomially many possible sequences n_1, \dots, n_s , we have

$$(A^n)_{i, j} \leq \text{const} n^r \theta^n,$$

where r is one less than the number of components C_t with $\theta_{C_t} = \theta$.

We show below the other direction $(A^n)_{i, j} \geq \text{const} n^r \theta^n$ by showing that n^r is the order of the number of valid sequences n_1, \dots, n_s such that the sum of n_t for $\theta_{C_t} = \theta$ is at a bounded distance from n .

Indeed, let us start with any walk with the corresponding sequence n_1, \dots, n_s . In order to prove the order n^r , it is sufficient to show that: For any t_1, t_2 such that C_{t_1}, C_{t_2} are both nonsingle and n_{t_1} is large enough, we can transform the original walk into another walk with the corresponding sequence n'_1, \dots, n'_s with $n_{t_1} - n'_{t_1} = n'_{t_2} - n_{t_2} = \Delta > 0$ while $n_t = n'_t$ for $t \neq t_1, t_2$; all possible values of Δ form a sequence so that $\min \Delta, n_{t_1} - \max \Delta$ and the difference between two consecutive Δ are all bounded. We can say roughly that it is the action of transfer Δ steps from the subwalk of C_{t_1} to the subwalk of C_{t_2} and there are at least $\text{const} n_{t_1}$ possible values of Δ .

At first, we assume that the subwalk of C_{t_1} begins with a circuit of length m from i_{t_1} to i_{t_1} , where i_{t_1} is the first vertex in the subwalk, so that $n_{t_1} - m$ is bounded, since otherwise we can transform the subwalk into the desired subwalk by Lemma 8.

Let d be the least common multiple of d_{t_1} and d_{t_2} . Consider all the values $\Delta = pd$ for some p such that the value Δ is large enough so that there is a circuit from i_{t_2} to i_{t_2} of length Δ for the beginning vertex i_{t_2} in the subwalk of C_{t_2} , and in the same time Δ is small enough so that there is circuit from i_{t_1} to i_{t_1} of length $m - \Delta$. The existence is ensured by Lemma 4. The new subwalk in C_{t_1} starts with a circuit of length $m - \Delta$ instead of m . In the new subwalk of C_{t_2} , a circuit of length Δ from i_{t_2} to i_{t_2} is now inserted to the beginning.

Also by Lemma 4, we can see that the first possible Δ is bounded, the last possible Δ is at a bounded distance from n_{t_1} and the difference between any two consecutive possible Δ is at most d .

We have shown that n^r is the order of the number of sequences n_1, \dots, n_s such that the difference between n and the sum of n_t over the components C_t with $\theta_{C_t} = \theta$ is bounded. It means

$$\text{const} n^r \theta^n \leq (A^n)_{i, j} \leq \text{const} n^r \theta^n.$$

□

Proof of Theorem 7. We assume that there is a path from i to j , as it is trivial otherwise.

Different walks of length n from i to j may go through the components in different paths in the condensation. For a path P from the component of i to the component of j in the condensation, let A_P be the matrix induced by only the vertices in the components along P .

We can see that

$$(A^n)_{i,j} = \sum_P (A_P^n)_{i,j},$$

where P is ranged over all possible paths between the two components.

By Theorem 6, $(A_P^n)_{i,j}$ is of order $n^{r_P}(\theta_P)^n$, where θ_P is the maximal θ_C over all the components C on P and r_P is one less than the number of components C on P with $\theta_C = \theta_P$.

Let $n^r\theta^n$ be the maximal order $n^{r_P}(\theta_P)^n$ of $(A_P^n)_{i,j}$ over all the paths P . Since the number of paths is finite, we have

$$(A^n)_{i,j} = \sum_P (A_P^n)_{i,j} \leq \text{const } n^r\theta^n.$$

The upper bound sometimes cannot be attained since a walk of length n from i to j may not be able to go along the components on a path P^* with the order $n^r\theta^n$ (see again the example in Figure 1). However, there are infinitely often many n so that there is such a walk. Indeed, take any n_0 so that there is such a walk. Let C be any nonsingle component on P^* and k a vertex of C in the walk. For every $n = n_0 + md_k$ for m large enough, there is a circuit of length of md_k from k to k (by Lemma 4), hence there is a walk of length n from i to j . □

Proof of Theorem 8. From Theorem 7, each pair i, j gives an order $n^r\theta^n$ and also a path in the condensation. Consider a pair i, j that maximizes the order $n^r\theta^n$ over all the pairs i, j . Let P be the corresponding path in the condensation. We can assume that the last component on P , which contains j , is nonsingle since otherwise removing that component and take any j in the next to last component does not reduce the order.

The latter inequality $\max_i \max_j (A^n)_{i,j} \leq \text{const } n^r\theta^n$ is clear from the choice of $n^r\theta^n$.

This order can be attained for every n . Indeed, by Theorem 7, there is a sequence $\{n_t\}_t$ with the same difference d between two consecutive elements and $(A^{n_t})_{i,j} \geq \text{const } n_t^r\theta^{n_t}$ for every t . For any $n = n_t + u$ for $0 \leq u < d$, consider a walk of length n_t from i to j . Since the last component is nonsingle, we can extend the walk by appending a walk of length u from j to j' for some j' in that component. We obtain the former inequality since

$$(A^n)_{i,j'} \geq (A^{n_t})_{i,j} (A^u)_{j,j'} \geq \text{const } n_t^r\theta^{n_t} \geq \text{const } n^r\theta^n.$$

□

6. COMPACT VERSION OF THE THIRD PROOF

The proof below would be a bit sketchy in the later half since most arguments there are rather similar to the other sections.

Suppose i, j (not necessarily different) are in the same component in the dependency graph, we have

$$\begin{aligned} (A^{\ell+m+2\delta(j,i)+\delta(i,j)})_{i,j} &\geq (A^{\ell+m+2\delta(j,i)})_{i,i} (A^{\delta(i,j)})_{i,j} \\ &\geq (A^{\ell+\delta(j,i)})_{i,i} (A^{m+\delta(j,i)})_{i,i} (A^{\delta(i,j)})_{i,j} \\ &\geq (A^\ell)_{i,j} (A^{\delta(j,i)})_{j,i} (A^m)_{i,j} (A^{\delta(j,i)})_{j,i} (A^{\delta(i,j)})_{i,j} \\ &= K_{i,j} (A^\ell)_{i,j} (A^m)_{i,j}, \end{aligned}$$

where $K_{i,j} = ((A^{\delta(j,i)})_{j,i})^2 (A^{\delta(i,j)})_{i,j}$ and $\delta(i,j)$ is the distance from i to j .

Apply the extension of Fekete's lemma (Lemma 1) to the nonnegative supermultiplicative sequence $x_n = K_{i,j} (A^{n-2\delta(j,i)-\delta(i,j)})_{i,j}$ for each i, j , we have the subsequence of positive $\sqrt[n]{(A^n)_{i,j}}$ converges to $\sup_n \sqrt[n]{K_{i,j} (A^{n-2\delta(j,i)-\delta(i,j)})_{i,j}}$.

Note that for any i, j, i', j' of the same component, we have $\sup_n \sqrt[n]{K_{i,j} (A^{n-2\delta(j,i)-\delta(i,j)})_{i,j}} = \sup_n \sqrt[n]{K_{i',j'} (A^{n-2\delta(j',i')-\delta(i',j')})_{i',j'}}$ since they are supermultiplicative and $(A^{n+\delta(i,i')+\delta(j',j)})_{i,j} \geq \text{const}(A^n)_{i',j'}$, $(A^{n+\delta(i',i)+\delta(j,j')})_{i',j'} \geq \text{const}(A^n)_{i,j}$. Denote by λ_C the limit for component C . That is $\lambda_C = \sup_n \sqrt[n]{(A^n)_{i,i}}$ for any $i \in C$. Note that for every n , there exists a pair i, j in C such that $(A^n)_{i,j} > 0$. In other words,

$$\lim_{n \rightarrow \infty} \sqrt[n]{\max_{i,j \in C} (A^n)_{i,j}} = \lambda_C.$$

Let C^* be a component such that $\lambda_{C^*} = \max_C \lambda_C = \max_i \sup_n \sqrt[n]{(A^n)_{i,i}}$. We have

$$\liminf_{n \rightarrow \infty} \sqrt[n]{\max_{i,j} (A^n)_{i,j}} \geq \lambda_{C^*}.$$

For any i, j ,

$$(A^n)_{i,j} = \sum_{\substack{k \text{ different components for some } k \\ \text{all partitions } n=(k-1)+n_1+\dots+n_k \\ \text{all possible } i_*, j_* \text{ of the same component} \\ i_1=i_*, j_k=j_*}} (A^{n_1})_{i_1, j_1} A_{j_1, i_2} (A^{n_2})_{i_2, j_2} \dots (A^{n_{k-1}})_{i_{k-1}, j_{k-1}} A_{j_{k-1}, i_k} (A^{n_k})_{i_k, j_k}.$$

Since there are only finitely many choices for components and endpoints i_*, j_* , and only polynomially many choices for n_1, \dots, n_k , we have

$$\limsup_{n \rightarrow \infty} \sqrt[n]{\max_{i,j} (A^n)_{i,j}} \leq \lambda_{C^*}.$$

We obtain the conclusion

$$\lim_{n \rightarrow \infty} \sqrt[n]{\max_{i,j} (A^n)_{i,j}} = \max_i \sup_n \sqrt[n]{(A^n)_{i,i}}.$$

REFERENCES

- [1] Michael Fekete. Über die Verteilung der Wurzeln bei gewissen algebraischen Gleichungen mit ganzzahligen Koeffizienten. *Mathematische Zeitschrift*, 17(1):228–249, 1923.
- [2] Semyon Aranovich Gershgorin. Über die Abgrenzung der Eigenwerte einer Matrix. *Известия Российской академии наук. Серия математическая*, (6):749–754, 1931.
- [3] RV Mises and Hilda Pollaczek-Geiringer. Praktische Verfahren der Gleichungsauflösung. *ZAMM-Journal of Applied Mathematics and Mechanics/Zeitschrift für Angewandte Mathematik und Mechanik*, 9(1):58–77, 1929.
- [4] Vuong Bui. Growth of bilinear maps. *Linear Algebra and its Applications*, 2021.
- [5] H. K. Wimmer. Spectral radius and radius of convergence. *The American Mathematical Monthly*, 81(6):625–627, 1974.

VUONG BUI, INSTITUT FÜR INFORMATIK, FREIE UNIVERSITÄT BERLIN, TAKUSTRASSE 9, 14195 BERLIN, GERMANY

Email address: bui.vuong@fu-berlin.de