

Primitive Forms without Higher Residue Structure and Integrable Hierarchies (I)

Konstantin Aleshkin and Kyoji Saito

ABSTRACT. We introduce *primitive forms with or without higher residue structure* and explore their connection with the flat structures with or without a metric [18] and integrable hierarchies of KdV type. Just as the classical case of primitive forms with metric [11], the primitive forms without metrics are constructed as the positive part of the Birkhoff decomposition of formal oscillatory integrals with respect to the descendent variable. The oscillating integrals of a primitive form without metric give rise to a hierarchy of commuting PDE of the KdV type as in the case of primitive forms with metric. This shall be studied in (II).

1. Introduction

We introduce *primitive forms without higher residue structure* (or the same to say, *primitive forms without metric structure*)¹ and explore their connection with the flat structures without a metric and integrable hierarchies of the KdV type.

We start with a historical overview on primitive forms. Primitive forms were introduced by the second named author as the integrands for vanishing cycles in a universal unfolding family of a hypersurface isolated singularity. They are formulated in terms of higher residue pairings [23]. Namely, a primitive form is a top degree relative de Rham cohomology class on a certain family of (open) hypersurfaces, satisfying a system of bilinear relations described by the higher residue pairings [23, 11]. In particular, the conditions include a purity condition of the higher residue pairing on the first derivatives of the primitive form. Then, those bilinear relations on a primitive form induce a geometric structure on the parameter space of the family, called the flat structure [25]. That is, the tangent bundle of the parameter space obtains an algebra structure compatible with a flat metric on it. This implies the existence of a flat coordinate system on the parameter space and a potential function with respect to the triple tensor product of vector fields. Actually, such structure on the parameter space has been found already before on

2000 *Mathematics Subject Classification*. Primary.

¹In the present paper, we shall use a convention that the expression “without metric” includes also the cases “with metric”.

the orbit space of a finite reflection group ([24, 26]) and was called the flat structure. Later a similar structure was found in 2D-topological field theories [4], and was axiomatized by Dubrovin as a Frobenius manifold structure [4, 14, 6].

Sabbah [18] and Manin [14] considered a generalized version of that flat

flat structure without metric got a new interest and impetus since the structure appeared on the deformation parameter space of Okubo-type ordinary differential equations, and consequently, one finds a Saito structure without metric on the orbit space of a unitary reflection group [7, 8, 9] (c.f. also [10]). Similar structures appear in the study of relative and open Gromov-Witten invariants [28, 3]

On the other hand, the primitive forms, as mentioned already, to be considered as integrands for vanishing cycles, can be re-considered also as the integrands over Lefschetz thimbles for oscillatory integrals for the unfolding family of defining function of the hypersurface singularity.² Then, primitive forms can be captured as the positive part in the *descendent variable*³ of the Birkhoff decomposition of the evolution image by an oscillatory integral factor action on the top degree form on the generating center belonging to a suitable section of filtered relative De Rham cohomology group of the generating center ([11]). This view point naturally reduces the defining conditions on a primitive form to the conditions on the pair of the top form and the section on the generating center to be *good*. In particular, the purity condition of higher residue pairings on the primitive forms is reduced to a purity condition of the classical residue pairing defined on the section.

The present paper is the first part in a series of two papers. In the present Part I, we introduce a concept of a primitive form without higher residue structure (that is, we remove a purity condition on the primitive forms)⁴ and discuss their connection to the flat structure without metric.

We first recall in §2-4 some basic frame works and techniques on primitive forms. Namely, in §2, for a given unfolding $F(\underline{x}, \underline{t})$ of a function $f(\underline{x})$ with an isolated critical point (=the generating center), we attach the filtered De Rham cohomology groups (that is, the cohomology theory, which contains, so called, the *descendent variable* z). Then in §3 the concepts of sections and opposite filtrations on the filtered de-Rham cohomology groups are recalled. In §4, we recall the technique of formal analysis (i.e. perturbation theory) of the deformation of oscillatory integrals, developed in [11]. That is, we formally study the actions of the oscillatory integral factor $e^{(f(x)-F(\underline{x}, \underline{t}))/z}$ which intertwines the twisted de Rham cohomology group for f with that for F . The main difficulty here is the *mixture of the descent variable* z and the *deformation variable* \underline{t} , since, in the oscillatory integral factor, the descendent variable z has essential singularity at $z = 0$. However, the perturbative approach allows us to treat only finite order of poles in the descent variable

²To be exact, there is a one dimensional difference between vanishing cycles and Lefschetz thimbles, and accordingly a degree one difference between the corresponding integrands. This can be explained by the relation of a top-form on an ambient space and its Poincare residues on hypersurface-fibers. Here, in the universal unfolding, the initial function f for the unfolding and its defining domain X , is called the *generating center* according to R. Thom [30].

³We mean by the *descendent variable* the denominator variable z in the exponential of the oscillatory integral factor.

⁴Actually this version of primitive forms without higher residue structure was already discussed by some authors, but was not written explicitly (Lectures by the second author in Ecole Polytechnique, Neimehen and Nice (1980). See also [2]).

z so far as we remain in a formal neighborhood of the origin of the deformation space S of bounded order.

After these preparations, the definitions of primitive forms without metric structures are given in §5. As in the classical case [11], their formal construction is achieved as follows. We first consider a good section without metric (which is equivalent data of a good opposite filtration without metric)⁵ to the filtered De Rham cohomology group of the generating center. Then a pair of such section without metric and a top degree De Rham cohomology class in the section is called a good pair without metric if they satisfy certain primitivity and homogeneity conditions. Then the primitive form without metric is given by the positive power part w.r.t. the descendent variable, i.e. the positive part of the Birkhoff decomposition, of the evolution image by the oscillatory integral factor action on the top form. We show in §6 that there is a one to one correspondence between (a) the set of primitive forms without metric structure and (b) the set of good pairs without metric condition.

The construction of a flat structure without metric from a primitive form without metric is an easy formal procedure which is parallel to the classical case [22] just taking care of the lack of the purity condition on the residue pairing, given in §7. We compare the flat structure without metric with the already known definitions of Saito structure without metric and F-manifolds.

In the coming Part II, we explore the connection of these primitive forms without metric structure with integrable hierarchies of the KdV type. Let us outline how the integrable hierarchy is constructed from a primitive form and the corresponding flat structure. We note that the evolution by the oscillatory integral factor preserves opposite filtration both in the filtered De Rham cohomology groups for the generating center and for the unfolding. Therefore, the inverse evolution of the primitive form gives the principal part of the Laurent expansion with respect to the descendent variable in the generating center. That is, the oscillatory integral of the primitive form given by the inverse evolution map applied to the primitive form is a formal power series in the inverse powers of the descendent variable z . As a consequence of the properties of the primitive form, those coefficients of the inverse oscillatory integral satisfies a certain system of differential equations. These equations are crucial in the construction of integrable hierarchies. The decomposition coefficients are called deformed flat coordinates, since the first coefficients in the descendent variable coincide with actual flat coordinates. The particular choice of such a set of deformed flat coordinates defines a calibration of the flat structure (without metric).

In the case where the primitive form has a compatible metric structure, the corresponding flat structure is a Frobenius manifold. There is a well-known construction that associates an integrable system to a Frobenius manifold with an additional data called calibration [4]. Such an integrable system is called the principle hierarchy associated to a given calibrated Frobenius manifold. The equations of the principle hierarchy are defined on an appropriate version of the loop space of the Frobenius manifold. The most notable example is the (dispersionless) KdV hierarchy which is constructed on the one-dimensional deformation space of the

⁵We mean by “without metric” that we remove the purity condition on the residue pairing on the section in the definition.

A_1 -singularity. The KdV hierarchy is a system of PDE on one function. The range of this function is identified with the deformation space of the singularity.

In the case of primitive forms without metric the flat structure on the deformation space is basically a Saito structure without metric or a conformal flat F-manifold. It is known in the literature [13, 12], that such structures give rise to integrable systems of PDE as well via a very similar construction. However, the resulting PDE are not necessarily Hamiltonian which is due to the fact that the flat structure has no natural metric. We remark that there is another construction of integrable systems from generalized Frobenius-type structures which arise from quantum K-theory [16]. In the quantum K-theory case a Frobenius structure has a flat metric but lacks flat identity and Euler vector fields (see the main text).

The second paper is devoted the explanation of the structure of the non-Hamiltonian infinite commuting PDE that can be constructed from primitive forms without metric structure. The defining equations of the integable hierarchy are also constructed using the oscillatory integrals of the primitive form. We also discuss generalizations of the properties of integrable hierarchies from the case of primitive forms with metric structure.

2. Filtered De Rham cohomology modules \mathcal{H}_F

In this section, we introduce the semi-infinite filtered De Rham cohomology modules associated with an isolated critical point of a holomorphic function (cf. [6, 21, 23, 25, 11]).

Let X be an open Stein and contractible neighborhood of the origin 0 in the complex space \mathbb{C}^{n+1} ($n \in \mathbb{Z}_{\geq 0}$). Let f be a holomorphic function on X whose critical point is only at the origin.

DEFINITION 2.1. A tuple (Z, S, p) is called a *frame* of an unfolding F of f (or, F is an unfolding of f over a frame (Z, S, p)), if it satisfies the following conditions 1., 2. and 3.

1. The space Z is a Stein open neighborhood of $0 \in \mathbb{C}^{n+m+1}$ ($m \in \mathbb{Z}_{\geq 0}$), S is a Stein open neighborhood of the base point $0 \in \mathbb{C}^m$, $p : Z \rightarrow S$ is the natural projection with $p^{-1}(0) = X$.

2. The F is a holomorphic function on Z such that the restriction $F|_{X=p^{-1}(0)}$ coincides with f .

3. The restriction $p|_{C_F}$ of the projection p to the relative critical set C_F (see (2.1) below) is finite.

In particular, $F = f$ itself is a trivial unfolding defined over the frame $(X, S = \{0\}, p_0)$ where p_0 is the trivial projection $X \rightarrow \{0\}$.

We allow to change and shrink X , S and Z suitably if necessary. For a convenience, we assume further an auxiliary data that there is a projection

$$\pi_X : Z \longrightarrow X$$

so that Z is embedded into $X \times S \subset \mathbb{C}^{n+1} \times \mathbb{C}^m$ as a domain by the map (π_X, p) . We shall denote by (x_0, \dots, x_n) the coordinate system of \mathbb{C}^{n+1} and by t_1, \dots, t_m the coordinate system of \mathbb{C}^m . Partial derivatives with respect to the coordinate system are denoted by $\partial_{x_0}, \dots, \partial_{x_n}$ and $\partial_{t_1}, \dots, \partial_{t_m}$, respectively. Under this notation, the relative critical set is defined by

$$(2.1) \quad C_F := \{\partial_{x_0} F = \dots = \partial_{x_n} F = 0\} \quad \text{and} \quad \mathcal{O}_{C_F} := \mathcal{O}_Z / (\partial_{x_0} F, \dots, \partial_{x_n} F).$$

The relative critical set C_F is Cohen-Macaulay variety such that the projection $p|_{C_F}$ is finite μ -ramified covering over S . Actually, the direct image $p_*(\mathcal{O}_{C_F})$ is locally free \mathcal{O}_S -module of rank $\mu := \dim_{\mathbb{C}} \mathcal{O}_X/(\partial f)$, called the Milnor number of f . We define a natural \mathcal{O}_S -homomorphism,

$$(2.2) \quad \mathcal{T}_S \rightarrow p_*\mathcal{O}_{C_F}, \quad v \mapsto \tilde{v}F|_{C_F}$$

where \mathcal{T}_S is the sheaf of holomorphic tangent vector fields on S and \tilde{v} is any lifting of a vector field v on $U \subset S$ to that on $p^{-1}(U) \subset Z$ such that $p_*(\tilde{v}) = v$. Then, the restriction $\tilde{v}F|_{C_F \cap p^{-1}(U)}$ does not depend on the choice \tilde{v} .

DEFINITION 2.2. An unfolding F is called *universal* if the morphism (2.2) is an \mathcal{O}_S -isomorphism ([30]).

$$(2.3) \quad \mathcal{T}_S \simeq p_*\mathcal{O}_{C_F}$$

The definition implies, in particular, the dimension $\dim S =: m$ is equal to the Milnor number μ of f . Using this isomorphism, we particularly introduce global vector fields on S

$$(2.4) \quad \partial_1 \quad \text{and} \quad E \in \Gamma(S, \mathcal{T}_S)$$

called the *unit vector field* and the *Euler vector field*, respectively, as those corresponding to the unit 1 and the class of the function F in the RHS of (2.3), respectively. That is, we have

$$(2.5) \quad \tilde{\partial}_1 F|_{C_F} = 1 \quad \text{and} \quad \tilde{E}F|_{C_F} = F|_{C_F}.$$

(See [19] for a justification of this notion).

We usually choose $\tilde{\partial}_1$ in such manner that $\tilde{\partial}_1 F = 1$. We shall often use coordinates system t_1, \dots, t_m such that $\partial_1 = \partial_{t_1}$. The tangent bundle \mathcal{T}_S of S gets an \mathcal{O}_S -algebra structure through the isomorphism (2.3), where we denote by $*$ the product, and ∂_1 plays the role of the unit element.

$$(2.6) \quad \begin{aligned} * & : \mathcal{T}_S \times \mathcal{T}_S \longrightarrow \mathcal{T}_S \\ (\widetilde{v_1 * v_2})(F)|_{C_F} & := \tilde{v}_1(F)|_{C_F} \cdot \tilde{v}_2(F)|_{C_F} \quad \text{for } v_1, v_2 \in \mathcal{T}_S. \end{aligned}$$

Next, we introduce the filtered de-Rham cohomology groups (2.10) and (2.11) associated with any unfolding F of f . Let $\Omega_{Z/S}^* = \bigoplus_{i=0}^{n+1} \Omega_{Z/S}^i$, where $\Omega_{Z/S}^i := \Omega_Z^i/p^*\Omega_S^1 \wedge \Omega_Z^{i-1}$ ($i = 0, \dots, n+1$), be the sheaf of the exterior algebra of holomorphic relative de Rham differentials on Z relative to the map p . We consider two structures of complex on it: the relative de Rham complex $(\Omega_{Z/S}^*, d = d_{Z/S})$ over the base space S and the Poincare complex $(\Omega_{Z/S}^*, dF \wedge)$, which are anti-commuting to each other. Combining the two complexes linearly, we introduce the key complex ([23]) of the present paper:

$$(2.7) \quad (\Omega_{Z/S}^*((z)), D_F).$$

Here (1) $\Omega_{Z/S}^*((z)) := \lim_{\infty \leftarrow k} (\Omega_{Z/S}^* \otimes \mathbb{C}[z, z^{-1}])/z^k(\Omega_{Z/S}^* \otimes \mathbb{C}[z])$ (z is a formal variable commuting with the differentials d and $dF \wedge$). See Footnote 1), and (2) the differential D_F of the complex is the linear combination

$$(2.8) \quad D_F := z d_{Z/S} + dF \wedge \quad \text{such that} \quad (D_F)^2 = 0$$

Since (i) the relative critical points of F are isolated, (ii) the following sequence (2.9) is exact

$$(2.9) \quad 0 \rightarrow \mathcal{O}_Z \xrightarrow{dF \wedge} \Omega_{Z/S}^1 \xrightarrow{dF \wedge} \Omega_{Z/S}^2 \xrightarrow{dF \wedge} \dots \xrightarrow{dF \wedge} \Omega_{Z/S}^n \xrightarrow{dF \wedge} \Omega_{Z/S}^{n+1} \rightarrow \Omega_{Z/S}^{n+1}/\Omega_{Z/S}^n \rightarrow 0$$

(De Rham Lemma [17][20]) and (iii) the morphism p is Stein, we see that the direct image $\mathbb{R}p_*$ of the complex (2.7) is pure $n + 1$ -dimensional over $\mathcal{O}_S((z))$ [23, 25]. That is, only $n + 1$ -th cohomology group is non-trivial. Thus we set

$$(2.10) \quad \mathcal{H}_F := \mathbb{R}^{n+1}p_*(\Omega_{Z/S}^*(z), D_F) = p_*(\Omega_{Z/S}^{n+1}(z))/D_F(p_*(\Omega_{Z/S}^n(z)))$$

which is naturally a $\mathcal{O}_S((z))$ -module. The cohomology class of $s \in p_*(\Omega_{Z/S}^n(z))$ is denoted by $[s]$.

We also define the k -th filter $(F^k\Omega_{Z/S}^*(z), D_F)$ of the complex (2.7) by $F^k\Omega_{Z/S}^*(z) := z^{-k}\Omega_{Z/S}^*[[z]]$ for $k \in \mathbb{Z}$.⁶ Their cohomology groups are also pure $n + 1$ -dimensional over $\mathcal{O}_S[[z]]$ ([ibid]), and we set

$$(2.11) \quad \mathcal{H}_F^{(k)} := \mathbb{R}^{n+1}p_*(F^k\Omega_{Z/S}^*(z), D_F) = z^{-k}p_*(\Omega_{Z/S}^{n+1}[[z]])/D_F(z^{-k}p_*(\Omega_{Z/S}^n[[z]]))$$

which is naturally a $\mathcal{O}_S[[z]]$ -module. Then, we see the following semi-infinitely filtered structure on \mathcal{H}_F together with the Gauss-Manin connection and the higher residue pairings K_F ([S3, 21])

PROPOSITION 2.1. 1. The inclusion $F^k\Omega_{Z/S}^*(z) \subset \Omega_{Z/S}^*(z)$ for $k \in \mathbb{Z}$ induces an embedding (injection) $\mathcal{H}_F^{(k)} \subset \mathcal{H}_F$ as $\mathcal{O}_S[[z]]$ -submodule so that we obtain an increasing filtration in \mathcal{H}_F :

$$(2.12) \quad \dots \subset \mathcal{H}_F^{(-1)} \subset \mathcal{H}_F^{(0)} \subset \mathcal{H}_F^{(1)} \subset \dots \subset \mathcal{H}_F$$

with the regularity property:

$$(2.13) \quad \cup_{k \in \mathbb{Z}} \mathcal{H}_F^{(k)} = \mathcal{H}_F \quad \text{and} \quad \cap_{k \in \mathbb{Z}} \mathcal{H}_F^{(k)} = \{0\}.$$

The multiplication by z induces $\mathcal{O}_S[[z]]$ -isomorphisms between the filters:

$$(2.14) \quad z : \mathcal{H}_F^{(k+1)} \simeq \mathcal{H}_F^{(k)}.$$

That is, z is homogeneous of degree -1 with respect to the filtration (see Footnote 1).

The graded pieces of the filtration are given by the exact sequence:

$$(2.15) \quad 0 \rightarrow \mathcal{H}_F^{(k-1)} \rightarrow \mathcal{H}_F^{(k)} \xrightarrow{r_F^{(k)}} \Omega_F \rightarrow 0$$

where

$$\Omega_F := p_*(\Omega_{Z/S}^{n+1}/dF \wedge \Omega_{Z/S}^n) \simeq p_*(\Omega_{Z/S}^{n+1})/p_*(dF) \wedge p_*(\Omega_{Z/S}^n)$$

⁶In the early formulations [21, 23, 25] of the filtered de-Rham module \mathcal{H}_F , instead of the variable z as in the present paper, we used the variable δ_1^{-1} in order to indicate that it is a pseudo differential operator (or Fourier integral operator) associated to the inverse of the primitive derivation $\delta_1 = \partial_1$. We switch the view point and regard \mathcal{H}_F as a module over the space of the Fourier dual variable z of the primitive derivation δ_1 (this is the same for the variable t in [11]). For this reason, we have an unfortunate opposite sign convention between the index k for the filtration $F^k\Omega_{Z/S}^*(z)$ and the exponent $-k$ of z in $z^{-k}\Omega_{Z/S}^*[[z]]$.

is an \mathcal{O}_S -free module of rank μ , and $r^{(k)}$ is the natural quotient morphism modulo $dF \wedge \Omega_{Z/S}^n$ on the leading term of $z^{-k} p_*(\Omega_{Z/S}^{n+1}[[z]])$ to Ω_F such that $r_F^{(k)} = r_F^{(0)} \circ z^{-k}$.

2. There is a covariant differentiations acting on \mathcal{H}_F , called the *Gauss-Manin connection*:

$$\nabla : \mathcal{T}_{\mathbf{A}_z^1 \times S} \times \mathcal{H}_F \longrightarrow \mathcal{H}_F$$

given, for $\zeta = [\phi(\underline{x}, z) dx_1 \wedge \cdots \wedge dx_n] \in \mathcal{H}_F$, by

$$(2.16) \quad \begin{aligned} \nabla_{\frac{d}{dz}} \zeta &= \left[\left(\frac{d\phi}{dz} - z^{-2} F \phi \right) dx_1 \wedge \cdots \wedge dx_n \right], \\ \nabla_v \zeta &= \left[\left(\tilde{v} \phi + z^{-1} \tilde{v} F \phi \right) dx_1 \wedge \cdots \wedge dx_n \right] \end{aligned}$$

for $\frac{d}{dz} \in \mathcal{T}_{\mathbf{A}_z^1}$ and $v \in \mathcal{T}_S$, such that

- i) The connection ∇ is integrable: $\nabla^2 = 0$. In particular, $\nabla_{\frac{d}{dz}}$ and ∇_v commute each other.
- ii) The connection ∇ over \mathcal{T}_S satisfies the transversality: $\nabla_{\mathcal{T}_S} : \mathcal{H}_F^{(k-1)} \rightarrow \mathcal{H}_F^{(k)}$.
- iii) The covariant action of ∇_{∂_z} has pole of order 2 at $z = 0$ (that is, $z^2 \nabla_{\partial_z} = -\nabla_{\partial_{z^{-1}}}$ preserves the filtration (2.12)).
- iv) The operator $\nabla_{z\partial_z + E}$ on \mathcal{H}_F preserves the filtration (2.12).

3. There is a skew- $\mathcal{O}_S((z))$ -bilinear form, called the higher residue pairing:

$$(2.17) \quad K_F : \mathcal{H}_F \times \mathcal{H}_F \longrightarrow \mathcal{O}_S((z))$$

such that

- i) For $\zeta_1, \zeta_2 \in \mathcal{H}_F$, we have $K_F(\zeta_1, \zeta_2) = K_F(\zeta_2, \zeta_1)^*$.
- ii) For $\zeta_1, \zeta_2 \in \mathcal{H}_F$ and $P \in \mathcal{O}_S((z))$, we have

$$PK_F(\zeta_1, \zeta_2) = K_F(P\zeta_1, \zeta_2) = K(\zeta_1, P^*\zeta_2).$$

where we denote by the upper script of $*$ an \mathcal{O}_S -involution of $\mathcal{O}_S((z))$ defined by $z \mapsto -z$.

- iii) For $\zeta_1, \zeta_2 \in \mathcal{H}_F$ and $v \in \mathcal{T}_S$, we have

$$(2.18) \quad \partial_v K_F(\zeta_1, \zeta_2) = K_F(\nabla_v \zeta_1, \zeta_2) + K_F(\zeta_1, \nabla_v \zeta_2).$$

- iv) For $\zeta_1, \zeta_2 \in \mathcal{H}_F$, we have

$$(2.19) \quad z\partial_z K_F(\zeta_1, \zeta_2) = K_F(\nabla_{z\partial_z} \zeta_1, \zeta_2) + K_F(\zeta_1, \nabla_{z\partial_z} \zeta_2).$$

v) The restriction $K_F|_{\mathcal{H}_F^{(0)} \times \mathcal{H}_F^{(0)}}$ gives an $\mathcal{O}_S[[z]]$ -linear map $\mathcal{H}_F^{(0)} \times \mathcal{H}_F^{(0)} \rightarrow z^{n+1}\mathcal{O}_S[[z]]$ which makes the following commutative diagram:

$$(2.20) \quad \begin{array}{ccc} \mathcal{H}_F^{(0)} \times \mathcal{H}_F^{(0)} & \xrightarrow{K_F} & z^{n+1}\mathcal{O}_S[[z]] \\ r_F^{(0)} \times r_F^{(0)} \downarrow & & \downarrow \text{mod } z^n\mathcal{O}_S[[z]] \\ \Omega_F \times \Omega_F & \xrightarrow{Res} & \mathcal{O}_S, \end{array}$$

where the *Res* in the second row is given by the residue symbol ([6])

$$\text{Res} \left[\frac{\phi_1(\underline{x}, z) \phi_2(\underline{x}, z) dx_1 \wedge \cdots \wedge dx_{n+1}}{\partial_{x_1} F, \cdots, \partial_{x_{n+1}} F} \right],$$

for $\zeta_i = [\phi_i(\underline{x}, z) dx_1 \wedge \cdots \wedge dx_{n+1}] \in \mathcal{H}_F^{(0)}$ ($i = 1, 2$).

Note. 1. The explicit formula (2.16) implies that the leading terms of the covariant actions of ∇_v for $v \in \mathcal{T}_S$ and of $\nabla_{z\partial_z}$ with respect to the filtration on \mathcal{H}_F are given by the following commutative diagrams, respectively.

$$(2.21) \quad \begin{array}{ccc} \mathcal{H}_F^{(k-1)} & \xrightarrow{\nabla_v} & \mathcal{H}_F^{(k)} \\ r_F^{(k-1)} \downarrow & & \downarrow r_F^{(k)} \\ \Omega_F & \xrightarrow{\bar{v}F|_{\mathcal{C}_F}} & \Omega_F \end{array} \quad \begin{array}{ccc} \mathcal{H}_F^{(k-1)} & \xrightarrow{\nabla_{z\partial_z}} & \mathcal{H}_F^{(k)} \\ r_F^{(k-1)} \downarrow & & \downarrow r_F^{(k)} \\ \Omega_F & \xrightarrow{-F|_{\mathcal{C}_F}} & \Omega_F \end{array}$$

2. The following elementary fact shall be used later in the uniqueness of the evolution map $e^{(f-F)/z}$ in §4. The restriction $\mathcal{H}_F \otimes_{\mathcal{O}_S} \mathcal{O}_S/m$ (m : the maximal ideal of $\mathcal{O}_{S,0}$) of \mathcal{H}_F at the origin $0 \in S$ is naturally isomorphic to \mathcal{H}_f since the homomorphism of $\mathcal{H}_F \rightarrow \mathcal{H}_F \otimes_{\mathcal{O}_S} \mathcal{O}_S/m$ is naturally induced from the pull-back morphism $(\Omega_{Z/S}^*(z), D_F) \rightarrow (\Omega_X^*(z), D_f)$ of the complexes induced from the inclusion morphism $X \subset Z$.

3. For a later use in §6, we give a sketch of a proof of 2. iv). We prove only that the filter $\mathcal{H}_F^{(0)}$ is preserved by $\nabla_{z\partial_z+E}$. Let \tilde{E} be a lifting of E . Then by Definition 2.2, there exists $g_i \in \Gamma(Z, \mathcal{O}_Z)$ ($i = 1, \dots, n$) such that $F - \tilde{E}F = \sum_i g_i \partial_{x_i} F$. Therefore, for any $[s] \in \mathcal{H}_F^{(0)}$, we calculate

$$(2.22) \quad \begin{aligned} \nabla_{z\partial_z+E}[s] &= [(z\partial_z - F/z + \tilde{E} + \tilde{E}F/z)s] \\ &= [(z\partial_z + \tilde{E} - (\sum_i g_i \partial_i F)/z)s] = [(z\partial_z + \tilde{E})s + (d_{Z/S} - z^{-1}D_F)(s')] = \\ &= [(z\partial_z + \tilde{E})s + d_{Z/S}s'] \in \mathcal{H}_F^{(0)} \end{aligned}$$

where $s' = (s/(dt_1 \wedge \dots \wedge t_{n+1})) \sum_i (-1)^i g_i dt_1 \wedge \dots \wedge \check{d}t_i \wedge \dots \wedge dt_{n+1} \in p_*(\Omega_{Z/S}^n)$.

REMARK 2.3. Let us give an analytic interpretation of the modules (2.10) and (2.11).

Choose $\alpha \in \Omega_X^{n+1}\{z\} \subset \Omega_X^{n+1}[[z]]$ and consider a z -dependent family of oscillatory integrals for $z \in \mathbb{C}^*$:

$$(2.23) \quad \int_{\Gamma_z} e^{f/z} \alpha$$

for a continuous family $\{\Gamma_z\}$ of relative cycles $\Gamma_z \in H_{n+1}(X, \text{Re}(f/z) \ll 0)$. If $\alpha = D_z(\beta)$ for $\beta \in \Omega_X^n\{z\}$ then

$$(2.24) \quad \int_{\Gamma_z} e^{f/z} \alpha = \int_{\Gamma_z} e^{f/z} (z d\beta + df \wedge \beta) = z \int_{\Gamma_z} d(e^{f/z} \beta) = 0,$$

and we obtain a perfect pairing:

$$(2.25) \quad H_{n+1}(X, \text{Re}(f/z) \ll 0) \times \Omega_X^{n+1}/D_z \Omega_X^n \rightarrow \mathbb{C}$$

[1]. In the present paper, we do not use such analytic theory, however we may regard the splitting factor $\hat{\mathcal{B}}[[z]]$ in (3.5) as the space of representatives of integrants for ‘‘formal oscillatory integrals’’ $\int_{\Gamma_z} e^{f/z} \alpha$ out of their equivalence classes.

REMARK 2.4. The notation \mathcal{H}_F and $\mathcal{H}_F^{(k)}$ were already used in [23] for slightly different objects. Therefore, let us clarify the relationship between them and (2.10) and (2.11).

First, let us give a heuristic interpretation of the variable z and the complex D_F . We regard F as a family of functions parametrized by S so that we have a fibration $\varphi = (F, p) : Z \rightarrow \mathbf{A}^1 \times S$. Then the equation $D_F = 0$ together with the relation (2.5) should “imply” $z = -\left(\frac{dZ/S}{dF}\right)^{-1} \equiv \frac{\partial}{\partial t_1} \bmod dF$.

Therefore, in [21, 22], the variable z was denoted by δ_1^{-1} in order to identify it with the pseudo differential operator $\partial_{t_1}^{-1}$ where t_1 the coordinate for the range of F so that the module is defined on the relative-cotangent space of the range space of F . In the present paper, we regard it simply a formal variable.

The modules \mathcal{H}_f and $\mathcal{H}_f^{(k)}$ were sometimes called the module of semi-infinite Hodge structure for the vanishing cycles of f . However, in the present paper, we regard them as the spaces of integrands for oscillatory integrands with respect to the exponential factor $\exp(f/z)$ (see Remark 2.3). It is of our main interests in §4 how they evolve to the spaces of integrands for the oscillatory integrals with respect to the exponential factor $\exp(F/z)$ for an unfolding F of f .

3. Section and opposite filtration of \mathcal{H}_F

Let the setting be as in §2, i.e. F is an unfolding define on $Z \subset X \times S$ of a function f having an isolated critical point at $0 \in X \subset \mathbb{C}^{n+1}$, where S is the parameter space for the unfolding. We analyze the filtered \mathcal{O}_S -module \mathcal{H}_F by introducing concepts of sections and opposite filtrations (see Remark 3.1.).

DEFINITION 3.1. 1. A \mathcal{O}_S -submodule \mathcal{B} of $\mathcal{H}_F^{(0)}$ is called a *section* of $\mathcal{H}_F^{(0)}$, if the restriction of the projection $r_F^{(0)}$ (recall (2.15)) induces an \mathcal{O}_S -isomorphism $\mathcal{B} \simeq \Omega_F$. In view of Proposition 2.1, the condition on \mathcal{B} is equivalent to that we have natural $\mathcal{O}_S[[z]]$ -isomorphisms

$$(3.1) \quad \mathcal{B}[[z]] \simeq \mathcal{H}_F^{(0)} \quad \text{and} \quad \mathcal{B}((z)) \simeq \mathcal{H}_F.$$

The set of all sections of $\mathcal{H}_F^{(0)}$ is denoted by $\mathcal{M}_{s,F}$.

2. A \mathcal{O}_S -submodule \mathcal{L} of \mathcal{H}_F is called an *opposite filtration* of \mathcal{H}_F , if it satisfies (1) the \mathcal{O}_S -splitting condition: $\mathcal{H}_F = \mathcal{H}_F^{(0)} \oplus \mathcal{L}$ and (2) the opposite condition: $z\mathcal{L} \supset \mathcal{L}$. The set of all opposite filtrations of \mathcal{H}_F is denoted by $\mathcal{M}_{o,F}$.

For a later application in the proof of Theorem 6.1, below we describe the isomorphism (3.1) for the case of the trivial unfolding down to earth.

Consider any μ -dimensional \mathbb{C} -subspace $\hat{\mathcal{B}}$ of $\Gamma(X, \Omega_X^{n+1})[[z]]$ such that it is isomorphic to Ω_f by the projection $\Omega_X^{n+1}[[z]] \bmod (dF \wedge \Omega_X^n[[z]] + z\Omega_X^{n+1}[[z]]) \rightarrow \Omega_f$. In particular, the condition implies

$$(3.2) \quad \hat{\mathcal{B}} \cap z\Gamma(X, \Omega_X^{n+1})[[z]] = 0.$$

I.e. the projection of $\hat{\mathcal{B}}$ into $\Gamma(X, \Omega_X^{n+1})$ is an embedding. This implies that we get a sequence of injective morphisms: $\hat{\mathcal{B}}[z]/(z^i) \rightarrow \Gamma(X, \Omega_X^{n+1})[z]/(z^i)$ for $i = 1, 2, 3, \dots$ (since a graded piece of the morphisms is equivalent to the embedding for $i =$

1). Taking the inverse limit of the sequence, we see that $\hat{\mathcal{B}}[[z]]$ is embedded into $\Gamma(X, \Omega_X^{n+1})[[z]]$. The condition on $\hat{\mathcal{B}}$ implies further the splitting:

$$(3.3) \quad \Gamma(X, \Omega_X^{n+1}) = \text{the projection image of } \hat{\mathcal{B}} \text{ in } \Gamma(X, \Omega_X^{n+1}) \oplus \Gamma(X, df \wedge \Omega_X^n).$$

Then, repeating the similar arguments above, we can show the following splitting.

$$(3.4) \quad \Gamma(X, \Omega_X^{n+1})[[z]] = \hat{\mathcal{B}}[[z]] \oplus \Gamma(X, df \wedge \Omega_X^n)[[z]].$$

Note also that the modules $\hat{\mathcal{B}}[[z]]$ and $\hat{\mathcal{B}}((z))$ are equal to $\hat{\mathcal{B}} \otimes_{\mathbb{C}} \mathbb{C}[[z]]$ and $\hat{\mathcal{B}} \otimes_{\mathbb{C}} \mathbb{C}((z))$, respectively, due to the fact $\dim_{\mathbb{C}} \hat{\mathcal{B}} = \mu < \infty$, and are free of rank μ over $\mathbb{C}[[z]]$ and $\mathbb{C}((z))$, respectively.

PROPOSITION 3.1. Let $\hat{\mathcal{B}} \subset \Gamma(X, \Omega_X^{n+1})[[z]]$ be as above. Then, there are splittings

$$(3.5) \quad \begin{aligned} \Gamma(X, \Omega_X^{n+1})((z)) &= \hat{\mathcal{B}}((z)) \oplus D_f(\Gamma(X, \Omega_X^n)((z))) \\ z^{-k}\Gamma(X, \Omega_X^{n+1})[[z]] &= z^{-k}\hat{\mathcal{B}}[[z]] \oplus D_f(z^{-k}\Gamma(X, \Omega_X^n)[[z]]). \end{aligned}$$

PROOF. It is sufficient to show the second formula.

1. Since each factor in the RHS of (3.5) is embedded in the LHS, we consider the natural homomorphism from the RHS to the LHS. We first show that it is surjective.

Pick an element α from the k th filter $z^{-k}\Gamma(X, \Omega_X^{n+1})[[z]]$. Due to (3.4), we find $v^1 \in z^{-k}\hat{\mathcal{B}}[[z]]$ and $\beta^1 \in z^{-k}\Gamma(X, \Omega_X^n)[[z]]$ such that

$$\alpha = df \wedge \beta^1 + v^1.$$

Setting $\alpha^1 := \alpha$ and $\alpha^2 := -d\beta^1$, we obtain

$$\alpha^1 = v^1 + D_f\beta^1 + z\alpha^2,$$

Since $\alpha^2 \in z^{-k}\Gamma(X, \Omega_X^{n+1})[[z]]$, we continue inductively

$$\alpha^2 = v^2 + D_f\beta^2 + z\alpha^3,$$

$$\alpha^3 = v^3 + D_f\beta^3 + z\alpha^4,$$

...

where $\alpha^i := -d\beta^{i-1} \in z^{-k}\Gamma(X, \Omega_X^{n+1})[[z]]$ for $i \geq 2$. Multiplying the i -th line by z^{i-1} and summing up all lines, we obtain the decomposition:

$$(3.6) \quad \alpha = v + D_f\beta.$$

where $v = \sum_{i=1}^{\infty} v^i z^{i-1} \in z^{-k}\hat{\mathcal{B}}[[z]]$ and $\beta = \sum_{i=1}^{\infty} \beta^i z^{i-1} \in z^{-k}\Omega_X^n[[z]]$. \square

2. Next, let us show that $\hat{\mathcal{B}}[[z]] \cap D_f(\Gamma(X, \Omega_X^n)[[z]]) = \{0\}$.

Suppose the contrary, and there exist $v = \sum_{i=0}^{\infty} v_i z^i \in \hat{\mathcal{B}}[[z]]$ ($v_i \in \hat{\mathcal{B}}$), $v_i = \sum_{j=0}^{\infty} v_{ij} z^j$ ($v_{ij} \in \Gamma(X, \Omega_X^{n+1})$) and $\alpha = \sum_{i=0}^{\infty} \alpha_i z^i \in \Gamma(X, \Omega_X^n)[[z]]$ ($\alpha_i \in \Gamma(X, \Omega_X^n)$) such that $v = D_f\alpha$. Let us show by induction on $i \in \mathbb{Z}_{\geq 0}$ the following assertion.

(*)_i: the vanishing $v_i = 0$ and existences of $\beta_i \in \Gamma(X, \Omega_X^{n-1})$ such that $\alpha_i = df \wedge \beta_i + d\beta_{i-1}$.

Let us write the equality $v = D_f\alpha$ term-wisely: $\sum_{k+l=i} v_{kl} = d\alpha_{i-1} + df \wedge \alpha_i$ for $i \in \mathbb{Z}_{\geq 0}$, where $\alpha_{-1} = 0$. The equality for the case $i = 0$ turns to be $v_{00} = df \wedge \alpha_0$. Then, in view of the splitting (3.3), we see $v_{00} = 0$ and $df \wedge \alpha_0 = 0$. Then,

(3.2) implies $v_0 = 0$, and applying Poincare Lemma, we see $\alpha_0 = df \wedge \beta_0$ for $\beta_0 \in \Gamma(X, \Omega_X^{n-1})$. That is $(*)_0$ is shown by setting $\beta_{-1} = 0$. Suppose that $(*)_i$ is shown for $i \in \mathbb{Z}_{\geq 0}$. Then we have

$$v_{i+1,0} = d(df \wedge \beta_i + d\beta_{i-1}) + df \wedge \alpha_{i+1} = df \wedge (-d\beta_i + \alpha_{i+1}).$$

Applying again the splitting (3.3) to this equality, we get $v_{i+1,0} = 0$ and hence $v_{i+1} = 0$, and get the existence of $\beta_{i+1} \in \Gamma(X, \Omega_X^{n-1})$ such that $\alpha_{i+1} = df \wedge \beta_{i+1} + d\beta_i$, implying $(*)_{i+1}$. The induction shows that $v = 0$ and $\alpha = D_f \beta$. \square

Above 1. and 2. together give the proof of the splittings (3.5). \square

In later sections, we shall abuse the notion of sections and opposite filtrations for the cases when the coefficient is no-longer \mathcal{O}_S but is either an Artinian ring $\mathcal{O}_{S,0}/m^{k+1}$ or the formal rings $\check{\mathcal{O}}_{S,0}$.

REMARK 3.2. Present definitions of section and opposite filtration and the coming Definition 6.1 of good pair are “weaker” copies of those Definition 2.16 and Definition 4.13 in [11] in the sense that we do not assume for them the metric condition discussed in the following Proposition 3.4. except we explicitly require it, whereas in [11] the metric condition is assumed for every section, opposite filtration and good pair. Nevertheless, we shall see that the most of statements in [11] are recovered.

PROPOSITION 3.2. There is a natural bijection:

$$(3.7) \quad \mathcal{M}_{o,F} \simeq \mathcal{M}_{s,F}, \quad \mathcal{L} \mapsto \mathcal{B}(\mathcal{L}) := \mathcal{H}_F^{(0)} \cap z\mathcal{L}.$$

The inverse correspondence is given by $\mathcal{B} \mapsto \mathcal{L}(\mathcal{B}) := z^{-1}\mathcal{B}[z^{-1}]$.

PROOF. (c.f. [11] 2.16) Let $\mathcal{L} \in \mathcal{M}_{o,F}$ be given. Multiplying the decomposition in Definition 3.1. 2. (1) by z^k for $k \in \mathbb{Z}$, we obtain $\mathcal{H}_F = \mathcal{H}_F^{(-k)} \oplus z^k\mathcal{L}$, and hence $\mathcal{H}_F^{(-k)} = \mathcal{H}_F^{(-k-1)} \oplus (\mathcal{H}_F^{(-k)} \cap z^{k+1}\mathcal{L})$. For $k = 0$, we obtain $\mathcal{H}_F^{(0)} = \mathcal{H}_F^{(-1)} \oplus \mathcal{B}(\mathcal{L})$, implying the isomorphism $\mathcal{B}(\mathcal{L}) \simeq \Omega_F$ due to (2.15) for $k = 0$.

Conversely, let $\mathcal{B} \in \mathcal{M}_{s,F}$ be given. Then, applying (2.14) and (2.15) repeatedly, we obtain $\mathcal{H}_F^{(k)} = \mathcal{H}_F^{(0)} \oplus \bigoplus_{i=1}^k z^{-i}\mathcal{B}$ for all $k \in \mathbb{Z}_{>0}$. In view of (2.13), we see that $\mathcal{L}(\mathcal{B}) := z^{-1}\mathcal{B}[z^{-1}] \in \mathcal{M}_{o,F}$.

By the construction and the property Definition 3.1. 2. (2) of \mathcal{L} , it is clear that $\mathcal{L}(\mathcal{B}(\mathcal{L})) \subset \mathcal{L}$. Since we have the decompositions $\mathcal{H}_F = \mathcal{H}_F^{(0)} \oplus \mathcal{L} = \mathcal{H}_F^{(0)} \oplus \mathcal{L}(\mathcal{B}(\mathcal{L}))$, we get $\mathcal{L}(\mathcal{B}(\mathcal{L})) = \mathcal{L}$.

Conversely, by the construction, it is clear that $\mathcal{B}(\mathcal{L}(\mathcal{B})) \supset \mathcal{B}$. Since we have the decompositions $\mathcal{H}_F^{(0)} = \mathcal{H}_F^{(-1)} \oplus \mathcal{B} = \mathcal{H}_F^{(-1)} \oplus \mathcal{B}(\mathcal{L}(\mathcal{B}))$, we get $\mathcal{B}(\mathcal{L}(\mathcal{B})) = \mathcal{B}$. \square

PROPOSITION 3.3. Let $\mathcal{B} \in \mathcal{M}_{s,F}$ and $\mathcal{L} \in \mathcal{M}_{o,F}$ correspond to each other by (3.7). Then the following conditions a), a') and b) are equivalent. We shall call these conditions the $z\partial_z$ -stability condition.

$$\text{a) } \nabla_{z\partial_z}\mathcal{B} \subset \mathcal{B} + z^{-1}\mathcal{B}, \text{ a') } \nabla_{z\partial_z}\mathcal{B} \subset \mathcal{B}[z^{-1}], \text{ b) } \nabla_{z\partial_z}\mathcal{L} \subset \mathcal{L}.$$

PROOF. Clearly, a) implies a').

a') \Rightarrow b): $\nabla_{z\partial_z}\mathcal{L} = \nabla_{z\partial_z}z^{-1}\mathcal{B}[z^{-1}] \subset (\nabla_{z\partial_z}\mathcal{B})z^{-1}\mathbb{C}[z^{-1}] + \mathcal{B}z\partial_z(z^{-1}\mathbb{C}[z^{-1}]) \subset \mathcal{B}z^{-1}\mathbb{C}[z^{-1}] = \mathcal{L}$.

b) \Rightarrow a): $\nabla_{z\partial_z}\mathcal{B} = \nabla_{z\partial_z}(\mathcal{H}_F^{(0)} \cap z\mathcal{L}) \subset \nabla_{z\partial_z}(\mathcal{H}_F^{(0)}) \cap \nabla_{z\partial_z}(z\mathcal{L}) \subset z^{-1}\mathcal{H}_F^{(0)} \cap z\mathcal{L} = \mathcal{B} + z^{-1}\mathcal{B}$. \square

PROPOSITION 3.4. Let $\mathcal{B} \in \mathcal{M}_{s,F}$ and $\mathcal{L} \in \mathcal{M}_{o,F}$ correspond to each other by (3.7). Then the following two conditions c) and d) are equivalent. We shall call these conditions the metric condition.

c) The set of values $K_F(\mathcal{B}, \mathcal{B}) := \{K_F(v_1, v_2) \mid v_1, v_2 \in \mathcal{B}\}$ is contained in $z^n \mathcal{O}_S$ (c.f. (2.20)).

d) The set of values $K_F(\mathcal{L}, \mathcal{L}) := \{K_F(l_1, l_2) \mid l_1, l_2 \in \mathcal{L}\}$ is contained in $z^{-2+n} \mathcal{O}_S[z^{-1}]$.

PROOF. c) \Rightarrow d): $K_F(\mathcal{L}, \mathcal{L}) \subset K_F(z^{-1}\mathcal{B}[z^{-1}], z^{-1}\mathcal{B}[z^{-1}]) \subset K_F(\mathcal{B}, \mathcal{B})z^{-2}\mathcal{O}_S[z^{-1}] \subset z^{-2+n}\mathcal{O}_S[z^{-1}]$.

d) \Rightarrow c): $K_F(\mathcal{B}, \mathcal{B}) \subset K_F(\mathcal{H}_F^{(0)} \cap z\mathcal{L}, \mathcal{H}_F^{(0)} \cap z\mathcal{L}) \subset K_F(\mathcal{H}_F^{(0)}, \mathcal{H}_F^{(0)}) \cap K_F(z\mathcal{L}, z\mathcal{L}) \subset z^n \mathcal{O}_S[[z]] \cap z^n \mathcal{O}_S[z^{-1}] = \mathcal{O}_S \cdot z^n$. \square

REMARK 3.3. According to Givental [5], we may regard \mathcal{L} as a \mathcal{O}_S -linear Lagrangean subspace by setting the symplectic form

$$\omega : \mathcal{H}_F \times \mathcal{H}_F \rightarrow \mathcal{O}_S, \quad \omega(s_1, s_2) := \text{Res}_{z=0}(z^{-n} K_F(s_1, s_2)).$$

Then, the initial property (2.20) implies $\omega(\mathcal{H}_F^{(0)}, \mathcal{H}_F^{(0)}) = 0$, and the metric condition in Proposition 3.4 is equivalent to $\omega(\mathcal{L}, \mathcal{L}) = 0$. That is, $\mathcal{H}_F = \mathcal{H}_F^{(0)} \oplus \mathcal{L}$ is a *holonomic decomposition* in the sense of Mikio Sato.

4. Formal analysis on deformation parameter space

From this paragraph, we take a perturbative approach with respect to the deformation variable \underline{t} in S at 0, as was done in [11]. Namely, we embed \mathcal{H}_F into its formal completion $\check{\mathcal{H}}_F$ with respect to the deformation variable \underline{t} . We show that the formal (infinite) bundle $\check{\mathcal{H}}_F$ over the formal space $\text{Spf } \check{\mathcal{O}}_{S,0}$ is trivialized (w.r.t. the Gauss-Manin connection ∇) by the evolution of \mathcal{H}_f by the oscillatory integral factor $\exp(f-F)/z$ (Proposition 4.1). In particular, sections \mathcal{B} in $\mathcal{H}_f^{(0)}$ and opposite filtrations \mathcal{L} in \mathcal{H}_f evolve to flat sections and flat opposite filtrations in $\check{\mathcal{H}}_F$, respectively. This fact leads to the construction of formal primitive forms (without metric) in the next section.

Consider any unfolding F defined over the frame (Z, S, p, F) in 2.1. Let $0 \in S$ be the base point and let m be the maximal ideal of the local ring $\mathcal{O}_{S,0}$. Then m^{k+1} for $k \in \mathbb{Z}_{\geq 0}$ form the k -th formal neighborhood of $0 \in \mathcal{O}_{S,0}$ with respect to the m -adic topology on $\mathcal{O}_{S,0}$. Set $\check{\mathcal{O}}_{S,0} := \varprojlim_k \mathcal{O}_{S,0}/m^{k+1}$ and $\check{\mathcal{F}} := \mathcal{F} \hat{\otimes}_{\mathcal{O}_S} \check{\mathcal{O}}_{S,0} := \varprojlim_k \mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{O}_{S,0}/m^{k+1}$ for a \mathcal{O}_S -module \mathcal{F} . Using a local coordinate system $\underline{t} = (t_1, \dots, t_m)$ of S at the origin 0, we sometimes denote $\check{\mathcal{O}}_{S,0}$ also by $\mathbb{C}[[\underline{t}]]$ and $\check{\mathcal{F}}$ by $\mathcal{F}[[\underline{t}]]$.

Recall Proposition 2.1 that the \mathcal{O}_S -module \mathcal{H}_F admits the Gauss-Manin connection ∇ and the higher residue pairings K_F . We extend them to $\check{\mathcal{H}}_F := \varprojlim_k \mathcal{H}_F \otimes_{\mathcal{O}_S} \mathcal{O}_{S,0}/m^{k+1}$ as follows. Namely, one easily shows (see [11], Lemma 4.3) that ∇ induces a connection

$$(4.1) \quad \check{\nabla} : \mathcal{T}_{\mathbb{A}_z^1 \times S} \times \check{\mathcal{H}}_F \longrightarrow \check{\mathcal{H}}_F$$

defined by the projective limit of $\mathcal{T}_S \times \mathcal{H}_F/m^{k+1}\mathcal{H}_F \rightarrow \mathcal{H}_F/m^k\mathcal{H}_F$ and $\mathcal{T}_{\mathbb{A}_z^1} \times \mathcal{H}_F/m^{k+1}\mathcal{H}_F \rightarrow \mathcal{H}_F/m^{k+1}\mathcal{H}_F$ for $k \in \mathbb{Z}_{>0}$, which we call again the Gauss-Manin

connection and denote again by ∇ . The higher residue pairings K_F on \mathcal{H}_F also induces an $(\mathcal{O}_S((z)))^\vee$ -skew-symmetric pairing:

$$(4.2) \quad \check{K}_F : \check{\mathcal{H}}_F \times \check{\mathcal{H}}_F \longrightarrow (\mathcal{O}_S((z)))^\vee.$$

They satisfy the properties parallel to those listed in 2. and 3. of Proposition 2.1, whose detailed formulations and verifications are left to the reader.

REMARK 4.1. Let us consider the formal unfolding \check{F} (associated with an unfolding F) defined on the formal frame $(\check{Z}, \check{S}, p, \check{F})$ (that is, $\check{S} = \mathrm{Spf}(\check{\mathcal{O}}_{S,0})$ is the formal neighborhood of the origin $0 \in S$, \check{Z} is a formal neighborhood of $X \times \{0\}$ in $\mathbb{C}^n \times \check{S}$ and \check{F} is the formal power series expansion of F with respect to a local coordinate system \underline{t} of S at 0. Detailed descriptions are left to the reader). Then, analogous to §2, one can define the $\check{\mathcal{O}}_S((z))$ -module $\mathcal{H}_{\check{F}} := p_*(\Omega_{\check{Z}/\check{S}}^n((z))/D_{\check{F}}(p_*(\Omega_{\check{Z}/\check{S}}^{n-1}((z))))$ which admits increasing filtration $\mathcal{H}_{\check{F}}^{(k)}$ satisfying the properties parallel to (2.10),(2.11),(2.14) and (2.15) (here, we note that $\Omega_{\check{F}} = \check{\Omega}_F = \Omega_F \otimes_{\mathcal{O}_S} \check{\mathcal{O}}_{S,0}$). In particular, we have isomorphisms $\mathcal{H}_{\check{F}} \simeq \mathcal{H}_F \otimes_{\mathcal{O}_S((z))} \check{\mathcal{O}}_{S,0}((z))$, which admits the filtration $\mathcal{H}_{\check{F}}^{(k)} \simeq \mathcal{H}_F^{(k)} \otimes_{\mathcal{O}_S[[z]]} \check{\mathcal{O}}_{S,0}[[z]] (= \check{\mathcal{H}}_F^{(k)})$ for $k \in \mathbb{Z}$ (regarded as the formal version of \mathcal{H}_F).

However, due to the “non-commutativity” $\check{\mathcal{O}}_{S,0}((z)) = \mathbb{C}[[\underline{t}]((z))] \not\subseteq \mathcal{O}_{S,0}((z))^\vee = \mathbb{C}((z))[[\underline{t}]]$ (since the RHS is a formal power series in (\underline{t}) of coefficients in Laurent series of z whose order of poles may not necessarily be bounded, e.g. $\sum_{k=0}^{\infty} z^{-k} t^k = z/(z-t) = -\sum_{m=1}^{\infty} t^{-m} z^m$ is in the RHS but not in the LHS), we have

$$(4.3) \quad \mathcal{H}_{F,0} \subsetneq \mathcal{H}_{\check{F}} \subsetneq \check{\mathcal{H}}_F.$$

So, even if the module $\check{\mathcal{H}}_F$ has the increasing sequence of $\mathbb{C}[[z]][[\underline{t}]] = \mathbb{C}[[z, \underline{t}]]$ -submodules $\check{\mathcal{H}}_F^{(k)} \simeq \mathcal{H}_F^{(k)}$ ($k \in \mathbb{Z}$), the union $\cup_{k \in \mathbb{Z}} \check{\mathcal{H}}_F^{(k)}$ covers only the submodule $\mathcal{H}_{\check{F}}$ consisting of elements which admit bounded Laurent series expansion in z with coefficients in $\mathbb{C}[[\underline{t}]]$ (w.r.t. a fixed basis), but does not cover the full-module $\check{\mathcal{H}}_F$ which consists of power series in \underline{t} whose coefficient in a monomial in \underline{t} is a Laurent series in z but the order of poles in z may grow to infinity as the order of the monomial increases. Thus, the analogy of the regularity (2.13) does not hold for $\check{\mathcal{H}}_F$.

We now compare the given unfolding F over a frame (Z, S, p) with the trivial unfolding f . Precisely, recall the projection $\pi_X : Z \rightarrow X$ (§2), and depending on that, we define the trivial unfolding

$$F_0 := \pi_X^{-1}(f)$$

over the frame (Z, S, p) so that $F - F_0 \in m\mathcal{O}_Z$. Then following Propositions 4.1 and 4.2, giving a trivialization of $\check{\mathcal{H}}_F$ and an opposite filtration of $\check{\mathcal{H}}_F$, are the key results in the formal analysis in the present paper.

PROPOSITION 4.1. ([11], Lemma/Definition 4.7) 1. Consider a correspondence

$$(4.4) \quad e^{(f-F)/z} : \mathcal{H}_f \longrightarrow \check{\mathcal{H}}_F.$$

given by associating, for each $[s] \in \mathcal{H}_f$, the projective system of elements

$$(4.5) \quad \left[\sum_{0 \leq l \leq k} \frac{(F_0 - F)^l}{z^l l!} \pi_X^{-1}(s) \right] \in \mathcal{H}_F \bmod m^{k+1} \mathcal{H}_F \quad \text{for } k \in \mathbb{Z}_{>0},$$

where $s \in p_{0*}(\Omega_X^n((z)))$ is any representative of $[s]$. Then, it defines a well-defined $\mathbb{C}((z))$ -module injective homomorphism, which we shall call the evolution map.⁷

2. The composition of the evolution map $e^{(f-F)/z}$ (4.4) with the restriction map $\check{\mathcal{H}}_F \rightarrow \check{\mathcal{H}}_F \otimes_{\check{\mathcal{O}}_{S,0}} (\check{\mathcal{O}}_{S,0}/m) \simeq \mathcal{H}_f$ (c.f. §2 Note. 2.) is the identity map on \mathcal{H}_f .

3. The image $e^{(f-F)/z}[s] \in \check{\mathcal{H}}_F$ for $[s] \in \mathcal{H}_f$ of the evolution map (4.4) is horizontal (flat) with respect to the Gauss-Manin connection on $\check{\mathcal{H}}_F$ over \mathcal{T}_S -direction (but not $\mathcal{T}_{\mathbb{A}_1^2}$ -direction). Precisely,

$$(4.6) \quad \ker(\nabla_{\mathcal{T}_S}) = e^{(f-F)/z}(\mathcal{H}_f).$$

4. The evolution map (4.4) induces an $\mathbb{C}((z)) \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0} := \varprojlim_k \mathbb{C}((z)) \otimes_{\mathbb{C}} \mathcal{O}_{S,0}/m^{k+1}$ -isomorphism:

$$(4.7) \quad \mathcal{H}_f \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0} \simeq \check{\mathcal{H}}_F$$

$$\mathcal{H}_f \otimes_{\mathbb{C}} \mathcal{O}_S/m^{k+1} \xrightarrow{e^{(F_0-F)/z}} \mathcal{H}_F/m^{k+1}\mathcal{H}_F, \quad [s] \otimes [r_k] \mapsto e^{(F_0-F)/z} \pi_X^{-1}(s) r_k \bmod m^{k+1}\mathcal{H}_F.$$

The inverse map of (4.7) is given by multiplying $e^{(F-F_0)/z} \bmod m^{k+1}$ at each k -th neighborhood, and is denoted by $e^{(F-f)/z}$. We shall call the extended isomorphism (4.7) also the evolution map.

5. (i) The evolution isomorphism (4.7) is equivariant with the Gauss-Manin connection $\nabla_{z\partial_z}$ (2.16), where its action in LHS is defined to be commutative with the action of $\check{\mathcal{O}}_{S,0}$.

(ii) The evolution isomorphism (4.7) is equivariant with the action of \mathcal{T}_S , where $v \in \mathcal{T}_S$ acts on LHS as the derivation on the coefficient $\check{\mathcal{O}}_{S,0}$ and on RHS via Gauss-Manin connection ∇_v (2.16).

PROOF. The proof given below is inspired from the oscillatory integral view point (see Remark 2.3).

1. First, remark that the k -th neighborhood $\mathcal{H}_F \otimes_{\mathcal{O}_S} \mathcal{O}_S/m^{k+1} = \mathcal{H}_F \bmod m^{k+1}\mathcal{H}_F$ in the RHS of (4.5) has an expression

$$(4.8) \quad \left(p_*(\Omega_{Z/S}^{n+1}((z))) \bmod m^{k+1} p_*(\Omega_{Z/S}^{n+1}((z))) \right) / \left(D_F(p_*(\Omega_{Z/S}^n((z))) \bmod m^{k+1} D_F(p_*(\Omega_{Z/S}^n((z)))) \right)$$

that is, quotient out by the co-boundary operator D_F and quotient out by the higher terms m^{k+1} are commutative to each other (this fact can be seen by the right-exactness of the tensor product $\otimes_{\mathcal{O}_S} \mathcal{O}_S/m^{k+1}$). Then, in order to show the well-definedness of the map (4.4), we need to show the following two facts.

1) For a given class $[s] \in \mathcal{H}_f$, fix its representative $s \in p_{0*}(\Omega_X^n((z)))$. Since the order of the pole in z of the expression in the bracket of (4.5) for each $k \in \mathbb{Z}_{>0}$ is bounded by k +the order of the pole in z of s , the expression defines an element in the numerator of (4.8). Further more, the sequence of the elements for $k \in \mathbb{Z}_{>0}$ forms a projective system, since we have $e^{(F_0-F)/z} = \sum_{l=0}^{\infty} \frac{(F_0-F)^l}{z^l l!}$ such that $\frac{(F_0-F)^l}{z^l l!} \in m^l p_* \mathcal{O}_Z((z))$ (recall Definition 2.1, 2. so that $F_0 - F \in m \cdot p_* \mathcal{O}_Z$).

⁷We use the notation $e^{(f-F)/z}$ instead of $e^{(F_0-F)/z}$, since we shall see immediately in Corollary 4.1 that the correspondence does not depend on the choice of the projection π_X and, hence, of F_0 .

2) The class of the image in (4.5) depends not on the choice of the representative $s \in p_{0*}(\Omega_X^n((z)))$ but only on the class $[s] \in \mathcal{H}_f$. (*Proof.* Consider s' such that $s' - s = D_f(r)$ for $r \in p_{0*}(\Omega_X^{n-1}((z)))$. Then, in the module of the numerator of (4.8), we calculate the difference of the representatives of $e^{(f-F)/z}[s'] - e^{(f-F)/z}[s]$:

$$(4.9) \quad e^{(f-F)/z}\pi_X^{-1}(D_f(r)) \equiv e^{(F_0-F)/z}D_{F_0}(\pi_X^{-1}(r)) \equiv D_F(e^{(F_0-F)/z}(\pi_X^{-1}(r))) \pmod{m^{k+1}}$$

where in the RHS, the order of poles in z is again bounded by k so that the expression defines an element in the denominator of (4.8), i.e. $e^{(f-F)/z}[s'] - e^{(f-F)/z}[s] = 0$. So the evolution map is well-defined. It commutes with the action of $\mathbb{C}((z))$ since it is so for π_X^{-1} and in k th neighborhood (4.5).

2. This is trivial due to the initial term of the explicit expression (4.5) of the evolution map.

3. We first show that RHS of (4.6) is included in LHS. This follows formally by the calculation:

$$\begin{aligned} (\nabla_v(e^{(f-F)/z}[s]) \pmod{m^k} &\equiv \nabla_v(e^{(F_0-F)/z}\pi_X^{-1}(s) \pmod{m^{k+1}}) \\ &\equiv [(L_{\tilde{v}} + \tilde{v}F/z)(e^{(F_0-F)/z}\pi_X^{-1}(s))] \pmod{m^k} \end{aligned}$$

where $L_{\tilde{v}}$ means the Lie derivative with respect to a lifting \tilde{v} of v and the lifting \tilde{v} may be chosen such that $\tilde{v}(\pi_X^{-1}(\mathcal{O}_X)) = 0$. Then, we have $L_{\tilde{v}}(e^{(F_0-F)/z}\pi_X^{-1}(s)) = (-\tilde{v}F/z)e^{(F_0-F)/z}\pi_X^{-1}(s)$ so that the two terms in the RHS of the above expression cancel to each other and the sum vanishes.

The opposite inclusion is shown as follows. Let ζ be any element of LHS. Consider the evolution $e^{(f-F)/z}(\zeta|_0)$ where $\zeta|_0 = \zeta \pmod{m} \in \mathcal{H}_f$ is the restriction of ζ at the origin $0 \in S$ (recall §2 Note.2.). Then the uniqueness of the solution of the linear equation $\nabla_{\mathcal{T}_S}\zeta = 0$ with the same initial value $\zeta|_0$ (recall 2. of present Proposition) implies the coincidence of ζ with $e^{(f-F)/z}(\zeta|_0) \in$ RHS.

4. The evolution map $\mathcal{H}_f \rightarrow \check{\mathcal{H}}_F$ (4.4) is a $\mathbb{C}((z))$ -morphism, and the correspondence (4.7) naturally extends the map to a $(\mathcal{O}_{S,0}((z)))^\vee$ -morphism $\mathcal{H}_f \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0} \rightarrow \check{\mathcal{H}}_F$ between the free modules. Its residue map modulo the maximal ideal $\mathbb{C}((T))\tilde{m}$ is the natural identity $\mathcal{H}_f \xrightarrow{\sim} \mathcal{H}_F|_{\underline{t}=0} = \mathcal{H}_f$. Then, due to Nakayama's lemma, the morphism is isomorphic.

It is formal to see that the formal multiplication by $e^{(F-F_0)/z}$ gives the inversion map to $e^{(f-F)/z}$.

5. (i) In the cochain complex level, the evolution morphism satisfies the following commutativity:

$$e^{(f-F)/z} \cdot (z\partial_z - f/z) = (z\partial_z - F/z) \cdot e^{(f-F)/z}$$

which is shown by a direct calculation. Then, in view of the formula (2.16) for $\nabla_{z\partial_z}$, we obtain the commutativity in the cohomological level.

(ii) This follows from the flatness 3. of present Proposition and the fact that the evolution morphism is an $(\mathcal{O}_{S,0}((z)))^\vee$ -linear morphism, since $\nabla_v e^{(f-F)/z}([s] \hat{\otimes} a) = \nabla_v(e^{(f-F)/z}[s]) \hat{\otimes} a = \nabla_v(e^{(f-F)/z}[s]) \hat{\otimes} a + (e^{(f-F)/z}[s]) \hat{\otimes} \nabla_v a = (e^{(f-F)/z}[s]) \hat{\otimes} \nabla_v a = e^{(f-F)/z}([s] \hat{\otimes} \nabla_v a)$ for $[s] \in \mathcal{H}_f$ and $a \in (\mathcal{O}_{S,0}((z)))^\vee$.

This completes a proof of Proposition 4.1 \square

COROLLARY 4.1. The evolution map $e^{(f-F)/z}$ (4.4) is defined independent of a choice of the projection π_X and, hence, of $F_0 = \pi_X^{-1}f$. This justifies the use of f in the notation $e^{(f-F)/z}$.

PROOF. For any element $[s] \in \mathcal{H}_f$, the image $e^{(f-F)/z}[s]$ is uniquely characterized by the two conditions: (1) it satisfies the system of linear equations $\nabla_v(e^{(f-F)/z}[s]) = 0$ for all $v \in \mathcal{T}_S$ (Proposition 4.1, 3.), and (2) its initial value $(e^{(f-F)/z}[s])|_0$ at the origin $0 \in S$ is equal to the original $[s] \in \mathcal{H}_f$ (Proposition 4.1) due to the uniqueness of the solution of the system of linear partial differential equation. Obviously, these two conditions do not depend on the choice of the projection π_S . So the solution $e^{(f-F)/z}[s]$, i.e. the image of $[s]$ by the evolution map, does not depend on the choice of π_X . \square

REMARK 4.2. The Proposition 4.1 says that the ‘‘infinite dimensional bundle’’ \mathcal{H}_f over a point $\{0\}$ flatly evolves to the trivial bundle $\check{\mathcal{H}}_F$ over the infinitesimal neighborhood $\check{S} = Spf(\check{\mathcal{O}}_{S,0})$ of $\{0\}$.

Note that the lattice $\mathcal{H}_f^{(0)} \subset \mathcal{H}_f$ does not evolve into the lattice $\check{\mathcal{H}}_F^{(0)} \subset \check{\mathcal{H}}_F$ by the formal evolution map (4.4) (since the oscillatory integral factor $e^{(f-F)/z}$ has pole in z). However, as we see in the next Proposition that an opposite filtration \mathcal{L} of \mathcal{H}_f evolves to a formal opposite filtration in $\check{\mathcal{H}}_F$.

We now want to apply this formal evolution map to the sections and opposite filtrations on \mathcal{H}_f .

Consider a section $\mathcal{B} \in \mathcal{M}_{s,f}$ and a opposite filtration $\mathcal{L} \in \mathcal{M}_{o,f}$ related as (3.7) for the trivial unfolding f . We first formally extend them as $\mathcal{B} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0} := \lim_{\leftarrow} \mathcal{B} \otimes_{\mathbb{C}} \mathcal{O}_S/m^k \subset \mathcal{H}_f^{(0)} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0}$ (actually, this is equal to $\mathcal{B} \otimes_{\mathbb{C}} \check{\mathcal{O}}_{S,0}$ due to the finite dimensionality of \mathcal{B}) and $\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0} := \lim_{\leftarrow} \mathcal{L} \otimes_{\mathbb{C}} \mathcal{O}_S/m^k \subset \mathcal{H}_f \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0}$. Then, we consider the images $e^{(f-F)/z}(\mathcal{B} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$ and $e^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$ in $\check{\mathcal{H}}_F$ by the evolution morphism (4.4), and call them the formal section and the formal opposite filtration on $\check{\mathcal{H}}_F$ associated with \mathcal{B} and \mathcal{L} , respectively.

Let us first justify the namings: the formal section and the formal opposite filtration.

PROPOSITION 4.2. ([11] Lemma 4.9) Let $\mathcal{L} \in \mathcal{M}_{o,f}$ be an opposite filtration. Then,

1. The formal evolution image $e^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$ of $\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0}$ gives a formal opposite filtration of $\check{\mathcal{H}}_F$ in the sense that it defines a splitting of $\check{\mathcal{O}}_{S,0}$ -module

$$(4.10) \quad \check{\mathcal{H}}_F = \check{\mathcal{H}}_F^{(0)} \oplus e^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$$

satisfying the formal opposite condition:

$$(4.11) \quad z^{-1}(e^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})) \subset e^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0}).$$

2. The intersection $\check{\mathcal{H}}_F^{(0)} \cap ze^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$ is a formal section of $\check{\mathcal{H}}_F^{(0)}$ in the sense that the natural projection to the z -graded piece, (2.15) for $k = 0$, induces an isomorphism:

$$(4.12) \quad \check{\mathcal{H}}_F^{(0)} \cap ze^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0}) \simeq \check{\mathcal{H}}_F^{(0)} / \check{\mathcal{H}}_F^{(-1)} \quad (\simeq \check{\Omega}_F).$$

PROOF. 1. Let a \mathbb{C} -basis of $\mathcal{B} := \mathcal{H}_f^{(0)} \cap z\mathcal{L}$ be presented by forms $\phi_j \in p_{0*}(\Omega_X^n)[[z]]$ ($j = 1, \dots, \mu$). Consider the \mathcal{O}_S -morphism $r_F^{(0)} : \mathcal{B}_F := \bigoplus_{j=1}^{\mu} \mathcal{O}_S[\pi_X^{-1}(\phi_j)] \rightarrow \Omega_F$ given in (2.15). Its restriction $r_f^{(0)}$ to the origin $0 \in S$ (i.e. to tensor with $\mathcal{O}_{S,0}/m$) is the isomorphism $\mathcal{B} \simeq \Omega_f$ (since \mathcal{B} is a section). Then by Nakayama’s

lemma, the morphism is isomorphic at the stalks at the origin. Then the \mathcal{O}_S -isomorphism between coherent sheaves at the origin extends to an \mathcal{O}_S -isomorphism in a neighborhood of the origin of S . This implies that $\mathcal{B}_F = \bigoplus_{j=1}^{\mu} \mathcal{O}_S[\pi_X^{-1}(\phi_j)]$ ($\simeq \mathcal{B} \otimes_{\mathbb{C}} \mathcal{O}_S$ depending on ϕ_j 's) on the neighborhood is a section of $\mathcal{H}_F^{(0)}$. So, $z^{-1}\mathcal{B}_F[z^{-1}]$ is an opposite filtration of \mathcal{H}_F in a neighborhood of $0 \in S$. In particular, we get a splitting

$$(4.13) \quad \mathcal{H}_F \otimes_{\mathcal{O}_S} \mathcal{O}_S/m^{k+1} = \left(\mathcal{H}_F^{(0)} \otimes_{\mathcal{O}_S} \mathcal{O}_S/m^{k+1} \right) \oplus \left(z^{-1}\mathcal{B}_F[z^{-1}] \otimes_{\mathcal{O}_S} \mathcal{O}_S/m^{k+1} \right) \quad k \in \mathbb{Z}_{>0}$$

On the other hand, recall the isomorphism: $e^{(f-F)/z} : \mathcal{B}((z)) \otimes_{\mathbb{C}} \mathcal{O}_S/m^{k+1} \rightarrow \mathcal{H}_F \otimes_{\mathcal{O}_S} \mathcal{O}_S/m^{k+1}$ (Proposition 4.1). Let us show that the composition of this isomorphism with the embedding $\mathcal{L} \otimes_{\mathbb{C}} \mathcal{O}_S/m^{k+1} (= z^{-1}\mathcal{B}[z^{-1}] \otimes_{\mathbb{C}} \mathcal{O}_S/m^{k+1}) \subset \mathcal{B}((z)) \otimes_{\mathbb{C}} \mathcal{O}_S/m^{k+1}$ and with the projection to the second factor $z^{-1}\mathcal{B}_F[z^{-1}] \otimes_{\mathcal{O}_S} \mathcal{O}_S/m^{k+1}$ of (4.13) (i.e. quotient by the submodule $\mathcal{H}_F^{(0)} \otimes_{\mathcal{O}_S} \mathcal{O}_S/m^{k+1}$, see the following Remark 4.3) is an $\mathcal{O}_S/m^{k+1}[z^{-1}]$ -isomorphism

$$(4.14) \quad e^{(f-F)/z} \mathcal{L} \otimes_{\mathbb{C}} \mathcal{O}_S/m^{k+1} \simeq z^{-1}\mathcal{B}_F[z^{-1}] \otimes_{\mathcal{O}_S} \mathcal{O}_S/m^{k+1}.$$

(Proof. We note the facts

- i) The map is an $\mathcal{O}_S/m^{k+1}[z^{-1}]$ -homomorphism.
- ii) Both hand sides of the map are $\mathcal{O}_S/m^{k+1}[z^{-1}]$ -free modules.
- iii) The map induces an isomorphism if one consider the map modulo the maximal ideal m .

Then it is clear that the map itself is an $\mathcal{O}_S/m^{k+1}[z^{-1}]$ -isomorphism. \square)

Thus, the image $e^{(f-F)/z}(\mathcal{L} \otimes_{\mathbb{C}} \mathcal{O}_S/m^{k+1})$ in $\mathcal{H}_F \otimes_{\mathcal{O}_S} \mathcal{O}_S/m^{k+1}$ defines a splitting

$$\mathcal{H}_F \otimes_{\mathcal{O}_S} \mathcal{O}_S/m^{k+1} = (\mathcal{H}_F^{(0)} \otimes_{\mathcal{O}_S} \mathcal{O}_S/m^{k+1}) \oplus e^{(f-F)/z}(\mathcal{L} \otimes_{\mathbb{C}} \mathcal{O}_S/m^{k+1}).$$

Clearly these splittings for $k \in \mathbb{Z}_{\geq 0}$ form a projective system, and hence taking the limit for $k \rightarrow \infty$, we obtain the splitting (4.10). The opposite condition (i.e. z^{-1} -stability) on $e^{(f-F)/z}(\mathcal{L} \otimes_{\mathbb{C}} \mathcal{O}_S/m^{k+1})$ obviously follows from that on \mathcal{L} .

2. The formula (4.12) follow from the same logic in the first half of Proposition 3.2. \square

REMARK 4.3. This quotient process in the proof of (4.14) is essentially necessary, since the evolution map $e^{(f-F)/z}$ may not bring elements of $\mathcal{L} \otimes_{\mathbb{C}} \mathcal{O}_S/m^{k+1}$ into $z^{-1}\mathcal{B}_F[z^{-1}] \otimes_{\mathcal{O}_S} \mathcal{O}_S/m^{k+1}$. This is caused, since, in order to express elements $(F_0 - F)^l \phi_j$'s in a linear combination of elements in $\phi_j \otimes \mathcal{O}_S/m^{k+1}$, one need to calculate modulo the image of D_F , which may create some terms having coefficients of positive powers in z .

REMARK 4.4. The evolution image $e^{(f-F)/z}(\mathcal{B} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$ of the section \mathcal{B} for \mathcal{L} does not give the section $\check{\mathcal{H}}_F^{(0)} \cap ze^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$ for the opposite filtration $e^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$, since the image as a submodule of the direct sum (4.10) may not be contained in the first factor, but just truncating its second factor part, it does give the section $\check{\mathcal{H}}_F^{(0)} \cap ze^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$. Actually, on the contrary, the evolution image of \mathcal{B} contains highly interesting information in the second factor.

REMARK 4.5. The naive relation $z^{-1}\mathcal{B}[z^{-1}] = \mathcal{L}$ inside \mathcal{H}_f does not extends to the formal level, but we only have $z^{-1}(\check{\mathcal{H}}_F^{(0)} \cap ze^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0}))[z^{-1}] \subsetneq \neq$

$e^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0})$ inside $\check{\mathcal{H}}_F$ due to the phenomenon in Remark 4.1. However, the opposite filtration $e^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0})$ can be recovered from the section $\check{\mathcal{H}}_F^{(0)} \cap ze^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0})$ by a ‘‘formal completion’’ (see §6 proof of Theorem 6.1).

The flatness of the formal evolution morphism $e^{(f-F)/z}$ (Proposition 4.1, 2.) implies the following important stability property of the formal section and the formal opposite filtration.

PROPOSITION 4.3. ([11] Lemma 4.11.) The Gauss-Manin connection ∇ on $\check{\mathcal{H}}_F$ over \mathcal{T}_S preserves the formal opposite filtration given in Proposition 4.2. That is,

$$\nabla_{\mathcal{T}_S}(e^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0})) \subset e^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0}).$$

We call this property \underline{t} -stability of the formal opposite filtration.

PROOF. Let an element s of $e^{(f-F)/z}(\mathcal{B}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0})$ or $e^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0})$ be given by a sequence $s_k = \sum_i e^{(f-F)/z} a_i \otimes b_i \bmod m^{k+1}$ for $a_i \in \mathcal{B}$ or $\in \mathcal{L}$ and $b_i \in \mathcal{O}_{S,0}/m^{k+1}$ for $k \in \mathbb{Z}_{\geq 0}$. Since $e^{(f-F)/z} a_i$ is flat (4.6), we have $\nabla_v s \equiv \sum_i e^{(f-F)/z} a_i \otimes \partial_{\bar{v}} b_i \bmod m^k$ where the RHS converges to an element in $e^{(f-F)/z}(\mathcal{B}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0})$ or in $e^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0})$, respectively. \square

The $z\partial_z$ -stability property and the metric property of sections \mathcal{B} and opposite filtrations \mathcal{L} in \mathcal{H}_f are inherited to formal section and formal opposite filtrations in $\check{\mathcal{H}}_F$ as follows.

PROPOSITION 4.4. Let $\mathcal{L} \in \mathcal{M}_{o,f}$ be an opposite filtration associated with a section $\mathcal{B} \in \mathcal{M}_{s,f}$.

1. If \mathcal{L} is $z\partial_z$ -stable (recall Proposition 3.3), then the formal opposite filtration $e^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0})$ is formally $z\partial_z$ -stable in the sense:

$$(4.15) \quad \nabla_{z\partial_z}(e^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0})) \subset e^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0}).$$

2. If \mathcal{L} satisfies the metric condition (see Proposition 3.4), then the formal opposite filtration $e^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0})$ satisfies the formal metric condition:

$$(4.16) \quad \check{K}_F(e^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0}), e^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0})) \subset z^{-2+n}(\mathcal{O}_S[z^{-1}])^\vee$$

and the formal section $\check{\mathcal{H}}_F^{(0)} \cap ze^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0})$ satisfies the formal purity condition:

$$(4.17) \quad \check{K}_F(\check{\mathcal{H}}_F^{(0)} \cap ze^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0}), \check{\mathcal{H}}_F^{(0)} \cap ze^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0})) \subset z^n \check{\mathcal{O}}_{S,0}.$$

PROOF. Those follow formally from the definitions.

1. Let us express an element of $\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0}$ by the limit of a sequence of elements of the form $\sum_i a_i^k \otimes b_i^k \bmod m^{k+1}$ for $a_i^k \in \mathcal{L}$ and $b_i^k \in \mathcal{O}_S$ for $k = 1, 2, 3, \dots$. Then, using the first line of (2.16),

$$\begin{aligned} & \nabla_{z\partial_z}(e^{(f-F)/z} \sum_i a_i \otimes b_i \bmod m^{k+1}) \\ &= e^{(f-F)/z}(\sum_i -((f-F)/z)a_i \otimes b_i + (z\partial_z a_i) \otimes b_i - z^{-1}F a_i \otimes b_i) \bmod m^{k+1} \\ &= e^{(f-F)/z}(\sum_i -(f/z)a_i + (z\partial_z a_i)) \otimes b_i \bmod m^{k+1} \\ &= e^{(f-F)/z}(\nabla_{z\partial_z} a_i) \otimes b_i \bmod m^{k+1} \end{aligned}$$

By assumption, $\nabla_{z\partial_z} a_i \in \mathcal{L}$ so that the last line belongs to $e^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0})$, implying (4.15).

2. Since K_F is \mathcal{O}_S -bilinear (2.17), it induces a \mathcal{O}_S/m^{k+1} -bilinear form such that $K_F(e^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0}) \bmod m^{k+1}, e^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0}) \bmod m^{k+1}) \subset z^{-2+n}\mathcal{O}_{S,0}[z^{-1}] \bmod m^{k+1}$. This implies clearly (4.16). Hence, combining with (2.20), we get (4.17). \square

REMARK 4.6. Suppose the opposite filtration \mathcal{L} in \mathcal{H}_f satisfy the metric condition. Then, Proposition 4.4, 2. implies $\omega(e^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0}), e^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0})) = 0$. Hence the decomposition (4.10) is holonomic decomposition of $\check{\mathcal{H}}_F$, since $\omega(\check{\mathcal{H}}_F^{(0)}, \check{\mathcal{H}}_F^{(0)}) = 0$ is automatic due to the formal version of (2.20) (see Remark 3.3). But we shall not use this fact explicitly.

REMARK 4.7. As in the latter half of Remark 4.1, one should note again that the range $z^{-2+n}(\mathcal{O}_S[z^{-1}])^\vee$ (see (4.16)) of the residue pairings \check{K}_F on $e^{(f-F)/z}(\mathcal{L}\hat{\otimes}_{\mathbb{C}}\check{\mathcal{O}}_{S,0})$ is strictly larger than $z^{-2+n}\check{\mathcal{O}}_{S,0}[z^{-1}]$. That is, the order of the pole in z of the coefficient of t^k may increase infinitely as $k \rightarrow \infty$.

5. Primitive forms without metric structure

In this section, we introduce the notion of (formal or non-formal) primitive forms and then (formal or non-formal) primitive forms without metric structure.

Let us recall the definition of primitive forms [21] in slightly reformed way.⁸

DEFINITION 5.1. **Primitive Form.** An element $\zeta \in \Gamma(S, \mathcal{H}_F^{(0)})$ is called a *primitive form* if it satisfies the following conditions (P0)-(P4).

(P0) (Primitivity): The covariant derivations of ζ induces an \mathcal{O}_S -isomorphism:

$$z\nabla\zeta : \mathcal{T}_S \rightarrow \mathcal{H}_F^{(0)}/\mathcal{H}_F^{(-1)} \simeq \Omega_F, \quad v \mapsto z\nabla_v\zeta \bmod \mathcal{H}_F^{(-1)}.$$

(P1) (Triviality in primitive direction): The ζ is trivial in the primitive direction ∂_1 . That is,

$$z\nabla_{\partial_1}\zeta = \zeta.$$

(P2) (Purity of higher residue pairings): The covariant derivations of ζ has the following purity w.r.t. the residues.

$$K_F(\nabla_{\mathcal{T}_S}\zeta, \nabla_{\mathcal{T}_S}\zeta) \subset z^{n-1}\mathcal{O}_S.$$

(P3) (Holonomicity): The second covariant derivatives of ζ satisfies the followings.

$$\begin{aligned} K_F(\nabla_{\mathcal{T}_S}\nabla_{\mathcal{T}_S}\zeta, \nabla_{\mathcal{T}_S}\zeta) &\in z^{n-2}\mathcal{O}_S + z^{n-1}\mathcal{O}_S \\ K_F(\nabla_{z\partial_z}\nabla_{\mathcal{T}_S}\zeta, \nabla_{\mathcal{T}_S}\zeta) &\in z^{n-2}\mathcal{O}_S + z^{n-1}\mathcal{O}_S. \end{aligned}$$

(P4) (Homogeneity): There is a constant $r \in \mathbb{C}$ such that

$$\nabla_{z\partial_z+E}\zeta = r\zeta.$$

where E is the Euler vector field (2.5).

⁸Compared with [21], in the present formulation as in [25] and [11], we included the descendent parameter z explicitly so that the base space $\mathbb{C} \times S$ of the family of vanishing cycles is one dimensional higher than the base space S of the unfolding in [21]. This makes essentially two changes of the formulation: 1. F_1 action is changed to $z\partial_z$ -action in the second condition of (P3) and in (P4). 2. To kill the redundancy of the base space, we need to add one more relation (P1) (which was missing in [25] and [11], see [29]), formulated as the triviality of primitive forms in primitive direction (actually, this formula (P1) is the reason of calling ∂_1 primitive derivation or primitive direction).

REMARK 5.2. The definition is analytic in S direction. However, it is still formal in z -variable.

REMARK 5.3. The condition (P0) is equivalent to that the image of covariant differentiations of ζ

$$\mathcal{B}_\zeta := \text{Im}\left(z\nabla\zeta : \mathcal{T}_S \rightarrow \mathcal{H}_F^{(0)}\right) = z\nabla_{\mathcal{T}_S}\zeta$$

form a section of $\mathcal{H}_F^{(0)}$. Using this notation, the condition (P2) is equivalent to the following

$$(P2)^* \text{ (Orthogonality)} \quad K_F(\mathcal{B}_\zeta, \mathcal{B}_\zeta) \subset z^{n+1}\mathcal{O}_S.$$

Then, under the assumptions (P0) and (P2) (\simeq (P2)*), the condition (P3) is equivalent to

(P3)* (Holonomicity)*: The second covariant derivatives of ζ satisfies the followings.

$$\begin{aligned} \nabla_{\mathcal{T}_S}\nabla_{\mathcal{T}_S}\zeta &\subset z^{-2}\mathcal{B}_\zeta + z^{-1}\mathcal{B}_\zeta \\ \nabla_{z\partial_z}\nabla_{\mathcal{T}_S}\zeta &\subset z^{-2}\mathcal{B}_\zeta + z^{-1}\mathcal{B}_\zeta. \end{aligned}$$

We now, removing the metric condition (P2) from the conditions (P0)-(P4) for a primitive form, introduce a concept of *primitive forms without metric condition* (see Remark 5.4 for this terminology), where we replace the condition (P3) by (P3)* (see Remark 5.1 for this replacement).

DEFINITION 5.4. **Primitive Form without metric structure.** An element $\zeta \in \Gamma(S, \mathcal{H}_F^{(0)})$ is called a *primitive form without metric structure* if it satisfies the conditions (P0), (P1), (P3)* and (P4) but may not necessarily satisfy (P2)*.

Next, let us recall the definition of formal primitive forms [11], Definition 4.6.

DEFINITION 5.5. **Formal Primitive Form.** An element $\zeta \in \check{\mathcal{H}}_F^{(0)}$ is called a *formal primitive form*, if it satisfies (P0) $^\vee$, (P1), (P2) $^\vee$, (P3) $^\vee$ and (P4), where

(P0) $^\vee$: Replace \mathcal{T}_S and Ω_F in (P0) by $\check{\mathcal{T}}_S = \mathcal{T}_S \otimes_{\mathcal{O}_S} \check{\mathcal{O}}_{S,0}$ and by $\check{\Omega}_F = \Omega_F \otimes_{\mathcal{O}_S} \check{\mathcal{O}}_{S,0}$, respectively.

(P2) $^\vee$: Replace \mathcal{T}_S , K_F and \mathcal{O}_S in (P2) by $\check{\mathcal{T}}_S$, \check{K}_F and by $\check{\mathcal{O}}_{S,0}$, respectively.

(P3) $^\vee$: Replace \mathcal{T}_S , K_F and \mathcal{O}_S in (P3) by $\check{\mathcal{T}}_S$, \check{K}_F and by $\check{\mathcal{O}}_{S,0}$, respectively.

Finally, let us introduce a concept of formal primitive forms without metric structure

DEFINITION 5.6. **Formal Primitive Form without metric structure.** An element $\zeta \in \check{\mathcal{H}}_F^{(0)}$ is called a *formal primitive form without metric structure*, if it satisfies (P0) $^\vee$, (P1), (P3)* $^\vee$ and (P4) but may not necessarily satisfy (P2) $^\vee$, where

(P3)* $^\vee$: Replace \mathcal{T}_S and \mathcal{B}_ζ in (P3)* by $\check{\mathcal{T}}_S$ and by $\check{\mathcal{B}}_\zeta$, respectively, where we set the formal section by

$$\check{\mathcal{B}}_\zeta := \text{Im}(z\nabla\zeta : \check{\mathcal{T}}_S \rightarrow \check{\mathcal{H}}_F^{(0)}) = z\nabla_{\check{\mathcal{T}}_S}\zeta.$$

REMARK 5.7. An obvious remark is that the embedding (4.3) send a primitive form ζ without metric structure in $\Gamma(S, \mathcal{H}_F^{(0)})$ to a formal primitive form ζ without metric structure in $\check{\mathcal{H}}_F^{(0)}$. Is the converse, i.e. all formal primitive forms without metric structure are induced from some analytic primitive form without metric structure, true?

REMARK 5.8. Note that the condition (P3) alone cannot be equivalent to (P3)* or (P3)** , but the combination (P2) and (P3) is equivalent to the combination (P2) and (P3)* (see Proof 4. of Theorem 6.3).

REMARK 5.9. In Definitions 5.4 and 5.6, the terminology “without metric structure” is used in a weak sense “with or without metric structure”. That is, a primitive form ζ without metric structure may eventually satisfy the metric condition (P2) so that it can actually be a primitive form, unless it is explicitly stated that ζ does not satisfy (P2).

6. Construction of formal primitive forms with or without metric structure

In order to formulate the first main result, Theorem 6.1, of the present paper, we prepare a notion of a *good pair* ([11] Def. 4.15, see Remark 6.1).

DEFINITION 6.1. A pair (\mathcal{L}, ζ_0) of a $z\partial_z$ -stable opposite filtration \mathcal{L} of \mathcal{H}_f and an element $\zeta_0 \in \mathcal{B} := \mathcal{H}_f^{(0)} \cap z\mathcal{L}$ is called a *good pair* (without metric condition) if it satisfies the following (1) and (2).

(1) (primitivity) The image of ζ_0 in Ω_f by $r_f^{(0)}$ (see (2.15)) generates Ω_f over \mathcal{O}_{C_f} .

(2) (homogeneity) There is a constant $r \in \mathbb{C}$ such that $\nabla_{z\partial_z} \zeta_0 - r\zeta_0 \in \mathcal{L}$.

If \mathcal{L} satisfies further the metric condition (c.f. Proposition 3.4), the pair (\mathcal{L}, ζ_0) is called a *good pair with metric condition* (see Remark 3.1).

REMARK 6.2. According to the correction of Definition of primitive forms (see 2. of the Footnote 3 in §5), we correct the definition of a good pair from [11]. Namely, we took ζ_0 from $\mathcal{H}_f^{(0)}$ in [11], but, in the present paper, we assume more strongly that ζ_0 belongs to $\mathcal{B} := \mathcal{H}_f^{(0)} \cap z\mathcal{L}$.

Theorem 4.16 in [11] gives a construction of formal primitive forms from a good pair with metric condition. We confirm in the following Theorem that the same construction works for the case without metric condition.

THEOREM 6.3. Let F be a universal unfolding of a function f with an isolated critical point at the origin (See Definitions 2.1 and 2.2). Then, we have bijections

(6.1)

$$\{\text{good pairs } (\mathcal{L}, \zeta_0) \text{ on } \mathcal{H}_f\} \simeq \{\text{formal primitive forms in } \check{\mathcal{H}}_F^{(0)} \text{ without metric structure}\}$$

(6.2)

$$\{\text{good pairs } (\mathcal{L}, \zeta_0) \text{ on } \mathcal{H}_f \text{ with metric structure}\} \simeq \{\text{formal primitive forms in } \check{\mathcal{H}}_F^{(0)}\}$$

The correspondence $(\mathcal{L}, \zeta_0) \mapsto \zeta_+$ from LHS to RHS is given as follows: apply the evolution morphism to ζ_0 so that we obtain an element $e^{(f-F)/z}\zeta_0$ in $\check{\mathcal{H}}_F$. Then, apply the splitting (4.10) to this element so that we have the decomposition:

$$(6.3) \quad e^{(f-F)/z}\zeta_0 = \zeta_+ + \zeta_-$$

for $\zeta_+ \in \check{\mathcal{H}}_F^{(0)}$ and $\zeta_- \in e^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$.

PROOF. In the following, the proof is divided in Steps 1-13. Compared with the proof for Theorem 4.16 in [11], since we added a new axiom, say (P1) (∂_1 -triviality), on the primitive form side, and an additional condition $\zeta_0 \in z\mathcal{L}$ on the primitive element side, we need some additional arguments, in particular, the Steps 2, and 13. We added also Proposition 6.1 to complete the Step 13 of the proof. In order to clarify the role of metric conditions, we take caution on the treatment of the conditions (P3) and (P3)* and distinguish the places where and when the metric condition is necessary or not.

Before coming to the proof of Theorem 6.3, we prepare below a proposition on $\zeta_- \in e^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$ for a use in the proof. Actually, beyond this use, ζ_- itself is of interest and we shall come back to ζ_- in §10.

PROPOSITION 6.1. Let (\mathcal{L}, ζ_0) be a good pair, and consider the decomposition (6.3). Set

$$(6.4) \quad \zeta_- = e^{(f-F)/z} \left(\sum_{j=1}^{\infty} z^{-j} u_j \right)$$

for $\sum_{j=1}^{\infty} z^{-j} u_j \in \mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0}$ with $u_j \in \mathcal{B} \otimes_{\mathbb{C}} \check{\mathcal{O}}_{S,0}$ ($j \in \mathbb{Z}_{>0}$).⁹ Then,

$$(6.5) \quad u_j \in \mathcal{B} \otimes_{\mathbb{C}} m^j \check{\mathcal{O}}_{S,0}.$$

PROOF. Applying the inversion $e^{(F-f)/z}$ of the evolution map to the formula (6.3), we obtain

$$(6.6) \quad \zeta_0 = e^{(F-f)/z} \zeta_+ + \sum_{j=1}^{\infty} z^{-j} u_j$$

In $\mathcal{H}_f \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0}$, we rewrite: $\sum_{j=1}^{\infty} z^{-j} u_j = \zeta_0 - e^{(F-f)/z} \zeta_+$. We want to represent the projection image of the RHS of this element in $\mathcal{H}_f \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0}/m^{k+1}$ for $k \geq 0$ by an element of $\bigoplus_{l=1}^k z^{-l} \hat{\mathcal{B}} \otimes_{\mathbb{C}} m^l \check{\mathcal{O}}_{S,0}$ modulo the image of $D_f(\Gamma(X, \Omega_X^n)(z)) \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0}$, where we recall that $\hat{\mathcal{B}} \subset \Gamma(X, \Omega_X^{n+1})[[z]]$ is a μ -dimensional \mathbb{C} -subspace representing \mathcal{B} (see paragraphs following §3 Definition 3.1) and note that the action of D_f commutes with the coefficients in $\check{\mathcal{O}}_{S,0}$.

Let $\hat{\zeta}_0 \in \Gamma(X, \Omega_X^n)[[z]]$ and $\hat{\zeta}_+ \in \Gamma(Z, \Omega_{Z/S}^n)[[z]]$ be representatives of ζ_0 and ζ_+ , respectively, where we may assume that $\hat{\zeta}_0 = \hat{\zeta}_+|_X$ (see Proposition 4.1 and its Corollary). Therefore, we have an expression $\hat{\zeta}_+ = \pi_X^{-1}(\hat{\zeta}_0)(1 + a(\underline{x}, \underline{t}, z))$ where $a = \sum_{|I|>0} a_I(\underline{x}, z) \underline{t}^I$ is a power series in \underline{t} with respect to the multi-index $I = (i_1, \dots, i_\mu) \in (\mathbb{Z}_{\geq 0})^\mu$ ($|I| = \sum_{j=1}^\mu i_j$) and $a_I(\underline{x}, z) \in \Gamma(X, \mathcal{O}_X)[[z]]$ (by shrinking X and Z if necessary). Owing to the expression (4.8), we lift $\zeta_0 - e^{(F-f)/z} \zeta_+ \pmod{m^{k+1}}$ to an element in $\Gamma(X, \Omega_X^n)[[z]] \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0} \pmod{m^{k+1}}$:

$$\hat{\zeta}_0 - \sum_{l=0}^{\infty} \frac{(F(\underline{x}, \underline{t}) - F_0(\underline{x}))^l}{l! z^l} \hat{\zeta}_0 (1 + a(\underline{x}, \underline{t}, z)) \pmod{m^{k+1}}$$

⁹For the same reason as in Remark 4.1, the elements in $\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0}$ may not always be expanded in Laurent series in z of bounded order of poles. Actually, we shall see in §10 that the coefficients in u_j are non-trivial polynomials of degree j with respect to the flat coordinates introduced in §8.

Since $F - F_0$ vanishes along X , i.e. it is contained in $m\Gamma(Z, \mathcal{O}_Z)$, the terms of summation index l larger than k vanish mod m^{k+1} . Hence, the expression is a finite sum till $l = k$. Let us develop this expression in the form $\sum_{K \in (\mathbb{Z}_{\geq 0})^\mu, 0 < |K| \leq k} \alpha_K(\underline{x}, z) \underline{t}^K$. The coefficient of this lifting is given by

$$\alpha_K(\underline{x}, z) = - \sum_{\substack{I, J \in (\mathbb{Z}_{\geq 0})^\mu, \\ I+J=K}} \binom{k-|I|}{l=0} \left(\text{the coefficient of } \underline{t}^J \text{ in } \frac{(F(\underline{x}, \underline{t}) - F_0(\underline{x}))^l}{l! z^l} \hat{\zeta}_0 \right) \times a_I(\underline{x}, z)$$

where we set $a_0 := 0$. In view that a_I and $\hat{\zeta}_0$ have no-poles in z , we see that the order of pole in z in the term for a fixed J and $I = K - J$ is bounded by $l \leq |J| \leq |K|$. That is, $\alpha_K(\underline{x}, z) \in z^{-|K|} \Gamma(X, \Omega_X^{n+1})[[z]]$. We apply the decomposition (3.5) to α_K so that it belongs to the direct sum $z^{-|K|} \hat{\mathcal{B}}[[z]] \oplus D_f(z^{-|K|} \Gamma(X, \Omega_X^n)[[z]])$. Separating the principal part of the Laurent series part in $z^{-|K|} \hat{\mathcal{B}}[[z]] = \bigoplus_{m=1}^{|K|} z^{-m} \mathcal{B} \oplus \mathcal{B}[[z]]$ of α_K , we decompose $\alpha_K \equiv \sum_{m=1}^{|K|} z^{-m} \beta_{K,m} + \beta_{K,0} \pmod{D_f(z^{-|K|} \Gamma(X, \Omega_X^n)[[z]])}$ for $\beta_{K,m} \in \hat{\mathcal{B}}$ ($m = 1, \dots, |K|$) and $\beta_{K,0} \in \hat{\mathcal{B}}[[z]]$. Multiply the monomial \underline{t}^K to $\alpha_K(\underline{x}, z)$ and sum up by the index K . Since it is a finite sum, we resum it according to the order l of the pole in z , and we obtain the expression

$$\sum_{0 < |K| \leq k} \alpha_K(\underline{x}, z) \underline{t}^K = \sum_{l=1}^k z^{-l} \left(\sum_{|K| \leq k} \beta_{K,l} \underline{t}^K \right) + \sum_{0 < |K| \leq k} \beta_{K,0} \underline{t}^K \pmod{D_f(z^{-k} \Gamma(X, \Omega_X^n)[[z]]) \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_S / m^{k+1}}$$

in the module $z^{-k} \Gamma(X, \Omega_X^{n+1})[[z]] \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_S / m^{k+1}$. Here, the l -th ($l = 1, \dots, k$) component of the first term $\sum_{|K| \leq k} \beta_{K,l} \underline{t}^K$ belongs to $\sum_{|K| \leq k} \mathcal{B} \otimes_{\mathbb{C}} m^{|K|} \check{\mathcal{O}}_S / m^{k+1} = \mathcal{B} \otimes_{\mathbb{C}} m^l \check{\mathcal{O}}_S / m^{k+1}$ and the second term $\sum_{0 < |K| \leq k} \beta_{K,0} \underline{t}^K$ belongs to $\mathcal{B}[[z]] \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_S / m^{k+1}$. On the other hand, we recall that the sum $\sum_{0 < |K| \leq k} \alpha_K(\underline{x}, z) \underline{t}^K$ is a representative of the class $\sum_{j=1}^{\infty} z^{-j} u_j \pmod{m^{k+1}}$. This implies

$$u_j \equiv \sum_{j \leq |K| \leq k} \beta_{K,j} \underline{t}^K \quad \text{and} \quad \sum_{0 < |K| \leq k} \beta_{K,0} \underline{t}^K \equiv 0 \pmod{m^{k+1}}$$

(actually, this is the constraint on $\hat{\zeta}_+$). In particular, we see that, for any infinitesimal neighborhood $\mathcal{O}_{S,0} / m^{k+1}$ ($k \in \mathbb{Z}_{\geq 0}$), u_j belongs to $\mathcal{B} \otimes_{\mathbb{C}} m^j \check{\mathcal{O}}_{S,0}$. \square

Let us return to the proof of Theorem 6.3. The proof consists of the following steps 0-13. In the first steps 0 - 6., we show that the correspondence

$$(6.7) \quad \Phi : (\mathcal{L}, \zeta_0) \mapsto \zeta_+$$

defines a map from the LHS to the RHS of (6.1) and (6.2). In the steps 7 - 13., we construct the inverse map Ψ of Φ .

Step 0. *For any good pair (\mathcal{L}, ζ_0) , the associated ζ_+ satisfies the primitivity $(P0)^\vee$.*

PROOF. We need to show that the $\mathcal{O}_{S,0} / m^{k+1}$ -morphism $z \nabla \zeta_+ : \mathcal{T}_S / m^{k+1} \mathcal{T}_S \rightarrow \Omega_F / m^{k+1} \Omega_F$ between the $\mathcal{O}_{S,0} / m^{k+1}$ -free modules is an isomorphism for any $k \in \mathbb{Z}_{\geq 0}$. Recall the description of the Gauss-Manin connection (2.16) so that we have $z \nabla_v \zeta_+ \pmod{\check{\mathcal{H}}_F^{(-1)}} \equiv [\partial_{\check{v}} F \hat{\zeta}_+]$ where $\hat{\zeta}_+ \in (p_*(\Omega_{Z/S}^{n+1}[[z]]))^\vee$ is a lifting of ζ_+ to a relative differential form. However, this morphism for $k = 0$ is an isomorphism since 1) $e^{(f-F)/z} \zeta_0 \pmod{m} \equiv \zeta_0$ implying that $\zeta_+ \pmod{m} \equiv \zeta_0$, and 2) ζ_0 satisfies the

primitivity (1) in Definition 6.1. Then, Nakayama Lemma implies the isomorphism over \mathcal{O}_S/m^{k+1} for any $k \in \mathbb{Z}_{>0}$ and hence over $\check{\mathcal{O}}_{S,0}$. \square

Step 1. *For any good pair (\mathcal{L}, ζ_0) , the associated ζ_+ gives the following $\check{\mathcal{O}}_{S,0}$ -isomorphism*

$$(6.8) \quad z\nabla\zeta_+ : \check{\mathcal{T}}_S \simeq \check{\mathcal{H}}_F^{(0)} \cap ze^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$$

where we recall that the RHS is the formal section of $\check{\mathcal{H}}_F^{(0)}$ given in Proposition 4.2, 2.

PROOF. Let $v \in \check{\mathcal{T}}_S$ ($i = 1, 2$) and let $e^{(f-F)/z}\zeta_0 = \zeta_+ + \zeta_-$ be the splitting (4.10). Since $e^{(f-F)/z}\zeta_0$ is flat (Proposition 4.1, 3.), we have $z\nabla_v\zeta_+ = -z\nabla_v\zeta_-$, where the LHS belongs to $\check{\mathcal{H}}_F^{(0)}$ (transversality: 2. ii) of Proposition 2.1) and the RHS belongs to $ze^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$ (see Proposition 4.3). That is,

$$(6.9) \quad z\nabla_v\zeta_+ \in \check{\mathcal{H}}_F^{(0)} \cap ze^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$$

Combining isomorphisms (P0) $^\vee$ (above Step 0.) and (4.12), we obtain the isomorphism (6.8). \square

Step 2. *For any good pair (\mathcal{L}, ζ_0) , the associated ζ_+ satisfies the ∂_1 -triviality (P1).*

PROOF. Since $\zeta_0 \in \mathcal{B} \subset z\mathcal{L}$, one has $\zeta_+ + \zeta_- = e^{(f-F)/z}\zeta_0 \in ze^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$. On the other hand, by definition, $\zeta_- \in e^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$. Then, in view of the opposite condition (4.11) of the formal opposite filtration, we see that ζ_- and hence $\zeta_+ = e^{(f-F)/z}\zeta_0 - \zeta_-$ belongs to $ze^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$. Thus, ζ_+ belongs to the RHS of (6.8), implying that there exists a unique element $v \in \check{\mathcal{T}}_S$ such that $z\nabla_v\zeta_+ = \zeta_+$. This means, by the morphism $r_F^{(0)}$ (2.15), we obtain the relation $(\tilde{v}F|_{C_F}) \cdot \zeta_+ = \zeta_+$ in $\check{\Omega}_F$. On the other hand, since $\zeta_+ \bmod m = \zeta_0$ is a generating element in Ω_f (Definition 6.1 (1)), ζ_+ is also a generating element in $\check{\Omega}_F$. Then the relation $(\tilde{v}F|_{C_F} - 1)\zeta_+ = 0$ implies $\tilde{v}F|_{C_F} - 1 = 0$. That is, by Definition 2.2, v is the primitive vector field ∂_1 . \square

Step 3. *Suppose that a good pair (\mathcal{L}, ζ_0) satisfies the metric condition (see Definition 6.1 and Proposition 3.4), then the associated ζ_+ satisfies the orthogonality (P2) $^\vee$.*

PROOF. This is an immediate consequence of above Step 1. combined with the purity formula (4.17). \square

Step 4. *For any good pair (\mathcal{L}, ζ_0) , the associated ζ_+ satisfies the holonomicity (P3) $^{*\vee}$.*

PROOF. For $v_1, v_2 \in \check{\mathcal{T}}_S$, we have $\nabla_{v_1}\nabla_{v_2}\zeta_+ = -\nabla_{v_1}\nabla_{v_2}\zeta_-$. Applying the transversality of ∇ (2. i) of Proposition 2.1) on the LHS and the \underline{t} -stability of $e^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$ (Proposition 4.3) on the RHS, we see that it is contained in $z^{-2}\check{\mathcal{H}}_F^{(0)}$ and $e^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$, whose intersection decomposes

$$z^{-2}\check{\mathcal{H}}_F^{(0)} \cap e^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0}) = z^{-2}(\check{\mathcal{H}}_F^{(0)} \cap ze^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})) \oplus z^{-1}(\check{\mathcal{H}}_F^{(0)} \cap ze^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0}))$$

into two graded pieces. Then, applying Step 1. to each piece, this module is expressed as $z^{-1}\nabla_{\check{\mathcal{T}}_S}\zeta_+ \oplus \nabla_{\check{\mathcal{T}}_S}\zeta_+$. Then the first formula of (P3) $^{*\vee}$ follows.

Next, take $v \in \mathcal{T}_S$. Then, $\nabla_{z\partial_z} \nabla_v \zeta_+ = -\nabla_{z\partial_z} \nabla_v \zeta_-$ belongs to $z^{-2} \check{\mathcal{H}}_F^{(0)} \cap e^{(f-F)/z} (\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$ (recall irregularity of ∇_{∂_z} : 2. iii) of Proposition 2.1 and the $z\partial_z$ -stability of $e^{(f-F)/z} (\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$ Proposition 4.4). As in the first half, the intersection decomposes, and we obtain the second formula of (P3)^{*v}. \square

Step 5. *Suppose that a good pair (\mathcal{L}, ζ_0) satisfies the metric condition (see Definition 6.1), then the associated ζ_+ satisfies the holonomicity (P3)^v.*

PROOF. The metric condition implies the purity (orthogonality) of higher residue pairings (4.17). Then, the conditions in (P3)^{*v} are reformulated into those in (P3)^v. \square

Step 6. *For any good pair (\mathcal{L}, ζ_0) , the associated ζ_+ satisfies the homogeneity (P4).*

PROOF. We calculate $(\nabla_{z\partial_z+E} - r)e^{(f-F)/z} \zeta_0$ by two different ways.

1. The flatness of the evolution map and the commutativity of $\nabla_{z\partial_z}$ with the evolution map (3. and 5. of Proposition 4.1) implies $(\nabla_{z\partial_z+E} - r)e^{(f-F)/z} \zeta_0 = e^{(f-F)/z} ((\nabla_{z\partial_z} - r)\zeta_0)$. Then, the homogeneity of ζ_0 ((2) of Definition 6.1) implies that the last term is contained in $e^{(f-F)/z} \mathcal{L} \subset e^{(f-F)/z} (\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$. That is, the image is contained in the second splitting factor of (4.10).

2. We remark that the splitting (4.10) is preserved by the action of $\nabla_{z\partial_z+E} - r$, since (1) the filter $\check{\mathcal{H}}_F^{(0)}$ is preserved due to 2. iv) of Proposition 2.1, and (2) $e^{(f-F)/z} (\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$ is preserved due to the $z\partial_z$ -stability (1. of Proposition 4.4) and \underline{t} -stability (Proposition 4.3). In particular, the action of $\nabla_{z\partial_z+E} - r$ on the decomposition (6.3) preserves each factor to each factor.

Comparing 1. and 2., we see that the action of $\nabla_{z\partial_z+E} - r$ on the first factor ζ_+ should be zero. \square

Note. According to the homogeneity of ζ_0 (Definition 6.1), set $G := \nabla_{z\partial_z} \zeta_0 - r\zeta_0 \in \mathcal{L}$. Then, the above argument in 6. shows that $\nabla_{E+z\partial_z} \zeta_- = r\zeta_- + e^{(f-F)/z} G$.

Above 0.-6. altogether show that one direction Φ (6.7) of the correspondences (6.1) and (6.2) are well-defined. In the following 7-13., we construct the inverse correspondence $\Psi : \zeta_+ \mapsto (\mathcal{L}, \zeta_0)$ for a given formal primitive form $\zeta_+ \in \check{\mathcal{H}}_F^{(0)}$ which may or may not be equipped with metric structure (recall Definitions 5.5 and 5.6).

Step 7. We first consider a set $\check{\mathcal{B}}_{\zeta_+}$ of covariant differentiations of ζ_+ .

$$(6.10) \quad \check{\mathcal{B}}_{\zeta_+} := \text{Im}(z\nabla_{\zeta_+} : \check{\mathcal{T}}_S \rightarrow \check{\mathcal{H}}_F^{(0)}) = z\nabla_{\check{\mathcal{T}}_S} \zeta_+.$$

By the primitivity (P0)^v, $\check{\mathcal{B}}_{\zeta_+}$ is a section to the projection $\check{\mathcal{H}}_F^{(0)} \rightarrow \check{\Omega}_F$ (2.15) in the sense that $\check{\mathcal{B}}_{\zeta_+}$ is a $\check{\mathcal{O}}_{S,0}$ -submodule of $\check{\mathcal{H}}_F^{(0)}$ which is isomorphic to $\check{\Omega}_F$ by the projection $\check{r}_F^{(0)}$.

Therefore, for each $k \in \mathbb{Z}_{\geq 0}$, we have identifications (c.f. Definition 3.1,1)

$$(6.11) \quad \mathcal{H}_F/m^{k+1}\mathcal{H}_F = (\check{\mathcal{B}}_{\zeta_+}/m^{k+1}\check{\mathcal{B}}_{\zeta_+})(z) \quad \text{and} \quad \mathcal{H}_F^{(0)}/m^{k+1}\mathcal{H}_F^{(0)} = (\check{\mathcal{B}}_{\zeta_+}/m^{k+1}\check{\mathcal{B}}_{\zeta_+})[[z]].$$

Then, we consider the formal opposite filtration of $\check{\mathcal{H}}_F$ (this terminology shall be justified in the following Step 9.) by considering the projective system of $\mathcal{O}_S/m^{k+1}\mathcal{O}_S$ -submodule of $\mathcal{H}_F/m^{k+1}\mathcal{H}_F$ for $k \in \mathbb{Z}_{\geq 0}$

$$(6.12) \quad \mathcal{L}_{\zeta_+}^k := z^{-1}(\check{\mathcal{B}}_{\zeta_+}/m^{k+1}\check{\mathcal{B}}_{\zeta_+})[z^{-1}]$$

which is a splitting factor in $\mathcal{H}_F/m^{k+1}\mathcal{H}_F = \mathcal{H}_F^{(0)}/m^{k+1}\mathcal{H}_F^{(0)} \oplus \mathcal{L}_{\zeta_+}^k$ for $k \in \mathbb{Z}_{\geq 0}$.

Before going further to study the limit of this projective system, we formulate, using the particular case $k = 0$ of (6.12), the statement of the proof of Theorem 6.3.

Step 8. *Let ζ_+ be a formal primitive form without metric structure. Consider the pair*

$$\mathcal{L}_{\zeta_+}^0 := (6.12) \text{ for } k = 0 \quad \text{and} \quad \zeta_+^0 := \zeta_+ \bmod m \in \mathcal{H}_F^{(0)}/m\mathcal{H}_F^{(0)} \simeq \mathcal{H}_f^{(0)}$$

(recall Note after the proof of Theorem 4.1). Then, the correspondence

$$\Psi : \zeta_+ \mapsto (\mathcal{L}_{\zeta_+}^0, \zeta_+^0)$$

gives a map from the RHS to the LHS of (6.1) and (6.2), which is the inverse of Φ .

PROOF. The proof is divided into Steps 9-13. Precisely, the fact that $\mathcal{L}_{\zeta_+}^0$ is a $z\partial_z$ -stable opposite filtration of \mathcal{H}_f without metric condition is a particular case of Step 9. The facts that ζ_+^0 belongs to $z\mathcal{L}_{\zeta_+}^0$ and is primitive and homogeneous are shown in Steps 10 and 11, respectively. The fact that Ψ is the inverse to Φ is shown in Step 13. \square

Step 9. *The $\mathcal{O}_S/m^{k+1}\mathcal{O}_S$ -modules (6.12) for $k \in \mathbb{Z}_{\geq 0}$ form a projective system so that the limit*

$$\check{\mathcal{L}}_{\zeta_+}^\infty := \varprojlim \mathcal{L}_{\zeta_+}^k$$

is a formal opposite submodule in $\check{\mathcal{H}}_F$ in the sense of (6.13) below such that $\check{\mathcal{B}}_{\zeta_+} = \check{\mathcal{H}}_F^{(0)} \cap z\check{\mathcal{L}}_{\zeta_+}^\infty$. The $\check{\mathcal{L}}_{\zeta_+}^\infty$ is $z\partial_z$ -stable and \mathfrak{t} -stable. If, further, ζ_+ is a formal primitive form (that is, it further satisfies the metric condition (P2) $^\vee$), then the $\check{\mathcal{L}}_{\zeta_+}^\infty$ satisfies the formal metric condition (see Proposition 3.4).

PROOF. The fact that $\mathcal{L}_{\zeta_+}^k$ is a splitting factor $\mathcal{O}_{S,0}/m^{k+1}[z^{-1}]$ -submodule of $\mathcal{H}_F/m^{k+1}\mathcal{H}_F$ forming a projective system is obvious from the descriptions (6.11). These imply that $\mathcal{L}_{\zeta_+}^\infty$ is a opposite submodule in $\check{\mathcal{H}}_F$ in the sense that

$$(6.13) \quad \check{\mathcal{H}}_F = \check{\mathcal{H}}_F^{(0)} \oplus \check{\mathcal{L}}_{\zeta_+}^\infty \quad \text{and} \quad z^{-1}\check{\mathcal{L}}_{\zeta_+}^\infty \subset \check{\mathcal{L}}_{\zeta_+}^\infty.$$

The $z\partial_z$ -stability (see Proposition 3.3) and the \mathfrak{t} -stability (see Proposition 4.3) of $\check{\mathcal{L}}_{\zeta_+}^\infty$ follow from those of $\mathcal{L}_{\zeta_+}^k$, which follow from the condition (P3) * on the primitive form ζ_+ .

Suppose further that ζ_+ satisfies (P2) $^\vee$. This means, by definition of $\check{\mathcal{B}}_{\zeta_+}$, immediately

$$(6.14) \quad \check{K}_F(\check{\mathcal{B}}_{\zeta_+}, \check{\mathcal{B}}_{\zeta_+}) \subset z^{n+1}\check{\mathcal{O}}_{S,0} \quad \square$$

Step 10. *The initial term ζ_+^0 of ζ_+ belong to $z\mathcal{L}_{\zeta_+}^0$.*

PROOF. The condition (P1) on ζ_+ implies that $\zeta_+ \in \check{\mathcal{B}}_{\zeta_+}$ (6.10). In particular, this means $\zeta_+ \bmod m^{k+1} \in \check{\mathcal{B}}_{\zeta_+}/m^{k+1}\check{\mathcal{B}}_{\zeta_+} \subset z\check{\mathcal{L}}_{\zeta_+}^k$. Then, taking modulo the maximal ideal m , we obtain the result. \square

Step 11. *The initial term ζ_+^0 of ζ_+ is primitive and homogeneous.*

PROOF. Primitivity: Recall that the section $\check{\mathcal{B}}_{\zeta_+}$ is given by the covariant differentiation of ζ_+ (6.10). By definition of the section, its projection to $\check{\Omega}_F$ is surjective, i.e. $\check{\Omega}_F \subset \{\check{v}F\zeta_+ \mid v \in \check{\mathcal{T}}_S\} \subset \check{\mathcal{O}}_{C_F}\zeta_+ \subset \check{\Omega}_F$. Specializing this to $t = 0$ (i.e. mod m), we obtain the primitivity of ζ_0 .

Homogeneity : The condition (P4) on ζ_+ implies

$$\nabla_{z\partial_0}\zeta_+^0 - r\zeta_+^0 = \nabla_{z\partial_z}(\zeta_+|_X) - r(\zeta_+|_X) = (-\nabla_E\zeta_+)|_X \in z^{-1}(\check{\mathcal{B}}_{\zeta_+} \bmod m) \subset \mathcal{L}_{\zeta_+}^0.$$

□

Step 12. *The initial data $\mathcal{L}_{\zeta_+}^0$ (i.e. (6.12) for $k = 0$) is a $z\partial_z$ -stable opposite filtration of \mathcal{H}_f such that the formal opposite filtration $\check{\mathcal{L}}_{\zeta_+}^\infty$ is recovered from the initial data $\mathcal{L}_{\zeta_+}^0$ by the evolution map (4.7):*

$$(6.15) \quad e^{(f-F)/z}(\mathcal{L}_{\zeta_+}^0 \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0}) = \check{\mathcal{L}}_{\zeta_+}^\infty.$$

PROOF. We know already that $\mathcal{L}_{\zeta_+}^0$ is a $z\partial_z$ -stable opposite filtration in $\mathcal{H}_F/m\mathcal{H}_F = \check{\mathcal{H}}_f$ (see *Proof.* of Step 9.). Let us show (6.15). We first note the following.

a) For any given $a \in \mathcal{L}_{\zeta_+}^0$, the equation $\nabla_{\mathcal{T}_S}\alpha = 0$ for $\alpha \in \check{\mathcal{H}}_F$ with the initial condition $\alpha \bmod m = a$ has a unique solution in $\check{\mathcal{L}}_{\zeta_+}^\infty$.

PROOF. This essentially reduces to the existence and the uniqueness of the (formal) solution to the system of the first order linear partial differential equation $\nabla_{\mathcal{T}_S}\alpha = 0$ (for the coefficients of α in the Laurent expansion in z , c.f. (6.12)) over S for the given initial value a at $\underline{t} = 0$, where one should note that the equation is non-singular (i.e. the coefficient matrices are regular) on S . For details, see [11] Claim B in *Proof* of **Lemma/Definition 4.18**. □

Let us return to a proof of (6.15). Let $a \in \mathcal{L}_{\zeta_+}^0$ as above. The flatness (see (4.6)) of element $e^{(f-F)/z}a \in e^{(f-F)/z}\mathcal{H}_f \subset \check{\mathcal{H}}_F$ implies that it is a solution of the equation in a) satisfying the initial condition $e^{(f-F)/z}a \bmod m = a$. Due to the uniqueness of the solution, we observe that $e^{(f-F)/z}\mathcal{L}_{\zeta_+}^0$ is contained in $\check{\mathcal{L}}_{\zeta_+}^\infty$. More precisely, for any $k \in \mathbb{Z}_{\geq 0}$, we have inclusions $e^{(f-F)/z}\mathcal{L}_{\zeta_+}^0/m^{k+1} \subset (\check{\mathcal{L}}_{\zeta_+}^\infty/m^{k+1}\check{\mathcal{L}}_{\zeta_+}^\infty)$ and hence $e^{(f-F)/z}(\mathcal{L}_{\zeta_+}^0 \otimes_{\mathbb{C}} \mathcal{O}_{S,0}/m^{k+1}) \subset (\check{\mathcal{L}}_{\zeta_+}^\infty/m^{k+1}\check{\mathcal{L}}_{\zeta_+}^\infty)$. Then, taking the limit w.r.t. $k \in \mathbb{Z}_{>0}$, we obtain an inclusion $e^{(f-F)/z}(\mathcal{L}_{\zeta_+}^0 \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0}) \subset \check{\mathcal{L}}_{\zeta_+}^\infty$.

On the other hand, we have splittings (4.10) and (6.13) of the same module $\check{\mathcal{H}}_F^{(0)}$ of the same first factor $\check{\mathcal{H}}_F^{(0)}$, so that the inclusion is actually the equality (6.15). □

The following Step 13. is the final step of the proof of Theorem 6.3.

Step 13. *The compositions $\Psi \circ \Phi$ and $\Phi \circ \Psi$ are identities on each side of (6.1) and (6.2), respectively.*

PROOF. Let (\mathcal{L}, ζ_0) be a good pair. Set $\zeta_+ := \Phi(\mathcal{L}, \zeta_0)$ so that $\Psi(\zeta_+) = (\mathcal{L}_{\zeta_+}^0, \zeta_+ \bmod m)$. Recall (6.10) and (6.8), and we get $\check{\mathcal{B}}_{\zeta_+} = \check{\mathcal{H}}_F^{(0)} \cap ze^{(f-F)/z}(\mathcal{L} \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$. Restricting this equality on the subspace $X \subset Z$ (i.e. taking mod m , or setting $k = 0$), we obtain $\mathcal{B}_{\zeta_+}^0 = \mathcal{H}_f^{(0)} \cap \mathcal{L}$, where the LHS $\mathcal{B}_{\zeta_+}^0 := \check{\mathcal{B}}_{\zeta_+}/m\check{\mathcal{B}}_{\zeta_+}$ is the section of $\mathcal{H}_f^{(0)}$ corresponding to the opposite filtration $\mathcal{L}_{\zeta_+}^0$ in the sense of Proposition 3.2 and the RHS is the section corresponding to \mathcal{L} . Again due to Proposition 3.2, we get

$\mathcal{L}_{\zeta_+}^0 = \mathcal{L}$. On the other hand, $\zeta_+ \bmod m = \zeta_+ + \zeta_- \bmod m = e^{(f-F)/z} \zeta_0 \bmod m = \zeta_0$ (here, we use Proposition 6.1 for the vanishing of $\zeta_- \bmod m$). That is, $\Psi(\zeta_+) = (\mathcal{L}, \zeta_0)$, and hence $\Psi \circ \Phi$ is the identity on the set of good pairs.

Conversely, let us start with a formal primitive form ζ_+ . Then $\Phi \circ \Psi(\zeta_+) = \Phi(\mathcal{L}_{\zeta_+}^0, \zeta_+^0)$ where $\zeta_+^0 := \zeta_+ \bmod m$, by definition, is the first splitting factor of $e^{(f-F)/z} \zeta_+^0$ in $\check{\mathcal{H}}_F = \check{\mathcal{H}}_F^{(0)} \oplus e^{(f-F)/z} (\mathcal{L}_{\zeta_+}^0 \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0})$, and $e^{(f-F)/z} (\mathcal{L}_{\zeta_+}^0 \hat{\otimes}_{\mathbb{C}} \check{\mathcal{O}}_{S,0}) = \check{\mathcal{L}}_{\zeta_+}^\infty$ (6.15). Therefore, in order to show that it is actually equal to ζ_+ , i.e. $\Phi \circ \Psi$ is an identity, we have to show that $\xi := e^{(f-F)/z} \zeta_+^0 - \zeta_+ \in \check{\mathcal{L}}_{\zeta_+}^\infty$. We know that ξ is an element of $z\check{\mathcal{L}}_{\zeta_+}^\infty \subset \check{\mathcal{H}}_F$ since $\zeta_+^0 \in z\mathcal{L}_{\zeta_+}^0$ and $\zeta_+ \in \check{\mathcal{B}}_{\zeta_+}$ (see Step 10. and its proof). On the other hand, we see 1) ξ satisfies differential equation $\nabla_{\mathcal{T}_S} \xi (= -\nabla_{\mathcal{T}_S} \zeta_+) \subset z^{-1}\check{\mathcal{B}}_{\zeta_+} \subset \check{\mathcal{L}}_{\zeta_+}^\infty$ (recall (4.6) and (6.10)), and 2) the initial value: $\xi \bmod m$ is equal to zero (by the definition of ζ_+^0). The solution for such system of equations belongs to $\check{\mathcal{L}}_{\zeta_+}^\infty$ (*Proof.* this can be shown similar to a) in the proof of Step 12. Namely, let $\xi_0 \in \check{\mathcal{B}}_{\zeta_+}$ be the degree zero term of the Laurent expansion of ξ in z , where we note that there is positive degree term in z . Then it satisfies 1) a regular linear equation $\nabla_{\mathcal{T}_S} \xi_0 = 0 \bmod \check{\mathcal{L}}_{\zeta_+}^\infty$, and 2) the initial value $\xi_0 \bmod m$ is equal to 0, and then the unique solution is $\xi_0 = 0$). \square

This completes a proof of Theorem (6.1). \square

7. Flat structure without metric structure

It is well-known that a primitive form induces a flat structure (Frobenius manifold structure [6, 4, 14]) on S ([25]). In the present section we confirm that a primitive form without a metric structure induces a flat structure without a metric structure in the sense of Sabbah (cf. [7, 8, 9]), in particular, flat coordinate system on S are introduced. This is achieved by writing down explicitly the holonomicity property (P3)* of a primitive form ζ as in the following proposition.

PROPOSITION 7.1. (Flat structure without metric structure.)

1. Let ζ be a primitive form without higher residue structure (Definition 5.4). Then it satisfies the following (P3)**, which is an explicit form of (P3) and (P3)* (Holonomicity).

(P3)** (Holonomicity)**: The second covariant derivatives of ζ have the followings expressions.

1. There exists an affine torsion free connection $\nabla/ : \mathcal{T}_S \times \mathcal{T}_S \rightarrow \mathcal{T}_S$ on S such that

$$(7.1) \quad \nabla_{v_1} \nabla_{v_2} \zeta = z^{-1} \nabla_{v_1 * v_2} \zeta + \nabla_{\nabla/v_1 v_2} \zeta \quad \text{for } v_1, v_2 \in \mathcal{T}_S.$$

where “*” means the product structure on \mathcal{T}_S given in (2.6).

2. There is an \mathcal{O}_S -endomorphism $N \in \text{End}_{\mathcal{O}_S}(\Omega_F)$ such that

$$(7.2) \quad \nabla_{z\partial_z} \nabla_v \zeta = -z^{-1} \nabla_{E*v} \zeta + \nabla_{N(v)} \zeta \quad \text{for } v \in \mathcal{T}_S,$$

where E is the Euler vector field (see Definition 2.2).

3. The connection $\nabla/$ and the endomorphism N satisfy the following relations.

- i) Flatness and charge of Primitive vector field: $\nabla/\partial_1 = 0$ and $N(\partial_1) = r\partial_1$.
- ii) Integrability of $\nabla/$: $[\nabla/, \nabla/] = 0$,

iii) Symmetry of $\nabla/*$: Set $T(u, v, w) := \nabla/u(v*w) - (\nabla/u v)*w - v*(\nabla/u w)$ for $u, v, w \in \mathcal{T}_S$.

Then, T is a symmetric \mathcal{O}_S -tensor of type $(3, 1)$, i.e. $T \in \text{Hom}_{\mathcal{O}_S}(S^3(\mathcal{T}_S), \mathcal{T}_S)$.

iv) Horizontality of N : $\nabla/N = 0$.

v) Exponents: $N(v) = (r+1) \cdot v - \nabla/v E$ for $v \in \mathcal{T}_S$,

vi) Flatness of E : $\nabla/w \nabla/v E = \nabla/\nabla/w v E$ for $v, w \in \mathcal{T}_S$.

vii) Homogeneity of $*$ -product: $[E, v*w] = v*[E, w] + [E, v]*w + v*w$ for $v, w \in \mathcal{T}_S$.

2. Let ζ be a formal primitive form without metric structure (Definition 5.6). Then it satisfies the following $(P3)^{**\vee}$ (a explicit form of $(P3)^\vee$) and $(P3)^{* \vee}$ (Holonomicity)), where

$(P3)^{**\vee}$ (Holonomicity) $^{**\vee}$: Replace \mathcal{T}_S in $(P3)^{**}$ by $\check{\mathcal{T}}_S$.

PROOF. We prove only **1.**, since **2.** can be proven parallery to the case for **1.**
Proofs of 1. and 2.: According to $(P3)^*$, we set two expansions:

$$\nabla_{v_1} \nabla_{v_2} \zeta = z^{-1} \nabla_{A(v_1, v_2)} \zeta + \nabla_{B(v_1, v_2)} \zeta \quad \text{for } v_1, v_2 \in \mathcal{T}_S$$

$$\nabla_{z \partial_z} \nabla_v \zeta = z^{-1} \nabla_{M(v)} \zeta + \nabla_N(v) \zeta \quad \text{for } v \in \mathcal{T}_S,$$

where $A, B : \mathcal{T}_S \times \mathcal{T}_S \rightarrow \mathcal{T}_S$ are \mathbb{C} -bilinear morphism in the variables v_1, v_2 and \mathcal{O}_S -linear morphisms in the variable v_1 , and $M, N : \mathcal{T}_S \rightarrow \mathcal{T}_S$ are \mathcal{O}_S -linear morphisms in the variable v .

In view of (2.21) and Kodaira-Spencer morphism (2.3) with the $*$ -product (2.6), the first terms of the above expansions are calculated as $A(v_1, v_2) = v_1 * v_2$ and $M(v) = -E * v$, respectively. Further more inserting the above expansions to the integrability relation and the Leibniz rule of the Gauss-Manin connection ∇ , we see that $B(v_1, v_2)$ is given as the covariant differentiation $\nabla/v_1 v_2$ of the tangent vector v_2 by v_1 for an affine torsion free connection $\nabla/$ on S :

$$\nabla/v w - \nabla/w v = [v, w] \quad \text{for } v, w \in \mathcal{T}_S.$$

Thus we obtain the expressions (7.1) and (7.2).

Proof of 3. : The covariant derivation by $v \in \mathcal{T}_S$ of the ∂_1 -triviality (P1) $z \nabla_{\partial_1} \zeta = \zeta$ implies the horizontality $\nabla/v \partial_1 = 0$ of the primitive vector field. The covariant derivation by $z \partial_z$ of the ∂_1 -triviality (P1) $z \nabla_{\partial_1} \zeta = \zeta$ implies the equality $z \nabla_{\partial_z} \zeta = -\nabla_E \zeta + z \nabla_{N(\partial_1)} \zeta$. Comparing this with the homogeneity axiom (P3) of ζ , we get $\nabla_{r \partial_1} \zeta = \nabla_{N(\partial_1)} \zeta$ and, therefore, $N(\partial_1) = r \partial_1$. That is, ∂_1 is an eigenvector of N belonging to the eigenvalue r , and **(1. 3. i)** is shown.

$$(7.3) \quad \nabla/\partial_1 = 0 \quad \text{and} \quad N(\partial_1) = r \partial_1$$

The integrability relation $[\nabla_{v_1}, \nabla_{v_2}] = \nabla_{[v_1, v_2]}$ acting on $\nabla_w \zeta$ for $w \in \mathcal{T}_S$ implies the integrability $[\nabla/v_1, \nabla/v_2] = \nabla_{[v_1, v_2]}$ of $\nabla/$ (**1. 3. ii**) together with the relation:

$$(7.4) \quad [v_1, v_2] * v_3 = \nabla/v_1(v_2 * v_3) + v_1 * \nabla/v_2 v_3 - \nabla/v_2(v_1 * v_3) - v_2 * \nabla/v_1 v_3.$$

Let us show that the expression:

$$(7.5) \quad T(u, v, w) := \nabla_u(v * w) - (\nabla/u v) * w - v * (\nabla/u w) \quad \text{for } u, v, w \in \mathcal{T}_S$$

is a symmetric \mathcal{O}_S -tensor of type $(3, 1)$ (i.e. a \mathfrak{S}_3 -symmetric \mathcal{O}_S -tri-linear map $\mathcal{T}_S^{\otimes 3} \rightarrow \mathcal{T}_S$). The \mathcal{O}_S -linearity on the variable u is obvious. The symmetry by

the permutation of v and w is also obvious. Therefore, we have only to show that the symmetry by the permutation of u and v . However, it is immediate to see $T(v_1, v_2, v_3) - T(v_2, v_1, v_3) = 0$ due to (7.4). Thus, (1. 3. iii) is shown.

The integrability (commutativity) relation $[\nabla_{z\partial_z}, \nabla_v] = 0$ acting on $\nabla_w \zeta$ for $w \in \mathcal{T}_S$ implies the horizontality $\nabla/N = 0$ of N (1. 3. iv) together with the relation:

$$(7.6) \quad N(v * w) - v * N(w) = E * \nabla/v w - \nabla/v(E * w) + v * w$$

Inserting $w = \partial_1$ in (7.6) and applying (7.3), we obtain the expression (1. 3. v) of N :

$$(7.7) \quad N(v) = (r + 1) \cdot v - \nabla/v E.$$

Applying the expression (7.7) back to (7.6), we obtain the relation

$$(7.8) \quad \nabla/v * w E - v * \nabla/w E + E * \nabla/v w + v * w = \nabla/v(E * w).$$

In (7.4), insert $v_1 = E$, $v_2 = v$ and $v_3 = w$ and take the difference with (7.8) so that we obtain

$$(\nabla/E(v * w) - \nabla/v * w E) - v * (\nabla/E w - \nabla/w E) - [E, v] * w = v * w.$$

Remembering that $\nabla/$ is torsion free, this means that the homogeneity (1. 3. vii) of the $*$ -product.

$$(7.9) \quad [E, v * w] = v * [E, w] + [E, v] * w + v * w.$$

This completes a proof of Proposition 7.1. \square

REMARK 7.1. Let the Euler vector field E be generically invertible as an element of the Frobenius algebra \mathcal{T}_S and we denote the inverse element $E^{-1} \in \mathcal{T}_S$. Then the structure in the Proposition 7.1 is actually a bi-flat structure in a sense that there is another affine torsion free compatible connection $\nabla/^* : \mathcal{T}_S \times \mathcal{T}_S \rightarrow \mathcal{T}_S$ (defined by the Euler vector field) such that $\nabla/^* w = \nabla/v E^{-1} * w$.

REMARK 7.2. Including the case when ζ_+ is a formal primitive form (with or without higher residue structure), recall (4.1) that the Gauss-Manin connection $\check{\nabla}$ acts on the trivial bundle $\check{\mathcal{H}}_F$. However, the formulae (7.1) and (7.2) imply that the $\mathcal{O}_S[z^{\pm 1}]$ -free submodule $\mathcal{B}_{\zeta_+}[z^{\pm 1}]$, where we set $\mathcal{B}_{\zeta_+} := \text{Im}(z\nabla\zeta_+ : \mathcal{T}_{S,0} \rightarrow \check{\mathcal{H}}_F^{(0)}) = z\nabla_{\mathcal{T}_{S,0}}\zeta_+$, is invariant under the actions of ∇_{∂_z} and ∇_v for $v \in \mathcal{T}_{S,0}$. In this sense, the connection is extended to a meromorphic connection on $\mathbf{P}^1 \times S'$ where S' is a neighborhood of the origin $0 \in S$ and \mathbf{P}^1 is the projective line of inhomogeneous coordinate z . For brevity, we shall call S' again S . So we obtain

$$(7.10) \quad \nabla : \mathcal{T}_{\mathbf{P}^1 \times S} \times \mathcal{B}_{\zeta_+}[z^{\pm 1}] \longrightarrow \mathcal{B}_{\zeta_+}[z^{\pm 1}]$$

where we may regard $\mathcal{B}_{\zeta_+}[z^{\pm 1}]$ as a finite rank bundle $\pi^* \mathcal{T}_S$ on $\mathbf{P}^1 \times S$ for $\pi : \mathbf{P}^1 \times S \rightarrow S$.

Flat coordinates of S . The integrability of the connection $\nabla/$ implies that the unfolding parameter space S carries an affine linear structure ([21]). Namely, *torsion-freeness of $\nabla/$ implies that the horizontal sections of the dual connection $\nabla/^*$ of $\nabla/$*

on the cotangent bundle Ω_S^1 of S are closed forms. Therefore, the following second order differential equation on a function u on S :¹⁰

$$(7.11) \quad \nabla^* du = 0$$

is integrable and has (locally) $\mu + 1$ linearly independent solution (including constant functions). We shall call them the (local affine linear) *flat coordinate system* of S w.r.t. the primitive form ζ . Using the natural pairing between tangent and cotangent bundles, the equation (7.11) is equivalent to the vanishing of $\langle v_2, \nabla^* du \rangle = v_1 \langle v_2, du \rangle - \langle \nabla^* v_1 v_2, du \rangle$ for all $v_1, v_2 \in \mathcal{T}_S$. That is,

$$(7.12) \quad v_1 v_2 u - (\nabla^* v_1 v_2) u = 0 \quad \forall v_1, v_2 \in \mathcal{T}_S.$$

Actually, the system (7.12) is integrable due to the torsion freeness of ∇^* : $(v_1 v_2 - (\nabla^* v_1 v_2)) - (v_2 v_1 - (\nabla^* v_2 v_1)) = [v_1, v_2] - (\nabla^* v_1 v_2 - \nabla^* v_2 v_1) = 0$.

For a sake of completeness, we add a few classical well-known results on the case when ζ is a primitive form with metric structure. Recall that Kodaira-Spencer morphism (2.2) has identified the tangent bundle of S with $\pi_* \mathcal{O}_C$. A choice of a primitive form (without metric structure) identify them with the vector bundle Ω_F over S due to (P0), where we recall that the bundle Ω_F admits an inner product η (2.20). We denote the pull back metric $\eta_\zeta := (z \nabla \zeta)^{-1} \zeta$ again simply by η and regard it as a metric on the tangent bundle \mathcal{T}_S of S .

PROPOSITION 7.2. Let ζ be a primitive form satisfying the metric condition (P2) and let η be the metric on \mathcal{T}_S induced from the residue pairing (2.20). Then, the flat connection ∇^* and the endomorphism N in Proposition 7.1 satisfy further the following relations:

- i) Horizontality of η : $\nabla^* \eta = 0$.
- ii) Self-adjointness of $*$ -product: $\eta(u * v, w) = \eta(u, v * w)$ for $u, v, w, \in \mathcal{T}_S$.
- iii) Duality of exponents: $N + N^* = 1$ where N^* is the adjoint of N w.r.t. η .

PROOF. According to the above convention of the metric η , the condition (P2) implies the equality: $K_F(\nabla_v \zeta, \nabla_w \zeta) = \eta(v, w) z^{n-2}$ for $v, w \in \mathcal{T}_S$. Let us apply this to the formulae (2.18) and (2.19), where we choose $\zeta_i = \nabla_{v_i} \zeta$ for $v_i \in \mathcal{T}_S$ ($i = 1, 2$) and use the expansion formulae (7.1) and (7.2). Then, for the first case (2.18), we obtain:

$$(7.13) \quad v \eta(v_1, v_2) z^{n-2} = \eta(v * v_1, v_2) z^{n-3} + \eta(\nabla^* v_1, v_2) z^{n-2} - \eta(v_1, v * v_2) z^{n-3} + \eta(v_1, \nabla^* v_2) z^{n-2}.$$

The relation of coefficients of z^{n-2} implies i) the horizontality $\nabla^* \eta = 0$, and the relation of coefficients of z^{n-3} implies ii) the self-adjointness of the $*$ -product. For the second case (2.19), we obtain:

$$(7.14) \quad (n-2+*) \eta(v_1, v_2) z^{n-2} = \eta(E * v_1, v_2) z^{n-3} + \eta(N(v_1), v_2) z^{n-2} - \eta(v_1, E * v_2) z^{n-3} + \eta(v_1, N(v_2)) z^{n-2}.$$

The relation of coefficients of z^{n-2} implies iii) the duality $N + N^* = 1$, and the relation of coefficients of z^{n-3} is a special case of ii) and does not give any new information. \square

¹⁰Using tangent-cotangent duality, the equation is equivalent to a system of equations $(v w - \nabla^* v w) u = 0$ for $v, w \in \mathcal{T}_S$.

Actually, the residue pairing defines a metric on \mathcal{T}_S via Kodaira-Spencer map even when the metric condition is not satisfied. But this metric is not flat in general.

PROPOSITION 7.3. Let ζ be a primitive form without metric structure. Let η be a metric induced on \mathcal{T}_S via the map $v \rightarrow z\nabla_v\zeta \in \mathcal{B} \simeq \Omega_F$. Let $(\mathcal{T}_S)^0 \subset \mathcal{T}_S$ be a subbundle of vector fields for which the condition (P2) is satisfied, that is if $v, w \in (\mathcal{T}_S)^0$ then $K_F(\nabla_v\zeta, \nabla_w\zeta) \in \mathbb{C}z^{n-2}$. Then the $*$ -product is symmetric with respect to η : $\eta(u * v, w) = \eta(u, v * w)$ for $u, v, w \in \mathcal{T}_S$.

PROOF. In general we have $K_F(\nabla_v\zeta, \nabla_w\zeta) = \eta(v, w)z^{n-2} + O(z^{n-1})$. We consider the relation (7.13) modulo the terms of order $O(z^{n-2})$:

$$(7.15) \quad 0 = \eta(v * v_1, v_2)z^{n-3} - \eta(v_1, v * v_2)z^{n-3},$$

which proves the statement. \square

Flat potential on S. As a consequence the metric property of $\nabla/$, we are now able to define a potential function [21]. Namely, *the following third order differential equation on a function F on S*

$$(7.16) \quad vvwF = \eta(u * v, w) \quad \text{for } u, v, w \in \text{Kernel}(\nabla/)$$

is solvable. Here we denote by $\text{Kernel}(\nabla/)$ the space of horizontal sections of $\nabla/$ (note that elements in $\text{Kernel}(\nabla/)$ are commutative as derivations due to the torsion-free-ness of $\nabla/$). (*Proof of the integrability of (7.16):* The 3-form in RHS of the equation (7.16) is symmetric in three vector fields u, v, w on S (Proposition 7.2 ii)). Let show that the 4-form $x\eta(u * v, w)$ for 4 flat vector fields $u, v, w, x \in \text{Kernel}(\nabla/)$ is symmetric. Let us specialize the symmetric tensor T (Proposition 7.1 1. iii) to flat vector fields $u, v, w \in \text{Kernel}(\nabla/)$. Then, we see that $\nabla_u(v * w)$ is symmetric in the three variables. On the other hand, using the horizontality of η (Proposition 7.2 i)), we see $x\eta(u * v, w) = \eta(\nabla_x(u * v), w) + \eta(u * v, \nabla_x w) = \eta(\nabla_x(u * v), w)$. LHS of this equality was already symmetric in the variables u, v, w , and RHS of the equality is now symmetric in three variables x, u, v . These together imply that the 4-tensor on $u, v, w, x \in \text{Kernel}(\nabla/)$ is symmetric. \square)

REMARK 7.3. Morihiko Saito [27] introduced the concept of very good section. Recently, Todor Milnov gave a characterization of a very good section in terms of the flat structure as follows.

PROPOSITION 7.4. ([15]) A section \mathcal{L} is very good if and only if

$$(7.17) \quad [N, E] = -E.$$

One observes that the formula (7.17) does not involve the metric structure (either the higher or the first residue pairings). Therefore, one can formally define a very good section without metric structure, if a good section without the metric condition satisfies (7.17). Using the flat structure in Proposition 7.1, we can rewrite the condition (7.17) only in terms of E as follows.

$$(7.18) \quad \nabla/E(E * a) - E * \nabla/E a = -E * a \quad \text{for } a \in \mathcal{T}_S.$$

PROOF. Applying Proposition 1. 3. v), LHS of the condition (7.17) implies

$$N(E * a) - E * N(a) = \nabla/E * a E - E * \nabla/a E = [E * a, E] + \nabla/E(E * a) - E * ([a, E] + \nabla/E a) = \nabla/E(E * a) - E * \nabla/E a$$

\square

Saito structures without metric and flat F-manifolds. Now we can compare our construction with the definition of the Saito structure without metric by Sabbah [18].

Let M be a complex analytic manifold, TM its tangent bundle, Θ_M the sheaf of holomorphic vector fields and Ω_M^1 the sheaf of holomorphic 1-forms.

DEFINITION 7.4. (Sabbah) A *Saito structure without metric* on M consists of the following data:

- (1) A torsionless flat connection ∇ on TM .
- (2) A 1-form Φ with values in $\text{End}(TM)$, namely a section of the sheaf $\text{End}(\Theta_M) \otimes_{\mathcal{O}_M} \Omega_M^1$, which is symmetric when considered as a bilinear map $\Theta_M \otimes_{\mathcal{O}_S} \Theta_M \rightarrow \Theta_M$;
- (3) Two global sections of Θ_M (vector fields) e and \mathfrak{E} respectively called unit field and Euler field such that the following two conditions are satisfied:
 - (1) The meromorphic connection ∇ defined on the vector bundle π^*TM on $\mathbb{A}^1 \times M$ by

$$(7.19) \quad \nabla = \pi^*\nabla + \frac{\pi^*\Phi}{\tau} - \left(\frac{\Phi(\mathfrak{E})}{\tau} + \nabla\mathfrak{E} \right) \frac{d\tau}{\tau}$$

is integrable.

- (2) The vector field e is ∇ -horizontal and $\Phi(e) = -Id$.

We consider a tuple $(M, \nabla, \Phi, e, \mathfrak{E})$ satisfying the conditions above as a Saito structure without metric.

Given a universal unfolding F over a frame (Z, S, p) the multiplication structure $*$: $\mathcal{T}_S \times \mathcal{T}_S \rightarrow \mathcal{T}_S$ defines a symmetric tensor C in $\text{End}(\mathcal{T}_S) \otimes_{\mathcal{O}_S} \Omega_S^1$ via $C(v, w) = v * w$, $v, w \in \mathcal{T}_S$, since the multiplication is \mathcal{O}_S -linear.

PROPOSITION 7.5. Let ζ be a primitive form without metric in a sense of Definition 5.4 .Then the tuple $(S, \nabla, -C, \partial_1, E)$ defines a Saito structure without metric.

PROOF. 1. The connection ∇ : $\mathcal{T}_S \times \mathcal{T}_S \rightarrow \mathcal{T}_S$ is flat and torsionless by the proposition 7.1.

2. The 1-form $-C$ is symmetric since the multiplication $*$ is commutative.

Condition 1: The connection ∇ reads:

$$(7.20) \quad \nabla = \pi^*\nabla - \frac{\pi^*C}{\tau} + \left(\frac{C(E)}{\tau} + \nabla E \right) \frac{d\tau}{\tau}.$$

The formula (7.20) means that for $v, w \in \mathcal{T}_S \otimes_{\mathcal{O}_S} \mathcal{O}_{S \times \mathbb{A}^1}$:

$$(7.21) \quad \begin{aligned} \nabla_v w &= \nabla'_v w - \tau^{-1} v * w, \\ \nabla_{\tau \partial_\tau} w &= \tau \partial_\tau w - \nabla'_w E + \tau^{-1} E * w. \end{aligned}$$

The integrability conditions read for all $v, u, w \in \mathcal{T}_S \subset \mathcal{T}_S \otimes_{\mathcal{O}_S} \mathcal{O}_{S \times \mathbb{A}^1}$, $[v, w] = 0$:

$$(7.22) \quad \begin{aligned} (\nabla'_v \nabla'_w - \nabla'_w \nabla'_v)u &= 0, \quad \nabla'_v(w * u) + v * \nabla'_w u = \\ &= \nabla'_w(v * u) + w * \nabla'_v u, \quad v * (w * u) - w * (v * u) = 0 \end{aligned}$$

from the compatibility of the first line of (7.21) with itself. The first equation holds since ∇ is flat (1. from the proposition 7.1), the second equation is the symmetry

condition 3. iii) from 7.1 and the third one is associativity and commutativity of the $*$ -product.

The following condition (together with the already checked) imply the compatibility of the first line of (7.21) with the second one:

$$(7.23) \quad w * \nabla_u E + \nabla/w(E * u) = E * \nabla/w u + \nabla/w(E * u) + w * u + \nabla/w * u E.$$

This equation is nothing but the equation (7.8).

To show property 2 we note that $-C(\partial_1) = -Id$ by the primitivity property of ∂_1 . Flatness of ∂_1 is 3. i) of the proposition 7.1. \square

REMARK 7.5. The connection (7.20) from the Saito structures without metric is essentially the Gauss-Manin connection. More precisely, a primitive form ζ defines a \mathcal{O}_S -module isomorphism $\mathcal{T}_S \rightarrow \mathcal{B}$ via $v \rightarrow zv\zeta$. Let $\tau = -z$. Then the connection (7.20) defines a connection on $\mathcal{B}((z)) \simeq \mathcal{H}_F$ via the isomorphism $\mathcal{T}_S \rightarrow \mathcal{B}$. Using the formula (7.21) we compute

$$(7.24) \quad \begin{aligned} \nabla_v \nabla_w \zeta &= \nabla_{\nabla/v w} \zeta + z^{-1} \nabla_{v * w} \zeta, \\ \nabla_{z \partial_z} \nabla_w \zeta &= z \partial_z \nabla_w \zeta - \nabla_{\nabla/w E} \zeta - z^{-1} \nabla_{E * w} \zeta. \end{aligned}$$

The first line is (7.1) if we identify $\nabla = \nabla$. The right hand side of the second line differs from the right hand side of (7.2) by the summand $r \cdot \nabla_w \zeta$.

A flat structure without metric also defines a structure of the flat F-manifold [14] on S which is a similar notion.

DEFINITION 7.6. (Manin) An *analytic F-manifold with compatible flat structure* (or *flat F-manifold*) is a tuple (M, \circ, ∇, e) , where M is a complex manifold, $\circ : \mathcal{T}_M \times_{\mathcal{O}_M} \mathcal{T}_M \rightarrow TM$ is a product on the tangent spaces depending holomorphically on the point in M , ∇ is a connection on TM and e is a distinguished vector field satisfying the properties:

- (1) The one-parameter family of connections $\nabla - \lambda \circ$ is flat and torsionless for any $\lambda \in \mathbb{C}$.
- (2) The vector field e is a unit of the product and is flat $\nabla e = 0$.

An F-manifold (M, \circ, ∇, e) is called conformal if it is equipped with a vector field $E \in \mathcal{T}_M$ called the Euler vector field such that $\nabla \nabla E = 0$, $[e, E] = e$ and $\mathcal{L}_E(\circ) = \circ$.

LEMMA 7.7. Let ζ be a primitive form without metric structure. Then $(S, *, \nabla, \partial_1, E)$ where $*$ is the multiplication structure on \mathcal{T}_S defined in (2.6) is a conformal flat F-manifold.

PROOF. Let us first that a primitive form without metric defines a flat F-structure.

1. Flatness and torsionless of the connection $\nabla - \lambda *$ follow from the flatness and torsionless of the \mathcal{T}_S -part of the Gauss-Manin connection or from the first line of (7.21).

2. Vector field ∂_1 is a unit of the product $*$ and is flat due to the properties of the flat structure without metric.

Properties $\nabla \nabla E = 0$ and homogeneity of the $*$ -product are is 3. vi) and 3. vii) of the proposition 7.1 correspondingly.

To show the property $[\partial_1, E] = \partial_1$ we notice that $[\partial_1, E] = \nabla_{\partial_1} E - \nabla_E \partial_1 = \nabla_{\partial_1} E$. Now we use

$$(7.25) \quad r\partial_1 = N(\partial_1) = (r+1)\partial_1 - \nabla_{\partial_1} E \implies \nabla_{\partial_1} E = \partial_1.$$

□

Acknowledgements

This work was partially supported by JP18H01116 (the second named author), OIST unit budget (Mathematical and Theoretical Physics Unit). The first named author is grateful to IPMU, OIST and Shinobu Hikami for hospitality.

References

- [1] Vladimir Arnold, Alexander Varchenko, and Sabir Gusein-Zade. *Singularities of Differentiable Maps*. Birkhauser Basel, 1985.
- [2] Sergey Barannikov and Maxim Kontsevich. “Frobenius manifolds and formality of Lie algebras of polyvector fields”. In: *Internat. Math. Res. Notices* 4 (1998), pp. 201–215. ISSN: 1073-7928. DOI: 10.1155/S1073792898000166. URL: <https://doi.org/10.1155/S1073792898000166>.
- [3] Alexey Basalaev and Alexandr Buryak. “Open Saito theory for A and D singularities”. In: *arXiv e-prints*, arXiv:1909.00598 (Sept. 2019), arXiv:1909.00598. arXiv: 1909.00598 [math-ph].
- [4] Boris Dubrovin. “Geometry of 2D topological field theories”. In: *Integrable systems and quantum groups (Montecatini Terme, 1993)*. Vol. 1620. Lecture Notes in Math. Springer, Berlin, 1996, pp. 120–348. DOI: 10.1007/BFb0094793. URL: <https://doi.org/10.1007/BFb0094793>.
- [5] Alexander B. Givental. “Symplectic geometry of Frobenius structures”. In: *Frobenius manifolds*. Aspects Math., E36. Friedr. Vieweg, Wiesbaden, 2004, pp. 91–112.
- [6] Claus Hertling. *Frobenius manifolds and moduli spaces for singularities*. Vol. 151. Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 2002, pp. x+270. ISBN: 0-521-81296-8. DOI: 10.1017/CB09780511543104. URL: <https://doi.org/10.1017/CB09780511543104>.
- [7] Mitsuo Kato, Toshiyuki Mano, and Jiro Sekiguchi. “Flat structures without potentials”. In: *Rev. Roumaine Math. Pures Appl.* 60.4 (2015), pp. 481–505. ISSN: 0035-3965.
- [8] Mitsuo Kato, Toshiyuki Mano, and Jiro Sekiguchi. “Flat structures and algebraic solutions to Painlevé VI equation”. In: *Analytic, algebraic and geometric aspects of differential equations*. Trends Math. Birkhäuser/Springer, Cham, 2017, pp. 383–398.
- [9] Mitsuo Kato, Toshiyuki Mano, and Jiro Sekiguchi. “Flat structure and potential vector fields related with algebraic solutions to Painlevé VI equation”. In: *Opuscula Math.* 38.2 (2018), pp. 201–252. ISSN: 1232-9274. DOI: 10.7494/opmath.2018.38.2.201. URL: <https://doi.org/10.7494/opmath.2018.38.2.201>.
- [10] Yukiko Konishi, Satoshi Minabe, and Yuuki Shiraiishi. “Almost duality for Saito structure and complex reflection groups”. In: *J. Integrable Syst.* 3.1 (2018), xyy003, 48. DOI: 10.1093/integr/xyy003. URL: <https://doi.org/10.1093/integr/xyy003>.
- [11] Changzheng Li, Si Li, and Kyoji Saito. “Primitive forms via polyvector fields”. In: *arXiv e-prints*, arXiv:1311.1659 (Nov. 2013), arXiv:1311.1659. arXiv: 1311.1659 [math.AG].

- [12] Paolo Lorenzoni and Marco Pedroni. “Natural connections for semi-Hamiltonian systems: the case of the ϵ -system”. In: *Lett. Math. Phys.* 97.1 (2011), pp. 85–108. ISSN: 0377-9017. DOI: 10.1007/s11005-011-0469-x. URL: <https://doi.org/10.1007/s11005-011-0469-x>.
- [13] Paolo Lorenzoni, Marco Pedroni, and Andrea Raimondo. “ F -manifolds and integrable systems of hydrodynamic type”. In: *Arch. Math. (Brno)* 47.3 (2011), pp. 163–180. ISSN: 0044-8753.
- [14] Yuri I. Manin. *Frobenius manifolds, quantum cohomology, and moduli spaces*. Vol. 47. American Mathematical Society Colloquium Publications. American Mathematical Society, Providence, RI, 1999, pp. xiv+303. ISBN: 0-8218-1917-8. DOI: 10.1090/coll/047. URL: <https://doi.org/10.1090/coll/047>.
- [15] Todor Milanov. *Singularity theory and mirror symmetry*. 2018.
- [16] Todor Milanov and Valentin Tonita. “K-theoretic Gromov-Witten invariants in genus 0 and integrable hierarchies”. In: *Lett. Math. Phys.* 108.6 (2018), pp. 1455–1483. ISSN: 0377-9017. DOI: 10.1007/s11005-018-1075-y. URL: <https://doi.org/10.1007/s11005-018-1075-y>.
- [17] Georges de de Rham. “Sur la division de formes et de courants par une forme linéaire.” In: *Commentarii mathematici Helvetici* 28 (1954), pp. 346–352. URL: <http://eudml.org/doc/139090>.
- [18] Claude Sabbah. *Isomonodromic deformations and frobenius manifolds: an introduction*. Universitext. London: Springer, 2008. DOI: 10.1007/978-1-84800-054-4. URL: <https://cds.cern.ch/record/1639313>.
- [19] Kyoji Saito. “Quasihomogene isolierte Singularitäten von Hyperflächen”. In: *Invent. Math.* 14 (1971), pp. 123–142. ISSN: 0020-9910. DOI: 10.1007/BF01405360. URL: <https://doi.org/10.1007/BF01405360>.
- [20] Kyoji Saito. “On a generalization of de-Rham lemma”. In: *Ann. Inst. Fourier (Grenoble)* 26.2 (1976), pp. vii, 165–170. ISSN: 0373-0956. URL: http://www.numdam.org/item?id=AIF_1976__26_2_0_0.
- [21] Kyoji Saito. *On the periods of primitive integrals I*. preprint Harvard. 1980.
- [22] Kyoji Saito. “Period mapping associated to a primitive form”. In: *Publ. Res. Inst. Math. Sci.* 19.3 (1983), pp. 1231–1264. ISSN: 0034-5318. DOI: 10.2977/prims/1195182028. URL: <https://doi.org/10.2977/prims/1195182028>.
- [23] Kyoji Saito. “The higher residue pairings $K_F^{(k)}$ for a family of hypersurface singular points”. In: *Singularities, Part 2 (Arcata, Calif., 1981)*. Vol. 40. Proc. Sympos. Pure Math. Amer. Math. Soc., Providence, RI, 1983, pp. 441–463.
- [24] Kyoji Saito. “On a linear structure of the quotient variety by a finite reflexion group”. In: *Publ. Res. Inst. Math. Sci.* 29.4 (1993), pp. 535–579. ISSN: 0034-5318. DOI: 10.2977/prims/1195166742. URL: <https://doi.org/10.2977/prims/1195166742>.
- [25] Kyoji Saito and Atsushi Takahashi. “From primitive forms to Frobenius manifolds”. In: *From Hodge theory to integrability and TQFT tt*-geometry*. Vol. 78. Proc. Sympos. Pure Math. Amer. Math. Soc., Providence, RI, 2008, pp. 31–48. DOI: 10.1090/pspum/078/2483747. URL: <https://doi.org/10.1090/pspum/078/2483747>.
- [26] Kyoji Saito, Tamaki Yano, and Jiro Sekiguchi. “On a certain generator system of the ring of invariants of a finite reflection group”. In: *Comm. Algebra* 8.4 (1980), pp. 373–408. ISSN: 0092-7872. DOI: 10.1080/00927878008822464. URL: <https://doi.org/10.1080/00927878008822464>.
- [27] Morihiko Saito. “On the structure of Brieskorn lattice”. In: *Ann. Inst. Fourier (Grenoble)* 39.1 (1989), pp. 27–72. ISSN: 0373-0956. URL: http://www.numdam.org/item?id=AIF_1989__39_1_0_0.
- [28] Jake P. Solomon and Sara B. Tukachinsky. “Relative quantum cohomology”. In: (2019). arXiv: 1906.04795 [math.SG].

- [29] Atsushi Takahashi and Yuuki Shiraishi. “On the Frobenius manifolds for cusp singularities”. In: *Adv. Math.* 273 (2015), pp. 485–522. ISSN: 0001-8708. DOI: 10.1016/j.aim.2014.12.019. URL: <https://doi.org/10.1016/j.aim.2014.12.019>.
- [30] René Thom. *Structural stability and morphogenesis*. Advanced Book Classics. An outline of a general theory of models, Translated from the French by D. H. Fowler, With a foreword by C. H. Waddington. Addison-Wesley Publishing Company, Advanced Book Program, Redwood City, CA, 1989, pp. xxxvi+348. ISBN: 0-201-09419-3.

COLUMBIA UNIVERSITY, 2990 BROADWAY NEW YORK, NY 10027

Email address: aleshkin@math.columbia.edu

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY, SAKYOKU KITASHIRAKAWA, KYOTO, 606-8502, JAPAN, INSTITUTE FOR PHYSICS AND MATHEMATICS OF THE UNIVERSE, THE UNIVERSITY OF TOKYO, 5-1-5 KASHIWANOHA, KASHIWA, 277-8583, JAPAN, LABORATORY OF AGHA, MOSCOW INSTITUTE OF PHYSICS AND TECHNOLOGY, 9 INSTITUTSKIY PER., DOLGOPRUDNY, MOSCOW REGION, 141700, RUSSIAN FEDERATION.

Email address: kyoji.saito@ipmu.jp