



arXiv:2012.01157v1 [math.CV] 2 Dec 2020

On the decay of singular inner functions

Thomas Ransford*

Abstract. It is known that, if $S(z)$ is a non-constant, singular inner function defined on the unit disk, then $\min_{|z|\leq r} |S(z)| \rightarrow 0$ as $r \rightarrow 1^-$. We show that the convergence may be arbitrarily slow.

A *singular inner function* on the unit disk \mathbb{D} is a function of the form

$$S(z) := \exp\left(-\int_{\mathbb{T}} \frac{e^{i\theta} + z}{e^{i\theta} - z} d\mu(\theta)\right) \quad (z \in \mathbb{D}), \tag{1}$$

where μ is a positive finite measure on the unit circle \mathbb{T} that is singular with respect to linear Lebesgue measure. Such functions play an important role in the theory of Hardy spaces H^p . For example, according to the so-called canonical factorization theorem, every function in H^p ($0 < p \leq \infty$) that is not identically zero can be factorized in an essentially unique way as the product of an outer function in H^p , a Blaschke product and a singular inner function (see e.g. [2, Theorem 1.12]).

As the name suggests, a singular inner function S is inner, i.e., it is a bounded holomorphic function whose radial limits satisfy $|S^*(e^{i\theta})| = 1$ Lebesgue-a.e. on \mathbb{T} . In addition, the fact of being singular implies that S also satisfies $S^*(e^{i\theta}) = 0$ μ -a.e. on \mathbb{T} (see e.g. [2, Lemma 1.8]). In particular, if S is non-constant (so $\mu \neq 0$), then S has at least one radial limit equal to zero, and hence

$$\min_{|z|\leq r} |S(z)| \rightarrow 0 \quad (r \rightarrow 1^-). \tag{2}$$

Using the formula (1), it is easy to show that

$$\inf_{0 \leq r < 1} \left(\min_{|z|\leq r} |S(z)| \right)^{1-r} > 0. \tag{3}$$

This imposes a constraint on how rapidly the minimum in (2) can converge to zero, and in fact this constraint is optimal (consider the case when μ is a point mass). In this note, we consider how *slowly* the minimum in (2) can converge to zero. The following theorem shows that the convergence may be arbitrarily slow.

Theorem *Let $\omega : [0, 1) \rightarrow (0, 1)$ be a function such that $\sup_{r \in [0, 1)} \omega(r) < 1$ and $\lim_{r \rightarrow 1^-} \omega(r) = 0$. Then there exists a non-constant singular inner function S on \mathbb{D} such that*

$$\min_{|z|\leq r} |S(z)| \geq \omega(r) \quad (r \in [0, 1)). \tag{4}$$

2020 Mathematics Subject Classification: primary 30J15, secondary 28A78.

Keywords: singular inner function, Hausdorff measure.

*Research supported by grants from NSERC and the Canada research chairs program

Proof Replacing ω by a slightly larger, piecewise-linear function, if necessary, we may assume that, in addition to its other properties, ω is a continuous, decreasing function on $[0, 1)$.

Define two auxiliary functions $g, h : (0, \pi] \rightarrow (0, \infty)$ by

$$g(t) := \frac{1}{3\pi} \log \frac{1}{\omega(1-t/\pi)} \quad \text{and} \quad h(t) := \min \left\{ \sqrt{t}, \inf_{s \in [t, \pi]} sg(s) \right\}.$$

Then g is a positive, continuous, decreasing function on $(0, \pi]$ with $\lim_{t \rightarrow 0^+} g(t) = \infty$, and h is a positive, continuous, increasing function on $(0, \pi]$ such that $\lim_{t \rightarrow 0^+} h(t) = 0$. In addition, we clearly have $h(t) \leq tg(t)$ for all $t \in (0, \pi]$. Notice also that, for $t \in (0, 1]$,

$$\frac{h(t)}{t} = \min \left\{ \frac{\sqrt{t}}{t}, \inf_{s \in [t, \sqrt{t}]} \frac{sg(s)}{t}, \inf_{s \in [\sqrt{t}, \pi]} \frac{sg(s)}{t} \right\} \geq \min \left\{ \frac{1}{\sqrt{t}}, g(\sqrt{t}), \frac{g(\pi)}{\sqrt{t}} \right\},$$

and so $\lim_{t \rightarrow 0^+} h(t)/t = \infty$.

The properties of h (continuous, increasing, $\lim_{t \rightarrow 0^+} h(t) = 0$) make it a so-called measure function, which can be used to define an h -Hausdorff measure, Λ_h (see e.g. [1, Chapter II]). We shall consider Λ_h on the unit circle \mathbb{T} , thought of as a metric space with the arclength distance.

Since $\lim_{t \rightarrow 0^+} h(t)/t = \infty$ and \mathbb{T} has positive linear measure, we have $\Lambda_h(\mathbb{T}) > 0$ (in fact even $\Lambda_h(\mathbb{T}) = \infty$). By [1, Chapter II, Theorem 3], there exists a closed subset E of \mathbb{T} such that $0 < \Lambda_h(E) < \infty$. Again using the fact that $\lim_{t \rightarrow 0^+} h(t)/t = \infty$, the inequality $\Lambda_h(E) < \infty$ implies that E is of linear measure zero.

Now we apply a theorem of Frostman, [1, Chapter II, Theorem 1]. The fact that $\Lambda_h(E) > 0$ implies that there is a finite positive measure μ on E such that $\mu(E) > 0$ and $\mu([\theta - t, \theta + t]) \leq h(t)$ for all $\theta \in [-\pi, \pi]$ and $t \in (0, \pi]$. Since E is of linear measure zero, μ is singular with respect to Lebesgue measure. Define S by (1). Then S is a non-constant, singular inner function, and we shall show that it satisfies the estimate (4).

Fix $r \in [0, 1)$ and $\theta \in [-\pi, \pi]$. We have

$$-\log |S(re^{i\theta})| = \int_{\mathbb{T}} P_r(t - \theta) d\mu(t),$$

where $P_r(\cdot)$ denotes the Poisson kernel, namely

$$P_r(t - \theta) = \frac{1 - r^2}{1 + r^2 - 2r \cos(t - \theta)}.$$

We estimate the integral by splitting it into two, according to whether $|t - \theta|$ is smaller than or greater than ϵ . Here $\epsilon \in (0, \pi]$ is a constant, to be chosen later. Firstly, we have

$$\begin{aligned} \int_{|t-\theta| \leq \epsilon} P_r(t - \theta) d\mu(t) &\leq \int_{|t-\theta| \leq \epsilon} \frac{1 - r^2}{(1 - r)^2} d\mu(t) \\ &\leq \frac{1 - r^2}{(1 - r)^2} \mu([\theta - \epsilon, \theta + \epsilon]) \\ &\leq \frac{1 + r}{1 - r} h(\epsilon) \leq \frac{2}{1 - r} \epsilon g(\epsilon). \end{aligned}$$

Secondly, using integration by parts, we have

$$\begin{aligned}
\int_{|t-\theta|>\epsilon} (P_r(t-\theta) - P_r(\pi)) d\mu(t) &= \int_{|t-\theta|>\epsilon} \int_{s=|t-\theta|}^{\pi} -P_r'(s) ds d\mu(t) \\
&= \int_{s=\epsilon}^{\pi} \int_{|t-\theta|\leq s} -P_r'(s) d\mu(t) ds \\
&= \int_{s=\epsilon}^{\pi} \mu([\theta-s, \theta+s]) (-P_r'(s)) ds \\
&\leq \int_{s=\epsilon}^{\pi} h(s) (-P_r'(s)) ds \\
&\leq \int_{s=\epsilon}^{\pi} g(s) (-sP_r'(s)) ds \\
&\leq g(\epsilon) \int_{s=0}^{\pi} -sP_r'(s) ds \\
&= -\pi P_r(\pi)g(\epsilon) + g(\epsilon) \int_0^{\pi} P_r(s) ds \\
&= -\pi P_r(\pi)g(\epsilon) + \pi g(\epsilon),
\end{aligned}$$

whence

$$\begin{aligned}
\int_{|t-\theta|>\epsilon} P_r(t-\theta) d\mu(t) &\leq -\pi P_r(\pi)g(\epsilon) + \pi g(\epsilon) + P_r(\pi)\mu(\mathbb{T}) \\
&\leq -\pi P_r(\pi)g(\epsilon) + \pi g(\epsilon) + P_r(\pi)\pi g(\pi) \leq \pi g(\epsilon).
\end{aligned}$$

Putting these estimates together, we obtain

$$-\log |S(re^{i\theta})| \leq \left(\frac{2\epsilon}{1-r} + \pi \right) g(\epsilon).$$

In particular, taking $\epsilon = \pi(1-r)$, we get

$$-\log |S(re^{i\theta})| \leq 3\pi g(\pi(1-r)) = \log \frac{1}{\omega(r)}.$$

Rearranging this equality and minimizing over θ yields

$$\min_{|z|=r} |S(z)| \geq \omega(r),$$

which is equivalent to (4), by the minimum principle. ■

Acknowledgement. The theorem above was proved in order to solve a problem posed to me by Joe Cima. I am grateful to Joe for drawing my attention to this question.

References

- [1] L. Carleson, *Selected Problems on Exceptional Sets*, Van Nostrand, Princeton NJ, 1967.
- [2] J. Mashreghi, *Derivatives of Inner Functions*, Springer, New York NY, 2013.

Département de mathématiques et de statistique, Université Laval, Québec, QC, G1V 0A6
e-mail: ransford@mat.ulaval.ca.