

On the Brauer groups of fibrations

Yanshuai Qin

October 10, 2021

Contents

1	Introduction	1
1.1	Tate conjecture and Brauer group	2
1.2	Brauer groups for fibrations	3
1.3	The finiteness of the Galois invariant part of geometric Brauer group	4
1.4	Sketch of the proof of Theorem 1.4	4
1.5	Notation and Terminology	5
2	Geometric Brauer groups and Tate-Shafarevich groups	8
2.1	Preliminary on Tate's conjecture and Brauer groups	8
2.2	The Galois invariant part of geometric Brauer group	8
2.3	Tate-Shafarevich group	12
2.4	Cofiniteness of Brauer groups and Tate-Shafarevich groups	14
3	Proof of the main theorem	15
3.1	The left exactness in Theorem 1.4	15
3.2	The pull-back trick	18
3.3	Proof of Theorem 1.4	20
4	Applications	21
4.1	Proof of Theorem 1.8	21
4.2	Proof of Theorem 1.2	22
4.3	Brauer groups of integral models of abelian varieties and $K3$ surfaces	22

1 Introduction

Let C be the spectrum of the ring of integers in a number field or a smooth projective geometrically connected curve over a finite field with function field K . Let \mathcal{X} be a 2-dimensional regular scheme and $\pi : \mathcal{X} \rightarrow C$ be a proper flat morphism such that

the generic fiber X of π is smooth and geometrically connected over K . Artin and Grothendieck (cf. [Gro3, § 4] or [Ulm, Prop. 5.3]) proved that there is an isomorphism

$$\mathrm{III}(\mathrm{Pic}_{X/K}^0) \cong \mathrm{Br}(\mathcal{X})$$

up to finite groups. Assuming the finiteness of $\mathrm{III}(\mathrm{Pic}_{X/K}^0)$ or $\mathrm{Br}(\mathcal{X})$, Milne [Mil4], Gonzales-Aviles [Goa] and Geisser [Gei1] gave a precise relation between their orders.

In this paper, we generalize the original result of Artin and Grothendieck to fibrations of high relative dimensions over arbitrary finitely generated fields. As a consequence, we reduce the Tate conjecture for divisors on smooth projective varieties over a finitely generated field K to the Tate conjecture for divisors on smooth projective varieties over the prime subfield of K .

For any noetherian scheme X , the *cohomological Brauer group*

$$\mathrm{Br}(X) := H^2(X, \mathbb{G}_m)_{\mathrm{tor}}$$

is defined to be the torsion part of the étale cohomology group $H^2(X, \mathbb{G}_m)$.

1.1 Tate conjecture and Brauer group

Let us first recall the Tate conjecture for divisors over finitely generated fields.

Conjecture 1.1. (*Conjecture $T^1(X, \ell)$*) *Let X be a projective and smooth variety over a finitely generated field k of characteristic $p \geq 0$, and $\ell \neq p$ be a prime number. Then the cycle class map*

$$\mathrm{Pic}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_{\ell} \longrightarrow H^2(X_{k^s}, \mathbb{Q}_{\ell}(1))^{G_k}$$

is surjective.

The conjecture is proved for abelian varieties by Tate [Tat3] (over finite fields), Zarhin [Zar1, Zar2] (over positive characteristics) and Faltings [Fal1, Fal2] (over characteristic zero), for K3 surfaces over characteristic zero by André [And] and Tankeev [Tan4, Tan5], for K3 surfaces over positive characteristics by Nygaard [Nyg], Nygaard-Ogus [NO], Artin-Swinnerton-Dyer [ASD], Maulik [Mau], Charles [Cha] and Madapusi-Pera [MP].

By a well-known result (see Proposition 2.1), $T^1(X, \ell)$ is equivalent to the finiteness of $\mathrm{Br}(X_{k^s})^{G_k}(\ell)$.

Theorem 1.2. *Let k be a prime field and K be a finitely generated field over k . Let ℓ be a prime different from $\mathrm{char}(k)$. Assuming that $T^1(\mathcal{X}, \ell)$ is true for all smooth projective varieties \mathcal{X} over k , then $T^1(X, \ell)$ holds for all smooth projective varieties X over K .*

Remark 1.3. $T^1(X, \ell)$ for smooth projective varieties over finite fields has been reduced to $T^1(X, \ell)$ for smooth projective surfaces X over finite fields with $H^1(X, \mathcal{O}_X) = 0$ by the work of de Jong [deJ2], Morrow [Mor, Thm. 4.3] and Yuan [Yua1, Thm. 1.6].

The above reduction theorem was proved by Ambrosi [Amb]. The idea of our proof is to work on the obstructions of the Tate conjecture, i.e. Brauer groups directly. We obtain the following relation between Brauer groups for fibrations.

1.2 Brauer groups for fibrations

For a torsion abelian group M and a prime number p , let $M(\text{non-}p)$ denote the subgroup of elements of order prime to p . We also define $M(\text{non-}p) := M$ for $p = 0$. For two abelian groups M and N , we say that they are *almost isomorphic* if there exists a subquotient M_1/M_0 of M (resp. N_1/N_0 of N) such that M_0 (resp. N_0) and M/M_1 (resp. N/N_1) are finite and $M_1/M_0 \cong N_1/N_0$. Let $f : M \rightarrow N$ be a homomorphism with a finite cokernel and a kernel almost isomorphic to an abelian group H , in this situation, we say that the sequence $0 \rightarrow H \rightarrow M \rightarrow N \rightarrow 0$ is exact up to finite groups.

Theorem 1.4. *Let k be a finitely generated field of characteristic $p \geq 0$. Let \mathcal{X} be a smooth geometrically connected variety over k and C be a smooth projective geometrically connected curve over k with function field K . Let $\pi : \mathcal{X} \rightarrow C$ be a dominant morphism such that the generic fiber X is smooth projective geometrically connected over K . Let K' denote Kk^s and write P for $\text{Pic}_{X/K, \text{red}}^0$. Define*

$$\mathbb{H}_{K'}(P) := \text{Ker}(H^1(K', P) \rightarrow \prod_{v \in C_{k^s}} H^1(K_v^{sh}, P)).$$

Then there is an exact sequence up to finite groups

$$0 \rightarrow \mathbb{H}_{K'}(P)^{G_k(\text{non-}p)} \rightarrow \text{Br}(\mathcal{X}_{k^s})^{G_k(\text{non-}p)} \rightarrow \text{Br}(X_{K^s})^{G_K(\text{non-}p)} \rightarrow 0.$$

In the case that k is a finite field and \mathcal{X} is projective over k , the following result generalizes Artin-Grothendieck's theorem to high relative dimensions.

Corollary 1.5. *Let $\pi : \mathcal{X} \rightarrow C$ be a proper flat morphism, where C is a smooth projective geometrically connected curve over a finite field k of characteristic p with function field K . Assuming that \mathcal{X} is smooth projective over k and the generic fiber X of π is smooth projective geometrically connected over K , then there is an exact sequence up to finite groups*

$$0 \rightarrow \mathbb{H}(\text{Pic}_{X/K, \text{red}}^0)(\text{non-}p) \rightarrow \text{Br}(\mathcal{X})(\text{non-}p) \rightarrow \text{Br}(X_{K^s})^{G_K(\text{non-}p)} \rightarrow 0.$$

In Theorem 1.4, when k is finite, the canonical maps $\mathbb{H}(P)(\text{non-}p) \rightarrow \mathbb{H}_{K'}(P)^{G_k(\text{non-}p)}$ and $\text{Br}(\mathcal{X}) \rightarrow \text{Br}(\mathcal{X}_{k^s})^{G_k}$ have finite kernels and cokernels (cf. Proposition 2.8 and [Yua2, Cor. 1.4]). Thus, Corollary 1.5 follows from Theorem 1.4. It is well-known that the BSD conjecture for an abelian variety over a global function field is equivalent to the finiteness of the Tate-Shafarevich group (cf. [Mil5] and [Sch]). Thus, Corollary 1.5 implies the following result:

Corollary 1.6. *Notations as in Corollary 1.5, then $T^1(\mathcal{X}, \ell)$ is equivalent to the BSD conjecture for $\text{Pic}_{X/K, \text{red}}^0$ and $T^1(X, \ell)$.*

Remark 1.7. Geisser[Gei2] first studied fibrations of high relative dimensions over finite fields by using etale motivic cohomology theory. He proved Corollary 1.6 and gave a

precise relation between the orders of the Tate-Shafarevich group of the Albanese variety of the generic fiber and the Brauer group of \mathcal{X} under the assumption of the finiteness of $\text{Br}(\mathcal{X})$. For fibrations over the spectrum of the ring of integers in a number field of relative dimension > 1 , the above question was first studied by Tankeev (cf.[Tan1, Tan2, Tan3]). In our forthcoming work, we will prove a version of Corollary 1.5 over the spectrum of the ring of integers in a number field.

1.3 The finiteness of the Galois invariant part of geometric Brauer group

In [SZ1], Skorobogatov and Zarkhin conjectured that $\text{Br}(X_{k^s})^{G_k}$ is finite for any smooth projective variety X over a finitely generated field k . By the following result, assuming Tate conjecture for divisors, we have that $\text{Br}(X_{k^s})^{G_k(\text{non-}p)}$ is finite if k is a finitely generated field of characteristic $p > 0$.

Theorem 1.8. *Let k be a prime field and K be a field finitely generated over k . Then the following statements are true.*

- (1) *If $k = \mathbb{Q}$ and $\text{Br}(X_{k^s})^{G_k}$ is finite for any smooth projective variety X over k , then $\text{Br}(X_{K^s})^{G_K}$ is finite for any smooth projective variety X over K .*
- (2) *If $k = \mathbb{F}_p$ and the Tate conjecture for divisors holds for any smooth projective variety over k , then $\text{Br}(X_{K^s})^{G_K(\text{non-}p)}$ is finite for any smooth projective variety X over K .*

Remark 1.9. In [SZ1, SZ2], Skorobogatov and Zarkhin proved the finiteness of $\text{Br}(X_{k^s})^{G_k(\text{non-}p)}$ for abelian varieties and K3 surfaces ($\text{char}(k) \neq 2$).

1.4 Sketch of the proof of Theorem 1.4

For the proof of Theorem 1.4, the left exactness will follow from Grothendieck's argument in [Gro3, § 4]. The tricky part is to show that the natural map

$$\text{Br}(\mathcal{X}_{k^s})^{G_k(\text{non-}p)} \longrightarrow \text{Br}(X_{K^s})^{G_K(\text{non-}p)}$$

has a finite cokernel. The key idea is to use a pull-back trick developed by Colliot-Thélène and Skorobogatov (cf.[CTS] or [Yua2]) to reduce it to cases of relative dimension 1. Choose an open dense subset $U \subseteq C$ such that $\pi^{-1}(U) \rightarrow U$ is smooth and proper. Write \mathcal{V} for $\pi^{-1}(U)$. By purity for Brauer groups, the natural map

$$\text{Br}(\mathcal{X}_{k^s})^{G_k(\text{non-}p)} \longrightarrow \text{Br}(\mathcal{V}_{k^s})^{G_k(\text{non-}p)}$$

is injective and has a finite cokernel. It suffices to show that

$$\text{Br}(\mathcal{V})(\text{non-}p) \longrightarrow \text{Br}(X_{K^s})^{G_K(\text{non-}p)}$$

has a finite cokernel. For simplicity, we assume that $X(K)$ is not empty. By the spectral sequence

$$E_2^{p,q} = H^p(U, R^q \pi_* \mathbb{G}_m) \Rightarrow H^{p+q}(\mathcal{V}, \mathbb{G}_m),$$

we get an exact sequence

$$H^2(\mathcal{V}, \mathbb{G}_m) \longrightarrow H^0(U, R^2 \pi_* \mathbb{G}_m) \xrightarrow{d_2^{0,2}} H^2(U, R^1 \pi_* \mathbb{G}_m).$$

And we will show there is a canonical isomorphism

$$H^0(U, R^2 \pi_* \mathbb{G}_m)(\text{non-}p) \cong \text{Br}(X_{K^s})^{G_K}(\text{non-}p).$$

So the question is reduced to show the vanishing of $d_2^{0,2}$. In the case of relative dimension 1, this is obvious since $R^2 \pi_* \mathbb{G}_m = 0$ by Artin's theorem [Gro3, Cor. 3.2]. In general, taking a smooth projective curve $Y \subseteq X$, then spreading out, we get a relative curve $\pi' : \mathcal{Y} \longrightarrow C$ and a C -morphism $\mathcal{Y} \longrightarrow \mathcal{X}$. This gives a commutative diagram

$$\begin{array}{ccc} H^0(U, R^2 \pi_* \mathbb{G}_m) & \longrightarrow & H^2(U, R^1 \pi_* \mathbb{G}_m) \\ \downarrow & & \downarrow \\ H^0(U, R^2 \pi'_* \mathbb{G}_m) & \longrightarrow & H^2(U, R^1 \pi'_* \mathbb{G}_m) \end{array}$$

We hope that the second column is injective. In fact, we need finitely many $\pi^i : \mathcal{Y}_i \longrightarrow C$ such that the induced morphism of abelian sheaves

$$R^1 \pi_* \mathbb{G}_m \longrightarrow \bigoplus_i R^1 \pi_*^i \mathbb{G}_m$$

is split up to abelian sheaves of finite exponent. The splitting implies that the natural map

$$H^2(U, R^1 \pi_* \mathbb{G}_m) \longrightarrow \bigoplus_i H^2(U, R^1 \pi_*^i \mathbb{G}_m)$$

has a kernel of finite exponent. It follows that $d_2^{0,2}$ has an image of finite exponent.

To get this splitness, we will choose Y_i as the complete intersection of very ample divisors on X such that there exists an abelian variety A/K and a G_K -equivariant map

$$\text{Pic}(X_{K^s}) \times A(K^s) \longrightarrow \bigoplus_i \text{Pic}(Y_{i,K^s})$$

with finite kernel and cokernel (cf. [Yua2, §3.3, p17]).

1.5 Notation and Terminology

Fields

By a *finitely generated field*, we mean a field which is finitely generated over a prime field. For any field k , denote by k^s the separable closure. Denote by $G_k = \text{Gal}(k^s/k)$ the absolute Galois group of k .

Henselization

Let R be a noetherian local ring, denote by R^h (resp. R^{sh}) the henselization (resp. strict henselization) of R at the maximal ideal. If R is a domain, denote by K^h (resp. K^{sh}) the fraction field of R^h (resp. R^{sh}).

Varieties

By a *variety* over a field k , we mean a scheme which is separated and of finite type over k .

For a smooth proper geometrically connected variety X over a field k , we use $\text{Pic}_{X/k}^0$ to denote the identity component of the Picard scheme $\text{Pic}_{X/k}$. Denote by $\text{Pic}_{X/k,\text{red}}^0$ the underlying reduced closed subscheme of $\text{Pic}_{X/k}^0$ (which is known to be an abelian variety cf. FGA 236-17, Cor. 3.2.)

Cohomology

The default sheaves and cohomology over schemes are with respect to the small étale site. So H^i is the abbreviation of $H_{\text{ét}}^i$.

Brauer groups

For any noetherian scheme X , denote the *cohomological Brauer group*

$$\text{Br}(X) := H^2(X, \mathbb{G}_m)_{\text{tor}}.$$

Tate-Shafarevich group

For an abelian variety A defined over a global field K , denote the *Tate–Shafarevich group*

$$\text{III}(A) := \text{Ker}(H^1(G_K, A) \longrightarrow \prod_v H^1(G_{K_v}, A)),$$

where K_v denotes the completion of K at a place v .

Abelian group

For any abelian group M , integer m and prime ℓ , we set

$$\begin{aligned} M[m] &= \{x \in M \mid mx = 0\}, & M_{\text{tor}} &= \bigcup_{m \geq 1} M[m], & M(\ell) &= \bigcup_{n \geq 1} M[\ell^n], \\ T_\ell M &= \text{Hom}_{\mathbb{Z}}(\mathbb{Q}_\ell/\mathbb{Z}_\ell, M) = \varprojlim_n M[\ell^n], & V_\ell M &= T_\ell(M) \otimes_{\mathbb{Z}_\ell} \mathbb{Q}_\ell. \end{aligned}$$

For any torsion abelian group M and prime number p , we set

$$M(\text{non-}p) = \bigcup_{p \nmid m} M[m].$$

For two abelian groups M and N , we say that they are *almost isomorphic* if there exist filtrations $M_0 \subseteq M_1 \subseteq M$ and $N_0 \subseteq N_1 \subseteq N$ such that M_0 (resp. N_0) and M/M_1 (resp. N/N_1) are finite and M_1/M_0 is isomorphic to N_1/N_0 .

Acknowledgements:

I would like to thank my advisor Xinyi Yuan for suggesting this question to me and for telling me the pull-back trick. I also want to thank Jean-Louis Colliot-Thélène and Alexei N. Skorobogatov for helpful suggestions on references.

2 Geometric Brauer groups and Tate-Shafarevich groups

2.1 Preliminary on Tate's conjecture and Brauer groups

Proposition 2.1. *Let X be a projective and smooth variety over a finitely generated field k of characteristic $p \geq 0$, and $\ell \neq p$ be a prime number. Then the cycle class map*

$$\text{Pic}(X) \otimes_{\mathbb{Z}} \mathbb{Q}_{\ell} \longrightarrow H^2(X_{k^s}, \mathbb{Q}_{\ell}(1))^{G_k}$$

is surjective if and only if $\text{Br}(X_{k^s})^{G_k}(\ell)$ is finite.

Proof. See [Yua2, Thm. 1.2]. □

Lemma 2.2. *Let X be a projective and smooth variety over a field k of characteristic $p \geq 0$, and $\ell \neq p$ be a prime number. Then there is a canonical G_k -splitting exact sequence*

$$0 \longrightarrow \text{NS}(X_{k^s}) \otimes_{\mathbb{Z}} \mathbb{Q}_{\ell} \longrightarrow H^2(X_{k^s}, \mathbb{Q}_{\ell}(1)) \longrightarrow V_{\ell}\text{Br}(X_{k^s}) \longrightarrow 0$$

Proof. See [Yua2, §2.2]. □

2.2 The Galois invariant part of geometric Brauer group

In this section, we will show that the Galois invariant part of the geometric Brauer group of a smooth variety is a birational invariant up to finite groups. This will allow us to shrink the base of a fibration without changing the question.

The following results in the case of characteristic 0 were proved in [CTS] (cf. [CTS, Prop. 6.1 and Thm. 6.2 (iii)]).

Proposition 2.3. *Let X be a smooth geometrically connected variety over a finitely generated field k of characteristic $p \geq 0$. Let $U \subseteq X$ be an open dense subset. Then the natural map*

$$\text{Br}(X_{k^s})^{G_k}(\text{non-}p) \longrightarrow \text{Br}(U_{k^s})^{G_k}(\text{non-}p)$$

is injective and has a finite cokernel.

Proof. The injectivity follows from that X_{k^s} is regular and irreducible. To show that the cokernel is finite, we need the lemma below. Let Y denote $X - U$ with reduced scheme structure. By purity for Brauer group (cf. [Čes, Thm. 1.1]), removing a close subset of codimension ≥ 2 will not change the Brauer group. Thus, by shrinking X , we might assume that Y is regular and of codimension 1 in X . Let $\ell \neq p$ be a prime. By purity for Brauer group (cf. [Gro3, §6] and [Fuj]), there is a canonical exact sequence

$$0 \longrightarrow \text{Br}(X_{k^s})(\ell) \longrightarrow \text{Br}(U_{k^s})(\ell) \longrightarrow H^1(Y_{k^s}, \mathbb{Q}_{\ell}/\mathbb{Z}_{\ell}).$$

Thus, it suffices to show that the group

$$H^1(Y_{k^s}, \mathbb{Q}_{\ell}/\mathbb{Z}_{\ell})^{G_k}$$

is finite and vanishes for all but finitely many ℓ . This follows from the lemma below. □

Lemma 2.4. *Let U be a regular variety over a finitely generated field k . Let $\ell \neq \text{char}(k)$ be a prime. Then*

$$H^1(U_{k^s}, \mathbb{Q}_\ell/\mathbb{Z}_\ell)^{G_k}$$

is finite and vanishes for all but finitely many ℓ .

Proof. Firstly, we will show that it suffices to prove the claim for smooth varieties over k . Notice that we can replace k by a finite separable extension. Thus, we might assume that U_{k^s} is irreducible. Let V be an open dense subset of U_{k^s} . By semi-purity (cf. [Fuj, §8]),

$$H^1(U_{k^s}, \mathbb{Z}/\ell^n) \longrightarrow H^1(V, \mathbb{Z}/\ell^n)$$

is injective. We might assume that V is defined over k . Therefore, it suffices to prove the claim for an open dense subset. Since U_{k^s} and $U_{\bar{k}}$ have the same underlying topological space, by shrinking U_{k^s} , we might assume that $(U_{\bar{k}})_{\text{red}}$ is irreducible and smooth over \bar{k} . So there exists a finite extension l/k such that $(U_l)_{\text{red}}$ is irreducible and smooth over l . We might assume that l/k is purely inseparable. Then $l^s = l \otimes_k k^s$ and $G_l = G_k$. Let V denote $(U_l)_{\text{red}}$. Thus $V_{l^s} = V \times_{\text{Spec } k} \text{Spec } k^s$. The k -morphism $V \rightarrow U$ induces a G_k -equivariant isomorphism

$$H^1(U_{k^s}, \mathbb{Z}/\ell^n) \cong H^1(V_{l^s}, \mathbb{Z}/\ell^n),$$

so it suffices to prove the claim for V . Thus we might assume that U is smooth and geometrically connected over k .

By above arguments, we can always replace k by a finite extension and shrink U . By de Jong's alteration theorem, we might assume that there is a finite flat morphism $f : V \rightarrow U$ such that V admits a smooth projective geometrically connected compactification over k . Since the kernel of

$$H^1(U_{k^s}, \mathbb{Z}/\ell^n) \longrightarrow H^1(V_{k^s}, \mathbb{Z}/\ell^n)$$

is killed by the degree of f (cf. Lemma 2.6 below), it suffices to prove the claim for V . Thus we might assume that U is an open subvariety of a smooth projective geometrically connected variety X over k . Let Y_i be irreducible components of $X - U$ of codimension 1. Let D_i be the regular locus of Y_i . By extending k to a finite separable extension, we might assume that D_i is geometrically irreducible. By purity, there is a canonical exact sequence

$$0 \longrightarrow H^1(X_{k^s}, \mathbb{Z}/\ell^n) \longrightarrow H^1(U_{k^s}, \mathbb{Z}/\ell^n) \longrightarrow \bigoplus_i H^0(D_{i,k^s}, \mathbb{Z}/\ell^n(-1))$$

Taking G_k -invariants, we get an exact sequence

$$0 \longrightarrow H^1(X_{k^s}, \mathbb{Z}/\ell^n)^{G_k} \longrightarrow H^1(U_{k^s}, \mathbb{Z}/\ell^n)^{G_k} \longrightarrow \bigoplus_i H^0(D_{i,k^s}, \mathbb{Z}/\ell^n(-1))^{G_k}.$$

It suffices to show that the size of the first and third group are bounded independent of n and is equal to 1 for all but finitely many ℓ . Since D_{i,k^s} is connected, we have

$$H^0(D_{i,k^s}, \mathbb{Z}/\ell^n(-1)) = \text{Hom}(\mu_{\ell^n}, \mathbb{Z}/\ell^n).$$

By the lemma below, the size of the third group is bounded independent of n and is equal to 1 for all but finitely many ℓ . Since

$$H^1(X_{k^s}, \mathbb{Z}/\ell^n) \cong \text{Pic}_{X/K}[\ell^n](-1)$$

and

$$0 \longrightarrow \text{Pic}_{X/K, \text{red}}^0[\ell^n] \longrightarrow \text{Pic}_{X/K}[\ell^n] \longrightarrow \text{NS}(X_{k^s})[\ell^n]$$

is exact, it suffices to show that the size of $(\text{Pic}_{X/K, \text{red}}^0[\ell^n](-1))^{G_k}$ is bounded independent of n and is equal to 1 for all but finitely many ℓ . Let A be the dual of $\text{Pic}_{X/K, \text{red}}^0$. By the Weil pairing,

$$\text{Pic}_{X/K, \text{red}}^0[\ell^n](-1) \cong \text{Hom}(A[\ell^n], \mathbb{Z}/\ell^n).$$

Taking G_k -invariants, we get

$$(\text{Pic}_{X/K, \text{red}}^0[\ell^n](-1))^{G_k} \cong \text{Hom}(A[\ell^n], \mathbb{Z}/\ell^n)^{G_k}.$$

By the lemma below, the claim follows. \square

Lemma 2.5. *Let A be an abelian variety over a finitely generated field K of characteristic $p \geq 0$. Let $\ell \neq p$ be a prime. Then the sizes of*

$$\text{Hom}(A[\ell^n], \mathbb{Z}/\ell^n)^{G_K} \quad \text{and} \quad \text{Hom}(\mu_{\ell^n}, \mathbb{Z}/\ell^n)^{G_K}$$

are bounded independent of n . Moreover, for all but finitely many ℓ , these two groups vanish for any n .

Proof. We will only prove the claim for the first group, since the second one follows from the same arguments. Firstly, we assume that K is finite, then we will use a specialization technique to reduce the general case to the finite field case.

If K is finite, then we have

$$\text{Hom}(A[\ell^n], \mathbb{Z}/\ell^n)^{G_K} = \text{Hom}(A[\ell^n]_{G_K}, \mathbb{Z}/\ell^n),$$

which has the same size as $A[\ell^n]^{G_K}$. Then the claim follows from the finiteness of $A(K)$.

In general, choose an integral regular scheme S of finite type over $\text{Spec } \mathbb{Z}$ with function field K . By Shrinking S , we might assume A/K extends to an abelian scheme \mathcal{A}/S . Fix a closed point $s \in S$. For any $\ell \neq \text{char}(k(s))$, we can shrink S such that ℓ is invertible on S and $s \in S$. Then the etale sheaf $\mathcal{A}[\ell^n]$ is a locally constant sheaf of \mathbb{Z}/ℓ^n -module since $\mathcal{A}[\ell^n]$ is finite etale over S . Thus

$$\mathcal{H}om(\mathcal{A}[\ell^n], \mathbb{Z}/\ell^n)$$

is also a locally constant sheaf of \mathbb{Z}/ℓ^n -module and its stalk at the generic point can be identified with

$$\text{Hom}(A[\ell^n], \mathbb{Z}/\ell^n).$$

Set $\mathcal{F} = \mathcal{H}om(\mathcal{A}[\ell^n], \mathbb{Z}/\ell^n)$, we have

$$\mathrm{Hom}(A[\ell^n], \mathbb{Z}/\ell^n)^{G_K} = H^0(S, \mathcal{F}) \hookrightarrow H^0(s, \mathcal{F}).$$

Since $H^0(s, \mathcal{F}) = \mathrm{Hom}(\mathcal{A}_s[\ell^n], \mathbb{Z}/\ell^n)^{G_{k(s)}}$ and $k(s)$ is finite, thus the claim holds for $\ell \neq \mathrm{char}(k(s))$. In the case $\mathrm{char}(K) = 0$, we can choose another closed point s' such that $\mathrm{char}(k(s')) \neq \mathrm{char}(k(s))$, then by the same argument, the claim also holds for $\ell = \mathrm{char}(k(s))$. This completes the proof. \square

Lemma 2.6. *Let $f : X \rightarrow Y$ be a dominant morphism between regular integral schemes. Assuming that $K(X)/K(Y)$ is a finite extension of degree d , then the induced map*

$$\mathrm{Br}(Y) \rightarrow \mathrm{Br}(X)$$

has a kernel killed by d . Moreover, if ℓ is a prime invertible on Y , then the induced map

$$H^1(Y, \mu_{\ell^n}) \rightarrow H^1(X, \mu_{\ell^n})$$

also has a kernel killed by d .

Proof. We will only prove the first claim since the second one follows from the same arguments. Since X and Y are regular, we have

$$\mathrm{Br}(X) \rightarrow \mathrm{Br}(K(X)) \quad \text{and} \quad \mathrm{Br}(Y) \rightarrow \mathrm{Br}(K(Y))$$

are injective. It suffices to show

$$\mathrm{Br}(K(Y)) \rightarrow \mathrm{Br}(K(X))$$

has a kernel killed by d . Let L (resp. K) denote $K(X)$ (resp. $K(Y)$). Since $f : \mathrm{Spec} L \rightarrow \mathrm{Spec} K$ is finite, we have

$$H^2(\mathrm{Spec} L, \mathbb{G}_m) = H^2(\mathrm{Spec} K, f_* \mathbb{G}_m) = H^2(K, (K^s \otimes_K L)^\times).$$

There is a norm map $(K^s \otimes_K L)^\times \rightarrow (K^s)^\times$ such that the composition of $(K^s)^\times \rightarrow (K^s \otimes_K L)^\times \rightarrow (K^s)^\times$ is the d -th power map. Taking cohomology, the composition of

$$H^2(K, (K^s)^\times) \rightarrow H^2(K, (K^s \otimes_K L)^\times) \rightarrow H^2(K, (K^s)^\times)$$

is equal to multiplication by d . It follows that

$$\mathrm{Ker}(\mathrm{Br}(K) \rightarrow \mathrm{Br}(L))$$

is killed by d . \square

2.3 Tate-Shafarevich group

In this section, first, we will study a geometric version of the Tate-Shafarevich group for an abelian variety over a function field with a base field k . Then, we will prove that the Galois fixed part of the geometric Tate-Shafarevich group is canonically isomorphic to Tate-Shafarevich group up to finite groups in the case that k is a finite field. The idea is reducing the question to the relation between arithmetic Brauer groups and geometric Brauer groups which was studied in [CTS](cf. [Yua2]).

Proposition 2.7. *Let C be a smooth projective geometrically connected curve defined over a finitely generated field k of characteristic $p \geq 0$. Let K be the function field of C and A be an abelian variety over K . Denote Kk^s by K' . Let $U \subseteq C$ be an open dense subscheme. Define*

$$\text{III}_{U_{k^s}}(A) := \text{Ker}(H^1(K', A) \longrightarrow \prod_{v \in U_{k^s}} H^1(K_v^{sh}, A))$$

Then the natural map

$$\text{III}_{C_{k^s}}(A)^{G_k(\text{non-}p)} \longrightarrow \text{III}_{U_{k^s}}(A)^{G_k(\text{non-}p)}$$

is injective and has a cokernel of finite exponent.

Proof. By definitions, the injectivity is obvious. It suffices to show that the cokernel is of finite exponent. Since there exists an abelian variety B/K such that $A \times B$ is isogenous to $\text{Pic}_{X/K}^0$ for some smooth projective geometrically connected curve X over K , it suffices to prove the claim for $A = \text{Pic}_{X/K}^0$. Without loss of generality, we might replace k by a finite extension l/k . This is obvious if l/k is separable. For l/k a purely inseparable extension of degree p^n , the quotient $A(K^{sl^s})/A(K^s)$ is killed by p^n . Then $H^1(K', A) \longrightarrow H^1(Kl^s, A)$ has a kernel and a cokernel killed by p^n . It follows that $\text{III}_{U_{k^s}}(A) \longrightarrow \text{III}_{U_{l^s}}(A)$ has a kernel and a cokernel killed by some power of p . By resolution of singularity of surfaces, $X \longrightarrow \text{Spec } K$ admits a proper regular model $\pi : \mathcal{X} \longrightarrow C$. By extending k , we might assume that \mathcal{X} is smooth proper geometrically connected over k . Write V for $\pi^{-1}(U)$. By the Leray spectral sequence

$$E_2^{p,q} = H^p(U_{k^s}, R^q \pi_* \mathbb{G}_m) \Rightarrow H^{p+q}(V_{k^s}, \mathbb{G}_m),$$

we get a long exact sequence

$$\begin{aligned} H^2(U_{k^s}, \mathbb{G}_m) &\longrightarrow \text{Ker}(H^2(V_{k^s}, \mathbb{G}_m) \longrightarrow H^0(U_{k^s}, R^2 \pi_* \mathbb{G}_m)) \\ &\longrightarrow H^1(U_{k^s}, R^1 \pi_* \mathbb{G}_m) \longrightarrow H^3(U_{k^s}, \mathbb{G}_m). \end{aligned}$$

By [Gro3, Cor. 3.2 and Lem. 3.2.1], $R^2 \pi_* \mathbb{G}_m = 0$ and $H^i(U_{k^s}, \mathbb{G}_m)(\text{non-}p) = 0$ for $i \geq 2$. Thus, we have a canonical isomorphism

$$H^2(V_{k^s}, \mathbb{G}_m)(\text{non-}p) \cong H^1(U_{k^s}, R^1 \pi_* \mathbb{G}_m)(\text{non-}p).$$

Let $j : \text{Spec } K' \longrightarrow C_{k^s}$ be the generic point. By the spectral sequence

$$H^p(U_{k^s}, R^q j_* (j^* R^1 \pi_* \mathbb{G}_m)) \Rightarrow H^{p+q}(K', \text{Pic}(X_{K^s})),$$

we have

$$H^1(U_{k^s}, j_* j^* R^1 \pi_* \mathbb{G}_m) = \text{Ker}(H^1(K', \text{Pic}(X_{K^s})) \longrightarrow \prod_{v \in U_{k^s}} H^1(K_v^{sh}, \text{Pic}(X_{K^s}))).$$

Since $\text{Pic}(X_{K^s}) = \text{Pic}_{X/K}^0(K^s) \oplus \mathbb{Z}$ as G_K -modules, we get a natural isomorphism

$$H^1(U_{k^s}, j_* j^* R^1 \pi_* \mathbb{G}_m) \cong \text{III}_{U_{k^s}}(\text{Pic}_{X/K}^0).$$

Without loss of generality, we might shrink U such that π is smooth on $\pi^{-1}(U)$. Then, by Lemma 3.1, the natural map

$$R^1 \pi_* \mathbb{G}_m \longrightarrow j_* j^* R^1 \pi_* \mathbb{G}_m$$

is an isomorphism on U . It follows that

$$\begin{aligned} H^2(V_{k^s}, \mathbb{G}_m)(\text{non-}p) &\cong H^1(U_{k^s}, R^1 \pi_* \mathbb{G}_m)(\text{non-}p) \\ &\cong H^1(U_{k^s}, j_* j^* R^1 \pi_* \mathbb{G}_m)(\text{non-}p) \cong \text{III}_{U_{k^s}}(\text{Pic}_{X/K}^0)(\text{non-}p). \end{aligned}$$

Consider the following commutative diagram

$$\begin{array}{ccccc} \text{Br}(\mathcal{X}_{k^s})(\text{non-}p) & \longrightarrow & H^1(C_{k^s}, j_* j^* R^1 \pi_* \mathbb{G}_m)(\text{non-}p) & \longrightarrow & \text{III}_{C_{k^s}}(\text{Pic}_{X/K}^0)(\text{non-}p) \\ \downarrow & & \downarrow & & \downarrow \\ \text{Br}(V_{k^s})(\text{non-}p) & \longrightarrow & H^1(U_{k^s}, j_* j^* R^1 \pi_* \mathbb{G}_m)(\text{non-}p) & \longrightarrow & \text{III}_{U_{k^s}}(\text{Pic}_{X/K}^0)(\text{non-}p) \end{array}$$

Since maps on the bottom are isomorphisms, it suffices to show that the natural map

$$\text{Br}(\mathcal{X}_{k^s})^{G_k}(\text{non-}p) \longrightarrow \text{Br}(V_{k^s})^{G_k}(\text{non-}p)$$

has a finite cokernel. This follows from Proposition 2.3. \square

Proposition 2.8. *Notations as in the above proposition. Assuming that k is a finite field, define*

$$\text{III}_{K'}(A) := \text{Ker}(H^1(K', A) \longrightarrow \prod_{v \in C_{k^s}} H^1(K_v^{sh}, A)).$$

Then the natural map

$$\text{III}(A)(\text{non-}p) \longrightarrow \text{III}_{K'}(A)^{G_k}(\text{non-}p)$$

has a kernel and a cokernel of finite exponent.

Proof. We use the same arguments as in the proof of the previous lemma. It suffices to prove the claim for $A = \text{Pic}_{Y/K}^0$ where Y is a smooth projective geometrically connected curve over k . Y admits a projective regular model $\mathcal{Y} \rightarrow C$. Then we have

$$\text{Br}(\mathcal{Y}) \cong \text{III}(\text{Pic}_{Y/K}^0)$$

and

$$\text{Br}(\mathcal{Y}_{k^s}) \cong \text{III}_{K'}(\text{Pic}_{Y/K}^0)$$

up to finite groups. Thus the question is reduced to show that

$$\text{Br}(\mathcal{Y}) \rightarrow \text{Br}(\mathcal{Y}_{k^s})^{G_k}$$

has a finite kernel and a finite cokernel. This follows from [Yua2, Cor. 1.4]. \square

2.4 Cofiniteness of Brauer groups and Tate-Shafarevich groups

Let ℓ be a prime number. Recall that a ℓ -torsion abelian group M is of *cofinite type* if $M[\ell]$ is finite. This is also equivalent to that M can be written as $(\mathbb{Q}_\ell/\mathbb{Z}_\ell)^r \oplus M_0$ for some integer $r \geq 0$ and some finite group M_0 . We also say that a torsion abelian group M is of cofinite type if $M(\ell)$ is of cofinite type for all primes ℓ . It is easy to see that a morphism $M \rightarrow N$ between abelian groups of cofinite type has a kernel and a cokernel of finite exponent if and only if the kernel and cokernel are finite.

In the following, we will show the cofiniteness of geometric Tate-Shafarevich groups and geometric Brauer groups defined in previous sections.

Lemma 2.9. *The group $\text{III}_{U_{k^s}}(A)(\ell)$ defined in the Proposition 2.7 is of cofinite type for any $\ell \neq p$.*

Proof. We might shrink U such that A extends to an abelian scheme \mathcal{A} over U . By the exact sequence

$$0 \rightarrow \mathcal{A}[\ell] \rightarrow \mathcal{A} \xrightarrow{\ell} \mathcal{A} \rightarrow 0$$

we get a surjection

$$H^1(U_{k^s}, \mathcal{A}[\ell]) \rightarrow H^1(U_{k^s}, \mathcal{A})[\ell].$$

Since $H^1(U_{k^s}, \mathcal{A}[\ell])$ is finite, $H^1(U_{k^s}, \mathcal{A})[\ell]$ is also finite. Thus, it suffices to show

$$H^1(U_{k^s}, \mathcal{A}) \cong \text{III}_{U_{k^s}}(A).$$

This follows from $\mathcal{A} \cong j_* A$, since \mathcal{A} is a Néron model of A over U , where $j : \text{Spec } K \rightarrow U$ is the generic point. \square

Lemma 2.10. *Let X be a smooth variety over a separable closed field k of characteristic $p \geq 0$. Let $\ell \neq p$ be a prime. Then $\text{Br}(X)(\ell)$ is of cofinite type.*

Proof. The Kummer exact sequence

$$0 \longrightarrow \mu_\ell \longrightarrow \mathbb{G}_m \xrightarrow{\ell} \mathbb{G}_m \longrightarrow 0$$

induces a surjection

$$H^2(X, \mu_\ell) \longrightarrow \mathrm{Br}(X)[\ell].$$

Then the claim follows from the finiteness of $H^2(X, \mu_\ell)$. □

3 Proof of the main theorem

3.1 The left exactness in Theorem 1.4

In this section, we will prove the left exactness of the sequence in Theorem 1.4 by following Grothendieck's arguments in [Gro3, §4].

Lemma 3.1. *Let U be an irreducible regular scheme of dimension 1 with function field K . Let $\pi : \mathcal{X} \rightarrow U$ be a smooth proper morphism with a generic fiber geometrically connected over K . Let $j : \mathrm{Spec} K \rightarrow U$ be the generic point of U . Then we have*

(a) *the natural map*

$$R^1\pi_*\mathbb{G}_m \longrightarrow j_*j^*R^1\pi_*\mathbb{G}_m$$

is an isomorphism,

(b) *the natural map*

$$R^2\pi_*\mathbb{G}_m(\ell) \longrightarrow j_*j^*R^2\pi_*\mathbb{G}_m(\ell)$$

is an isomorphism for any prime ℓ invertible on U .

Proof. It suffices to show that the induced maps on stalks are isomorphism. Thus, we might assume that $U = \mathrm{Spec} R$ where R is a strictly henselian DVR.

Let X denote the generic fiber. Let $s \in U$ be the closed point. Then we have

$$(R^1\pi_*\mathbb{G}_m)_{\bar{s}} = \mathrm{Pic}(\mathcal{X}) \quad \text{and} \quad (j_*j^*R^1\pi_*\mathbb{G}_m)_{\bar{s}} = \mathrm{Pic}_{X/K}(K).$$

Since \mathcal{X}_s admits a section $s \rightarrow \mathcal{X}_s$ and π is smooth, the section can be extended to a section $U \rightarrow \mathcal{X}$. Thus $X(K)$ is not empty. So

$$\mathrm{Pic}_{X/K}(K) = \mathrm{Pic}(X).$$

Since \mathcal{X} is regular, the natural map

$$\mathrm{Pic}(\mathcal{X}) \longrightarrow \mathrm{Pic}(X)$$

is surjective and has a kernel generated by vertical divisors. It suffices to show that \mathcal{X}_s is connected. This actually follows from $\pi_*\mathcal{O}_{\mathcal{X}} = \mathcal{O}_U$ (cf. [Har, Chap. III, Cor. 11.3]). This proves (a).

Let I denote G_K . For (b), the induced map on the stalk at s is

$$\mathrm{Br}(\mathcal{X})(\ell) \longrightarrow \mathrm{Br}(X_{K^s})^I(\ell).$$

Since π is smooth and proper, we have

$$H^2(\mathcal{X}, \mu_{\ell^\infty}) \cong H^2(X_{K^s}, \mu_{\ell^\infty}) = H^2(X_{K^s}, \mu_{\ell^\infty})^I.$$

Consider

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathrm{Pic}(\mathcal{X}) \otimes \mathbb{Q}_\ell/\mathbb{Z}_\ell & \longrightarrow & H^2(\mathcal{X}, \mu_{\ell^\infty}) & \longrightarrow & \mathrm{Br}(\mathcal{X})(\ell) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \mathrm{NS}(X_{K^s}) \otimes \mathbb{Q}_\ell/\mathbb{Z}_\ell & \longrightarrow & H^2(X_{K^s}, \mu_{\ell^\infty}) & \longrightarrow & \mathrm{Br}(X_{K^s})(\ell) & \longrightarrow & 0 \end{array}$$

Since $\mathrm{NS}(X_{K^s}) \otimes \mathbb{Q}_\ell/\mathbb{Z}_\ell$ is I -invariant and $\mathrm{Pic}(\mathcal{X}) = \mathrm{Pic}(X)$. It suffices to show that

$$\mathrm{Pic}(X) \otimes \mathbb{Q}_\ell/\mathbb{Z}_\ell \longrightarrow (\mathrm{NS}(X_{K^s}) \otimes \mathbb{Q}_\ell/\mathbb{Z}_\ell)^I$$

is surjective. Write $\mathrm{NS}(X_{K^s})_{free}$ for $\mathrm{NS}(X_{K^s})/\mathrm{NS}(X_{K^s})_{tor}$. The action of I on $\mathrm{NS}(X_{K^s})$ factors through a finite quotient I' . Consider the exact sequence

$$\begin{aligned} 0 &\longrightarrow (\mathrm{NS}(X_{K^s})_{free})^{I'} \otimes \mathbb{Z}_\ell \longrightarrow \mathrm{NS}(X_{K^s})^{I'} \otimes \mathbb{Q}_\ell \longrightarrow (\mathrm{NS}(X_{K^s}) \otimes \mathbb{Q}_\ell/\mathbb{Z}_\ell)^I \\ &\longrightarrow H^1(I', \mathrm{NS}(X_{K^s})_{free} \otimes \mathbb{Z}_\ell). \end{aligned}$$

$H^1(I', \mathrm{NS}(X_{K^s})_{free} \otimes \mathbb{Z}_\ell)$ is killed by the order of I' . Since $(\mathrm{NS}(X_{K^s}) \otimes \mathbb{Q}_\ell/\mathbb{Z}_\ell)^I = \mathrm{NS}(X_{K^s}) \otimes \mathbb{Q}_\ell/\mathbb{Z}_\ell$ is divisible, so the image of the last map is zero. Since

$$\mathrm{Pic}(X) \otimes \mathbb{Q}_\ell \longrightarrow \mathrm{NS}(X_{K^s})^I \otimes \mathbb{Q}_\ell$$

is surjective (cf. [Yua2, §2.2]), the claim follows. By Snake Lemma, the natural map

$$\mathrm{Br}(\mathcal{X})(\ell) \longrightarrow \mathrm{Br}(X_{K^s})^I(\ell)$$

is an isomorphism. □

Lemma 3.2. *Let U be a smooth geometrically connected curve over a field k of characteristic $p \geq 0$ with function field K . Let $\pi : \mathcal{X} \rightarrow U$ be a smooth proper morphism with a generic fiber X geometrically connected over K . Let K' denote Kk^s . Define*

$$\mathbb{H}_{U_{k^s}}(\mathrm{Pic}_{X/K}) := \mathrm{Ker}(H^1(K', \mathrm{Pic}(X_{K^s})) \longrightarrow \prod_{v \in U_{k^s}} H^1(K_v^{sh}, \mathrm{Pic}(X_{K^s}))).$$

Then we have a canonical exact sequence

$$0 \rightarrow (\mathbb{H}_{U_{k^s}}(\mathrm{Pic}_{X/K}))^{G_k}(\mathrm{non-}p) \rightarrow \mathrm{Br}(\mathcal{X}_{k^s})^{G_k}(\mathrm{non-}p) \rightarrow \mathrm{Br}(X_{K^s})^{G_K}(\mathrm{non-}p).$$

Moreover, the natural map

$$(\mathbb{H}_{U_{k^s}}(\mathrm{Pic}_{X/K, \mathrm{red}}^0))^{G_k} \longrightarrow (\mathbb{H}_{U_{k^s}}(\mathrm{Pic}_{X/K}))^{G_k}$$

has a kernel and cokernel of finite exponent.

Proof. We might assume that $k = k^s$. By the Leray spectral sequence

$$E_2^{p,q} = H^p(U, R^q \pi_* \mathbb{G}_m) \Rightarrow H^{p+q}(\mathcal{X}, \mathbb{G}_m),$$

we get a long exact sequence

$$\begin{aligned} H^2(U, \mathbb{G}_m) &\longrightarrow \text{Ker}(H^2(\mathcal{X}, \mathbb{G}_m) \longrightarrow H^0(U, R^2 \pi_* \mathbb{G}_m)) \\ &\longrightarrow H^1(U, R^1 \pi_* \mathbb{G}_m) \longrightarrow H^3(U, \mathbb{G}_m) \end{aligned}$$

By [Gro3, Lem. 3.2.1], $H^i(U, \mathbb{G}_m)(\text{non-}p) = 0$ for $i \geq 2$. Thus, we get a canonical exact sequence

$$0 \rightarrow H^1(U, R^1 \pi_* \mathbb{G}_m)(\text{non-}p) \rightarrow H^2(\mathcal{X}, \mathbb{G}_m)(\text{non-}p) \rightarrow H^0(U, R^2 \pi_* \mathbb{G}_m)(\text{non-}p).$$

Let $j : \text{Spec } K \rightarrow U$ be the generic point. By Lemma 3.1, we have canonical isomorphisms

$$H^1(U, R^1 \pi_* \mathbb{G}_m)(\text{non-}p) \cong H^1(U, j_* j^* R^1 \pi_* \mathbb{G}_m)(\text{non-}p)$$

and

$$H^0(U, R^2 \pi_* \mathbb{G}_m)(\text{non-}p) \cong H^0(U, j_* j^* R^2 \pi_* \mathbb{G}_m)(\text{non-}p).$$

Since $j^* R^i \pi_* \mathbb{G}_m$ corresponds to the G_K -module $H^i(X_{K^s}, \mathbb{G}_m)$, we have

$$H^1(U, j_* j^* R^1 \pi_* \mathbb{G}_m) = \text{III}_U(\text{Pic}_{X/K}) \quad \text{and} \quad H^0(U, j_* j^* R^2 \pi_* \mathbb{G}_m) = \text{Br}(X_{K^s})^{G_K}.$$

This proves the first claim.

For the second claim, consider the exact sequence

$$0 \longrightarrow \text{Pic}^0(X_{K^s}) \longrightarrow \text{Pic}(X_{K^s}) \longrightarrow \text{NS}(X_{K^s}) \longrightarrow 0.$$

Since $\text{NS}(X_{K^s})$ is finitely generated, there exists a finite Galois extension L/K such that $\text{Pic}(X_L) \rightarrow \text{NS}(X_{K^s})$ is surjective. Taking Galois cohomology, we get a long exact sequence

$$\begin{aligned} 0 &\longrightarrow H^0(K, \text{Pic}^0(X_{K^s})) \longrightarrow H^0(K, \text{Pic}(X_{K^s})) \longrightarrow H^0(K, \text{NS}(X_{K^s})) \\ &\xrightarrow{a_K} H^1(K, \text{Pic}^0(X_{K^s})) \longrightarrow H^1(K, \text{Pic}(X_{K^s})) \longrightarrow H^1(K, \text{NS}(X_{K^s})). \end{aligned}$$

We have a similar long exact sequence for $H^i(L, -)$. Since $\text{Pic}(X_L) \rightarrow \text{NS}(X_{K^s})$ is surjective, we have $a_L = 0$ and

$$H^1(L, \text{NS}(X_{K^s})) = \text{Hom}(G_L, \text{NS}(X_{K^s})) = \text{Hom}(G_L, \text{NS}(X_{K^s})_{\text{tor}}).$$

$a_L = 0$ implies that the image of a_K is contained in

$$\text{Ker}(H^1(K, \text{Pic}^0(X_{K^s})) \longrightarrow H^1(L, \text{Pic}^0(X_{K^s})))$$

which is killed by $[L : K]$. Similarly, one can show that $H^1(K, \text{NS}(X_{K^s}))$ is killed by $[L : K]|\text{NS}(X_{K^s})_{\text{tor}}|$. Therefore, the kernel and cokernel of

$$H^1(K, \text{Pic}^0(X_{K^s})) \longrightarrow H^1(K, \text{Pic}(X_{K^s}))$$

are killed by $[L : K]|\text{NS}(X_{K^s})_{\text{tor}}|$. The claim also holds for

$$H^1(K_v^{sh}, \text{Pic}^0(X_{K^s})) \longrightarrow H^1(K_v^{sh}, \text{Pic}(X_{K^s})).$$

By diagram chasings, the kernel and cokernel of

$$\text{III}_U(\text{Pic}_{X/K, \text{red}}^0) \longrightarrow \text{III}_U(\text{Pic}_{X/K})$$

is of finite exponent. This completes the proof. \square

3.2 The pull-back trick

In this section, we will introduce a pull-back trick initially developed by Colliot-Thélène and Skorobogatov in [CTS] which allows us to reduce the question to cases of relative dimension 1.

Let U be a regular integral excellent scheme of dimension 1 with function field K . Let $\pi : \mathcal{X} \longrightarrow U$ be a smooth projective morphism with the generic fiber X geometrically connected over K . The Leray spectral sequence

$$E_2^{p,q} = H^p(U, R^q\pi_*\mathbb{G}_m) \Rightarrow H^{p+q}(\mathcal{X}, \mathbb{G}_m)$$

induces canonical maps

$$\begin{aligned} d_2^{1,1} : E_2^{1,1} &\longrightarrow E_2^{3,0}, \\ d_3^{0,2} : E_3^{0,2} &\longrightarrow E_3^{3,0}, \\ d_2^{0,2} : E_2^{0,2} &\longrightarrow E_2^{2,1}. \end{aligned}$$

Lemma 3.3. *Assuming that $X(K)$ is not empty, then the canonical maps $d_2^{1,1}$ and $d_3^{0,2}$ vanish and $E_3^{3,0} = E_2^{3,0}$. Moreover, there exists an open dense subscheme $V \subseteq U$ such that the canonical map $d_2^{0,2}$ has an image of finite exponent when replacing U by V . As a result, the natural map*

$$H^2(\pi^{-1}(V), \mathbb{G}_m) \longrightarrow H^0(V, R^2\pi_*\mathbb{G}_m)$$

has a cokernel of finite exponent.

Proof. Let $s \in X(K)$. Since π is proper, it extends to a section $s : U \longrightarrow \mathcal{X}$. Let $\tilde{E}_2^{p,q}$ denote the Leray spectral sequence for the identity map $U \longrightarrow U$. Then s induces a commutative diagram

$$\begin{array}{ccc} E_2^{1,1} & \xrightarrow{d_2^{1,1}} & E_2^{3,0} \\ \downarrow & & \downarrow \\ \tilde{E}_2^{1,1} & \longrightarrow & \tilde{E}_2^{3,0} \end{array}$$

The second column is an isomorphism since $E_2^{3,0} = H^3(U, \mathbb{G}_m) = \tilde{E}_2^{3,0}$. Since $\tilde{E}_2^{1,1} = 0$, thus $d_2^{1,1} = 0$. By definition, $d_2^{3,0} = 0$, it follows that $E_3^{3,0} = E_2^{3,0}$. By the same arguments, we have $d_3^{0,2} = 0$.

For the proof of the second claim, we will use the pullback method for finitely many U -morphisms $\mathcal{Y}_i \rightarrow \mathcal{X}$ where \mathcal{Y}_i is of relative dimension 1 over U . Let ${}^i E_2^{p,q}$ denote the Leray spectral sequence for $\pi_i : \mathcal{Y}_i \rightarrow U$. There is a commutative diagram

$$\begin{array}{ccc} E_2^{0,2} & \xrightarrow{d_2^{0,2}} & E_2^{2,1} \\ \downarrow & & \downarrow \\ \bigoplus_i {}^i E_2^{0,2} & \longrightarrow & \bigoplus_i {}^i E_2^{2,1} \end{array}$$

By shrinking U , we can assume that π_i is smooth and projective for all i . By Artin's theorem [Gro3, Cor. 3.2], $R^2 \pi_{i,*} \mathbb{G}_m = 0$. Thus, to show that $d_2^{0,2}$ has an image of finite exponent, it suffices to show that the second column has a kernel of finite exponent. The second column is the map

$$H^2(U, R^1 \pi_* \mathbb{G}_m) \longrightarrow \bigoplus_i H^2(U, R^1 \pi_{i,*} \mathbb{G}_m).$$

By Lemma 3.1(a), we have a canonical isomorphism

$$R^1 \pi_{i,*} \mathbb{G}_m \cong j_* j^* R^1 \pi_{i,*} \mathbb{G}_m,$$

where $j : \text{Spec } K \rightarrow U$ is the generic point. It suffices to show that there exists an étale sheaf \mathcal{F} on $\text{Spec } K$ and a morphism $\mathcal{F} \rightarrow \bigoplus_i j^* R^1 \pi_{i,*} \mathbb{G}_m$ such that the induced map

$$(1) \quad j^* R^1 \pi_* \mathbb{G}_m \oplus \mathcal{F} \longrightarrow \bigoplus_i j^* R^1 \pi_{i,*} \mathbb{G}_m$$

has a kernel and a cokernel killed by some positive integer. This will imply that the induced map

$$H^2(U, j_* j^* R^1 \pi_* \mathbb{G}_m) \oplus H^2(U, j_* \mathcal{F}) \longrightarrow \bigoplus_i H^2(U, j_* j^* R^1 \pi_{i,*} \mathbb{G}_m)$$

has a kernel and a cokernel of finite exponent. Since $j^* R^1 \pi_* \mathbb{G}_m$ corresponds to the G_K -module $\text{Pic}(X_{K^s})$, (1) can be interpreted as that there is a G_K -module M and a G_K -equivariant map $M \rightarrow \bigoplus_i \text{Pic}(Y_{i,K^s})$ such that

$$\text{Pic}(X_{K^s}) \oplus M \longrightarrow \bigoplus_i \text{Pic}(Y_{i,K^s})$$

has a kernel and a cokernel of finite exponent. In [Yua2, §3.3, p17], Yuan proved that there exist smooth projective curves Y_i in X which are obtained by taking intersection

of very ample divisors and an abelian subvariety A of $\prod_i \text{Pic}_{Y_i/K}^0$ such that the induced morphism

$$\text{Pic}(X_{K^s}) \times A(K^s) \longrightarrow \prod_i \text{Pic}(Y_{i,K^s})$$

has a finite kernel and a finite cokernel. Taking \mathcal{Y}_i to be the Zariski closure of Y_i in \mathcal{X} , then shrinking U , we get smooth and proper morphisms $\pi : \mathcal{Y}_i \longrightarrow U$. This proves the second claim.

For the last claim, consider the following canonical exact sequences induced by the Leray spectral sequence,

$$\begin{aligned} E^2 &\longrightarrow E_4^{0,2} \longrightarrow 0, \\ 0 &\longrightarrow E_4^{0,2} \longrightarrow E_3^{0,2} \xrightarrow{d_3^{0,2}} E_3^{3,0}, \\ 0 &\longrightarrow E_3^{0,2} \longrightarrow E_2^{0,2} \xrightarrow{d_2^{0,2}} E_2^{2,1}. \end{aligned}$$

Since $d_3^{0,2}$ vanishes and $d_2^{0,2}$ has an image of finite exponent, it follows that

$$E^2 \longrightarrow E_2^{0,2}$$

has a cokernel of finite exponent. This completes the proof. \square

3.3 Proof of Theorem 1.4

Now we prove Theorem 1.4. Let U be an open dense subscheme of C such that π is smooth and projective over U . By Lemma 3.2, the natural map

$$(\text{III}_{U_{K^s}}(\text{Pic}_{X/K,\text{red}}^0))^{G_k(\text{non-}p)} \rightarrow \text{Ker}(\text{Br}(\pi^{-1}(U_{K^s}))^{G_k(\text{non-}p)} \rightarrow \text{Br}(X_{K^s})^{G_K(\text{non-}p)})$$

has a kernel of finite exponent. By Lemma 2.9 and 2.10, all groups here are of cofinite type, thus the kernel is actually finite. By Proposition 2.3 and 2.7, it suffices to show that the natural map

$$\text{Br}(\pi^{-1}(U_{K^s}))^{G_k(\text{non-}p)} \longrightarrow \text{Br}(X_{K^s})^{G_K(\text{non-}p)}$$

has a cokernel of finite exponent.

Firstly, we will prove this under the assumption that $X(K)$ is not empty. We still use \mathcal{X} to denote $\pi^{-1}(U)$. By Proposition 2.3, without loss of generality, we can always replace U by an open dense subset. By Lemma 3.1(b), we have

$$H^0(U, R^2\pi_*\mathbb{G}_m)(\text{non-}p) \cong H^0(U, j_*j^*R^2\pi_*\mathbb{G}_m)(\text{non-}p) = \text{Br}(X_{K^s})^{G_K(\text{non-}p)},$$

so it suffices to show that the natural map induced by the Leray spectral sequence for π

$$H^2(\mathcal{X}, \mathbb{G}_m) \longrightarrow H^0(U, R^2\pi_*\mathbb{G}_m)$$

has a cokernel of finite exponent. By shrinking U , the claim follows from Lemma 3.3.

Secondly, we will show that the question can be reduced to the case that $X(K)$ is not empty. There exists a finite Galois extension L/K such that $X(L)$ is not empty. Let W be a smooth curve over k with function field L . By shrinking U and W , the map $\text{Spec } L \rightarrow \text{Spec } K$ extends to a finite etale Galois covering map $W \rightarrow U$. Let $\mathcal{X}_W \rightarrow W$ be the base change of $\mathcal{X} \rightarrow U$ to W . By the above arguments, the natural map

$$\text{Br}(\mathcal{X}_W)(\text{non-}p) \longrightarrow \text{Br}(X_{K^s})^{G_L}(\text{non-}p)$$

has a cokernel of finite exponent. Let G denote $\text{Gal}(L/K)$. The above map is compatible with the G -action. Taking G -invariant, we have that

$$\text{Br}(\mathcal{X}_W)^G(\text{non-}p) \longrightarrow \text{Br}(X_{K^s})^{G_K}(\text{non-}p)$$

has a cokernel of finite exponent. Then the question is reduced to show that the natural map

$$\text{Br}(\mathcal{X}) \longrightarrow \text{Br}(\mathcal{X}_W)^G$$

has cokernel of finite exponent. Consider the spectral sequence

$$H^p(G, H^q(\mathcal{X}_W, \mathbb{G}_m)) \Rightarrow H^{p+q}(\mathcal{X}, \mathbb{G}_m).$$

Since $H^p(G, -)$ is killed by the order of G , by similar arguments as in Lemma 3.3, we can conclude that the cokernel of

$$H^2(\mathcal{X}, \mathbb{G}_m) \longrightarrow H^2(\mathcal{X}_W, \mathbb{G}_m)^G$$

is of finite exponent. This completes the proof of the theorem.

4 Applications

4.1 Proof of Theorem 1.8

Firstly, note that in the case that k is finite, by Milne's theorems in [Mil5, Mil6], the finiteness of $\text{Br}(\mathcal{X}_{k^s})^{G_k}(\text{non-}p)$ is equivalent to Tate conjecture for divisors on \mathcal{X} .

We will prove that the finiteness of $\text{Br}(X_{K^s})^{G_K}(\text{non-}p)$ by induction on the transcendence degree of K/k . If K/k is of transcendence degree 0, a smooth projective variety over K can be view as a smooth projective variety over k . So the claim is true by assumptions.

Assume that the claim is true for all extensions l/k of transcendence degree n . Let K/k be a finitely generated extension of transcendence degree $n + 1$. Let X/K be a smooth projective connected variety over K , by extending K , we might assume that X/K is geometrically connected. K can be regarded as the function field of a smooth projective geometrically connected curve C over a field l of transcendence degree n over k . $X \rightarrow \text{Spec } K$ can be extended to a smooth morphism $\pi : \mathcal{X} \rightarrow C$. By Theorem 1.4,

$$\text{Br}(\mathcal{X}_{l^s})^{G_k}(\text{non-}p) \longrightarrow \text{Br}(\mathcal{X}_{K^s})^{G_K}(\text{non-}p)$$

has a finite cokernel. It suffices to show that $\mathrm{Br}(\mathcal{X}_{l^s})^{G_l}(\mathrm{non-}p)$ is finite. By Proposition 2.3, we might shrink \mathcal{X} such that there is an alteration $f : \mathcal{X}' \rightarrow \mathcal{X}$ such that \mathcal{X}' admits a smooth projective compactification over a finite extension of l . Replacing l by a finite extension will not change the question, so we might assume that \mathcal{X}' admits a smooth projective compactification over l . By shrinking \mathcal{X} , we might assume that f is finite flat. Then the kernel of

$$\mathrm{Br}(\mathcal{X}_{l^s})(\mathrm{non-}p) \longrightarrow \mathrm{Br}(\mathcal{X}'_{l^s})(\mathrm{non-}p)$$

is killed by the degree of f (cf. Lemma 2.6) and therefore is finite. Since $\mathrm{Br}(\mathcal{X}'_{l^s})^{G_l}(\mathrm{non-}p)$ is finite by induction, so $\mathrm{Br}(\mathcal{X}_{l^s})^{G_l}(\mathrm{non-}p)$ is finite. This completes the proof.

4.2 Proof of Theorem 1.2

Let $\ell \neq p$ be a prime, the the claim in Theorem 1.8 also holds when replacing the prime to p part by the ℓ primary part in the statement. So by the same argument as in the proof of Theorem 1.8, Theorem 1.2 is true.

4.3 Brauer groups of integral models of abelian varieties and K3 surfaces

Proposition 4.1. *Let C be a smooth projective geometrically connected curve over a finite field k of characteristic $p > 2$. Let $\pi : \mathcal{X} \rightarrow C$ be a projective flat morphism. Let X denote the generic fiber of π . Assuming that \mathcal{X} is regular and X_{K^s} is an abelian variety over K^s , then there is an isomorphism up to finite groups*

$$\mathrm{III}(\mathrm{Pic}_{X/K, \mathrm{red}}^0)(\mathrm{non-}p) \cong \mathrm{Br}(\mathcal{X})(\mathrm{non-}p).$$

Proof. By [SZ1, Thm. 1.1], $\mathrm{Br}(X_{K^s})^{G_K}(\mathrm{non-}p)$ is finite, the claim follows directly from Corollary 1.5. \square

Proposition 4.2. *Let C be a smooth projective geometrically connected curve over a finite field k of characteristic p . Let $\pi : \mathcal{X} \rightarrow C$ be a projective flat morphism. Assuming that \mathcal{X} is regular and the generic fiber X of π is a smooth projective geometrically connected surface over K with $H^1(X_{K^s}, \mathbb{Q}_\ell) = 0$ for some prime $\ell \neq p$, then the natural map*

$$\mathrm{Br}(\mathcal{X})(\mathrm{non-}p) \longrightarrow \mathrm{Br}(X_{K^s})^{G_K}(\mathrm{non-}p)$$

has a finite kernel and cokernel. As a result, $\mathrm{Br}(X_{K^s})^{G_K}(\mathrm{non-}p)$ is finite if and only if $\mathrm{Br}(X_{K^s})^{G_K}(\ell)$ is finite for some prime $\ell \neq p$.

Proof. $\mathrm{Pic}_{X/K, \mathrm{red}}^0 = 0$ since $H^1(X_{K^s}, \mathbb{Q}_\ell) = 0$. It follows that

$$\mathrm{III}(\mathrm{Pic}_{X/K, \mathrm{red}}^0) = 0.$$

Then the claim follows from Corollary 1.5. \square

Corollary 4.3. *Notations as above, if X/K is a K3 surface and $p > 2$, then $\mathrm{Br}(\mathcal{X})$ is finite.*

Proof. Since Tate conjecture for X is known by work of lots of people (cf. Conjecture 1.1), so $\mathrm{Br}(X_{K^s})^{G_K}(\ell)$ is finite for some prime $\ell \neq p$. By the above proposition, $\mathrm{Br}(\mathcal{X})(\ell)$ is also finite. Then by Milne's theorems [Mil5, Mil6], $\mathrm{Br}(\mathcal{X})$ is finite. \square

Proposition 4.4. *Let X be a K3 surface over a global field K of characteristic $p > 2$. Then $\mathrm{Br}(X_{K^s})^{G_K(\mathrm{non-}p)}$ is finite.*

Proof. By previous results, it suffices to show that X admits a projective regular model. By resolution of singularity of threefolds(cf. [CP]), X does admit a projective regular model. \square

References

- [Amb] Ambrosi, Emiliano. A note on the behaviour of the Tate conjecture under finitely generated field extensions. *Pure and Applied Mathematics Quarterly* 14.3 (2018): 515-527.
- [And] Y. André, On the Shafarevich and Tate conjectures for hyper-Kähler varieties, *Math. Ann.* 305 (1996), no. 2, 205-248.
- [ASD] M. Artin, H. P. F. Swinnerton-Dyer, The Shafarevich-Tate conjecture for pencils of elliptic curves on K3 surfaces, *Invent. Math.* 20 (1973), 249-266.
- [Čes] Česnavičius, K., 2019. Purity for the Brauer group. *Duke Mathematical Journal*, 168(8), pp.1461-1486.
- [Cha] F. Charles, The Tate conjecture for K3 surfaces over finite fields, *Invent. Math.* 194, 119-145 (2013).
- [CP] Cossart, V., Piltant, O. (2009). Resolution of singularities of threefolds in positive characteristic II. *Journal of Algebra*, 321(7), 1836-1976.
- [CTS] Colliot-Thélène, J.-L.; Skorobogatov, A.N. Descente galoisienne sur le groupe de Brauer. *J. reine angew. Math.* 682 (2013) 141-165.
- [Del] Deligne, P. “SGA 4 1/2.” *Lect. Notes Math* 569 (1977).
- [deJ1] de Jong, A., 1997. Families of curves and alterations. In *Annals of the Fourier Institute* (Vol. 47, No. 2, pp. 599-621).
- [deJ2] de Jong, A. J.: Tate conjecture for divisors. Unpublished note.
- [EGA IV] GROTHENDIECK, A. Éléments de géométrie algébrique : IV. Étude locale des schémas et des morphismes de schémas, Troisième partie Publications mathématiques de l’I.H.É.S., tome 28 (1966), p. 5-255.
- [Fal1] G. Faltings, Endlichkeitssätze für abelsche Varietäten über Zahlkörpern. *Invent. Math.* 73 (1983), no. 3, 349-366.
- [Fal2] G. Faltings, Complements to Mordell. In: G. Faltings, G. Wüstholz et al., *Rational points*, 2nd edition, F. Vieweg & Sohn, 1986.
- [Fuj] Fujiwara, K. (2002). A proof of the absolute purity conjecture (after Gabber). In *Algebraic geometry 2000, Azumino* (pp. 153-183). Mathematical Society of Japan.
- [Fu] Fu, L., 2011. *Etale cohomology theory* (Vol. 13). World Scientific.
- [Gei1] Geisser, T. H. (2018). Comparing the Brauer group to the Tate–Shafarevich group. *Journal of the Institute of Mathematics of Jussieu*, 1-6.

- [Gei2] Geisser, Thomas H. “TATE’S CONJECTURE AND THE TATE–SHAFAREVICH GROUP OVER GLOBAL FUNCTION FIELDS.” *Journal of the Institute of Mathematics of Jussieu* (2018): 1-22.
- [Goa] Gonzalez-Aviles C D. Brauer Groups and Tate-Shafarevich Groups. *J. Math. Sci. Univ. Tokyo*, 2003, 10: 391-419.
- [Gro1] A. Grothendieck, Le groupe de Brauer. I. Algèbres d’Azumaya et interprétations diverses, *Dix Exposés sur la Cohomologie des Schémas*, North-Holland, Amsterdam, 1968, pp. 46–66.
- [Gro2] A. Grothendieck, Le groupe de Brauer. II. Théorie cohomologique, *Dix Exposés sur la Cohomologie des Schémas*, North-Holland, Amsterdam, 1968, pp. 67–87.
- [Gro3] A. Grothendieck, Le groupe de Brauer. III. Exemples et compléments, *Dix Exposés sur la Cohomologie des Schémas*, North-Holland, Amsterdam, 1968, pp. 88–188.
- [Har] Hartshorne, Robin. *Algebraic geometry*. Vol. 52. Springer Science & Business Media, 2013.
- [Mau] D. Maulik, Supersingular K3 surfaces for large primes, With an appendix by Andrew Snowden, *Duke Math. J.* 163, 2357-2425 (2014).
- [Mil1] Milne, James S, *Arithmetic duality theorems*, Vol. 20. Charleston, SC: BookSurge, 2006.
- [Mil2] Milne, James S, *Abelian Varieties*, Chapter V of *Arithmetic geometry* (Storrs, Conn., 1984), 103–150, Springer, New York, 1986.
- [Mil3] Milne, J.S., 1980. *Etale cohomology* (PMS-33) (Vol. 5657). Princeton university press.
- [Mil4] Milne, J. S., Comparison of the Brauer Group with the Tate-Safarevic group, *J. Fac. Science Univ. Tokyo, Sec. IA* 28 (1981), 735–743.
- [Mil5] Milne, J. S. (1975). On a conjecture of Artin and Tate. *Annals of Mathematics*, 517-533.
- [Mil6] Milne, J. S. (1986). Values of zeta functions of varieties over finite fields. *American Journal of Mathematics*, 108(2), 297-360.
- [MP] K. Madapusi Pera, The Tate conjecture for K3 surfaces in odd characteristic, *Invent. Math.* 201, 625-668 (2015).
- [Mor] Morrow, M. (2019). A Variational Tate Conjecture in crystalline cohomology. *Journal of the European Mathematical Society*, 21(11), 3467-3511.

- [NO] N. Nygaard, A. Ogus, Tate's conjecture for K3 surfaces of finite height, *Ann. of Math.* 122, 461-507 (1985).
- [Nyg] N. O. Nygaard, The Tate conjecture for ordinary K3 surfaces over finite fields, *Invent. Math.* 74, 213-237 (1983).
- [Sch] Schneider, P. (1982). On the conjecture of Birch and Swinnerton-Dyer about global function fields. *Mathematische Annalen* , 260 (4), 495-510.
- [SZ1] Skorobogatov, A. N., & Zarkhin, Y. G. (2008). A finiteness theorem for the Brauer group of Abelian varieties and KS surfaces. *Journal of Algebraic Geometry*, 17(3), 481-502.
- [SZ2] Skorobogatov, Alexei N., and Yuri G. Zarhin. A finiteness theorem for the Brauer group of K3 surfaces in odd characteristic. *International Mathematics Research Notices* 2015.21 (2015): 11404-11418.
- [Tan1] Tankeev, S. G. (2001). On the Brauer group of an arithmetic scheme. *Izvestiya: Mathematics*, 65(2), 357.
- [Tan2] Tankeev, S. G. (2003). On the Brauer group of an arithmetic scheme. II. *Izvestiya: Mathematics*, 67(5), 1007.
- [Tan3] Tankeev, S. G. (2014). On the finiteness of the Brauer group of an arithmetic scheme. *Mathematical Notes*, 95(1-2), 121-132.
- [Tan4] S. G. Tankeev, Surfaces of K3 type over number fields and the Mumford-Tate conjecture. *Izv. Akad. Nauk SSSR Ser. Mat.* 54 (1990), no. 4, 846-861; translation in *Math. USSR-Izv.* 37 (1991), no. 1, 191- 208.
- [Tan5] S. G. Tankeev, Surfaces of K3 type over number fields and the Mumford-Tate conjecture. II. *Izv. Ross. Akad. Nauk Ser. Mat.* 59 (1995), no. 3, 179-206; translation in *Izv. Math.* 59 (1995), no. 3, 619-646.
- [Tat1] Tate J. Algebraic cycles and poles of zeta functions. *Arithmetical algebraic geometry*. 1965:93-110.
- [Tat2] Tate, John. Conjectures on algebraic cycles in l -adic cohomology. *Motives* (Seattle, WA, 1991) 55 (1994): 71-83.
- [Tat3] J. Tate, Endomorphisms of abelian varieties over finite fields, *Invent. Math.* 2, 134-144 (1966).
- [Ulm] D.Ulmer, Curves and Jacobians over function fields, *Arithmetic geometry over global function fields*, 2014, pp.281-337.
- [Yua1] Yuan, X. (2018). Positivity of Hodge bundles of abelian varieties over some function fields. arXiv preprint arXiv:1808.03960.

- [Yua2] Yuan, Xinyi. Comparison of arithmetic Brauer groups with geometric Brauer group. arXiv preprint arXiv:2011.12897v2
- [Zar1] Yu.G. Zarhin, Endomorphisms of Abelian varieties over fields of finite characteristic. *Izv. Akad. Nauk SSSR Ser. Mat.* 39 (1975), 272-277; *Math. USSR Izv.* 9 (1975), 255-260.
- [Zar2] Yu.G. Zarhin, Abelian varieties in characteristic p . *Mat. Zametki* 19 (1976), 393-400; *Math. Notes* 19 (1976), 240-244.