

# BESOV-HANKEL NORMS IN TERMS OF THE CONTINUOUS BESSEL WAVELET TRANSFORM

ASHISH PATHAK AND DILEEP KUMAR  
 DEPARTMENT OF MATHEMATICS  
 INSTITUTE OF SCIENCE  
 BANARAS HINDU UNIVERSITY  
 VARANASI-221005, INDIA.

ABSTRACT. In this paper, we extend the concept of continuous Bessel wavelet transform in  $L^p$ -space and derived the Parseval's as well as the inversion formulas. By using Bessel wavelet coefficients we characterized the Besov- Hankel space.

## 1. INTRODUCTION

In this paper, as usual  $L^{p,\sigma}(\mathbb{R}^+ = (0, \infty))$  denotes the weighted  $L^p$ - space with norm

$$\|f\|_{L^{p,\sigma}} = \|f\|_{p,\sigma} = \left( \int_0^\infty |f(x)|^p d\sigma(x) \right)^{\frac{1}{p}}, \quad (1 \leq p < \infty), \quad (1.1)$$

$$\|f\|_{\infty,\sigma} = \text{ess sup}_{0 < x < \infty} |f(x)| < \infty. \quad (1.2)$$

The Hankel transformation of the function  $f \in L^{1,\sigma}(\mathbb{R}^+)$  is defined by

$$\hat{f}(x) = \int_0^\infty j(xt)f(t)d\sigma(t), \quad 0 \leq x < \infty, \quad (1.3)$$

where  $\sigma(t) = \frac{t^{2\nu+1}}{2^{\nu+\frac{1}{2}}\Gamma(\nu+\frac{3}{2})}$ ,  $j(\cdot) = C_\nu x^{\frac{1}{2}-\nu} J_{\nu-\frac{1}{2}}$ ,  $C_\nu = 2^{\nu+\frac{1}{2}}\Gamma(\nu + \frac{1}{2})$  and  $J_{\nu-\frac{1}{2}}$  denote the Bessel function of first kind of order  $\nu - \frac{1}{2}$ .

---

2010 *Mathematics Subject Classification.* 33A40; 42C10;44A05;42C40.

*Key words and phrases.* Besov-Hankel space;Continuous Bessel wavelet transform; Hankel transform; Hankel convolution.

<sup>1</sup> E-mail: ashishpathak@bhu.ac.in, <sup>2</sup> E-mail: dkbhu07@gmail.com .

If  $\hat{f} \in L^{1,\sigma}(\mathbb{R}^+)$ , then the inverse of Hankel transformations is given by

$$f(x) = \int_0^{\infty} j(xt)\hat{f}(t)d\sigma(t), \quad 0 < x < \infty. \quad (1.4)$$

Also, Parseval's formula of the Hankel transformation for  $f, g \in L^{1,\sigma} \cap L^{2,\sigma}$  is given by

$$\int_0^{\infty} \hat{f}(x)\hat{g}(x)d\sigma(x) = \int_0^{\infty} f(u)g(u)d\sigma(u). \quad (1.5)$$

By denseness and continuity the Parseval's formula can be extended to all  $f, g \in L^{2,\sigma}(\mathbb{R}^+)$ . Hence Hankel transform is isometry on  $L^{2,\sigma}(\mathbb{R}^+)$ .

If  $f, g \in L^{1,\sigma}(\mathbb{R}^+)$ , then the convolution associated with the Hankel is defined as (see [10])

$$(f\#g)(x) = \int_0^{\infty} f(x,y)g(y)d\sigma(y), \quad (1.6)$$

where the Hankel translation is given by

$$f(x,y) = \tau_y f(x) =: \int_0^{\infty} f(z)D(x,y,z)d\sigma(z), \quad 0 < x, y < \infty, \quad (1.7)$$

and

$$\begin{aligned} D(x,y,z) &= \int_0^{\infty} j(xu)j(yu)j(zu)d\sigma(u) \\ &= 2^{3\nu-\frac{5}{2}}[\Gamma(\nu+\frac{1}{2})]^2 \left(\Gamma(\nu)\pi^{\frac{1}{2}}\right)^{-1} (xyz)^{-2\nu-1} [\Delta(xyz)]^{2\nu-2} \end{aligned} \quad (1.8)$$

where  $\Delta$  denotes the area of a triangle.  $D(x,y,z)$  is symmetric in  $x, y, z$ .

From (1.4) and (1.8), we have

$$\int_0^{\infty} j(zu)D(x,y,z)d\sigma(z) = j(xu)j(yu), \quad 0 < x, y < \infty, \quad 0 \leq u < \infty, \quad (1.9)$$

and for  $u = 0$ , we get

$$\int_0^{\infty} D(x, y, z) d\sigma(z) = 1, \quad (1.10)$$

and

$$(f \# g)(x) = \hat{f}(x) \hat{g}(x), \quad 0 \leq x < \infty. \quad (1.11)$$

Now, we recall some properties of Hankel convolution ( see [7],[2],[9], [6] ) which are useful throught the paper.

**Lemma 1.1.** *Let  $f \in L^{p,\sigma}(\mathbb{R}^+)$ ,  $1 \leq p < \infty$ . Then we have*

$$\|\tau_y f(x)\|_{p,\sigma} \leq \|f\|_{p,\sigma}. \quad (1.12)$$

**Lemma 1.2.** *Let  $f \in L^{p,\sigma}(\mathbb{R}^+)$  and  $g \in L^{q,\sigma}(\mathbb{R}^+)$ ,  $\frac{1}{p} + \frac{1}{q} = 1 + \frac{1}{r}$ . Then we have*

$$\|f \# g\|_{r,\sigma} \leq \|f\|_{p,\sigma} \|g\|_{q,\sigma}. \quad (1.13)$$

Betancor and Rodriguez-Mesa defined the new function spaces called Besov-Hankel by using Hankel transform and its properties (see [9])

**Definition 1.3. (Besov-Hankel Space):** Let measurable function  $\phi$  on  $(0, \infty)$  belongs to  $BH_{\alpha,\sigma}^{p,q}$  if  $\phi \in L^{p,\sigma}(\mathbb{R}^+)$  and

$$\int_0^{\infty} (h^{-\alpha} w_p(\phi)(h))^q \frac{dh}{h} < \infty \quad \text{for } \alpha > 0, 1 \leq p, q < \infty, \quad (1.14)$$

where  $w_p(\phi)(h) =: \|\tau_h \phi - \phi\|_{p,\sigma}$ ,  $h \in (0, \infty)$ .

**1.1. Bessel Wavelet.** Using the properites of Hankel transform Pathak and Dixit (see [2]) define the continuous Bessel wavelet for  $\psi \in L^{p,\sigma}(\mathbb{R}^+)$ ,  $1 \leq p < \infty$ ,  $b \geq 0$  and  $a > 0$  as

$$\begin{aligned} \psi_{b,a}(x) &:= D_a \tau_b \psi(x) \\ &= a^{-2\nu-1} \int_0^{\infty} \psi(z) D\left(\frac{b}{a}, \frac{x}{a}, z\right) d\sigma(z) \end{aligned} \quad (1.15)$$

where  $D_a$  denote the dilation operator.

The continuous Bessel wavelet transform of  $f \in L^{2,\sigma}(\mathbb{R}^+)$  with respect to a wavelet  $\psi \in L^{2,\sigma}(\mathbb{R}^+)$  is defined as

$$\begin{aligned} (B_\psi f)(b, a) &= \int_0^\infty f(x) \overline{\psi_{b,a}(x)} d\sigma(x) \\ &= a^{-2\nu-1} \int_0^\infty \int_0^\infty f(x) \overline{\psi(z)} D\left(\frac{b}{a}, \frac{x}{a}, z\right) d\sigma(z) d\sigma(x). \end{aligned} \quad (1.16)$$

Moreover, using (1.6), we have

$$(B_\psi f)(b, a) = (f \# \psi_a)(b), \quad (1.17)$$

where  $\psi_a(t) = a^{-2\nu-1} \overline{\psi(t/a)}$ .

Betancor and Rodriguez-Mesa [9] first time introduce Besov-Hankel spaces and characterized by mean of the Bochner-Riesz mean and the partial Hankel integrals. Perrier and Basdevant [5] established the characterization of Besov spaces by means of continuous wavelet transform. Motivated by these two works, we characterized Besov-Hankel spaces by the continuous Bessel wavelet transform.

Present paper is organized in following manner: section 1 is introductory, in which we recall some properties of Hankel transform, Besov-Hankel space and continuous Bessel wavelet transform. Section 2 is related to continuous the continuous Besse wavelet transform in  $L^{p,\sigma}(\mathbb{R}^+)$ . In the section 3, characterize Besov-Hankel norms in terms of continuous Bessel wavelet transform.

## 2. THE CONTINUOUS BESSEL WAVELET TRANSFORM IN $L^{p,\sigma}(\mathbb{R}^+)$

In this section we extend the concept of Bessel wavelet transform on  $L^{p,\sigma}(\mathbb{R}^+)$ .

**Theorem 2.1.** *Suppose that the Bessel wavelet  $\psi \in L^{p,\sigma}(\mathbb{R}^+)$  satisfies the admissibility condition*

$$A_\psi = \int_0^\infty \omega^{-2\nu-1} |\hat{\psi}(\omega)|^2 d\omega > 0,$$

where  $\hat{\psi}$  denote the Hankel transform of  $\psi$  then continuous Bessel wavelet transform is a bounded linear operator

$$L^{p,\sigma}(\mathbb{R}^+) \rightarrow L^{2,\sigma}(\mathbb{R}^+, \frac{d\sigma(a)}{a^{2\nu+1}}) \times L^{p,\sigma}(\mathbb{R}^+),$$

moreover, for any  $f \in L^{p,\sigma}(\mathbb{R}^+)$ ,  $1 < p < \infty$

$$\|f\|_{L^{p,\sigma}(\mathbb{R}^+)} = \left( \int_0^\infty \left( \int_0^\infty |B_\psi f(b, a)|^2 \frac{d\sigma(a)}{a^{2\nu+1}} \right)^{\frac{p}{2}} d\sigma(b) \right)^{\frac{1}{p}}. \quad (2.1)$$

*Proof.* Let  $S_p$  denote the space  $L^{2,\sigma}(\mathbb{R}^+, \frac{d\sigma(a)}{a^{2\nu+1}}) \times L^{p,\sigma}(\mathbb{R}^+)$  associated to the norm

$$\|f\|_{S_p} = \left\{ \int_0^\infty \left( \int_0^\infty |f(b, a)|^2 \frac{d\sigma(a)}{a^{2\nu+1}} \right)^{\frac{p}{2}} d\sigma(b) \right\}^{\frac{1}{p}}.$$

If we take  $p = 2$ , then from Plancherel's theorem:

$$\begin{aligned} \|B_\psi f\|_{S_2} &= \left\{ \int_0^\infty \left( \int_0^\infty |B_\psi f(b, a)|^2 \frac{d\sigma(a)}{a^{2\nu+1}} \right) d\sigma(b) \right\}^{\frac{1}{2}} \\ \|B_\psi f\|_{S_2} &= \sqrt{A_\psi} \|f\|_{L^{2,\sigma}}, \end{aligned}$$

where  $A_\psi = \int_0^\infty \omega^{-2\nu-1} |\hat{\psi}(\omega)|^2 d\omega > 0$ , if  $\psi$  is real. From singular integral theorem, the operators on  $L^{2,\sigma}(\mathbb{R}^+, \frac{d\sigma(a)}{a^{2\nu+1}})$  holds inequality:

$$\|B_\psi f\|_{S_p} \leq C_p \|f\|_{L^{p,\sigma}(\mathbb{R}^+)} \text{ for } 1 < p \leq 2,$$

where the constant  $C_p$  depends only on  $p$  and  $\psi$  (see [12]). Due to duality the inequality is also valid for  $1 < p < \infty$ . It follows that

$$\left\{ \int_0^\infty \left( \int_0^\infty |B_\psi f(b, a)|^2 \frac{d\sigma(a)}{a^{2\nu+1}} \right)^{\frac{p}{2}} d\sigma(b) \right\}^{\frac{1}{p}} \leq C_p \|f\|_{L^{p,\sigma}(\mathbb{R}^+)}. \quad (2.2)$$

Conversely suppose that  $f \in L^{2,\sigma}(\mathbb{R}^+) \cap L^{p,\sigma}(\mathbb{R}^+)$ . Since continuous Bessel wavelet transform is isometry for every  $g \in L^{2,\sigma}(\mathbb{R}^+) \cap L^{q,\sigma}(\mathbb{R}^+)$ , we can write

$$\begin{aligned} \int_0^\infty \int_0^\infty B_\psi f(b, a) \overline{B_\psi g(b, a)} a^{-2\nu-1} d\sigma(a) d\sigma(b) &= A_\psi \langle f, g \rangle \\ \frac{1}{A_\psi} \int_0^\infty \int_0^\infty B_\psi f(b, a) \overline{B_\psi g(b, a)} a^{-2\nu-1} d\sigma(a) d\sigma(b) &= \int_0^\infty f(x) \overline{g(x)} d\sigma(x). \end{aligned} \quad (2.3)$$

Now,

$$\begin{aligned} \left| \int_0^\infty f(x)g(x)d\sigma(x) \right| &= \frac{1}{A_\psi} \left| \int_0^\infty \int_0^\infty B_\psi f(b,a) \overline{B_\psi g(b,a)} a^{-2\nu-1} d\sigma(a) d\sigma(b) \right| \\ &\leq \frac{1}{A_\psi} \int_0^\infty \int_0^\infty |B_\psi f(b,a) \overline{B_\psi g(b,a)}| a^{-2\nu-1} d\sigma(a) d\sigma(b), \end{aligned}$$

using Schwarz inequality and then Holder's inequality, we have

$$\begin{aligned} &\leq \frac{1}{A_\psi} \left( \int_0^\infty \left( \int_0^\infty |B_\psi f(b,a)|^2 a^{-2\nu-1} d\sigma(a) \right)^{\frac{p}{2}} d\sigma(b) \right)^{\frac{1}{p}} \\ &\quad \times \left( \int_0^\infty \left( \int_0^\infty |B_\psi g(b,a)|^2 a^{-2\nu-1} d\sigma(a) \right)^{\frac{q}{2}} d\sigma(b) \right)^{\frac{1}{q}}, \end{aligned}$$

where  $\frac{1}{p} + \frac{1}{q} = 1$ .

From equation (2.2), we get

$$\leq \frac{A_q}{A_\psi} \left( \int_0^\infty \left( \int_0^\infty |B_\psi f(b,a)|^2 a^{-2\nu-1} d\sigma(a) \right)^{\frac{p}{2}} d\sigma(b) \right)^{\frac{1}{p}} \|g\|_{L^{q,\sigma}(\mathbb{R}^+)}.$$

By Density theorem

$$\|f\|_{L^{p,\sigma}(\mathbb{R}^+)} \leq A \left( \int_0^\infty \left( \int_0^\infty |B_\psi f(b,a)|^2 a^{-2\nu-1} d\sigma(a) \right)^{\frac{p}{2}} d\sigma(b) \right)^{\frac{1}{p}},$$

where  $A = \frac{A_q}{A_\psi}$ . □

**Theorem 2.2. (Parseval's formula)** *Let us assume  $\phi_1 \in L^{p,\sigma}(\mathbb{R}^+)$ ,  $\phi_2 \in L^{q,\sigma}(\mathbb{R}^+)$  with  $1 \leq p, q < \infty$  and  $\frac{1}{p} + \frac{1}{q} = 1$ . If  $\psi$  is a real wavelet then*

$$\frac{1}{A_\psi} \int_0^\infty \int_0^\infty B_\psi \phi_1(b,a) \overline{B_\psi \phi_2(b,a)} a^{-2\nu-1} d\sigma(a) d\sigma(b) = \int_0^\infty \phi_1(x) \overline{\phi_2(x)} d\sigma(x), \quad (2.4)$$

where  $A_\psi = \int_0^\infty \omega^{-2\nu-1} |\hat{\psi}(\omega)|^2 d\omega > 0$  and  $\hat{\psi}$  denote the Hankel transform.

*Proof.* Let us define bilinear transform  $T : L^{p,\sigma}(\mathbb{R}^+) \times L^{q,\sigma}(\mathbb{R}^+) \rightarrow \mathbb{R}^+$  by

$$T(\phi_1, \phi_2) = \langle B_\psi \phi_1(b,a), B_\psi \phi_2(b,a) \rangle_{\left(\frac{d\sigma(a)}{a^{2\nu+1}} d\sigma(b)\right)}.$$

Now, applying Holder's inequality two times we obtain

$$\begin{aligned}
 |T(\phi_1, \phi_2)| &= |\langle B_\psi \phi_1(b, a), B_\psi \phi_2(b, a) \rangle_{\frac{d\sigma(a)}{a^{2\nu+1}}, d\sigma(b)}| \\
 &\leq \int_0^\infty \left( \int_0^\infty |B_\psi \phi_1(b, a)|^2 \frac{d\sigma(a)}{a^{2\nu+1}} \right)^{\frac{1}{2}} \left( \int_0^\infty |B_\psi \phi_2(b, a)|^2 d\sigma(b) \right)^{\frac{1}{2}} \\
 &\leq \left( \int_0^\infty \left( \int_0^\infty |B_\psi \phi_1(b, a)|^2 \frac{d\sigma(a)}{a^{2\nu+1}} \right)^{\frac{p}{2}} \right)^{\frac{1}{p}} \times \left( \int_0^\infty \left( \int_0^\infty |B_\psi \phi_2(b, a)|^2 d\sigma(b) \right)^{\frac{q}{2}} \right)^{\frac{1}{q}}
 \end{aligned}$$

using Theorem 2.1. we have

$$|T(\phi_1, \phi_2)| \leq C \|\phi_1\|_{L^{p,\sigma}(\mathbb{R}^+)} \|\phi_2\|_{L^{q,\sigma}(\mathbb{R}^+)}. \quad (2.5)$$

Moreover for all  $\phi_1 \in L^{2,\sigma}(\mathbb{R}^+) \cap L^{p,\sigma}(\mathbb{R}^+)$  and  $\phi_2 \in L^{2,\sigma}(\mathbb{R}^+) \cap L^{q,\sigma}(\mathbb{R}^+)$  we get

$$T(\phi_1, \phi_2) = \langle B_\psi \phi_1(b, a), B_\psi \phi_2(b, a) \rangle_{\frac{d\sigma(a)}{a^{2\nu+1}}, d\sigma(b)} = A_\psi \langle \phi_1, \phi_2 \rangle. \quad (2.6)$$

From equations (3.5), (3.6) and density of spaces  $L^{2,\sigma}(\mathbb{R}^+) \cap L^{p,\sigma}(\mathbb{R}^+)$  in  $L^{p,\sigma}(\mathbb{R}^+)$  gives the result.  $\square$

### 2.1. An inversion formula.

**Theorem 2.3.** *Let us consider  $\phi \in L^{p,\sigma}(\mathbb{R}^+)$  with  $1 < p < \infty$  and  $\psi$  is a real wavelet. Then*

$$\phi(x) = \frac{1}{A_\psi} \int_0^\infty \int_0^\infty B_\psi \phi(b, a) \psi_{b,a}(x) \frac{d\sigma(a)}{a^{2\nu+1}} d\sigma(b). \quad (2.7)$$

*The equality holds in  $L^{p,\sigma}(\mathbb{R}^+)$  sense and the integral of right hand side have to be taken in the sense of distributions.*

*Proof.* The proof followed from Theorem 2.2.  $\square$

## 3. CHARACTERIZATION OF BESOV-HANKEL NORMS

In present section, By using the above results, we characterize the Besov-Hankel norms associated the Bessel wavelet transform.

**Theorem 3.1.** *Let  $f \in B_{\alpha,\sigma}^{p,q}(\mathbb{R}^+)$  ( $p, q > 1, \alpha \notin \mathbb{Z}$ ) and analysing wavelet  $\psi$  has  $[\alpha]+1$  cancellations and  $(z^{\alpha-[\alpha]}\psi) \in L^{1,\sigma}(\mathbb{R}^+)$ , then the wavelet coefficient of function  $f$  holds following conditions:*

$$\begin{aligned} \text{if } q < \infty, \quad & \int_0^\infty [a^{-\alpha} \|B_\psi(\cdot, a)\|_{L^{p,\sigma}}]^q \frac{da}{a} < \infty \\ \text{if } q = \infty, \quad & a \rightarrow a^{-\alpha} \|B_\psi(\cdot, a)\|_{L^{p,\sigma}} \in L^\infty(\mathbb{R}^+). \end{aligned}$$

Moreover the function  $a \rightarrow a^{-\alpha} \|B_\psi(\cdot, a)\|_{L^{p,\sigma}} \in L^q(\mathbb{R}^+, \frac{da}{a^{2\nu+1}})$  and we have:

$$\|a^{-\alpha} \|B_\psi(\cdot, a)\|_{L^{p,\sigma}}\|_{L^q, \sigma(\frac{da}{a})} \leq \|z^{\alpha-[\alpha]}\psi\|^{L_{1,\sigma}} \times \|h^{\alpha-[\alpha]}\omega_p(f, h)\|_{L^q, \sigma(\frac{dh}{h})} \quad (3.1)$$

*Proof.* By the definition of continuous Bessel wavelet transform, we have

$$\begin{aligned} B_\psi(b, a) &= \int_0^\infty f(x) \overline{\psi_{b,a}(x)} d\sigma(x) \\ &= \int_0^\infty f(x) \left( \int_0^\infty a^{-2\nu-1} D\left(\frac{b}{a}, \frac{x}{a}, z\right) \overline{\psi(z)} d\sigma(z) \right) d\sigma(x) \\ &= \int_0^\infty \overline{\psi(z)} \left( \int_0^\infty a^{-2\nu-1} D\left(\frac{b}{a}, \frac{x}{a}, z\right) f(x) d\sigma(x) \right) d\sigma(z) \\ &= \int_0^\infty \overline{\psi(z)} \left( \int_0^\infty D(b, x, az) f(x) d\sigma(x) \right) d\sigma(z) \\ &= \left\{ \int_0^\infty (\tau_{az}f)(b) \overline{\psi(z)} d\sigma(z) - \int_0^\infty f(b) \overline{\psi(z)} d\sigma(z) \right\} \\ &= \int_0^\infty \overline{\psi(z)} ((\tau_{az}f)(b) - f(b)) d\sigma(z). \end{aligned}$$

Taking  $L^{p,\sigma}$  – norm of the wavelet coefficient

$$\|B_\psi(b, a)\|_{L^{p,\sigma}} = \int_0^\infty \left\{ \left| \int_0^\infty \overline{\psi(z)} ((\tau_{az}f)(b) - f(b)) d\sigma(z) \right|^p \right\}^{\frac{1}{p}} d\sigma(b).$$

Using Minkowski inequality of integrability for  $p \neq \infty$

$$\|B_\psi(b, a)\|_{L^{p,\sigma}} \leq \int_0^\infty \left\{ \int_0^\infty |(\tau_{az}f)(b) - f(b)| d\sigma(b) \right\}^{\frac{1}{p}} |\psi(z)| d\sigma(z). \quad (3.2)$$

Suppose that  $q < \infty$  and integrating w.r.t.  $a$ , we get

$$\int_0^\infty [a^{-\alpha} \|B_\psi(b, a)\|_{L^{p,\sigma}}]^q \frac{da}{a} \leq \int_0^\infty \left[ a^{-\alpha} \int_0^\infty |\psi(z)| \omega_p(f, az) d\sigma(z) \right]^q \frac{da}{a}.$$

Again using Minkowski integrabilty inequality

$$\int_0^\infty [a^{-\alpha} \|B_\psi(b, a)\|_{L^{p,\sigma}}]^q \frac{da}{a} \leq \left[ \int_0^\infty |\psi(z)| d\sigma(z) \left\{ \int_0^\infty (a^{-\alpha} \omega_p(f, az))^q \frac{da}{a} \right\}^{\frac{1}{q}} \right]^q.$$



Applying change of variable  $h = az$

$$\begin{aligned}
 &= \left[ \int_0^\infty z^{-\alpha} |\psi(z)| d\sigma(z) \left\{ \int_0^\infty (h^{-\alpha} \omega_{p,\sigma}(f, h))^q \frac{dh}{h} \right\}^{\frac{1}{q}} \right]^q \\
 &\leq \left\{ \int_0^\infty z^{-\alpha} |\psi(z)| d\sigma(z) \right\}^q \times \left\{ \int_0^\infty (h^{-\alpha} \omega_{p,\sigma}(f, h))^q \frac{dh}{h} \right\} \\
 &= \left\{ \int_0^\infty |z^{-\alpha} \psi(z)| d\sigma(z) \right\}^q \times \left\{ \int_0^\infty (h^{-\alpha} \omega_{p,\sigma}(f, h))^q \frac{dh}{h} \right\} \\
 &< \infty.
 \end{aligned} \tag{3.3}$$

If  $q = \infty$  the hypothesis on  $f$  says that  $h^{-\alpha} \omega_{p,\sigma}(f, h) \in L_{\infty,\sigma}(\mathbb{R}^+)$ , so

$$\|B_\psi(b, a)\|_{L_{p,\sigma}} \leq a^\alpha \|h^{-\alpha} \omega_{p,\sigma}(f, h)\|_{L_{\infty,\sigma}(\mathbb{R}^+)} \int_0^\infty |z^{-\alpha} \psi(z)| d\sigma(z). \tag{3.4}$$

The theorem has been proved for  $0 < \alpha < 1$ . For  $1 < \alpha < 2$ ,  $\alpha$  is not an integer, by hypothesis  $f'$  belongs to  $BH_{\alpha-1,\sigma}^p$ . Since  $\psi$  has cancellation up to order 2 therefore  $(z - \int_{-\infty}^z \psi(t) d\sigma(t))$  is an analysing wavelet and we can apply inequality (3.3) to the coefficient  $C(f', \int \psi, a, b)$

$$\begin{aligned}
 \int_0^\infty \left( \|C\left(f', \int \psi, a, \cdot\right)\|_{L_{\infty,\sigma}(\mathbb{R}^+)} \right)^q \frac{da}{a} &\leq \left\{ \int_0^\infty |z^{\alpha-1} \psi(z)| d\sigma(z) \right\}^q \\
 &\times \left\{ \int_0^\infty (h^{-\alpha-1} \omega_{p,\sigma}(f', h))^q \frac{dh}{h} \right\}.
 \end{aligned}$$

Now,

$$\begin{aligned}
 \int_0^\infty (a^{-\alpha} \|B_\psi(b, a)\|_{L_{p,\sigma}})^q \frac{d\sigma(a)}{a^{2\nu}} &\leq \left\{ \int_0^\infty |z^{\alpha-1} \psi(z)| d\sigma(z) \right\}^q \\
 &\times \left\{ \int_0^\infty (h^{-\alpha-1} \omega_{p,\sigma}(f', h))^q \frac{dh}{h} \right\} < \infty.
 \end{aligned}$$

This prove the result for  $1 < \alpha < 2$ . Similarly for  $q = \infty$ . Hence by recurrence on  $[\alpha]$  proved the theorem.  $\square$

Next theorem is the converse of the above theorem. The Bessel wavelet coefficients at small value of  $a$  is sufficient to characterizes Besov-Hankel spaces.

**Theorem 3.2.** *Suppose  $\alpha > 0$ ,  $\alpha$  not an integer and a function  $\psi$  is a real  $C^{[\alpha]+1}$ -regular analysing wavelet with all derivatives rapidly decreasing. If  $f, f', f'', f''', \dots, f^{[\alpha]} \in$*

$L^{p,\sigma}(\mathbb{R}^+)$  ( $1 < p < \infty$ ), and if  $a^{-\alpha}\|(B_\psi f)(a, \cdot)\|_{L^{p,\sigma}} \in L^{q,\sigma}(\mathbb{R}^+, \frac{da}{a})$ , then  $f \in BH_{\alpha,\sigma}^{p,q}$  and we have

$$\begin{aligned} \|h^{-(\alpha-[\alpha])}w_{p,\sigma}(f^{([\alpha])}, h)\|_{L^{q,\sigma}} &\leq \frac{1}{A_\psi} \left( \frac{2}{(\alpha - [\alpha])} \|\psi^{([\alpha])}\|_{L^{1,\sigma}} + \frac{1}{1 - (\alpha - [\alpha])} \|\psi^{([\alpha]+1)}\|_{L^{1,\sigma}} \right) \\ &\quad \times a^{-\alpha} \|(B_\psi f)(a, \cdot)\|_{L^{p,\sigma}} \end{aligned} \quad (3.5)$$

*Proof.* Let  $f \in L^{p,\sigma}(\mathbb{R}^+)$ . By inversion formula of Bessel wavelet transform

$$f(x) = \frac{1}{A_\psi} \int_0^\infty \frac{d\sigma(a)}{a^{2\nu+1}} \int_0^\infty (B_\psi f)(a, b) \psi_{a,b}(x) d\sigma(b) \quad (3.6)$$

and

$$\tau_h f(x) = \frac{1}{A_\psi} \int_0^\infty \frac{d\sigma(a)}{a^{2\nu+1}} \int_0^\infty (B_\psi f)(a, b) \tau_h \psi_{a,b}(x) d\sigma(b). \quad (3.7)$$

Then

$$\begin{aligned} \tau_h f(x) - f(x) &= \frac{1}{A_\psi} \int_0^\infty \frac{d\sigma(a)}{a^{2\nu+1}} \int_0^\infty (B_\psi f)(a, b) \{ \tau_h \psi_{a,b}(x) - \psi_{a,b}(x) \} d\sigma(b) \\ &= \frac{1}{A_\psi} \int_0^\infty \frac{d\sigma(a)}{a^{2\nu+1}} \int_0^\infty (B_\psi f)(a, b) \left\{ \tau_{\frac{h}{a}} \tau_{\frac{b}{a}} \psi\left(\frac{x}{a}\right) - \tau_{\frac{b}{a}} \psi\left(\frac{x}{a}\right) \right\} d\sigma(b) \\ &= \frac{1}{A_\psi} \int_0^\infty \frac{d\sigma(a)}{a^{2\nu+1}} \int_0^\infty (B_\psi f)(a, b) a^{-2\nu-1} D\left(\frac{b}{a}, \frac{x}{a}, y\right) d\sigma(b) \\ &\quad \times \int_0^\infty \left\{ \tau_{\frac{h}{a}} \psi(y) - \psi(y) \right\} d\sigma(y) \\ &= \frac{1}{A_\psi} \int_0^\infty \frac{d\sigma(a)}{a^{2\nu+1}} \tau_{\frac{h}{a}} (B_\psi f)\left(a, \frac{x}{a}\right) \int_0^\infty \left\{ \tau_{\frac{h}{a}} \psi(y) - \psi(y) \right\} d\sigma(y) \end{aligned}$$

Taking  $L^{p,\sigma}$ - norm on both side, we have

$$w_{p,\sigma}(f, h) = \frac{1}{A_\psi} \left\{ \int_0^\infty \left| \int_0^\infty \frac{d\sigma(a)}{a^{2\nu+1}} \tau_{\frac{h}{a}} (B_\psi f)\left(a, \frac{x}{a}\right) \int_0^\infty \left\{ \tau_{\frac{h}{a}} \psi(y) - \psi(y) \right\} d\sigma(y) \right|^p d\sigma(x) \right\}^{\frac{1}{p}}$$

applying Minkowski's inequality

$$\begin{aligned} &\leq \frac{1}{A_\psi} \int_0^\infty \frac{d\sigma(a)}{a^{2\nu+1}} \int_0^\infty \left| \tau_{\frac{h}{a}} \psi(y) - \psi(y) \right| d\sigma(y) \left\{ \int_0^\infty \left| \tau_{\frac{h}{a}} (B_\psi f)\left(a, \frac{x}{a}\right) \right|^p d\sigma(x) \right\}^{\frac{1}{p}} \\ &= \frac{1}{A_\psi} \int_0^\infty \frac{d\sigma(t)}{t^{2\nu}} \|(B_\psi f)\left(\frac{h}{t}, \cdot\right)\|_{L^{p,\sigma}} \int_0^\infty \left| \tau_{\frac{h}{a}} \psi(y) - \psi(y) \right| d\sigma(y) \end{aligned}$$

Now, consider  $0 < \alpha < 1$  and using Minkowski's inequality

$$\begin{aligned}
 \left\{ \int_0^\infty \frac{dh}{h} h^{-\alpha q} w_{p,\sigma}(f, h)^q \right\}^{\frac{1}{q}} &\leq \frac{1}{A_\psi} \int_0^\infty \frac{d\sigma(t)}{t^{2\nu}} \int_0^\infty |\tau_{\frac{h}{a}} \psi(y) - \psi(y)| d\sigma(y) \\
 &\times \left\{ \int_0^\infty \frac{dh}{h} h^{-\alpha q} \|(B_\psi f)\left(\frac{h}{t}, \cdot\right)\|_{L^{p,\sigma}}^q \right\}^{\frac{1}{q}} \\
 &= \frac{1}{A_\psi} \int_0^\infty \frac{d\sigma(t)}{t^{2\nu+\alpha}} \int_0^\infty |\tau_{\frac{h}{a}} \psi(y) - \psi(y)| d\sigma(y) \\
 &\times \left\{ \int_0^\infty \frac{da}{a} a^{-\alpha q} \|(B_\psi f)(a, \cdot)\|_{L^{p,\sigma}}^q \right\}^{\frac{1}{q}} \\
 &= \frac{C}{A_\psi} \int_0^\infty \frac{dt}{t^{1+\alpha}} \int_0^\infty |\tau_{\frac{h}{a}} \psi(y) - \psi(y)| d\sigma(y) \\
 &\times \left\{ \int_0^\infty \frac{da}{a} a^{-\alpha q} \|(B_\psi f)(a, \cdot)\|_{L^{p,\sigma}}^q \right\}^{\frac{1}{q}}.
 \end{aligned}$$

Using Lemma 1.1, we obtain

$$\int_0^\infty |\tau_{\frac{h}{a}} \psi(y) - \psi(y)| d\sigma(y) \leq 2\|\psi\|_{L^{1,\sigma}} \quad (3.8)$$

and

$$\begin{aligned}
 \int_0^\infty |\tau_{\frac{h}{a}} \psi(y) - \psi(y)| d\sigma(y) &= \int_0^\infty \left| \int_0^t (\tau_z \psi(y))' dz \right| d\sigma(y) \\
 &\leq \int_0^t \int_0^\infty |(\tau_z \psi(y))'| dz d\sigma(y) \\
 &\leq \|\psi'\|_{L^{1,\sigma}} t.
 \end{aligned} \quad (3.9)$$

Here, we observe that

$$\begin{aligned}
 \int_0^\infty \frac{dt}{t^{1+\alpha}} \int_0^\infty |\tau_{\frac{h}{a}} \psi(y) - \psi(y)| d\sigma(y) &= \int_0^1 \frac{dt}{t^{1+\alpha}} \int_0^\infty |\tau_{\frac{h}{a}} \psi(y) - \psi(y)| d\sigma(y) \\
 &+ \int_1^\infty \frac{dt}{t^{1+\alpha}} \int_0^\infty |\tau_{\frac{h}{a}} \psi(y) - \psi(y)| d\sigma(y) \\
 &\leq 2\|\psi\|_{L^{1,\sigma}} \int_1^\infty \frac{dt}{t^{1+\alpha}} + 2\|\psi'\|_{L^{1,\sigma}} \int_0^1 \frac{dt}{t^\alpha} \\
 &= \frac{2}{\alpha} \|\psi\|_{L^{1,\sigma}} + \frac{1}{1-\alpha} \|\psi'\|_{L^{1,\sigma}},
 \end{aligned}$$

which proved the result for  $0 < \alpha < 1$ .

If  $1 < \alpha < 2$ , by the hypothesis  $f' \in L^p(\mathbb{R}^+)$  and  $\psi$  is a  $C^1$ -regular function with  $\psi'$  rapidly decreasing at infinity. From equations (3.6) and (3.7), we have the equality

$$\tau_h f'(x) - f'(x) = \frac{1}{A_\psi} \int_0^\infty \frac{d\sigma(a)}{a^{2\nu+1}} \int_0^\infty (B_\psi f)(a, b) \{ \tau_h \psi'_{a,b}(x) - \psi'_{a,b}(x) \} d\sigma(b)$$

Calculate in similar manner as above for  $f'$  gives the following estimation

$$\begin{aligned} \left\{ \int_0^\infty \frac{dh}{h} h^{-(\alpha-1)q} w_{p,\sigma}(f, h)^q \right\}^{\frac{1}{q}} &\leq \frac{1}{A_\psi} \int_0^\infty \frac{dt}{t^{1+\alpha}} \int_0^\infty |\tau_{\frac{h}{a}} \psi(y) - \psi(y)| d\sigma(y) \\ &\quad \times \left\{ \int_0^\infty \frac{dh}{h} (a)^{-\alpha q} \|(B_\psi f)(a, \cdot)\|_{L^{p,\sigma}}^q \right\}^{\frac{1}{q}} \\ &\leq \frac{1}{A_\psi} \left( \frac{2}{(\alpha-1)} \|\psi'\|_{L^{1,\sigma}} + \frac{1}{1-(\alpha-1)} \|\psi''\|_{L^{1,\sigma}} \right) \\ &\quad \times \left\{ \int_0^\infty \frac{dh}{h} a^{-\alpha q} \|(B_\psi f)(a, \cdot)\|_{L^{p,\sigma}}^q \right\}^{\frac{1}{q}} \end{aligned}$$

this proves that  $f' \in BH_{\alpha-1,\sigma}^{p,q}$  the hypothesis  $a \rightarrow a^{-\alpha} \|(B_\psi f)(a, \cdot)\|_{L^{p,\sigma}}^q \in L^{q,\sigma}(\mathbb{R}^+)$  implies  $a \rightarrow a^{-(\alpha-1)} \|(B_\psi f)(a, \cdot)\|_{L^{p,\sigma}}^q \in L^{q,\sigma}(\mathbb{R}^+)$  then  $f \in BH_{\alpha-1,\sigma}^{p,q}$ . The theorem is established for  $1 < \alpha < 2$ , a recurrence on  $[\alpha]$  gives the final result.  $\square$

**Corollary 3.3.** *Let  $f \in B_{\alpha,\sigma}^{p,q}(\mathbb{R}^+)$  ( $p, q > 1, \alpha \neq \mathbb{Z}$ ), then*

$$\|f\|_{B_{\alpha,\sigma}^{p,q}} = \|f\|_{L^{p,\sigma}(\mathbb{R}^+)} + |f|_{B_{\alpha,\sigma}^{p,q}}$$

where  $|f|_{B_{\alpha,\sigma}^{p,q}}$  is equal to

$$|f|_{B_{\alpha,\sigma}^{p,q}}^q = \int_0^\infty (h^{-\alpha} w_p(\phi)(h))^q \frac{dh}{h} \approx \int_0^\infty [a^{-\alpha} \|B_\psi(\cdot, a)\|_{L^{p,\sigma}}]^q \frac{da}{a}.$$

**Remark 3.4.** *Considerable work has already been done on the characterization of Besov-k-Hankel norms by means of k-Hankel wavelet transform and Mehler-Besov-Fock spaces by using Mehler-Fock wavelet transform etc.. A further research in the context of different types of Besov space related to the different integral transform is needed.*

## ACKNOWLEDGEMENT

The research of the second author is supported by University Grants Commission (UGC), grant number: F.No. 16-6(DEC. 2017)/2018(NET/CSIR), New Delhi, India.

## REFERENCES

- [1] Pathak , R.S., Integral Transforms of Generalized Functions and Their Applications, Gordon and Breach Science Publishers, UK, 1997.
- [2] Pathak, R. S.and Dixit, M. M. , Continuous and discrete Bessel wavelet transforms, J. Comput. Appl. Math. 160 (2003), no. 1-2, 241-250.
- [3] Pathak, R. S., Upadhyay, S. K. and Pandey, R. S., The Bessel wavelet convolution product. Rend. Semin. Mat. Univ. Politec. Torino 69 (2011), no. 3, 267-279.
- [4] Pathak, R. S.and Upadhyay, S. K.  $L_p^\mu$ -boundedness of the pseudo-differential operator associated with the Bessel operator. J. Math. Anal. Appl. 257 (2001), no. 1, 141-153.
- [5] Perrier, Valérie; Basdevant, Claude, Besov norms in terms of the continuous wavelet transform. Application to structure functions. Math. Models Methods Appl. Sci. 6 (1996), no. 5, 649-664.
- [6] Upadhyay, S. K. and Singh, Reshma, Integrability of the continuum Bessel wavelet kernel. Int. J. Wavelets Multiresolut. Inf. Process. 13 (2015), no. 5, 1550032, 13 pp.
- [7] Hirschman, I. I., Jr. Variation diminishing Hankel transforms. J. Analyse Math. 8 (1960/61), 307-336.
- [8] Watson, G. N., A Treatise on the Theory of Bessel Functions, Cambridge University Press, Cambridge, 1958.
- [9] Betancor, Jorge J., Rodríguez-Mesa, L., On the Besov-Hankel spaces. J. Math. Soc. Japan 50 (1998), no. 3, 781-788.
- [10] Haimo, D.T. , Integral equations associated with Hankel convolutions, Trans. Amer. Math. Soc., 116 (1965), 330-375.
- [11] Trime'che, K., Generalized Wavelet and Hypergroup Gordon and Breach, Amsterdam,( 1997).
- [12] Stein, E. M., Singular Integral and Differentiability Properties of Functions, Princeton University Press, (1970).