

# THE $\mathbb{F}_p$ -SELBERG INTEGRAL OF TYPE $A_n$

RICHÁRD RIMÁNYI<sup>◊</sup> AND ALEXANDER VARCHENKO<sup>\*</sup>

<sup>◊,\*</sup> *Department of Mathematics, University of North Carolina at Chapel Hill  
Chapel Hill, NC 27599-3250, USA*

<sup>\*</sup> *Faculty of Mathematics and Mechanics, Lomonosov Moscow State University  
Leninskiye Gory 1, 119991 Moscow GSP-1, Russia*

<sup>\*</sup> *Moscow Center of Fundamental and Applied Mathematics  
Leninskiye Gory 1, 119991 Moscow GSP-1, Russia*

*Key words:* KZ and dynamical equations, reduction modulo  $p$ , Selberg integrals,  $\mathbb{F}_p$ -integrals

*2010 Mathematics Subject Classification:* 13A35 (33C60, 32G20)

ABSTRACT. We prove an  $\mathbb{F}_p$ -Selberg integral formula of type  $A_n$ , in which the  $\mathbb{F}_p$ -Selberg integral is an element of the finite field  $\mathbb{F}_p$  with odd prime number  $p$  of elements. The formula is motivated by analogy between multidimensional hypergeometric solutions of the KZ equations and polynomial solutions of the same equations reduced modulo  $p$ . For the type  $A_1$  the formula was proved in a previous paper by the authors.

## CONTENTS

|  |    |
|--|----|
| 1. Introduction  | 2  |
| 2. Preliminary remarks   | 3  |
| 2.1. Cancellation of factorials  | 3  |
| 2.2. Dyson's formula   | 4  |
| 2.3. $\mathbb{F}_p$ -Integrals   | 4  |
| 2.4. $\mathbb{F}_p$ -Beta integral   | 4  |
| 3. $\mathbb{F}_p$ -Selberg integral of type $A_n$  | 5  |
| 3.1. Admissible parameters   | 5  |
| 3.2. Main theorem  | 6  |
| 3.3. Relation between the $\mathbb{F}_p$ -Selberg integrals of types $A_{n-1}$ - and $A_n$ | 7  |
| 4. The $A_2$ -type Selberg integral over $\mathbb{C}$                                      | 8  |
| 4.1. The $A_2$ -formula over $\mathbb{C}$  | 8  |
| 4.2. Weight functions  | 9  |
| 4.3. Representations of $\mathfrak{sl}_3$  | 9  |
| 5. The $A_2$ -type Selberg integrals over $\mathbb{F}_p$                                   | 12 |
| 5.1. Relations between $\mathbb{F}_p$ -integrals   | 12 |
| 5.2. Theorem 3.3 for $n = 2$   | 14 |
| 5.3. Evaluation of $I_{0,0,0}(a, b_1, b_2, c)$   | 16 |
| 5.4. Proof of Theorem 3.3 for arbitrary $n > 2$  | 18 |
| References   | 19 |

<sup>◊</sup>*E-mail:* rimanyi@email.unc.edu, supported in part by Simons Foundation grant 523882

<sup>\*</sup>*E-mail:* anv@email.unc.edu, supported in part by NSF grant DMS-1954266

## 1. INTRODUCTION

In 1944 Atle Selberg proved the following integral formula:

$$(1.1) \quad \int_0^1 \cdots \int_0^1 \prod_{1 \leq i < j \leq k} (x_i - x_j)^{2\gamma} \prod_{i=1}^k x_i^{\alpha-1} (1-x_i)^{\beta-1} dx_1 \cdots dx_k \\ = \prod_{j=1}^k \frac{\Gamma(1+j\gamma)}{\Gamma(1+\gamma)} \frac{\Gamma(\alpha+(j-1)\gamma) \Gamma(\beta+(j-1)\gamma)}{\Gamma(\alpha+\beta+(k+j-2)\gamma)},$$

see [Se, AAR]. Hundreds of papers are devoted to the generalizations of the Selberg integral formula and its applications, see for example [AAR, FW] and references therein. There are  $q$ -analysis versions of the formula, the generalizations associated with Lie algebras, elliptic versions, finite field versions, see some references in [AAR, FW, As, Ha, Ka, Op, Ch, TV1, TV2, TV4, Wa1, Wa2, Sp, R, FSV, An, Ev]. In the finite field versions, one considers additive and multiplicative characters of a finite field, which map the field to the field of complex numbers, and forms an analog of equation (1.1), in which both sides are complex numbers. The simplest of such formulas is the classical relation between Jacobi and Gauss sums, see [AAR, An, Ev].

In [RV] we suggested another version of the Selberg integral formula, in which the  $\mathbb{F}_p$ -Selberg integral is an element of the finite field  $\mathbb{F}_p$  with an odd prime number  $p$  of elements.

Our motivation in [RV] came from the theory of Knizhnik-Zamolodchikov (KZ) equations, see [KZ, EFK]. These are the systems of linear differential equations, satisfied by conformal blocks on the sphere in the WZW model of conformal field theory. The KZ equations were solved in multidimensional hypergeometric integrals in [SV1], see also [V1, V2]. The following general principle was formulated in [MuV]: if an example of the KZ-type equations has a one-dimensional space of solutions, then the corresponding multidimensional hypergeometric integral can be evaluated explicitly. As an illustration of that principle in [MuV], an example of the  $\mathfrak{sl}_2$  differential KZ equations with a one-dimensional space of solutions was considered, the corresponding multidimensional hypergeometric integral was reduced to the Selberg integral and then evaluated by formula (1.1). See other illustrations in [FV, FSV, TV1, TV2, TV4, V3, RTVZ].

Recently in [SV2] the KZ equations were considered modulo a prime number  $p$  and polynomial solutions of the reduced equations were constructed, see also [SIV, V3, V4, V5, V6, V7]. The construction is analogous to the construction of the multidimensional hypergeometric solutions, and the constructed polynomial solutions were called the  $\mathbb{F}_p$ -hypergeometric solutions.

In [RV] we considered the reduction modulo  $p$  of the same example of the  $\mathfrak{sl}_2$  differential KZ equations, that led in [MuV] to the Selberg integral. We evaluated the corresponding  $\mathbb{F}_p$ -hypergeometric solution by analogy with the evaluation of the Selberg integral and obtained the  $\mathbb{F}_p$ -Selberg integral formula in [RV, Theorem 4.1].

In [TV4] the Selberg integral formula of type  $A_2$  was proved,

$$\begin{aligned}
(1.2) \quad & \int_{C_c^{k_1, k_2} [0, 1]} \prod_{i=1}^{k_1} t_i^{\alpha_1 - 1} (1 - t_i)^{\beta_1 - 1} \prod_{j=1}^{k_2} (1 - s_j)^{\beta_2 - 1} \prod_{i=1}^{k_1} \prod_{j=1}^{k_2} |s_j - t_i|^{-\gamma} \\
& \times \prod_{1 \leq i < i' \leq k_1} (t_i - t_{i'})^{2\gamma} \prod_{1 \leq j < j' \leq k_2} (s_j - s_{j'})^{2\gamma} dt_1 \dots dt_{k_1} ds_1 \dots ds_{k_2} \\
& = \prod_{i=1}^{k_1 - k_2} \frac{\Gamma(\beta_1 + (i - 1)\gamma)}{\Gamma(\alpha + \beta_1 + (i + k_1 - 2)\gamma)} \\
& \times \prod_{i=1}^{k_2} \frac{\Gamma(\beta_2 + (i - 1)\gamma)}{\Gamma(1 + \beta_2 + (i + k_2 - k_1 - 2)\gamma)} \frac{\Gamma(\beta_1 + \beta_2 + (i - 2)\gamma)}{\Gamma(\alpha + \beta_1 + \beta_2 + (i + k_2 - 3)\gamma)} \\
& \times \prod_{i=1}^{k_2} \frac{\Gamma(1 + (i - k_1 - 1)\gamma) \Gamma(i\gamma)}{\Gamma(\gamma)} \prod_{i=1}^{k_1} \frac{\Gamma(\alpha + (i - 1)\gamma) \Gamma(i\gamma)}{\Gamma(\gamma)}.
\end{aligned}$$

The starting point of this formula was an example of the joint system of the  $\mathfrak{sl}_3$  trigonometric differential KZ equations and associated dynamical difference equations, an example in which the space of solutions is one-dimensional. The  $A_n$ -type Selberg integral formula for arbitrary  $n$  was obtained in [Wa1, Wa2], see also [FW].

In this paper we consider the reduction modulo  $p$  of the same example of the joint system of the  $\mathfrak{sl}_{n+1}$  trigonometric differential KZ equations and associated dynamical difference equations, which led in [TV4, Wa1] to the  $A_n$ -type Selberg integral formula. Using the reduction modulo  $p$  of these differential and difference equations we obtain our  $A_n$ -type  $\mathbb{F}_p$ -Selberg integral formula in Theorem 3.3.

The paper is organized as follows. In Section 2 we collect useful facts. In Section 3 we introduce the notion of  $\mathbb{F}_p$ -integral and discuss the integral formula for the  $\mathbb{F}_p$ -beta integral. In Section 3 we formulate our main Theorem 3.3, containing the  $A_n$ -type  $\mathbb{F}_p$ -Selberg integral formula. For  $n = 1$  it is [RV, Theorem 4.1]. Theorem 3.3 is proved by induction on  $n$  in the remaining part of the paper. In Section 3 we also prove Theorem 3.6, which is used in the transition from the  $A_{n-1}$ -type formula to the  $A_n$ -type formula. In Section 4 we sketch the proof of formula (1.2) following [TV4]. In Section 5 we adapt this proof to prove the  $A_2$ -type  $\mathbb{F}_p$ -Selberg integral formula. The proof of the  $A_n$ -type  $\mathbb{F}_p$ -Selberg integral formula for  $n > 2$  is similar, see Section 5.4.

The authors thank I. Cherednik, P. Etingof, E. Rains for useful discussions.

## 2. PRELIMINARY REMARKS

Let  $p$  be an odd prime number throughout the paper.

### 2.1. Cancellation of factorials.

**Lemma 2.1.** *If  $a, b$  are nonnegative integers and  $a + b = p - 1$ , then in  $\mathbb{F}_p$  we have*

$$(2.1) \quad a! b! = (-1)^{a+1}.$$

*Proof.* We have  $a! = (-1)^a (p-1) \dots (p-a)$  and  $p-a = b+1$ . Hence  $a! b! = (-1)^a (p-1)! = (-1)^{a+1}$  by Wilson's Theorem.  $\square$

**2.2. Dyson's formula.** We shall use Dyson's formula

$$(2.2) \quad \text{C.T.} \quad \prod_{1 \leq i < j \leq k} (1 - x_i/x_j)^c (1 - x_j/x_i)^c = \frac{(kc)!}{(c!)^k},$$

where C.T. denotes the constant term. See the formula in [AAR, Section 8.8].

**2.3.  $\mathbb{F}_p$ -Integrals.** Let  $M$  an  $\mathbb{F}_p$ -module. Let  $P(x_1, \dots, x_k)$  be a polynomial with coefficients in  $M$ ,

$$(2.3) \quad P(x_1, \dots, x_k) = \sum_d c_d x_1^{d_1} \dots x_k^{d_k}.$$

Let  $l = (l_1, \dots, l_k) \in \mathbb{Z}_{>0}^k$ . The coefficient  $c_{l_1 p - 1, \dots, l_k p - 1}$  is called the  $\mathbb{F}_p$ -integral over the  $p$ -cycle  $[l_1, \dots, l_k]_p$  and is denoted by  $\int_{[l_1, \dots, l_k]_p} P(x_1, \dots, x_k) dx_1 \dots dx_k$ .

**Lemma 2.2.** For  $i = 1, \dots, k - 1$  we have

$$(2.4) \quad \int_{[l_1, \dots, l_{i+1}, l_i, \dots, l_k]_p} P(x_1, \dots, x_{i+1}, x_i, \dots, x_k) dx_1 \dots dx_k \\ = \int_{[l_1, \dots, l_k]_p} P(x_1, \dots, x_k) dx_1 \dots dx_k.$$

□

**Lemma 2.3.** For any  $i = 1, \dots, k$ , we have

$$\int_{[l_1, \dots, l_k]_p} \frac{\partial P}{\partial x_i}(x_1, \dots, x_k) dx_1 \dots dx_k = 0.$$

□

Let  $\mathbf{k} = (k_1, \dots, k_n) \in \mathbb{Z}_{>0}^n$  and

$$(2.5) \quad [\mathbf{k}]_p := [(1)_{k_1}; (k_1)_{k_2}; \dots; (k_{n-1})_{k_n}]_p,$$

where  $x_y$  denotes the  $y$ -tuple  $(x, \dots, x)$ . For example for  $n = 2$ ,  $\mathbf{k} = (3, 2)$ , we have  $[\mathbf{k}]_p = [1, 1, 1; 3, 3]_p$ .

**2.4.  $\mathbb{F}_p$ -Beta integral.** For nonnegative integers the classical beta integral formula says

$$(2.6) \quad \int_0^1 x^a (1-x)^b dx = \frac{a! b!}{(a+b+1)!}.$$

**Theorem 2.4** ([V7]). Let  $a < p$ ,  $b < p$ ,  $p - 1 \leq a + b$ . Then in  $\mathbb{F}_p$  we have

$$(2.7) \quad \int_{[1]_p} x^a (1-x)^b dx = -\frac{a! b!}{(a+b+1-p)!}.$$

If  $a + b < p - 1$ , then

$$(2.8) \quad \int_{[1]_p} x^a (1-x)^b dx = 0.$$

3.  $\mathbb{F}_p$ -SELBERG INTEGRAL OF TYPE  $A_n$ 

**3.1. Admissible parameters.** Let  $\mathbf{k} = (k_1, \dots, k_n) \in \mathbb{Z}_{>0}^n$  and  $k_i > k_{i+1}$ ,  $i = 1, \dots, n-1$ . Set  $k_0 = k_{n+1} = 0$ .

Let  $a, b_1, \dots, b_n, c \in \mathbb{Z}_{>0}$ . Denote  $b = (b_1, \dots, b_n)$  and

(3.1)

$$\begin{aligned} R_{\mathbf{k}}(a, b, c) &= \\ &= \prod_{1 \leq s \leq r \leq n} \prod_{i=1}^{k_r - k_{r+1}} \frac{(r - s + b_s + \dots + b_r + (i + s - r - 1)c)!}{(r - s + 1 + a_s + b_s + \dots + b_r + (i + s - r + k_s - k_{s-1} - 2)c - \delta_{s,1}p)!} \\ &\times (-1)^{\sum_{i=1}^n k_i} \left( \prod_{i=1}^{k_1} (a_1 + (i-1)c)! \right) \left( \prod_{r=1}^n \prod_{i=1}^{k_r} \frac{(ic)!}{c!} \right) \left( \prod_{r=2}^n \prod_{i=1}^{k_r} (p + (i - k_{r-1} - 1)c)! \right), \end{aligned}$$

where  $a_1 = a$ ,  $a_2 = \dots = a_n = 0$ ;  $\delta_{s,1}$  is 1 if  $s = 1$  and is zero otherwise.

We say that  $a, b_1, \dots, b_n, c \in \mathbb{Z}_{>0}$  are *admissible* if  $a + (k_1 - 1)c < p - 1$  and for any factorial  $x!$  in the right-hand side of (3.1) we have  $0 \leq x < p$ . The set of all admissible  $(a, b, c)$  is denoted by  $\mathcal{A}_{\mathbf{k}}$ .

**Lemma 3.1.** *The set  $\mathcal{A}_{\mathbf{k}}$  is defined in  $\mathbb{Z}_{>0}^{n+2}$  by the following system of inequalities:*

$$(3.2) \quad \begin{aligned} 0 &\leq r - s + b_s + \dots + b_r + (s - r)c, \\ r - s + b_s + \dots + b_r + (k_r - k_{r+1} + s - r - 1)c &\leq p - 1, \end{aligned}$$

for  $1 \leq s \leq r \leq n$ ;

$$(3.3) \quad \begin{aligned} 0 &\leq r - s + 1 + b_s + \dots + b_r + (s - r + k_s - k_{s-1} - 1)c, \\ r - s + 1 + b_s + \dots + b_r + (s - r + k_r - k_{r+1} + k_s - k_{s-1} - 2)c &\leq p - 1, \end{aligned}$$

for  $2 \leq s \leq r \leq n$ ;

$$(3.4) \quad \begin{aligned} p &\leq r + a + b_1 + \dots + b_r + (k_1 - r)c, \\ r + a + b_1 + \dots + b_r + (k_r - k_{r+1} + k_1 - r - 1)c &< 2p, \end{aligned}$$

for  $1 \leq r \leq n$ ;

$$(3.5) \quad a + (k_1 - 1)c < p - 1, \quad b_1 \geq p - 1 - (a + (k_1 - 1)c), \quad 0 < k_1 c < p.$$

□

**Lemma 3.2.** *Assume that  $(a, b, c) \in \mathcal{A}_{\mathbf{k}}$ . Then*

$$(3.6) \quad b_1 \geq p - 1 - (a + (k_1 - 1)c), \quad b_s \geq (k_{s-1} - k_s + 1)c - 1, \quad s = 2, \dots, n.$$

*Proof.* The inequality  $b_s \geq (k_{s-1} - k_s + 1)c - 1$  for  $s = 2, \dots, n$  follows from the first inequality in (3.3) for  $r = s$ . The inequality  $b_1 \geq p - 1 - (a + (k_1 - 1)c)$  follows from the first inequality in (3.4) for  $r = 1$ . □

**Example.** Let  $n = 1$ ,  $\mathbf{k} = (k_1)$ . Then

$$(3.7) \quad R_{(k_1)}(a, b_1, c) = \prod_{i=1}^{k_1} \frac{(ic)!}{c!} \frac{(a + (i-1)c)! (b_1 + (i-1)c)!}{(1 + a + b_1 + (i + k_1 - 2)c - p)!}$$

and  $\mathcal{A}_{(k_1)}$  consists of  $a, b, c \in \mathbb{Z}_{>0}$  such that

$$(3.8) \quad \begin{aligned} a + (k_1 - 1)c < p - 1, & \quad b_1 + (k_1 - 1)c \leq p - 1, & \quad k_1 c \leq p - 1, \\ p - 1 \leq a + b_1 + (k_1 - 1)c, & \quad a + b_1 + (2k_1 - 2)c < 2p - 1. \end{aligned}$$

**3.2. Main theorem.** Given  $\mathbf{k} = (k_1, \dots, k_n) \in \mathbb{Z}_{>0}^n$  introduce  $k_1 + \dots + k_n$  variables

$$(3.9) \quad t = (t^{(1)}, \dots, t^{(n)}), \quad \text{where} \quad t^{(i)} = (t_1^{(i)}, \dots, t_{k_i}^{(i)}), \quad i = 1, \dots, n.$$

Define the *master polynomial*

$$\Phi_{\mathbf{k}}(t; a, b, c) = \prod_{i=1}^n \left( \prod_{j=1}^{k_i} (t_j^{(i)})^{a_i} (1 - t_j^{(i)})^{b_i} \prod_{1 \leq j < j' \leq k_i} (t_j^{(i)} - t_{j'}^{(i)})^{2c} \right) \prod_{i=1}^{n-1} \prod_{j=1}^{k_{i+1}} \prod_{j'=1}^{k_i} (t_j^{(i+1)} - t_{j'}^{(i)})^{p-c}.$$

Denote

$$(3.10) \quad S_{\mathbf{k}}(a, b, c) = \int_{[\mathbf{k}]_p} \Phi_{\mathbf{k}} dt.$$

The  $\mathbb{F}_p$ -integral  $S_{\mathbf{k}}(a, b, c)$  is called the  $\mathbb{F}_p$ -Selberg integral of type  $A_n$ .

Here is our main result.

**Theorem 3.3.** *Let  $\mathbf{k} = (k_1, \dots, k_n) \in \mathbb{Z}_{>0}^n$ ,  $k_i > k_{i+1}$ ,  $i = 1, \dots, n - 1$ . Then for any  $(a, b, c) \in \mathcal{A}_{\mathbf{k}}$  we have the equality in  $\mathbb{F}_p$ :*

$$(3.11) \quad S_{\mathbf{k}}(a, b, c) = R_{\mathbf{k}}(a, b, c).$$

For  $n = 1$  this theorem is [RV, Theorem 4.1]. For  $n = 2$  the theorem is deduced from the case of  $n = 1$  in Section 5. More generally, for any  $k$  the theorem for  $n = k$  is deduced from the theorem for  $n = k - 1$  similarly, see Section 5.4.

**Remark.** Theorem 3.3 can be extended to the case of  $\mathbf{k}$  such that  $k_i \geq k_{i+1}$ ,  $i = 1, \dots, n - 1$ , but the structure of inequalities in Lemma 3.1 will depend on the appearance of equalities  $k_i = k_{i+1}$  in  $\mathbf{k}$ , and the proof of Theorem 3.3 will split into different sub-cases. To shorten the exposition we restrict ourselves with  $\mathbf{k}$  such that  $k_i > k_{i+1}$ ,  $i = 1, \dots, n - 1$ .

**Example.** Here is the simplest  $A_2$ -type  $\mathbb{F}_p$ -Selberg integral formula with  $k_1 = k_2 = 1$ .

**Theorem 3.4.** *Assume that  $a, b_1, b_2, c$  are integers such that*

$$\begin{aligned} 0 \leq a < p, \quad 0 < c \leq p, \quad 0 \leq b_2 - c + 1 < p, \\ 0 \leq b_1 + b_2 - c + 1 < p, \quad p - 1 \leq a + b_1 + b_2 - c + 1 < 2p - 1. \end{aligned}$$

*Then in  $\mathbb{F}_p$  we have*

$$(3.12) \quad \int_{[1;1]_p} t^a (1-t)^{b_1} (s-t)^{p-c} (1-s)^{b_2} dt ds = \frac{a! (b_1 + b_2 - c + 1)! (p-c)! (b_2)!}{(a + b_1 + b_2 - c + 2 - p)! (b_2 - c + 1)!}.$$

*Proof.* Change variables  $s = t + (1-t)v$ , then the  $\mathbb{F}_p$ -integral becomes equal to

$$\int_{[1,1]_p} t^a (1-t)^{b_1+b_2-c+1} v^{p-c} (1-v)^{b_2} dt dv.$$

Applying the  $\mathbb{F}_p$ -beta integral formula we obtain the theorem.  $\square$

The simplest  $A_3$ -type  $\mathbb{F}_p$ -Selberg integral formula  $k_1 = k_2 = k_3 = 1$  is given by the next theorem.

**Theorem 3.5.** *Let  $a, b_1, b_2, b_3, c$  be integers such that all factorials in the right-hand side of formula (3.13) are factorials of nonnegative integers less than  $p$ . Then in  $\mathbb{F}_p$  we have*

$$(3.13) \quad \int_{[1;1;1]_p} t^a (1-t)^{b_1} (s-t)^{p-c} (1-s)^{b_2} (u-s)^{p-c} (1-u)^{b_2} dt ds du \\ = - \frac{a! (b_1 + b_2 + b_3 - 2c + 2)!}{(a + b_1 + b_2 + b_3 - 2c + 3 - p)!} \frac{(p-c)! (b_2 + b_3 - c + 1)! (p-c)! (b_3)!}{(b_2 + b_3 - 2c + 2)! (b_3 - c + 1)!}.$$

*Proof.* The proof is the same as of the previous theorem.  $\square$

The versions of identities (3.12), (3.13) over complex numbers see in [MuV, Theorem 1].

### 3.3. Relation between the $\mathbb{F}_p$ -Selberg integrals of types $A_{n-1}$ - and $A_n$ .

**Theorem 3.6.** *Let  $n > 1$  and  $\mathbf{k} = (k_1, \dots, k_n)$ ,  $\mathbf{k}' = (k_1, \dots, k_{n-1})$ ,  $b = (b_1, \dots, b_n)$ ,  $b' = (b_1, \dots, b_{n-1})$ . Assume that formula (3.11) holds for the  $\mathbb{F}_p$ -Selberg integral  $S_{[\mathbf{k}']_p}(a, b', c)$  of type  $A_{n-1}$ . Also assume that  $b_n = (k_{n-1} - k_n + 1)c - 1$ . Then formula (3.11) holds for the  $\mathbb{F}_p$ -Selberg integral  $S_{[\mathbf{k}]_p}(a, b, c)$  of type  $A_n$ .*

*Proof.* Under the assumption  $b_n = (k_{n-1} - k_n + 1)c - 1$  all variables  $(t_j^{(n)})$  in  $\Phi_{[\mathbf{k}]_p}(\mathbf{t}; a, b, c)$  are used to reach the monomial  $\prod_{j=1}^{k_n} (t_j^{(n)})^{k_n - 1} p^{-1}$  in the calculation of the  $\mathbb{F}_p$ -integral  $S_{[\mathbf{k}]_p}(a, b, c)$ .

The remaining free variables  $(t_j^{(i)})$  with  $i < n$  all belong to the factor  $\Phi_{[\mathbf{k}']_p}(t^{(1)}, \dots, t^{(n-1)}; a, b', c)$  of  $\Phi_{[\mathbf{k}]_p}(\mathbf{t}; a, b, c)$  and are used to calculate the coefficient of  $\prod_{j=1}^{k_1} (t_j^{(1)})^{p-1} \prod_{i=2}^{n-1} \prod_{j=1}^{k_i} (t_j^{(i)})^{k_i - 1} p^{-1}$ .

More precisely, under the assumptions of the theorem we have

$$S_{\mathbf{k}}(a, b, c) = (-1)^{b_n k_n + c k_n (k_n - 1)/2} \frac{(k_n c)!}{(c!)^{k_n}} S_{\mathbf{k}'}(a', b', c),$$

where  $(-1)^{b_n k_n + c k_n (k_n - 1)/2} \frac{(k_n c)!}{(c!)^{k_n}}$  is the coefficient of  $\prod_{j=1}^{k_n} (t_j^{(n)})^{k_n - 1} p^{-1}$  in the expansion of

$$\prod_{j=1}^{k_n} \prod_{j'=1}^{k_{n-1}} (t_j^{(n)} - t_{j'}^{(n-1)})^{p-c} \prod_{1 \leq j < j' \leq k_n} (t_j^{(n)} - t_{j'}^{(n)})^{2c},$$

see Dyson's formula. We have  $(-1)^{b_n k_n + c k_n (k_n - 1)/2} = (-1)^{(k_{n-1} k_n - k_n (k_n + 1)/2)c - k_n}$ . Hence

$$S_{\mathbf{k}}(a, b, c) = (-1)^{(k_{n-1} k_n - k_n (k_n + 1)/2)c - k_n} \frac{(k_n c)!}{(c!)^{k_n}} S_{\mathbf{k}'}(a, b', c) \\ = (-1)^{(k_{n-1} k_n - k_n (k_n + 1)/2)c - k_n} \frac{(k_n c)!}{(c!)^{k_n}} R_{\mathbf{k}'}(a, b', c),$$

where  $S_{\mathbf{k}'}(a, b', c) = R_{\mathbf{k}'}(a, b', c)$  holds by assumptions. To prove the theorem we need to show that

$$R_{[\mathbf{k}]_p}(a, (b', b_n), c) = (-1)^{(k_{n-1} k_n - k_n (k_n + 1)/2)c - k_n} \frac{(k_n c)!}{(c!)^{k_n}} R_{\mathbf{k}'}(a, b', c).$$

Indeed we have

$$\begin{aligned}
R_{\mathbf{k}}(a, (b', b_n), c) &= R_{\mathbf{k}'}(a, b', c) \prod_{i=1}^{k_n} \frac{(ic)!}{c!} \prod_{i=1}^{k_n} (p + (i - k_{n-1} - 1)c)! \\
&\times \prod_{1 \leq s \leq n} \prod_{i=1}^{k_n} \frac{(n - s + b_s + \cdots + b_n + (i + s - n - 1)c)!}{(n - s + 1 + a_s + b_s + \cdots + b_n + (i + s - n + k_s - k_{s-1} - 2)c - \delta_{s,1}p)!} \\
&\times \prod_{1 \leq s \leq n-1} \prod_{i=1}^{k_n} \frac{(n - s + a_s + b_s + \cdots + b_{n-1} + (i + s - n + k_s - k_{s-1} - 3)c - \delta_{s,1}p)!}{(n - 1 - s + b_s + \cdots + b_{n-1} + (i + s - r - 1)c)!} \\
&= R_{\mathbf{k}'}(a, b', c) \prod_{i=1}^{k_n} \frac{(ic)!}{c!} \prod_{i=1}^{k_n} \frac{(b_n + (i - 1)c)! (p + (i - k_{n-1} - 1)c)!}{(1 + b_n + (i + k_n - k_{n-1} - 2)c)!} \\
&= R_{\mathbf{k}'}(a, b', c) \prod_{i=1}^{k_n} \frac{(ic)!}{c!} \prod_{i=1}^{k_n} \frac{((i + k_{n-1} - k_n)c - 1)! (p + (i - k_{n-1} - 1)c)!}{((i - 1)c)!} \\
&= R_{\mathbf{k}'}(a, b', c) (-1)^{(k_{n-1}k_n - k_n(k_n+1)/2)c - k_n} \frac{(k_n c)!}{(c!)^{k_n}},
\end{aligned}$$

where in the last step we use the cancellation Lemma 2.1. The theorem is proved.  $\square$

**Corollary 3.7.** *Let  $n > 1$ ,  $\mathbf{k} = (k_1, \dots, k_n)$ , and  $(a, b_1, c) \in \mathcal{A}_{(k_1)}$ . Let  $b = (b_1, \dots, b_n)$ , where  $b_i = (k_{i-1} - k_i + 1)c - 1$  for  $i = 2, \dots, n$ . Then formula (3.11) holds for the  $\mathbb{F}_p$ -Selberg integral  $S_{[\mathbf{k}]_p}(a, b, c)$  of type  $A_n$ .*

*Proof.* Formula (3.11) for the  $\mathbb{F}_p$ -Selberg integrals of type  $A_1$  is proved in [RV]. Hence the corollary follows from Theorem 3.6 by induction on  $n$ .  $\square$

#### 4. THE $A_2$ -TYPE SELBERG INTEGRAL OVER $\mathbb{C}$

In this section we formulate the  $A_2$ -type Selberg integral formula over  $\mathbb{C}$ , formulated and proved in [TV4], and sketch the proof of the formula, following [TV4]. In Section 5 we adapt this proof to prove the  $A_2$ -type  $\mathbb{F}_p$ -Selberg integral formula, that is, formula (3.11) for  $n = 2$ .

**4.1. The  $A_2$ -formula over  $\mathbb{C}$ .** For  $k_1 \geq k_2 \geq 0$  let  $t = (t_1, \dots, t_{k_1})$ ,  $s = (s_1, \dots, s_{k_2})$ . Define the *master function*

$$\begin{aligned}
(4.1) \quad \Phi(t; s) &= \prod_{i=1}^{k_1} t_i^{\alpha_1 - 1} (1 - t_i)^{\beta_1 - 1} \prod_{j=1}^{k_2} (1 - s_j)^{\beta_2 - 1} \prod_{i=1}^{k_1} \prod_{j=1}^{k_2} |s_j - t_i|^{-\gamma} \\
&\times \prod_{1 \leq i < i' \leq k_1} (t_i - t_{i'})^{2\gamma} \prod_{1 \leq j < j' \leq k_2} (s_j - s_{j'})^{2\gamma}
\end{aligned}$$

and the integral

$$(4.2) \quad \tilde{S}(\alpha, \beta_1, \beta_2, \gamma) = \int_{C^{k_1, k_2}[0, 1]} \Phi(t; s) dt ds,$$

where  $C^{k_1, k_2}[0, 1]$  is the integration cycle, defined in [TV4, Section 3], also see its definition in [Wa1, Wa2, FW]. The explicit description of this cycle is of no importance in this paper.

**Theorem 4.1** ([TV4, Theorem 3.3]). *Let  $a, b_1, b_2, c \in \mathbb{C}$  such that  $\operatorname{Re}(a), \operatorname{Re}(b_1), \operatorname{Re}(b_2) > -1$ ,  $\operatorname{Re}(c) < 0$  and  $|\operatorname{Re}(c)|$  sufficiently small. Then*

$$(4.3) \quad \begin{aligned} \tilde{S}(\alpha, \beta_1, \beta_2, \gamma) &= \prod_{i=1}^{k_1-k_2} \frac{\Gamma(\beta_1 + (i-1)\gamma)}{\Gamma(\alpha + \beta_1 + (i+k_1-2)\gamma)} \\ &\times \prod_{i=1}^{k_2} \frac{\Gamma(\beta_2 + (i-1)\gamma)}{\Gamma(1 + \beta_2 + (i+k_2-k_1-2)\gamma)} \frac{\Gamma(\beta_1 + \beta_2 + (i-2)\gamma)}{\Gamma(\alpha + \beta_1 + \beta_2 + (i+k_2-3)\gamma)} \\ &\times \prod_{i=1}^{k_2} \frac{\Gamma(1 + (i-k_1-1)\gamma) \Gamma(i\gamma)}{\Gamma(\gamma)} \prod_{i=1}^{k_1} \frac{\Gamma(\alpha + (i-1)\gamma) \Gamma(i\gamma)}{\Gamma(\gamma)}. \end{aligned}$$

In the next Sections 4.2, 4.3 we sketch the proof of formula (4.3) following [TV4].

**4.2. Weight functions.** To evaluate  $\tilde{S}(\alpha, \beta_1, \beta_2, \gamma)$  we introduce a collection of new integrals  $J_{l_1, l_2, m}(\alpha, \beta_1, \beta_2, \gamma)$ , which also can be evaluated explicitly, see [TV4].

For a function  $f(t_1, \dots, t_k)$  set

$$\operatorname{Sym}_{t_1, \dots, t_k} f(t_1, \dots, t_k) = \frac{1}{k!} \sum_{\sigma \in S_k} f(t_{\sigma_1}, \dots, t_{\sigma_k}).$$

Given  $k_1 \geq k_2 \geq 0$ , we say that a triple of nonnegative integers  $(l_1, l_2, m)$  is *allowable* if  $l_1 \leq k_1 - k_2 + l_2$ ,  $l_2 \leq k_2$  and  $m \leq \min(l_1, l_2)$ . For any allowable triple  $(l_1, l_2, m)$  define the weight function

$$\begin{aligned} W_{l_1, l_2, m}(t_1, \dots, t_{k_1}; s_1, \dots, s_{k_2}) &= \\ &= \operatorname{Sym}_{t_1, \dots, t_{k_1}} \operatorname{Sym}_{s_1, \dots, s_{k_2}} \left( \prod_{a=1}^{l_1} t_a \prod_{a=l_1+1}^{k_1} (1-t_a) \prod_{b=1}^m \frac{1-s_b}{s_b-t_b} \prod_{b=l_2+1}^{k_2} \frac{1-s_b}{s_b-t_{b+k_1-k_2}} \right) \end{aligned}$$

and the integral

$$J_{l_1, l_2, m}(\alpha, \beta_1, \beta_2, \gamma) = \int_{C^{k_1, k_2}[0,1]} \Phi(t; s) W_{l_1, l_2, m}(t; s) dt ds.$$

In particular,

$$(4.4) \quad J_{0, k_2, 0}(\alpha, \beta_1, \beta_2, \gamma) = \tilde{S}(\alpha, \beta_1 + 1, \beta_2, \gamma).$$

**4.3. Representations of  $\mathfrak{sl}_3$ .** Consider the complex Lie algebra  $\mathfrak{sl}_3$  with standard generators  $f_1, f_2, e_1, e_2, h_1, h_2$ , simple roots  $\sigma_1, \sigma_2$ , fundamental weights  $\omega_1, \omega_2$ . Let  $V_{\lambda_1}, V_{\lambda_2}$  be the irreducible  $\mathfrak{sl}_3$ -modules with highest weights

$$\lambda_1 = -\frac{\alpha}{\gamma} \omega_1, \quad \lambda_2 = -\frac{\beta_1}{\gamma} \omega_1 - \frac{\beta_2}{\gamma} \omega_2$$

and highest weight vectors  $v_1, v_2$ . For  $k_1 \geq k_2 \geq 0$  consider the weight subspace  $V_{\lambda_1} \otimes V_{\lambda_2}[\lambda_1 + \lambda_2 - k_1\sigma_1 - k_2\sigma_2]$  of the tensor product  $V_{\lambda_1} \otimes V_{\lambda_2}$  and the singular weight subspace  $\operatorname{Sing} V_{\lambda_1} \otimes V_{\lambda_2}[\lambda_1 + \lambda_2 - k_1\sigma_1 - k_2\sigma_2]$  consisting of the vectors  $w \in V_{\lambda_1} \otimes V_{\lambda_2}[\lambda_1 + \lambda_2 - k_1\sigma_1 - k_2\sigma_2]$

such that  $e_1 w = 0$ ,  $e_2 w = 0$ . A basis of  $V_{\lambda_1} \otimes V_{\lambda_2}[\lambda_1 + \lambda_2 - k_1 \sigma_1 - k_2 \sigma_2]$  is formed by the vectors

$$v_{l_1, l_2, m} = \frac{f_1^{k_1 - k_2 - l_1 + l_2} [f_1, f_2]^{k_2 - l_2} v_1 \otimes f_1^{l_1 - m} [f_1, f_2]^m f_2^{l_2 - m} v_2}{(k_1 - k_2 - l_1 + l_2)! (k_2 - l_2)! (l_1 - m)! m! (l_2 - m)!}$$

labeled by allowable triples  $(l_1, l_2, m)$ . It is known from the theory of KZ equations that the vector

$$J = \sum_{l_1, l_2, m} (-1)^{l_1} J_{l_1, l_2, m}(\alpha, \beta_1, \beta_2, \gamma) v_{l_1, l_2, m}$$

is a singular vector, see [M, Theorem 2.4], [MaV, Corollary 10.3], cf. [RSV].

The singular vector equations  $e_1 J = 0$ ,  $e_2 J = 0$  are calculated with the help of the formulas:

$$\begin{aligned} h_1 v_1 &= -\frac{\alpha}{\gamma} v_1, & h_2 v_1 &= 0, & h_1 v_2 &= -\frac{\beta_1}{\gamma} v_2, & h_2 v_2 &= -\frac{\beta_2}{\gamma} v_2, \\ [h_1, f_1] &= -2f_1, & [h_1, f_2] &= f_2, & [h_2, f_1] &= f_1, & [h_2, f_2] &= -2f_2, \\ [e_1, f_1] &= h_1, & [e_1, f_2] &= [e_2, f_1] = 0, & [e_2, f_2] &= h_2, \end{aligned}$$

$$[h_1, [f_1, f_2]] = -[f_1, f_2], \quad [h_2, [f_1, f_2]] = -[f_1, f_2], \quad [e_1, [f_1, f_2]] = f_2, \quad [e_2, [f_1, f_2]] = -f_1.$$

Here are some of the singular vector relations.

**Theorem 4.2** (cf. [TV4, Theorem 5.2]). *We have*

$$(4.5) \quad J_{0, l_2, 0} = (-1)^{l_2} J_{0, 0, 0} \prod_{i=0}^{l_2-1} \frac{(k_1 - k_2 + i + 1)\gamma}{\beta_2 + i\gamma}.$$

*Proof.* We have

$$\begin{aligned} e_2 \frac{f_1^{k_1 - k_2 + i} [f_1, f_2]^{k_2 - i} v_1 \otimes f_2^i v_2}{(k_1 - k_2 + i)! (k_2 - i)! i!} &= - (k_1 - k_2 + i + 1) \frac{f_1^{k_1 - k_2 + i + 1} [f_1, f_2]^{k_2 - i - 1} v_1 \otimes f_2^i v_2}{(k_1 - k_2 + i + 1)! (k_2 - i - 1)! i!} \\ &+ \left( -\frac{\beta_2}{\gamma} - i + 1 \right) \frac{f_1^{k_1 - k_2 + i} [f_1, f_2]^{k_2 - i} v_1 \otimes f_2^{i-1} v_2}{(k_1 - k_2 + i)! (k_2 - i)! (i-1)!}. \end{aligned}$$

Calculating the coefficient of  $\frac{f_1^{k_1 - k_2 + i + 1} [f_1, f_2]^{k_2 - i - 1} v_1 \otimes f_2^i v_2}{(k_1 - k_2 + i)! (k_2 - i)! i!}$  in  $e_2 J = 0$  we obtain

$$(4.6) \quad (k_1 - k_2 + i + 1)\gamma J_{0, i, 0} + (\beta_2 + i\gamma) J_{0, i+1, 0} = 0.$$

This implies the theorem. □

Hence

$$(4.7) \quad J_{0, k_2, 0}(\alpha, \beta_1, \beta_2, \gamma) = (-1)^{k_2} J_{0, 0, 0}(\alpha, \beta_1, \beta_2, \gamma) \prod_{i=0}^{k_2-1} \frac{(k_1 - k_2 + i + 1)\gamma}{\beta_2 + i\gamma}.$$

Combining (4.4) and (4.7) we observe that formula (4.3) is equivalent to the formula

$$(4.8) \quad \begin{aligned} J_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma) &= \prod_{i=1}^{k_1-k_2} \frac{\Gamma(1 + \beta_1 + (i-1)\gamma)}{\Gamma(1 + \alpha + \beta_1 + (i+k_1-2)\gamma)} \\ &\times \prod_{i=1}^{k_2} \frac{\Gamma(1 + \beta_2 + (i-1)\gamma)}{\Gamma(1 + \beta_2 + (i+k_2-k_1-2)\gamma)} \frac{\Gamma(1 + \beta_1 + \beta_2 + (i-2)\gamma)}{\Gamma(1 + \alpha + \beta_1 + \beta_2 + (i+k_2-3)\gamma)} \\ &\times \prod_{i=1}^{k_2} \frac{\Gamma((i-k_1-1)\gamma) \Gamma(i\gamma)}{\Gamma(\gamma)} \prod_{i=1}^{k_1} \frac{\Gamma(\alpha + (i-1)\gamma) \Gamma(i\gamma)}{\Gamma(\gamma)}. \end{aligned}$$

Denote by  $R_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma)$  the right-hand side of (4.8).

To prove (4.8) we use the following observation. The weight subspace  $V_{\lambda_1}[\lambda_1 - k_1\sigma_1 - k_2\sigma_2] \subset V_{\lambda_1}$  is one-dimensional with a basis vector

$$v_{0,0,0} = \frac{f_1^{k_1-k_2} [f_1, f_2]^{k_2} v_1 \otimes v_2}{(k_1 - k_2)! (k_2)!}.$$

By [MaV, Theorem 5.1] the vector-valued function  $J_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma)v_{0,0,0}$  satisfies the dynamical difference equations introduced in [TV3],

$$(4.9) \quad J_{0,0,0}(\alpha, \beta_1 - 1, \beta_2, \gamma)v_{0,0,0} = J_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma) \mathbb{B}_1 v_{0,0,0},$$

$$(4.10) \quad J_{0,0,0}(\alpha, \beta_1, \beta_2 - 1, \gamma)v_{0,0,0} = J_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma) \mathbb{B}_2 v_{0,0,0},$$

where  $\mathbb{B}_1, \mathbb{B}_2$  are certain linear operators acting on  $V_{\lambda_1}$  and preserving the weight decomposition of  $V_{\lambda_1}$ , see formulas for these operators in the example in [MaV, Section 7.1] and in [MaV, Section 3.1], also see [TV3, Formula (8)].

Written explicitly equations (4.9), (4.10) give us the difference equations for the scalar function  $J_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma)$  with respect to the shift of the variables  $\beta_1 \rightarrow \beta_1 - 1$  and  $\beta_2 \rightarrow \beta_2 - 1$ ,

$$(4.11) \quad \begin{aligned} J_{0,0,0}(\alpha, \beta_1 - 1, \beta_2, \gamma) &= J_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma) \prod_{i=1}^{k_1-k_2} \frac{\alpha + \beta_1 + (i+k_1-2)\gamma}{\beta_1 + (i-1)\gamma} \\ &\times \prod_{i=1}^{k_2} \frac{\alpha + \beta_1 + \beta_2 + (i+k_2-3)\gamma}{\beta_1 + \beta_2 + (i-2)\gamma}, \end{aligned}$$

$$(4.12) \quad \begin{aligned} J_{0,0,0}(\alpha, \beta_1, \beta_2 - 1, \gamma) &= J_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma) \\ &\times \prod_{i=1}^{k_2} \frac{\beta_2 + (i+k_2-k_1-2)\gamma}{\beta_2 + (i-1)\gamma} \frac{\alpha + \beta_1 + \beta_2 + (i+k_2-3)\gamma}{\beta_1 + \beta_2 + (i-2)\gamma}. \end{aligned}$$

The difference equations for  $J_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma)$  are the same as the difference equations for the function  $R_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma)$  with respect to the shift of the variables  $\beta_1 \rightarrow \beta_1 - 1$  and  $\beta_2 \rightarrow \beta_2 - 1$ . Therefore, the functions  $J_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma)$  and  $R_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma)$  are proportional up to a periodic function of  $\beta_1, \beta_2$ . The periodic function can be fixed by comparing asymptotics as  $\text{Re } \beta_1 \rightarrow \infty, \text{Re } \beta_2 \rightarrow \infty$ . This finishes the proof in [TV4] of formulas (4.8) and (4.3).

5. THE  $A_2$ -TYPE SELBERG INTEGRALS OVER  $\mathbb{F}_p$ 

5.1. **Relations between  $\mathbb{F}_p$ -integrals.** For  $\mathbf{k} = (k_1, k_2)$ ,  $k_1 > k_2 > 0$  and integers  $0 < a, b_1, b_2, c < p$  define the master polynomial

$$(5.1) \quad \Phi_{\mathbf{k}}(t; s; a, b_1, b_2, c) = \prod_{i=1}^{k_1} t_i^a (1-t_i)^{b_1} \prod_{j=1}^{k_2} (1-s_j)^{b_2} \prod_{i=1}^{k_1} \prod_{j=1}^{k_2} (s_j - t_i)^{p-c} \\ \times \prod_{1 \leq i < i' \leq k_1} (t_i - t_{i'})^{2c} \prod_{1 \leq j < j' \leq k_2} (s_j - s_{j'})^{2c}$$

and the  $\mathbb{F}_p$ -integral

$$(5.2) \quad S_{\mathbf{k}}(a, b_1, b_2, c) = \int_{[\mathbf{k}]_p} \Phi_{\mathbf{k}}(t; s; a, b_1, b_2, c) dt ds,$$

where the  $p$ -cycle  $[\mathbf{k}]_p$  is defined in (2.5). This is the  $A_2$ -type  $\mathbb{F}_p$ -Selberg integral, see (3.10).

For an allowable triple  $(l_1, l_2, m)$  define the  $\mathbb{F}_p$ -integral

$$(5.3) \quad I_{l_1, l_2, m}(a, b_1, b_2, c) = \int_{[\mathbf{k}]_p} \Phi_{\mathbf{k}}(t; s; a, b_1, b_2, c) \frac{W_{l_1, l_2, m}(t; s)}{\prod_{i=1}^{k_1} t_i (1-t_i) \prod_{j=1}^{k_2} (1-s_j)} dt ds,$$

where  $W_{l_1, l_2, m}(t; s)$  is the weight function defined in Section 4.2.

Clearly we have

$$(5.4) \quad I_{0, k_2, 0}(a, b_1, b_2, c) = S_{\mathbf{k}}(a-1, b_1, b_2-1, c).$$

Denote

$$(5.5) \quad B_0(a, b_1, b_2, c) = (-1)^{k_2} \prod_{i=0}^{k_2-1} \frac{(k_1 - k_2 + i + 1)c}{b_2 + ic}, \\ B_1(a, b_1, b_2, c) = \prod_{i=1}^{k_1-k_2} \frac{a + b_1 + (i + k_1 - 2)c}{b_1 + (i-1)c} \prod_{i=1}^{k_2} \frac{a + b_1 + b_2 + (i + k_2 - 3)c}{b_1 + b_2 + (i-2)c}, \\ B_2(a, b_1, b_2, c) = \prod_{i=1}^{k_2} \frac{b_2 + (i + k_2 - k_1 - 2)c}{b_2 + (i-1)c} \frac{a + b_1 + b_2 + (i + k_2 - 3)c}{b_1 + b_2 + (i-2)c}.$$

**Theorem 5.1.** *Assume that  $k_1 < p$ .*

(i) *Assume that every factor in  $B_0$  in the numerator or denominator is a nonzero element of  $\mathbb{F}_p$ . Then*

$$(5.6) \quad I_{0, k_2, 0}(a, b_1, b_2, c) = B_0(a, b_1, b_2, c) I_{0, 0, 0}(a, b_1, b_2, c).$$

(ii) *Assume that every factor in  $B_1$  in the numerator or denominator is a nonzero element of  $\mathbb{F}_p$  and  $b_1 > 1$ , then*

$$(5.7) \quad I_{0, 0, 0}(a, b_1 - 1, b_2, c) = B_1(a, b_1, b_2, c) I_{0, 0, 0}(a, b_1, b_2, c).$$

(iii) *Assume that every factor in  $B_2$  in the numerator or denominator is a nonzero element of  $\mathbb{F}_p$  and  $b_2 > 1$ , then*

$$(5.8) \quad I_{0, 0, 0}(a, b_1, b_2 - 1, c) = B_2(a, b_1, b_2, c) I_{0, 0, 0}(a, b_1, b_2, c).$$

*Proof.* Equation (5.6) is an  $\mathbb{F}_p$ -analog of equation (4.7) and its proof is analogous to the proof of equation (4.7).

More precisely, consider the complex Lie algebra  $\mathfrak{sl}_3$  with standard generators  $f_1, f_2, e_1, e_2, h_1, h_2$ , simple roots  $\sigma_1, \sigma_2$ , fundamental weights  $\omega_1, \omega_2$ . Let  $V_{\lambda_1}, V_{\lambda_2}$  be the irreducible  $\mathfrak{sl}_3$ -modules with highest weights

$$\lambda_1 = -\frac{a}{c+p}\omega_1, \quad \lambda_2 = -\frac{b_1}{c+p}\omega_1 - \frac{b_2}{c+p}\omega_2$$

and highest weight vectors  $v_1, v_2$ . The module  $V_{\lambda_1}$  has a basis  $(f_1^{r_1}[f_1, f_2]^{r_2}v_1)$  labeled by nonnegative integers  $r_1, r_2$  and the module  $V_{\lambda_2}$  has a basis  $(f_1^{r_1}[f_1, f_2]^{r_2}f_2^{r_3}v_2)$  labeled by nonnegative integers  $r_1, r_2, r_3$ . For every generator of  $\mathfrak{sl}_3$  the matrix of its action on  $V_{\lambda_1}$  or on  $V_{\lambda_2}$  in these bases is a polynomial in  $-\frac{a}{c+p}, -\frac{b_1}{c+p}, -\frac{b_2}{c+p}$  with integer coefficients.

Consider the Lie algebra  $\mathfrak{sl}_3$  over the field  $\mathbb{F}_p$ . Let  $V_{\lambda_1}^{\mathbb{F}_p}$  be the vector space over  $\mathbb{F}_p$  with basis  $(f_1^{r_1}[f_1, f_2]^{r_2}v_1)$  labeled by nonnegative integers  $r_1, r_2$  and with the action of  $\mathfrak{sl}_3$  defined by the same formulas as on  $V_{\lambda_1}$  but reduced modulo  $p$ . Similarly we define the  $\mathfrak{sl}_3$ -module  $V_{\lambda_2}^{\mathbb{F}_p}$ .

Recall  $\mathbf{k} = (k_1, k_2)$ ,  $k_1 > k_2 > 0$ . Consider the weight subspace  $V_{\lambda_1}^{\mathbb{F}_p} \otimes V_{\lambda_2}^{\mathbb{F}_p}[\lambda_1 + \lambda_2 - k_1\sigma_1 - k_2\sigma_2]$  of the tensor product  $V_{\lambda_1}^{\mathbb{F}_p} \otimes V_{\lambda_2}^{\mathbb{F}_p}$ . This weight subspace has a basis formed by the vectors

$$v_{l_1, l_2, m} = \frac{f_1^{k_1 - k_2 - l_1 + l_2}[f_1, f_2]^{k_2 - l_2}v_1 \otimes f_1^{l_1 - m}[f_1, f_2]^m f_2^{l_2 - m}v_2}{(k_1 - k_2 - l_1 + l_2)! (k_2 - l_2)! (l_1 - m)! m! (l_2 - m)!}$$

labeled by allowable triples  $(l_1, l_2, m)$ .

**Lemma 5.2.** *The vector*

$$I = \sum_{l_1, l_2, m} (-1)^{l_1} I_{l_1, l_2, l_m}(a, b_1, b_2, c) v_{l_1, l_2, m}$$

*is a singular vector of  $V_{\lambda_1}^{\mathbb{F}_p} \otimes V_{\lambda_2}^{\mathbb{F}_p}$ , that is,  $e_1 I = 0, e_2 I = 0$ .*

*Proof.* Equations  $e_1 I = 0, e_2 I = 0$  are  $\mathbb{F}_p$ -analogues of equations  $e_1 J = 0, e_2 J = 0$  over  $\mathbb{C}$ .

For  $i = 1, 2$ , the vector  $e_i J$  is the integral of a certain differential  $k_1 + k_2$ -form  $\mu_i$ . It is shown in [SV2, Theorems 6.16.2], [M, Theorem 2.4] that  $\mu_i = d\nu_i$ , where  $\nu_i$  is some explicitly written differential  $k_1 + k_2 - 1$ -form. This implies  $e_i J = 0$  by Stokes' theorem.

The vector  $e_i I$  is the  $\mathbb{F}_p$ -integral of the same  $\mu_i$  reduced modulo  $p$ . It is explained in [SV2, Section 4] that the differential form  $\nu_i$  also can be reduced modulo  $p$  and this implies that the  $\mathbb{F}_p$ -integral  $e_i I$  is zero by Lemma 2.3. Cf. the proof of [SV2, Theorem 2.4].  $\square$

Lemma 5.2 implies the equations

$$(5.9) \quad (k_1 - k_2 + i + 1)c I_{0, i, 0} + (b_2 + ic) I_{0, i+1, 0} = 0$$

for  $i = 0, \dots, k_2 - 1$ , similarly to the proof of equations (4.6). The iterated application of equation (5.9) implies equation (5.6).

The proof of equations (5.7), (5.8) is parallel to the proof of equations (4.11), (4.12). We prove (5.7). The proof of (5.8) is similar.

Equation (4.11) follows from equation (4.9):

$$(5.10) \quad J_{0,0,0}(\alpha, \beta_1 - 1, \beta_2, \gamma)v_{0,0,0} - J_{0,0,0}(\alpha, \beta_1, \beta_2, \gamma)\mathbb{B}_1v_{0,0,0} = 0$$

and equation  $\mathbb{B}_1v_{0,0,0} = B_1v_{0,0,0}$  in  $V_{\lambda_1}$ . The explicit formulas for  $\mathbb{B}_1$  show that under the assumptions of Theorem 5.1 the action of  $\mathbb{B}_1$  on  $v_{0,0,0}$  is well-defined modulo  $p$  and gives the same result  $\mathbb{B}_1v_{0,0,0} = B_1v_{0,0,0}$  but in  $V_{\lambda_1}^{\mathbb{F}_p}$ .

The proof of (5.10) in [MaV] goes as follows. The left-hand side of (5.10) is a vector-valued integral of a suitable differential  $k_1 + k_2$ -form  $\mu$ . It is shown in [M, Theorem 5.1] that  $\mu = d\nu$ , where  $\nu$  is some explicitly written differential  $k_1 + k_2 - 1$ -form. This implies (5.10) by Stokes' theorem.

The  $p$ -analog of the left-hand side of (5.10) is the element

$$(5.11) \quad I_{0,0,0}(a, b_1 - 1, b_2, c)v_{0,0,0} - I_{0,0,0}(a, b_1, b_2, c)\mathbb{B}_1v_{0,0,0} \in V_{\lambda_1}^{\mathbb{F}_p}.$$

This element is the  $\mathbb{F}_p$ -integral of the same  $\mu$  reduced modulo  $p$ . It is explained in [SV2, Section 4] that the differential form  $\nu$  also can be reduced modulo  $p$  and this implies that the  $\mathbb{F}_p$ -integral in the left-hand side of (5.11) equals zero by Lemma 2.3. Hence equation (5.7) is proved and Theorem 5.1 is proved.  $\square$

5.2. **Theorem 3.3 for  $n = 2$ .** Recall the set of admissible parameters  $\mathcal{A}_{\mathbf{k}}$  introduced in Section 3.1 for  $\mathbf{k} = (k_1, k_2)$ ,  $k_1 > k_2 > 0$ .

**Lemma 5.3.** *Assume that  $(a, b_1 - 1, b_2, c), (a, b_1, b_2, c) \in \mathcal{A}_{\mathbf{k}}$ . Then*

$$(5.12) \quad S_{\mathbf{k}}(a, b_1 - 1, b_2, c) = S_{\mathbf{k}}(a, b_1, b_2, c) \prod_{i=1}^{k_1 - k_2} \frac{1 + a + b_1 + (i + k_1 - 2)c - p}{b_1 + (i - 1)c} \\ \times \prod_{i=1}^{k_2} \frac{2 + a + b_1 + b_2 + (i + k_1 - 3)c - p}{1 + b_1 + b_2 + (i - 2)c}.$$

Assume that  $(a, b_1, b_2 - 1, c), (a, b_1, b_2, c) \in \mathcal{A}_{\mathbf{k}}$ . Then

$$(5.13) \quad S_{\mathbf{k}}(a, b_1, b_2 - 1, c) = S_{\mathbf{k}}(a, b_1, b_2, c) \prod_{i=1}^{k_2} \frac{1 + b_2 + (i + k_2 - k_1 - 2)c}{b_2 + (i - 1)c} \\ \times \prod_{i=1}^{k_2} \frac{2 + a + b_1 + b_2 + (i + k_1 - 3)c - p}{1 + b_1 + b_2 + (i - 2)c}.$$

*Proof.* The lemma follows from formulas (5.4) and (5.6) and Theorem 5.1.  $\square$

For  $n = 2$  formula (3.1) takes the form:

$$(5.14) \quad R_{\mathbf{k}}(a, b_1, b_2, c) = (-1)^{k_1+k_2} \prod_{i=1}^{k_1-k_2} \frac{(b_1 + (i-1)c)!}{(1+a+b_1+(i+k_1-2)c-p)!} \\ \times \prod_{i=1}^{k_2} \frac{(b_2 + (i-1)c)!}{(1+b_2+(i+k_2-k_1-2)c)!} \frac{(1+b_1+b_2+(i-2)c)!}{(2+a+b_1+b_2+(i+k_1-3)c-p)!} \\ \times \prod_{i=1}^{k_1} (a+(i-1)c)! \prod_{i=1}^{k_2} (p+(i-k_1-1)c)! \prod_{r=1}^2 \prod_{i=1}^{k_r} \frac{(ic)!}{c!}.$$

**Lemma 5.4.** *Assume that  $(a, b_1 - 1, b_2, c), (a, b_1, b_2, c) \in \mathcal{A}_{\mathbf{k}}$ . Then*

$$(5.15) \quad R_{\mathbf{k}}(a, b_1 - 1, b_2, c) = R_{\mathbf{k}}(a, b_1, b_2, c) \prod_{i=1}^{k_1-k_2} \frac{1+a+b_1+(i+k_1-2)c-p}{b_1+(i-1)c} \\ \times \prod_{i=1}^{k_2} \frac{2+a+b_1+b_2+(i+k_1-3)c-p}{1+b_1+b_2+(i-2)c}.$$

*Assume that  $(a, b_1, b_2 - 1, c), (a, b_1, b_2, c) \in \mathcal{A}_{\mathbf{k}}(a, b_1, b_2, c)$ . Then*

$$(5.16) \quad R_{\mathbf{k}}(a, b_1, b_2 - 1, c) = R_{\mathbf{k}}(a, b_1, b_2, c) \prod_{i=1}^{k_2} \frac{1+b_2+(i+k_2-k_1-2)c}{b_2+(i-1)c} \\ \times \prod_{i=1}^{k_2} \frac{2+a+b_1+b_2+(i+k_1-3)c-p}{1+b_1+b_2+(i-2)c}.$$

□

By Lemmas 5.3 and 5.4 the functions  $S_{\mathbf{k}}(a, b_1, b_2, c)$  and  $R_{\mathbf{k}}(a, b_1, b_2, c)$  defined on  $\mathcal{A}_{\mathbf{k}}$  satisfy the same difference equations with respects to the shifts of variables  $b_1 \rightarrow b_1 - 1$  and  $b_2 \rightarrow b_2 - 1$ .

**Lemma 5.5.** *Assume that  $a, c$  are positive integers such that  $0 < k_1 c \leq p - 1$ ,  $a + (k_1 - 1)c < p - 1$ . Then the point*

$$(5.17) \quad (a, b_1, b_2, c) = (a, p - 1 - (a + (k_1 - 1)c), (k_1 - k_2 + 1)c - 1, c)$$

*lies in  $\mathcal{A}_{\mathbf{k}}$ .*

*Proof.* If  $(a, b_1, b_2, c)$  is given by (5.17), then

$$R_{\mathbf{k}} = (-1)^{k_1+k_2} \prod_{i=1}^{k_1-k_2} \frac{(p-1-(a+(k_1-i)c))!}{((i-1)c)!} \\ \times \prod_{i=1}^{k_2} \frac{((k_1-k_2+i)c-1)!}{((i-1)c)!} \frac{(p-1-(a+(k_2-i)c))!}{((k_1-k_2+i-1)c)!} \\ \times \prod_{i=1}^{k_1} (a+(i-1)c)! \prod_{i=1}^{k_2} (p+(i-k_1-1)c)! \prod_{r=1}^2 \prod_{i=1}^{k_r} \frac{(ic)!}{c!}.$$

This proves the lemma.  $\square$

**Lemma 5.6.** *Assume that  $\tilde{a}, \tilde{c}$  are nonnegative integers such that  $0 < k_1 \tilde{c} \leq p-1$ ,  $\tilde{a} + (k_1 - 1)\tilde{c} \leq p-1$ . Denote by  $\mathcal{A}_{\mathbf{k}}(\tilde{a}, \tilde{c})$  the set of all  $(a, b_1, b_2, c) \in \mathcal{A}_{\mathbf{k}}$  such that  $a = \tilde{a}$ ,  $c = \tilde{c}$ . Then  $\mathcal{A}_{\mathbf{k}}(\tilde{a}, \tilde{c})$  consists of the pairs  $(b_1, b_2)$  of nonnegative integers satisfying the inequalities*

$$(5.18) \quad p-1 - (\tilde{a} + (k_1 - 1)\tilde{c}) \leq b_1, \quad (k_1 - k_2 + 1)\tilde{c} - 1 \leq b_2$$

and some other inequalities of the form

$$(5.19) \quad b_1 \leq A_1, \quad b_2 \leq A_2 \quad b_1 + b_2 \leq A_{12},$$

where  $A_1, A_2, A_{12}$  are some integers such that

$$\begin{aligned} A_1 &\geq p-1 - (\tilde{a} + (k_1 - 1)\tilde{c}), & A_2 &\geq (k_1 - k_2 + 1)\tilde{c} - 1, \\ A_{12} &\geq p-1 - (\tilde{a} + (k_1 - 1)\tilde{c}) + (k_1 - k_2 + 1)\tilde{c} - 1. \end{aligned}$$

*Proof.* The lemma follows from Lemmas 3.1 and 3.2.  $\square$

**Corollary 5.7.** *Any point  $(\tilde{a}, b_1, b_2, \tilde{c}) \in \mathcal{A}_{\mathbf{k}}(\tilde{a}, \tilde{c})$  can be connected with the point  $(\tilde{a}, p-1 - (\tilde{a} + (k_1 - 1)\tilde{c}), (k_1 - k_2 + 1)\tilde{c} - 1, \tilde{c}) \in \mathcal{A}_{\mathbf{k}}(\tilde{a}, \tilde{c})$  by a piece-wise linear path in  $\mathcal{A}_{\mathbf{k}}(\tilde{a}, \tilde{c})$  consisting of the vectors  $(0, -1, 0, 0)$  or  $(0, 0, -1, 0)$ .  $\square$*

*Proof of Theorem 3.3 for  $n = 2$ .* For  $n = 1$ ,  $\mathbf{k} = (k_1)$  and the point  $(\tilde{a}, p-1 - (\tilde{a} + (k_1 - 1)\tilde{c}), \tilde{c})$  formula (3.11) holds by [RV, Theorem 4.1].

For  $n = 2$ ,  $\mathbf{k} = (k_1, k_2)$  and the point  $(\tilde{a}, p-1 - (\tilde{a} + (k_1 - 1)\tilde{c}), (k_1 - k_2 + 1)\tilde{c} - 1, \tilde{c})$  formula (3.11) holds by Lemma 5.5 and Theorem 3.6.

For  $n = 2$ ,  $\mathbf{k} = (k_1, k_2)$  and arbitrary  $(\tilde{a}, b_1, b_2, \tilde{c}) \in \mathcal{A}_{\mathbf{k}}(\tilde{a}, \tilde{c})$  formula (3.11) holds by Lemmas 5.3, 5.4 and Corollary 5.7. Theorem 3.3 for  $n = 2$  is proved.  $\square$

**5.3. Evaluation of  $I_{0,0,0}(a, b_1, b_2, c)$ .** In this section we evaluate  $I_{0,0,0}(a, b_1, b_2, c)$  without using the evaluation of  $S_{\mathbf{k}}(a, b_1, b_2, c)$ .

**Theorem 5.8.** *Let  $\mathbf{k} = (k_1, k_2)$ ,  $k_1 > k_2 > 0$  and  $a, b_1, b_2, c \in \mathbb{Z}_{>0}$ . Assume that  $a + (k_1 - 1)c < p$  and all factorials in the right-hand side of the next formula are factorials of the nonnegative integers less than  $p$ . Then*

$$(5.20) \quad \begin{aligned} I_{0,0,0}(a, b_1, b_2, c) &= (-1)^{k_1+k_2} \prod_{i=1}^{k_1-k_2} \frac{(b_1 + (i-1)c)!}{(a + b_1 + (i+k_1-2)c - p)!} \\ &\times \prod_{i=1}^{k_2} \frac{(b_2 + (i-1)c)!}{(b_2 + (i+k_2-k_1-2)c)!} \frac{(b_1 + b_2 + (i-2)c)!}{(a + b_1 + b_2 + (i+k_1-3)c - p)!} \\ &\times \prod_{i=1}^{k_1} (a + (i-1)c - 1)! \prod_{i=1}^{k_2} (p + (i-k_1-1)c - 1)! \prod_{r=1}^2 \prod_{i=1}^{k_i} \frac{(ic)!}{c!}. \end{aligned}$$

*Proof.* The proof is parallel to the proof of Theorem 3.3 for  $n = 2$ .

Denote by  $\mathcal{A}_{\mathbf{k}}^I$  the set of all  $a, b_1, b_2, c \in \mathbb{Z}_{>0}$  satisfying the assumptions of Theorem 5.8. Notice that if  $(a, b_1, b_2, c) \in \mathcal{A}_{\mathbf{k}}^I$ , then

$$(5.21) \quad b_1 \geq p - (a + (k_1 - 1)c), \quad b_2 \geq (k_1 - k_2 + 1)c.$$

**Lemma 5.9.** *Formula (5.20) holds if  $b_2 = (k_1 - k_2 + 1)c$  and  $(a, b_1, (k_1 - k_2 + 1)c, c) \in \mathcal{A}_k^I$ .*

*Proof.* If  $b_2 = (k_1 - k_2 + 1)c$ , then all variables  $(s_j)$  in the integrand of  $I_{0,0,0}(a, b_1, (k_1 - k_2 + 1)c, c)$  are used to reach the monomial  $\prod_{j=1}^{k_1} s_j^{k_1 p - 1}$  in the calculation of the  $\mathbb{F}_p$ -integral  $I_{0,0,0}(a, b_1, (k_1 - k_2 + 1)c, c)$ . The remaining free variables  $(t_i)$  all belong to the factor

$$\Phi_{(k_1)}(t_1, \dots, t_{k_1}, a - 1, b_1, c) = \prod_{1 \leq i < i' \leq k_1} (t_i - t_{i'})^{2c} \prod_{i=1}^{k_1} t_i^{a-1} (1 - t_i)^{b_1}$$

of the integrand and are used to calculate the coefficient of the monomial  $\prod_{j=1}^{k_1} t_j^{p-1}$ .

More precisely, under the assumptions of the theorem we have

$$I_{0,0,0}(a, b_1, (k_1 - k_2 + 1)c, c) = (-1)^{b_2 k_2 + c k_2 (k_2 - 1)/2} \frac{(k_2 c)!}{(c!)^{k_2}} S_{(k_1)}(a - 1, b_1, c),$$

cf. the proof of Theorem 3.6. We have  $(-1)^{b_2 k_2 + c k_2 (k_2 - 1)/2} = (-1)^{(k_1 k_2 - k_2 (k_2 + 1)/2)c - k_2}$ .

By [RV, Theorem 4.1] we have  $S_{(k_1)}(a - 1, b_1, c) = R_{(k_1)}(a - 1, b_1, c)$ . Hence

$$(5.22) \quad \begin{aligned} I_{0,0,0}(a, b_1, (k_1 - k_2 + 1)c, c) &= (-1)^{(k_1 k_2 - k_2 (k_2 + 1)/2)c} (-1)^{k_1 + k_2} \\ &\quad \times \frac{(k_2 c)!}{(c!)^{k_2}} \prod_{j=1}^{k_1} \frac{(j c)!}{c!} \frac{(a + (j - 1)c - 1)! (b + (j - 1)c)!}{(a + b + (k_1 + j - 2)c - p)!}. \end{aligned}$$

Denote by  $R_{\mathbf{k}}^I(a, b_1, b_2, c)$  the right-hand side in (5.20). We have

$$(5.23) \quad \begin{aligned} R_{\mathbf{k}}^I(a, b_1, (k_1 - k_2 + 1)c, c) &= (-1)^{k_1 + k_2} \prod_{i=1}^{k_1 - k_2} \frac{(b_1 + (i - 1)c)!}{(a + b_1 + (i + k_1 - 2)c - p)!} \\ &\quad \times \prod_{i=1}^{k_2} \frac{((k_1 - k_2 + 1)c + (i - 1)c)!}{((k_1 - k_2 + 1)c + (i + k_2 - k_1 - 2)c)!} \\ &\quad \times \prod_{i=1}^{k_2} \frac{(b_1 + (k_1 - k_2 + 1)c + (i - 2)c)!}{(a + b_1 + (k_1 - k_2 + 1)c + (i + k_1 - 3)c - p)!} \\ &\quad \times \prod_{i=1}^{k_1} (a + (i - 1)c - 1)! \prod_{i=1}^{k_2} (p + (i - k_1 - 1)c - 1)! \prod_{r=1}^2 \prod_{i=1}^{k_i} \frac{(i c)!}{c!}. \\ &= (-1)^{k_1 + k_2} (-1)^{(k_1 k_2 - k_2 (k_2 + 1)/2)c} \frac{(k_2 c)!}{(c!)^{k_2}} \prod_{j=1}^{k_1} \frac{(j c)!}{c!} \frac{(a + (j - 1)c - 1)! (b + (j - 1)c)!}{(a + b + (k_1 + j - 2)c - p)!}, \end{aligned}$$

where we used the cancellation Lemma 2.1 in the last step. Hence

$I_{0,0,0}(a, b_1, (k_1 - k_2 + 1)c, c) = R_{\mathbf{k}}^I(a, b_1, (k_1 - k_2 + 1)c, c)$  and Lemma 5.9 is proved.  $\square$

Comparing equations (5.7), (5.8) and the formula for  $R_{\mathbf{k}}^I(a, b_1, b_2, c)$ , we conclude that the functions  $I_{0,0,0}(a, b_1, b_2, c)$  and  $R_{\mathbf{k}}^I(a, b_1, b_2, c)$  on  $\mathcal{A}_k^I$  satisfy the same difference equations with respect to the shifts of variables  $b_1 \rightarrow b_1 - 1$  and  $b_2 \rightarrow b_2 - 1$  and are equal if  $b_2$

takes its minimal value  $(k_1 - k_2 + 1)c$ . This implies Theorem 5.8, cf. Lemmas 5.5, 5.6 and Corollary 5.7.  $\square$

**5.4. Proof of Theorem 3.3 for arbitrary  $n > 2$ .** The proof is parallel to the proof of Theorem 3.3 for  $n = 2$ .

Analogously to the proof of Theorem 5.1, consider the Lie algebra  $\mathfrak{sl}_{n+1}$  and its representations  $V_{\lambda_1}^{\mathbb{F}_p}$  and  $V_{\lambda_2}^{\mathbb{F}_p}$  over  $\mathbb{F}_p$  with highest weights

$$\lambda_1 = -\frac{a}{c+p}\omega_1, \quad \lambda_2 = -\frac{b_1}{c+p}\omega_1 - \dots - \frac{b_n}{c+p}\omega_n$$

and highest weight vectors  $v_1, v_2$ . Consider the PBW basis  $\mathcal{B} = (u)$  of the weight subspace  $V_{\lambda_1}^{\mathbb{F}_p} \otimes V_{\lambda_2}^{\mathbb{F}_p}[\lambda_1 + \lambda_2 - \sum_{i=1}^n k_i \sigma_i]$  like in the proof of Theorem 5.1. We distinguish two elements of that basis:

$$u_1 = \frac{f_1^{k_1-k_2}[f_1, f_2]^{k_2-k_3} \dots [f_1, [f_2, \dots, [f_{n-1}, f_n] \dots]]^{k_n} v_1 \otimes v_2}{(k_1 - k_2)! (k_2 - k_3)! \dots (k_n)!},$$

$$u_2 = \frac{f_1^{k_1} v_1 \otimes f_2^{k_2} \dots f_n^{k_n} v_2}{(k_1)! (k_2)! \dots (k_n)!}.$$

For  $n = 2$  these vectors are the vectors  $v_{0,0,0}$  and  $v_{0,k_2,0}$  in the proof of Theorem 5.1.

To any basis vector  $u \in \mathcal{B}$  we assign the weight function  $W_u(t)$  defined in [RSV, Section 6.1], here  $t$  is the collection of variables defined in (3.9). Then we consider the  $\mathbb{F}_p$ -integrals

$$I_u(a, b, c) = \int_{[\mathbf{k}]_p} \Phi(t, a, b, c) W_u(t) dt.$$

It follows from the formulas for the weight functions that

$$I_{u_2}(a, b, c) = S_{\mathbf{k}}(a - 1, b_1, b_2 - 1, \dots, b_n - 1, c),$$

cf. (5.4). It is known from the theory of KZ equations that the vector

$$I(a, b, c) = \sum_{u \in \mathcal{B}} I_u(a, b, c) u$$

is a singular vector in  $V_{\lambda_1}^{\mathbb{F}_p} \otimes V_{\lambda_2}^{\mathbb{F}_p}[\lambda_1 + \lambda_2 - \sum_{i=1}^n k_i \sigma_i]$ . From the singular vector condition it follows that

$$(5.24) \quad I_{u_2}(a, b, c) = B_0(a, b, c) I_{u_1}(a, b, c),$$

where  $B_0(a, b, c)$  is an explicit expression like in (5.6).

Then consider the vector  $I_{u_1}(a, b, c)u_1$  of the one-dimensional weight subspace  $V_{\lambda_1}^{\mathbb{F}_p}[\lambda_1 + \lambda_2 - \sum_{i=1}^n k_i \sigma_i]$ . That vector satisfies the dynamical equations defined in [TV3]. The dynamical equations take the form

$$(5.25) \quad I_{u_1}(a, b_1, \dots, b_i - 1, \dots, b_n, c) = B_i(a, b, c) I_{u_1}(a, b, c), \quad i = 1, \dots, n,$$

where  $B_i(a, b, c)$  are explicit products like in (5.7) and (5.8).

Equation (5.24) and difference equations (5.25) imply that the two functions  $S_{\mathbf{k}}(a, b, c)$  and  $R_{\mathbf{k}}(a, b, c)$ , defined on  $\mathcal{A}_{\mathbf{k}}$ , satisfy the same difference equations with respect to the shift

of variables  $b_i \rightarrow b_i - 1$  for  $i = 1, \dots, n$ . By Corollary 3.7 we also know that the two functions are equal at the distinguished point

$$(a, p - 1 - (a + (k_1 - 1)c), (k_1 - k_2 + 1)c - 1, \dots, (k_{n-1} - k_n + 1)c - 1, c) \in \mathcal{A}_k.$$

This implies that the two functions are equal (cf. Corollary 5.7) and Theorem 3.3 is proved for any  $n$ . The details of the proof will be published elsewhere.

## REFERENCES

- [AAR] G.E. Andrews, R. Askey, R. Roy, *Special functions*, Cambridge University Press, 1999
- [An] Greg W. Anderson, *The evaluation of Selberg sums*, C. R. Acad. Sci. Paris Sér. I Math. **311** (1990), no. 8, 469–472
- [Ao] K. Aomoto, *Jacobi polynomials associated with Selberg integral*, Siam J. Math, **18** (1987), No. 2, 545–549
- [As] R. Askey, *Some basic hypergeometric extensions of integrals of Selberg and Andrews*, SIAM J. Math. Anal. **11** (1980), 938–951
- [Ch] I. Cherednik, *From double Hecke algebra to analysis*, Doc.Math.J.DMV, Extra Volume ICM 1998,II, 527–531
- [EFK] P. Etingof, I. Frenkel, A. Kirillov, *Lectures on representation theory and Knizhnik-Zamolodchikov equations*, Mathematical Surveys and Monographs, 58, AMS, Providence, RI, 1998. xiv+198 pp. ISBN: 0-8218-0496-0
- [Ev] R.J. Evans, *The evaluation of Selberg character sums*, L'Enseign. Math. **37** (1991), 235–248
- [FSV] G. Felder, L. Stevens, and A. Varchenko, *Elliptic Selberg integrals and conformal blocks*, Math. Res. Lett. 10 (2003), no. 5–6, 671–684
- [FV] G. Felder and A. Varchenko, *Integral representation of solutions of the elliptic Knizhnik-Zamolodchikov-Bernard equations*, Int. Math. Res. Notices, 1995, N. 5, 221–233
- [FW] P. J. Forrester and S. O. Warnaar, *The importance of the Selberg integral*, Bull. Amer. Math. Soc. (N.S.) **45** (2008), 489–534
- [Ha] L. Habsieger, *Une  $q$ -intégrale de Selberg et Askey*, SIAM J. Math. Anal. **19** (1988), 1475–1489
- [Ka] J. Kaneko,  *$q$ -Selberg integrals and Macdonald polynomials*, Ann. Sci. Ecole Norm. Sup. **29** (1996), 583–637
- [KZ] V. Knizhnik and A. Zamolodchikov, *Current algebra and the Wess-Zumino model in two dimensions*, Nucl. Phys. **B247** (1984), 83–103
- [M] A. Matsuo, *An application of Aomoto-Gelfand Hypergeometric Functions to the  $SU(n)$  Knizhnik-Zamolodchikov Equation*, Comm. Math. Phys. **134** (1990), 65–77 (1990).
- [MaV] Y. Markov and A. Varchenko, *Hypergeometric Solutions of Trigonometric KZ Equations satisfy Dynamical Difference Equations*, Adv. Math. **166** (2002), no. 1, 100–147
- [MuV] E. Mukhin and A. Varchenko, *Remarks on critical points of phase functions and norms of Bethe vectors*, Adv. Stud. Pure Math. **27** (2000), 239–246

- [Op] E.M. Opdam, *Some applications of hypergeometric shift operators*, Invent. Math. **98** (1989), 1–18
- [R] E. Rains, *Multivariate Quadratic Transformations and the Interpolation Kernel*, Contribution to the Special Issue on Elliptic Hypergeometric Functions and Their Applications, SIGMA **14** (2018), 019, 69 pages, <https://doi.org/10.3842/SIGMA.2018.019>
- [RSV] R. Rimányi, L. Stevens, A. Varchenko, *Combinatorics of rational functions and Poincare-Birchoff-Witt expansions of the canonical  $U(\mathfrak{n}_-)$ -valued differential form*, Ann. Comb. **9** (2005), no. 1, 57–74
- [RTVZ] R. Rimányi, V. Tarasov, A. Varchenko, P. Zinn-Justin, *Extended Joseph polynomials, quantized conformal blocks, and a  $q$ -Selberg type integral*, Journal of Geometry and Physics, **62** (2012), pp. 2188–2207
- [RV] R. Rimányi, A. Varchenko, *The  $\mathbb{F}_p$ -Selberg integral*, arXiv:2011.14248, 1–19
- [Se] A. Selberg, *Bemerkninger om et multipelt integral*, Norsk Mat. Tidsskr., **26** (1944), 71–78
- [SIV] A. Slinkin, A. Varchenko *Hypergeometric Integrals Modulo  $p$  and Hasse–Witt Matrices*, arXiv:2001.06869, 1–36
- [Sp] V. Spiridonov, *On the elliptic beta function*, (Russian) Uspekhi Mat. Nauk **56** (2001), no. 1, 181–182; translation in Russian Math. Surveys **56** (2001), no. 1, 185–186
- [SV1] V. Schechtman, A. Varchenko, *Arrangements of Hyperplanes and Lie Algebra Homology*, Invent. Math. **106** (1991), 139–194
- [SV2] V. Schechtman, A. Varchenko, *Solutions of KZ differential equations modulo  $p$* , The Ramanujan Journal, **48** (3), 2019, 655–683, <https://doi.org/10.1007/s11139-018-0068-x>
- [TV1] V. Tarasov and A. Varchenko, *Geometry of  $q$ -hypergeometric functions as a bridge between Yangians and quantum affine algebras*, Invent. Math. **128** (1997), 501–588
- [TV2] V. Tarasov and A. Varchenko, *Geometry of  $q$ -hypergeometric functions, quantum affine algebras and elliptic quantum groups*, Asterisque, **246** (1997), 1–135
- [TV3] V. Tarasov and A. Varchenko, *Difference Equations Compatible with Trigonometric KZ Differential Equations*, IMRN, **15** (2000), 801–829
- [TV4] V. Tarasov and A. Varchenko, *Selberg-type integrals associated with  $\mathfrak{sl}_3$* , Lett. Math. Phys. **65** (2003), 173–185
- [V1] A. Varchenko, *The Euler beta-function, the Vandermonde determinant, the Legendre equation, and critical values of linear functions on a configuration of hyperplanes*, I. (Russian) Izv. Akad. Nauk SSSR Ser. Mat. **53** (1989), no. **6**, 1206–1235, 1337; translation in Math. USSR-Izv. **35** (1990), no. 3, 543–571
- [V2] A. Varchenko, *Special functions, KZ type equations, and Representation theory*, CBMS, Regional Conference Series in Math., n. 98, AMS (2003)
- [V3] A. Varchenko, *Remarks on the Gaudin model modulo  $p$* , arXiv:1708.06264, Journal of Singularities, **18** (2018), 486–499
- [V4] A. Varchenko, *Solutions modulo  $p$  of Gauss-Manin differential equations for multidimensional hypergeometric integrals and associated Bethe ansatz*, arXiv:1709.06189, Mathematics 2017, **5**(4), 52; doi:10.3390/math5040052, 1–18

- [V5] A. Varchenko, *Hyperelliptic integrals modulo  $p$  and Cartier-Manin matrices*, [arXiv:1806.03289](#), 1–16
- [V6] A. Varchenko, *An invariant subbundle of the KZ connection mod  $p$  and reducibility of  $\widehat{\mathfrak{sl}}_2$  Verma modules mod  $p$* , [arXiv:2002.05834](#), 1–14
- [V7] A. Varchenko, *Determinant of  $\mathbb{F}_p$ -hypergeometric solutions under ample reduction*, [arXiv:2010.11275](#), 1–22
- [Wa1] S. Ole Warnaar, *A Selberg integral for the Lie algebra  $A_n$* , *Acta Math.* **203** (2009), no. 2, 269–304
- [Wa2] S. Ole Warnaar, *The  $\mathfrak{sl}_3$  Selberg integral*, *Adv. Math.* **224** (2010), no. 2, 499–524