

On Line Sum Optimization

Shmuel Onn *

Abstract

We show that the *column sum optimization problem*, of finding a $(0, 1)$ -matrix with prescribed row sums which minimizes the sum of evaluations of given functions at its column sums, can be solved in polynomial time, either when all functions are the same or when all row sums are bounded by any constant. We conjecture that the more general *line sum optimization problem*, of finding a matrix minimizing the sum of given functions evaluated at its row sums and column sums, can also be solved in polynomial time.

Keywords: majorization, column sum, row sum, matrix, degree sequence, graph

MSC: 05A, 15A, 51M, 52A, 52B, 52C, 62H, 68Q, 68R, 68U, 68W, 90B, 90C

1 Introduction

For a positive integer m let $[m] := \{1, 2, \dots, m\}$. For an $m \times n$ matrix A let $r_i(A) := \sum_{j=1}^n A_{i,j}$ for $i \in [m]$ and $c_j(A) := \sum_{i=1}^m A_{i,j}$ for $j \in [n]$ be its row sums and column sums.

We consider here the following algorithmic problem.

Column Sum Optimization. Given positive integers m, n and $r_1, \dots, r_m \leq n$, and functions $f_j : \{0, 1, \dots, m\} \rightarrow \mathbb{Z}$ for $j = 1, \dots, n$, find an $m \times n$ matrix A , where each entry is equal to 0 or 1, and with row sums $r_i(A) = r_i$ for $i = 1, \dots, m$, which minimizes $\sum_{j=1}^n f_j(c_j(A))$.

For instance, for $m = n = 4$, row sum tuple $(r_1, r_2, r_3, r_4) = (3, 3, 2, 2)$, and identical functions $f_1 = f_2 = f_3 = f_4 = f$ with $f(x) = (x - 1)^2(x - 3)^2$, an optimal solution is the following matrix, with column sum tuple $(c_1, c_2, c_3, c_4) = (3, 3, 3, 1)$ and objective value 0,

$$A = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix}.$$

We use a characterization of $(0, 1)$ -matrices by Ryser [7] and prove the following theorem.

Theorem 1.1 *The column sum optimization problem can be solved in polynomial time when:*

1. *either all functions are the same, that is, $f_1 = \dots = f_n = f$ for some f ;*
2. *or all row sums are bounded by some fixed integer b , that is, $r_1, \dots, r_m \leq b$.*

Note that even when the row sums are bounded, the number of possible $(0, 1)$ -matrices can be $\binom{n}{b}^m$ which is exponential in the data, hence exhaustive search is not polynomial.

The column sum optimization problem is a special case of the following more general *line sum optimization problem*: given m, n and $e \leq mn$, and functions $g_i : \{0, 1, \dots, n\} \rightarrow \mathbb{Z}$ for

*Technion - Israel Institute of Technology. Email: onn@ie.technion.ac.il

$i \in [m]$ and $f_j : \{0, 1, \dots, m\} \rightarrow \mathbb{Z}$ for $j \in [n]$, find $A \in \{0, 1\}^{m \times n}$ with e nonzero entries minimizing $\sum_{i=1}^m g_i(r_i(A)) + \sum_{j=1}^n f_j(c_j(A))$. Indeed, the column sum problem reduces to the line sum problem by setting $e := \sum_{i=1}^m r_i$ and $g_i(x) := a(x - r_i)^2$ for $i \in [m]$ and sufficiently large positive integer a . However, we do not know what is the complexity of this more general problem, and not even that of the column sum problem for arbitrary row sums and functions.

The line sum problem is in turn a special case of the following *degree sequence optimization problem*: given a graph $H = (V, E)$, integer $e \leq |E|$, and functions $f_v : \{0, 1, \dots, d_v(H)\} \rightarrow \mathbb{Z}$ for $v \in V$, where $d_v(H)$ is the degree of v in H , find a subgraph $G = (V, F) \subseteq H$ with e edges minimizing $\sum_{v \in V} f_v(d_v(G))$. Indeed, identifying matrices $A \in \{0, 1\}^{m \times n}$ with graphs $G = (V, F)$ where $V = \{u_1, \dots, u_m\} \uplus \{w_1, \dots, w_n\}$ and $F = \{\{u_i, w_j\} : A_{i,j} = 1\}$, the line sum problem reduces to the degree sequence problem with $H = K_{m,n}$ the complete bipartite graph. In the case of $H = K_n$ the complete graph and all functions the same, $f_v = f$ for all $v \in V$, the problem was recently shown in [3] to be polynomial time solvable, using the characterization of degree sequences by Erdős and Gallai [5]. For general graphs H , the problem was shown in [1] to be NP-hard already when $f_v(x) = -x^2$ for all $v \in V$, but is polynomial time solvable if the functions are convex [1, 4]. If m or n are fixed, the problem for any $H \subseteq K_{m,n}$, and hence in particular the line sum problem, for any functions, is polynomial time solvable [4]. We conjecture that the line sum problem, which is the degree sequence problem over $H = K_{m,n}$, as well as the degree sequence problem over $H = K_n$, are polynomial time solvable for arbitrary functions at the vertices, not necessarily identical.

2 Proof

Let the positive integers m, n and $r_1, \dots, r_m \leq n$ be given. Define the tuple $r := (r_1, \dots, r_m)$ and its *conjugate* tuple $d := (d_1, \dots, d_n)$ where $d_j := |\{i : r_i \geq j\}|$ for $j \in [n]$. Note that $\sum_{j=1}^n d_j = \sum_{i=1}^m r_i \leq mn$ and d is nonincreasing, that is, $d_1 \geq \dots \geq d_n$. A nonincreasing tuple $c = (c_1, \dots, c_n)$ is *majorized* by d if $\sum_{j=1}^k c_j \leq \sum_{j=1}^k d_j$ for $k \in [n]$ and $\sum_{j=1}^n c_j = \sum_{j=1}^n d_j$. (See [6] for more details on the theory of majorization and its applications.)

We use a characterization by Ryser [7] of $(0, 1)$ -matrices with prescribed line sums.

Proposition 2.1 [7] *There is a matrix $A \in \{0, 1\}^{m \times n}$ with row sum and column sum tuples r, c if and only if, permuting c to be nonincreasing, it is majorized by the conjugate tuple d .*

For instance, if $m = n = 4$ and $r = (3, 3, 2, 2)$, then $d = (4, 4, 2, 0)$, and $c = (3, 3, 3, 1)$ is majorized by d so there exists $A \in \{0, 1\}^{4 \times 4}$ with row sum tuple r and column sum tuple c .

Next we note that a matrix with given line sums r and c can be efficiently obtained.

Lemma 2.2 *Let $r = (r_1, \dots, r_m)$ and let $c = (c_1, \dots, c_n)$ be such that, permuting c to be nonincreasing, it is majorized by the tuple $d = (d_1, \dots, d_n)$ conjugate to r . Then a matrix A in $\{0, 1\}^{m \times n}$ with row sum tuple r and column sum tuple c is computable in polynomial time.*

Proof. The problem is solvable either by the efficient simple Gale-Ryser algorithm, see [2, Chapter 3], or by network flows as follows. Define a directed graph with capacities on the edges as follows. There are vertices s, t, u_1, \dots, u_m , and w_1, \dots, w_n . There are edges $[s, u_i]$ for $i \in [m]$ with capacity r_i , edges $[u_i, w_j]$ for $i \in [m]$ and $j \in [n]$ with capacity 1, and edges $[w_j, t]$ for $j \in [n]$ with capacity c_j . Then, as is well known, a maximum nonnegative integer flow from s to t can be computed in polynomial time, see e.g. [8]. Then A is read off from the maximum flow by taking $A_{i,j}$ to be the flow on edge $[u_i, w_j]$ for all $i \in [m]$ and $j \in [n]$. \square

We are now in position to prove our theorem.

Part 1. Since all functions at the column sums are now the same, that is, $f_1 = \dots = f_n = f$ for some f , the objective value $\sum_{j=1}^n f(c_j(A))$ is invariant under column permutations, and so without loss of generality we may and do search for a matrix A whose column sum tuple c is nonincreasing. (This is not true, however, for general objective $\sum_{j=1}^n f_j(c_j(A))$ with distinct f_j , and this is precisely the reason we do not know how to solve the general problem.)

We now reduce the column sum problem to the problem of finding a shortest directed path in a directed graph D where each edge has a length. We construct D as follows.

There are two special vertices u and v . In addition, there are vertices labeled by triplets of integers (k, c_k, s_k) where $1 \leq k \leq n$, $0 \leq c_k \leq d_1 \leq m$, and $0 \leq s_k \leq \sum_{j=1}^n d_j \leq mn$.

There are edges $[u, (1, c_1, s_1)]$ for $0 \leq c_1 \leq d_1$ and $s_1 = c_1$ of length $f(c_1)$. For $2 \leq k \leq n$ there are edges $[(k-1, c_{k-1}, s_{k-1}), (k, c_k, s_k)]$ for $c_k \leq c_{k-1}$, $s_k = s_{k-1} + c_k$, and $s_k \leq \sum_{j=1}^k d_j$, of length $f(c_k)$. Finally, there are edges $[(n, c_n, s_n), v]$ for $s_n = \sum_{j=1}^n d_j$ of length 0.

Consider any directed path from u to v in D , which by the definition of D looks like

$$u \longrightarrow (1, c_1, s_1) \longrightarrow \dots \longrightarrow (n, c_n, s_n) \longrightarrow v .$$

Intuitively, traveling on the path, being at vertex (k, c_k, s_k) indicates that the prefix (c_1, \dots, c_k) of the sequence had been determined, and it is nonincreasing and sums to s_k . Formally, by definition of the edges in D , we have $c_1 \geq \dots \geq c_n$ hence the corresponding tuple $c = (c_1, \dots, c_n)$ is nonincreasing. We also have that $\sum_{j=1}^k c_j = s_k \leq \sum_{j=1}^k d_j$ for $k \in [n]$ and $\sum_{j=1}^n c_j = s_n = \sum_{j=1}^n d_j$, and therefore c is majorized by the conjugate tuple d . So by Proposition 2.1, there is a matrix $A \in \{0, 1\}^{m \times n}$ with row sum tuple r and column sum tuple c . By the definition of the lengths, the length of this directed path is $\sum_{j=1}^n f(c_j) = \sum_{j=1}^n f(c_j(A))$.

Conversely, for any matrix $A \in \{0, 1\}^{m \times n}$ with row sum tuple r and nonincreasing column sum tuple $c = (c_1, \dots, c_n)$, the tuple c is majorized by $d = (d_1, \dots, d_n)$ by Proposition 2.1, hence the following is a directed path from u to v in D of length $\sum_{j=1}^n f(c_j) = \sum_{j=1}^n f(c_j(A))$,

$$u \longrightarrow (1, c_1, c_1) \longrightarrow (1, c_2, c_1 + c_2) \longrightarrow \dots \longrightarrow (n, c_n, \sum_{j=1}^n c_j) \longrightarrow v .$$

So the column sum problem indeed reduces to the shortest directed path problem.

Now, the number of vertices of D is bounded by $n(m+1)(mn+1) + 2 = O(m^2n^2)$ and hence is polynomial in the input. Therefore, as is well known, a shortest directed path from u to v in D can then be obtained in polynomial time, see e.g. [8]. Then an optimal nonincreasing tuple $c = (c_1, \dots, c_n)$ of column sums can be read off from this optimal directed path as explained above. Finally, an optimal matrix $A \in \{0, 1\}^{m \times n}$, with row sum tuple $r = (r_1, \dots, r_m)$ and column sum tuple $c = (c_1, \dots, c_n)$, can be computed in polynomial time by Lemma 2.2. \square

Part 2. Since all row sums are now bounded above by the constant b , by definition of the tuple $d = (d_1, \dots, d_n)$ conjugate to r we now have that $d_j = |\{i : r_i \geq j\}| = 0$ for all $j > b$. This implies that $\sum_{j=1}^b d_j = \sum_{j=1}^n d_j$ and hence a nonincreasing tuple $c = (c_1, \dots, c_n)$ is majorized by $d = (d_1, \dots, d_n)$ already if $\sum_{j=1}^i c_j \leq \sum_{j=1}^i d_j$ for $i = 1, \dots, b$ and $\sum_{j=1}^n c_j = \sum_{j=1}^n d_j$, since for $i > b$ we then automatically have $\sum_{j=1}^i c_j \leq \sum_{j=1}^n c_j = \sum_{j=1}^n d_j = \sum_{j=1}^i d_j$.

Note that, unlike the assumption in Part 1, the functions f_j can now be distinct, so we cannot assume that the matrix we are looking for has nonincreasing column sums.

Again we reduce the problem to the problem of finding a shortest directed path in a directed graph D where each edge has a length. We now construct D as follows.

There are two vertices u and v . There are also vertices labeled by $(b+3)$ -tuples of integers $(k, c_k, c_1^k, \dots, c_b^k, s_k)$ where $1 \leq k \leq n$, $0 \leq c_k \leq d_1 \leq m$, $m \geq d_1 \geq c_1^k \geq \dots \geq c_b^k \geq 0$, and $0 \leq s_k \leq \sum_{j=1}^n d_j \leq mn$. There are edges $[u, (1, c_1, c_1^1, \dots, c_b^1, s_1)]$ for $0 \leq c_1 \leq d_1$, $c_1^1 = c_1, c_2^1 = \dots = c_b^1 = 0$, and $s_1 = c_1$, of length $f_1(c_1)$. For $2 \leq k \leq n$ there are edges

$$[(k-1, c_{k-1}, c_1^{k-1}, \dots, c_b^{k-1}, s_{k-1}), (k, c_k, c_1^k, \dots, c_b^k, s_k)] ,$$

where $c_1^k \geq \dots \geq c_b^k$ are the b largest values among the $b+1$ values $c_1^{k-1}, \dots, c_b^{k-1}, c_k$, where $s_k = s_{k-1} + c_k$, and where $\sum_{j=1}^i c_j^k \leq \sum_{j=1}^i d_j$ for $i = 1, \dots, b$, of length $f_k(c_k)$. Finally, there are edges $[(n, c_n, c_1^n, \dots, c_b^n, s_n), v]$ for $s_n = \sum_{j=1}^n d_j$ of length 0.

Consider any directed path from u to v in D , which by the definition of D looks like

$$u \longrightarrow (1, c_1, c_1^1, \dots, c_b^1, s_1) \longrightarrow \dots \longrightarrow (n, c_n, c_1^n, \dots, c_b^n, s_n) \longrightarrow v .$$

Intuitively, traveling on the path, being at vertex $(k, c_k, c_1^k, \dots, c_b^k, s_k)$ indicates that the prefix (c_1, \dots, c_k) of the sequence had been determined, it sums to s_k , and its b largest values are $c_1^k \geq \dots \geq c_b^k$ and satisfy the majorization inequalities $\sum_{j=1}^i c_j^k \leq \sum_{j=1}^i d_j$ for $i = 1, \dots, b$. Formally, by definition of the edges in D , we have that the tuple $c = (c_1, \dots, c_n)$ corresponding to the path satisfies $\sum_{j=1}^n c_j = s_n = \sum_{j=1}^n d_j$, and, permuting it to be nonincreasing as $c_{\pi(1)} \geq \dots \geq c_{\pi(n)}$, it satisfies that $c_{\pi(j)} = c_j^n$ for $j = 1, \dots, b$ and hence $\sum_{j=1}^i c_{\pi(j)} \leq \sum_{j=1}^i d_j$ for $i = 1, \dots, b$. Since the row sums are bounded by b , as explained in the beginning of the proof of Part 2 above, it follows that $c_{\pi(1)} \geq \dots \geq c_{\pi(n)}$ is majorized by d . So by Proposition 2.1, there is a matrix $A \in \{0, 1\}^{m \times n}$ with row sum tuple r and column sum tuple c . By the definition of the lengths, the length of this directed path is $\sum_{j=1}^n f_j(c_j) = \sum_{j=1}^n f_j(c_j(A))$.

Conversely, for any matrix $A \in \{0, 1\}^{m \times n}$ with row sum tuple r and column sum tuple c , its nonincreasing permutation $c_{\pi(1)} \geq \dots \geq c_{\pi(n)}$ is majorized by d by Proposition 2.1, hence the following is a directed path from u to v in D of length $\sum_{j=1}^n f_j(c_j) = \sum_{j=1}^n f_j(c_j(A))$,

$$u \longrightarrow (1, c_1, c_1, 0, \dots, 0, c_1) \longrightarrow \dots \longrightarrow (n, c_n, c_{\pi(1)}, \dots, c_{\pi(b)}, \sum_{j=1}^n c_j) \longrightarrow v .$$

So the column sum problem indeed reduces to the shortest directed path problem.

Now, the number of vertices of D is bounded by $n(m+1)^{b+1}(mn+1) + 2 = O(m^{b+2}n^2)$ and since b is fixed it is polynomial in the input. So a shortest directed path from u to v in D can then be obtained in polynomial time, see e.g. [8]. Then an optimal nonincreasing tuple $c = (c_1, \dots, c_n)$ of column sums can be read off from this optimal directed path as explained above. Finally, an optimal matrix $A \in \{0, 1\}^{m \times n}$, with row sum tuple $r = (r_1, \dots, r_m)$ and column sum tuple $c = (c_1, \dots, c_n)$, can be computed in polynomial time by Lemma 2.2. \square

Acknowledgments

S. Onn was supported by a grant from the Israel Science Foundation and by the Dresner chair at the Technion. He thanks the referees for useful comments that improved the presentation.

References

- [1] Apollonio, N., Sebő, A.: Minconvex factors of prescribed size in graphs. *SIAM Journal on Discrete Mathematics* 23:1297–1310 (2009)
- [2] Brualdi, R.A.: *Combinatorial Matrix Classes*. Cambridge (2006)
- [3] Deza, A., Levin, A., Meesum, S.M., Onn, S.: Optimization over degree sequences. *SIAM Journal on Discrete Mathematics* 32:2067–2079 (2018)
- [4] Deza, G., Onn S.: Optimization over degree sequences of graphs. *Discrete Applied Mathematics* (2019). <http://doi.org/10.1016/j.dam.2019.12.016>
- [5] Erdős, P., Gallai, T.: Graphs with prescribed degrees of vertices (in Hungarian). *Matematikai Lapok* 11:264–274 (1960)
- [6] Marshall, A.W., Olkin, I., Arnold, B.C.: *Inequalities: Theory of Majorization and its Applications*. Springer (2011)
- [7] Ryser, H.J.: Combinatorial properties of matrices of zeroes and ones. *Canadian Journal of Mathematics* 9:371–377 (1957)
- [8] Schrijver A.: *Combinatorial Optimization*. Springer (2003)