

Infinite families of optimal systems of biangular lines related to representations of $SL(2, \mathbb{F}_q)$.

Mikhail Ganzhinov

Department of Communications and Networking
Aalto University School of Electrical Engineering
P.O. Box 15400, 00076 Aalto, Finland

Abstract

A line packing is optimal if its coherence is as small as possible. Most interesting examples of optimal line packings are achieving equality in some of the known lower bounds for coherence. In this paper two infinite families of real and complex biangular line packings are presented. New packings achieve equality in the real or complex second Levenshtein bound respectively. Both infinite families are constructed by analyzing well known representations of the finite groups $SL(2, \mathbb{F}_q)$. Until now the only known infinite families meeting the second Levenshtein bounds were related to the maximal sets of mutually unbiased bases (MUB). Similarly to the line packings related to the maximal sets of MUBs, the line packings presented here are related to the maximal sets of mutually unbiased weighing matrices. Another similarity is that the new packings are projective 2-designs. The latter property together with sufficiently large cardinalities of the new packings implies some improvement on largest known cardinalities of real and complex biangular tight frames (BTF).

1 Introduction

The Grassmannian line packing problem asks for line packings with minimal coherence. Packings with such properties are useful in signal processing applications [18] as they lead to improved error correction. The exact definition of coherence may vary in different applications.

Let \mathcal{L} be a set of lines represented by the set of unit vectors $\Phi_{\mathcal{L}} \subset \mathbb{F}^n$ where the field \mathbb{F} is \mathbb{R} or \mathbb{C} . The angle set of the line packing \mathcal{L} is defined as $\mathcal{A}_{\mathcal{L}} = \{|\langle x, y \rangle| : x, y \in \Phi_{\mathcal{L}}, x \neq y\}$. One natural way to define coherence $\mu_{\mathcal{L}}$ of the set \mathcal{L} is to put

$$\mu_{\mathcal{L}} = \max_{\alpha \in \mathcal{A}_{\mathcal{L}}} \alpha.$$

Line packings achieving minimal coherence among all packings of the same size in \mathbb{F}^n are called *optimal* or *Grassmannian line packings*. For $|\mathcal{A}_{\mathcal{L}}| = 1, 2$ and k corresponding line packings are called *equiangular*, *biangular* and *k-angular*, respectively.

Other properties of line packings may also be of great interest. A finite set of unit norm vectors $\Phi \subset \mathbb{F}^n$ is a *frame* if Φ spans \mathbb{F}^n . A frame Φ is *tight* if there exists a constant $a > 0$ such that $av = \sum_{\phi \in \Phi} \langle v, \phi \rangle \phi$ for any $v \in \mathbb{F}^n$. In other words, tight frames generalize some aspects of orthogonal bases. If $\Phi_{\mathcal{L}}$ is a tight frame, then the packing \mathcal{L} is also called a tight frame, in this case $a = |\mathcal{L}|/n$. The tightness of a frame is usually desired along other properties such as optimality and a small angle set.

Several lower bounds for minimal coherence are known. Packings with equality in those bounds are guaranteed to be optimal. One of these bounds is the Welch bound [21] which coincides with the first Levenshtein bound for line packings [16]. Line packings with equality in this bound are exactly equiangular tight frames (ETF). These are probably the most studied line systems [10, 11, 13, 19]. Maximal cardinality of real or complex equiangular line packings cannot exceed $\binom{n+1}{2}$ and n^2 lines, respectively [9]. Zauner's conjecture [23] predicts existence of complex ETFs of maximal cardinality in every dimension. According to [1, 15] existence of these packings seem to be related to abelian extensions of real quadratic fields.

Packings with cardinality larger than $\binom{n+1}{2}$ in the real case and n^2 in the complex case cannot be equiangular and cannot achieve equality in the Welch bound. When searching for larger line packings with good properties, one may turn the attention to systems of biangular lines. Biangular tight frames (BTFs) are investigated in [4, 6]. Introduction of the additional angle makes construction of tight frames with cardinalities $O(n^2)$ much easier than in the equiangular case. Despite that, construction of BTFs with cardinality exceeding the upper bound for equiangular lines proves to be challenging. To underline this fact, the following question is posed in [4].

Question 1. *Are there other "infinite families" of complex BTFs in addition to maximal sets of mutually unbiased bases (MUBs) [22] with cardinalities exceeding n^2 ?*

In contrary to the equiangular case, BTFs are not always optimal but may still possess good coherence.

To prove optimality of line packings with cardinality in the range $\binom{n+1}{2} < |\mathcal{L}| \leq \binom{n+2}{3}$ in the real case and $n^2 < |\mathcal{L}| \leq n\binom{n+1}{2}$ in the complex case, it is sufficient to achieve equality in the second Levenshtein bound [16]. Since packings of these sizes cannot be equiangular, it is natural to expect that some systems of biangular lines could be optimal. Linear programming bounds for systems of biangular lines with sufficiently small angles α, β such that $\mathcal{A}_{\mathcal{L}} = \{\alpha, \beta\}$ are given in [9].

Theorem 1 (Special bounds for biangular lines [9]). *If $3 - (n+2)(\alpha^2 + \beta^2) + n(n+2)\alpha^2\beta^2 > 0$ and $\alpha^2 + \beta^2 < \frac{6}{n+4}$, then the cardinality of an $\{\alpha, \beta\}$ -angular line system in \mathbb{R}^n cannot exceed*

$$N(\mathbb{R}^n, \alpha, \beta) = \frac{n(n+2)(1-\alpha^2)(1-\beta^2)}{3 - (n+2)(\alpha^2 + \beta^2) + n(n+2)\alpha^2\beta^2}.$$

If $2 - (n+1)(\alpha^2 + \beta^2) + n(n+1)\alpha^2\beta^2 > 0$ and $\alpha^2 + \beta^2 < \frac{4}{n+2}$, then the cardinality of an $\{\alpha, \beta\}$ -angular line system in \mathbb{C}^n cannot exceed

$$N(\mathbb{C}^n, \alpha, \beta) = \frac{n(n+1)(1-\alpha^2)(1-\beta^2)}{2 - (n+1)(\alpha^2 + \beta^2) + n(n+1)\alpha^2\beta^2}.$$

Systems of biangular lines achieving equality in the real or complex version of the special bound are known to be real or complex projective 2-designs [12]. One can verify that the real and complex special bounds for $\{0, \alpha\}$ -angular lines coincide with the real and complex second Levenshtein bounds, respectively. Thus, systems of $\{0, \alpha\}$ -angular lines attaining equality in the real or complex special bound are optimal and vice versa. In the real case a Gram matrix of a spherical code corresponding to a such optimal line system has the structure of Q -polynomial association scheme of class 3 [2]. Additionally, in the real case the angle α is necessarily a rational number of the form $\frac{1}{m}$ where $m \in \mathbb{N}$ [17]. Several examples of optimal $\{0, \alpha\}$ -angular packings are known, maximal sets of real and complex MUBs [14, 22] are some of these examples.

Several examples of $\{0, \alpha\}$ -angular line packings can be constructed as unions of nicely aligned disjoint full orthogonal bases. In this case it is said that these systems are related to mutually unbiased weighing matrices [3]. On the other hand, many exceptional line packings possess large automorphism groups and are line or basis transitive. In the latter case these bases may form a group G which is a subgroup of the full automorphism group of the line system. Original line packing may then be obtained from row/column vectors of matrices $M \in G$. The main results of this paper are two infinite families of real and complex BTFs given by Theorems 2 and 3, respectively. Both families produce optimal projective 2-designs because with angle sets of the form $\{0, \alpha\}$ and equalities in Theorem 1. Additionally, since the cardinalities of packings in these families exceed n^2 , Question 1 is answered in the affirmative.

The paper is organized as follows. Section 2 describes the mathematical machinery and the notations that will be used through the rest of the paper. In Section 3 we construct well-known $(q-1)$ -dimensional representations of the groups $\text{SL}(2, \mathbb{F}_q)$. In Section 4 we prove that certain representations of the groups $\text{SL}(2, \mathbb{F}_{2^{2k+1}})$ produce an infinite family of optimal $(2^{2k+1} - 1)$ -dimensional real line systems. Finally, in Section 5 we give an infinite family of optimal $(3^k - 1)/2$ -dimensional complex line systems related to representations of the groups $\text{SL}(2, \mathbb{F}_{3^k})$.

2 Preliminaries

Let \mathbb{F}_q be a finite field of order $q = p^n$ where p is a prime and $n \in \mathbb{N}$. Let \mathbb{L}/\mathbb{F}_q be a quadratic field extension of \mathbb{F}_q . If q is odd, \mathbb{L} can be defined as $\mathbb{F}_q(\rho)$ where $\rho \notin \mathbb{F}_q$ is a root of some quadratic polynomial of the form $x^2 - s$, $s \in \mathbb{F}_q$, that is irreducible over \mathbb{F}_q . If q is even, instead of $x^2 - s$, an irreducible polynomial (over \mathbb{F}_q) of the form $x^2 + x + s$ can be used.

Every element z of the field \mathbb{L} can be expressed in the form $z = x + y\rho$ and thus the additive group of \mathbb{L} is isomorphic to the vector space \mathbb{F}_q^2 . Let \mathbb{F}_q^* and \mathbb{L}^* be the multiplicative groups of the fields \mathbb{F}_q and \mathbb{L} , respectively; these groups are cyclic. If q is odd, \mathbb{F}_q^* can be decomposed into two sets of size $(q-1)/2$, QR_q and NQR_q , containing the squares (quadratic residues) and the nonsquares (quadratic nonresidues), respectively.

The field \mathbb{L} has a unique nontrivial automorphism σ fixing \mathbb{F}_q . If q is odd, then $\sigma(x + y\rho) = x - y\rho$ while if q is even then $\sigma(x + y\rho) = x + y + y\rho$ for all $x, y \in \mathbb{F}_q$. The functions $N : \mathbb{L} \rightarrow \mathbb{F}_q$ and $S : \mathbb{L} \rightarrow \mathbb{F}_q$ defined by $N(z) = z\sigma(z)$ and $S(z) = z + \sigma(z)$ for all $z \in \mathbb{L}$ are *relative norm* and *relative trace*, respectively. A *circle* C_d with $d \in \mathbb{F}_q^*$ is defined as the set $\{t \in \mathbb{L} \mid N(t) = d\}$. If $d = 1$, the notation $C_1 = C$ will be used. Each circle consists of $q+1$ elements, and C is a cyclic subgroup of \mathbb{L}^* . The *absolute trace* of the field \mathbb{F}_q is the function $Tr : \mathbb{F}_q \rightarrow \mathbb{F}_p$ defined by

$$Tr(x) = x + x^2 + \dots + x^{p^{n-1}} \text{ for all } x \in \mathbb{F}_q.$$

It is well known that while the norm is a multiplicative function, the trace is additive.

Let A be an abelian group. A function $\rho : A \rightarrow \mathbb{C} \setminus \{0\}$ is called a *character* of the group A if it is a homomorphism. The set \hat{A} formed by all characters of A is a group under pointwise multiplication. The group \hat{A} is always isomorphic to A . The character ρ_0 is said to be trivial if $\rho_0(a) = 1$ for all $a \in A$. The degree of a character ρ is defined to be the smallest integer k such that $\rho^k = \rho_0$. Additive characters of the field \mathbb{F}_q are homomorphisms from the additive group of the field \mathbb{F}_q to $\mathbb{C} \setminus \{0\}$. Similarly, the multiplicative character of a group $A \leq \mathbb{F}_q^*$ is a homomorphism from the subgroup A of the multiplicative group \mathbb{F}_q^* to $\mathbb{C} \setminus \{0\}$.

The special linear group $G = \text{SL}(2, \mathbb{F}_q)$ is a set of matrices $\{g = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \mid a, b, c, d \in \mathbb{F}_q, \det(g) = 1\}$ with the group operation being matrix multiplication. The group formed by invertible upper triangular matrices $g = \begin{bmatrix} a & b \\ 0 & d \end{bmatrix}$ where $a, d \in \mathbb{F}_q^*$ and $b \in \mathbb{F}_q$ will be denoted by B .

Finally, the real or complex Hilbert space (depending on the context) defined on a set S with the usual inner product $\langle \cdot, \cdot \rangle$ will be denoted by $H(S)$ and the associated norm of a vector $v \in H(S)$ will be denoted by $\|v\|_{H(S)}$.

3 Some representations of finite groups $\text{SL}(2, \mathbb{F}_q)$

In representation theory, elements of groups are represented as linear transformations of vector spaces. A representation of a group G is a pair (V, ϕ) where V is a vector space and $\phi : G \rightarrow \text{GL}(V)$ is a group homomorphism. In this paper we will be interested in $(q-1)$ -dimensional representations of the groups $G = \text{SL}(2, \mathbb{F}_q)$ and related line systems defined by the row and column vectors of matrices $\phi(g)$ where $g \in G$.

All irreducible representations of the groups $\text{SL}(2, \mathbb{F}_q)$ with q odd are covered (for example) in [20]. Throughout this section, the paper [20] will be used as the main source of information regarding representations of $\text{SL}(2, \mathbb{F}_q)$. Ultimately, some of these representations will produce optimal line packings.

As a starting point we introduce certain reducible representations of the groups $\text{SL}(2, \mathbb{F}_q)$ which later will be decomposed into irreducible ones. Let χ be a nontrivial additive character of \mathbb{F}_q defined by $\chi(x) = (-1)^{Tr(x)}$ for all $x \in \mathbb{F}_q$. Similarly as in [20], for all $g \in \text{SL}(2, \mathbb{F}_q)$ we define matrices $T_g : H(\mathbb{L}) \rightarrow H(\mathbb{L})$ by

$$T_g(v, u) = \begin{cases} -\frac{1}{q}\chi\left(\frac{aN(v)+dN(u)-S(u\sigma(v))}{c}\right), & \text{if } g \notin B \\ \chi(abN(v))\delta(u - va), & \text{otherwise,} \end{cases} \quad (1)$$

where δ is the delta function. We shall now show that the matrices T_g form a group.

Proposition 1. *The matrices T_g are unitary and form a group T . The group T is isomorphic to G with the group isomorphism $\phi : G \rightarrow T$ defined by $\phi(g) = T_g$ for all $g \in G$.*

Proof. For the case where q is odd, see [20]. If q is even, then the matrices T_g are real. The group property is equivalent to the equality

$$\langle T_{g_1}(v, \cdot), T_{g_2}(\cdot, u)^T \rangle = T_{g_1 g_2}(v, u) \text{ for all } g_1, g_2 \in G \text{ and } u, v \in \mathbb{L}. \quad (2)$$

When verifying the group property several cases arise. If at least one of matrices g_i is an element of the group B , the verification of the group property is trivial since one of matrices will be monomial. If $g_1 g_2 = 1$, then verification of the group property will only involve evaluation of the linear character sums. The most difficult case is when $g_1, g_2 \notin B$. In this case the problem can be reduced to computation of quadratic character sums of the form

$$I(v) = \sum_{u \in \mathbb{L}} \chi(N(u) + S(v\sigma(u))).$$

If $u = x + y\rho$ and $v = z + w\rho$, then the sum $I(v)$ can be rewritten as

$$I(v) = \sum_{x, y \in \mathbb{F}_q} \chi(x^2 + xy + sy^2 + xw + yz). \quad (3)$$

To evaluate $I(0)$ observe that when q is even the function $f : \mathbb{F}_q \rightarrow \mathbb{F}_q$, $f(x) = x^2$ for all $x \in \mathbb{F}_q$ is a bijection and thus $\sum_{x \in \mathbb{F}_q} \chi(x^2) = \sum_{x \in \mathbb{F}_q} \chi(x) = 0$. Each circle C_d contains $q + 1$ points one of which is on \mathbb{F}_q namely \sqrt{d} . Thus, the sum $I(0)$ can be rewritten as

$$I(0) = \sum_{u \in \mathbb{L}} \chi(N(u)) = \chi(0) + (q + 1) \sum_{u \in \mathbb{F}_q^*} \chi(N(u)) = (q + 1) \sum_{u \in \mathbb{F}_q} \chi(u^2) - q = -q.$$

All other sums $I(v)$ are related to $I(0)$. By performing a change of variables $u \rightarrow u + v$ in (3) we obtain

$$I(v) = \sum_{x, y \in \mathbb{F}_q} \chi(x^2 + xy + sy^2 + z^2 + zw + sw^2) = \sum_{u \in \mathbb{L}} \chi(N(v) + N(u)) = -\chi(N(v))q.$$

Now, verification of (2) will involve either calculation of sums $I(v)$, calculation of linear character sums or the evaluation of row vectors of the matrix T_{g_1} at a single element. \square

A group isomorphism ϕ defines a group homomorphism $\phi_1 : G \rightarrow \text{GL}(H(\mathbb{L}))$. Thus a pair $(H(\mathbb{L}), \phi_1)$ is a q^2 -dimensional representation of G . The next goal is to decompose a given q^2 -dimensional representation of G into smaller invariant subspaces. For $t \in C$ define an operator R_t by

$$R_t f(u) = f(tu) \text{ for all } f \in H(\mathbb{L}).$$

Now

$$\begin{aligned} T_g R_t f(u) &= T_g f(tu) = - \sum_{u \in \mathbb{L}} \chi\left(\frac{aN(v) + dN(u) - S(v\sigma(u))}{c}\right) f(tu)/q =_{u \rightarrow \sigma(t)u} \\ &= - \sum_{u \in \mathbb{L}} \chi\left(\frac{aN(tv) + dN(\sigma(t)u) - S(v\sigma(\sigma(t)u))}{c}\right) f(u)/q = R_t T_g f(u) \end{aligned}$$

for any $f \in H(\mathbb{L})$. Thus, the operators R_t commute with the matrices T_g . Let π be a character of the group C . Let $H(\mathbb{L})_\pi$ be a subspace of $H(\mathbb{L})$ consisting of elements that satisfy $f(tu) = \pi(t)f(u)$ for all $t \in C$. For all $f \in H(\mathbb{L})_\pi$

$$R_t T_g f(u) = \pi(t) T_g f(u) \in H(\mathbb{L})_\pi$$

which makes $H(\mathbb{L})_\pi$ an invariant subspace of the representation ϕ_1 .

If π_0 is the trivial (constant) character of C , then the subspace $H(\mathbb{L})_{\pi_0}$ is q -dimensional and consists of functions that are constant on circles C_d and $\{0\}$. If the character π is not trivial, the corresponding space $H(\mathbb{L})_\pi$ is now $(q - 1)$ -dimensional since necessarily $f(0) = 0$ for all $f \in H(\mathbb{L})_\pi$. Since subspaces $H(\mathbb{L})_\pi$ are

clearly orthogonal to each other, the whole q^2 -dimensional space $H(\mathbb{L})$ can be decomposed into a direct sum of $q + 1$ subspaces $H(\mathbb{L})_\pi$, indeed, $q^2 = q + q(q - 1)$.

Let $\theta = \{x_d \mid x_d \in C_d \text{ for every } d \in \mathbb{F}_q^*\}$ be a system of representatives of circles C_d . For each nontrivial character π , an orthonormal basis $B_{\pi,\theta} = \{e_d\}$ of the subspace $H(\mathbb{L})_\pi$ is obtained by choosing

$$e_d(x) = \begin{cases} \frac{1}{\sqrt{q+1}}\pi(x/x_d), & \text{if } x \in C_d \\ 0, & \text{otherwise.} \end{cases}$$

Let $F_{\pi,\theta} : H(\mathbb{L})_\pi \rightarrow H(\mathbb{F}_q^*)$ be a unitary operator defined by $F_{\pi,\theta}(e_d) = \delta(x - d)$ for all $e_d \in B_{\pi,\theta}$. Representations associated with subspaces $H(\mathbb{L})_\pi$ could be constructed explicitly as the pairs $(H(\mathbb{F}_q^*), \phi_\pi)$ where $\phi_\pi : G \rightarrow \text{GL}(H(\mathbb{F}_q^*))$ is a group homomorphism. An easy way to accomplish this is to simply put $\phi_\pi(g) = T_{g,\pi,\theta}$ where $T_{g,\pi,\theta} = F_{\pi,\theta}T_gF_{\pi,\theta}^{-1}$. Let $T_{\pi,\theta}$ be a set of all matrices $T_{g,\pi,\theta}$ with fixed π and θ . Explicitly, after performing matrix multiplications we get

$$T_{g,\pi,\theta}(v, u) = \begin{cases} \frac{-1}{q}\chi\left(\frac{av+du}{c}\right)\sum_{t \in C}\chi\left(\frac{-S(tx_u\sigma(x_v))}{c}\right)\pi(t), & \text{if } g \notin B \\ \pi(ax_v/x_u)\chi(abv)\delta(u - a^2v), & \text{otherwise} \end{cases} \quad (4)$$

for all $v, u \in \mathbb{F}_q^*$. Character sums of the form

$$J_\pi(z) := \frac{1}{q}\sum_{t \in C}\chi(-S(tz))\pi(t) \text{ where } z \in \mathbb{L}$$

are sometimes called Bessel functions of the field \mathbb{L} . By using these functions in the case $g \notin B$, the matrices $T_{g,\pi,\theta}$ can be expressed as

$$T_{g,\pi,\theta}(v, u) = -\chi\left(\frac{av + du}{c}\right)J_\pi(x_u\sigma(x_v)/c) \text{ for all } u, v \in \mathbb{F}_q^*. \quad (5)$$

At this point it is good to summarize what has been done so far. We began by introducing q^2 -dimensional representations $(H(\mathbb{L}), \phi_1)$ where matrices corresponding to elements $g \in \text{SL}(2, \mathbb{F}_q)$ were defined by (1). Later, representations $(H(\mathbb{L}), \phi_1)$ were decomposed into invariant subspaces $H(\mathbb{L})_\pi$ where π is a character of C . The representation corresponding to the subspace $H(\mathbb{L})_{\pi_0}$ where π_0 is trivial is q -dimensional and will not be needed in following sections. All other $(q - 1)$ -dimensional invariant subspaces $H(\mathbb{L})_\pi$ where π is nontrivial were used to construct representations $(H(\mathbb{F}_q^*), \phi_\pi)$. The homomorphism ϕ_π is given explicitly by (4) and (5) through Bessel functions of the field \mathbb{L} . Representation $(H(\mathbb{F}_q^*), \phi_\pi)$ depends on the choice of representatives θ . We conclude this section by remarks which clarify how exactly the choice of θ affects the corresponding representations.

Remark 1. It is clear that the particular form of the representations $(H(\mathbb{F}_q^*), \phi_\pi)$ depends on the particular choice of representatives θ . Let $\theta_i = \{x_{d,i} \mid x_{d,i} \in C_d \text{ for every } d \in \mathbb{F}_q^*\}$, $i = 1, 2$ be two different systems of representatives. From the definition of Bessel functions it is easy to see that

$$J_\pi(tz) = \bar{\pi}(t)J_\pi(z) \text{ for all } t \in C. \quad (6)$$

From (6) follows that

$$T_{g,\pi,\theta_2}(v, u) = T_{g,\pi,\theta_1}(v, u)\bar{\pi}(x_{v,2}/x_{v,1})\bar{\pi}(x_{u,2}/x_{u,1}). \quad (7)$$

when $g \notin B$ and (7) holds also in the case $g \in B$ which can be verified directly. In the following sections sets of biangular lines will be represented by row and column vectors of matrices $T_{g,\pi,\theta}$. Equation (7) implies that the choice of θ does not affect the structure of the resulting line system.

Remark 2. If q is even, by using perfectness of the field \mathbb{L} , it is possible to choose representatives of the circles C_d to be $\theta = \{\sqrt{d} \mid d \in \mathbb{F}_q^*\}$. With this particular choice

$$T_{g,\pi,\theta}(v, u) = \begin{cases} -\chi\left(\frac{av+du}{c}\right)J_\pi(\sqrt{vu}/c), & \text{if } g \notin B \\ \pi(a\sqrt{v/u})\chi(abv)\delta(u - a^2v), & \text{otherwise.} \end{cases}$$

In order to get rid of square roots in these expressions, the matrices $T_{g,\pi}$, will be defined for q even. Notice that $u^2 = N(u)$ for all $u \in \mathbb{F}_q$. Let $T_{g,\pi}(u, v) = T_{g,\pi,\theta}(N(v), N(u))$, explicitly

$$T_{g,\pi}(v, u) = \begin{cases} -\chi\left(\frac{aN(v)+dN(u)}{c}\right)J_\pi(vu/c), & \text{if } g \notin B \\ \pi(av/u)\chi(abN(v))\delta(N(u) - a^2N(v)), & \text{otherwise} \end{cases}$$

for all $v, u \in \mathbb{F}_q^*$. The matrices $T_{g,\pi}$ define representations $(H(\mathbb{F}_q^*), \phi_{2,\pi})$ where homomorphisms $\phi_{2,\pi}$ are defined by $\phi_{2,\pi}(g) = T_{g,\pi}$ for all $g \in G$.

4 Sets of biangular lines related to representations of $\mathbf{SL}(2, \mathbb{F}_{2^{2k+1}})$

This section is dedicated to real line systems \mathcal{L}_π represented by row or column vectors of matrices $T_{g,\pi}$ related to representations $(H(\mathbb{F}_q^*), \phi_{2,\pi})$. In the first part of this section we will derive Propositions 2-4 which express basic properties common to all such line systems. In particular, Proposition 4 gives a lower bound on coherence; line systems achieving this lower bound are always biangular. By Remark 3 these systems are also optimal. We will show then, that a particular infinite family achieving the lower bound in Proposition 4 actually exists. This is done by establishing a connection through Hasse-Davenport relations between some of the functions J_π and a certain additive character sums.

The usual p -norm of a vector $v \in H(\mathbb{F}_q^*)$ is denoted by $\|v\|_p$ where $1 \leq p \leq \infty$. Some important basic properties of the functions J_π which will be needed later are described in following proposition.

Proposition 2. *For q even and π a nontrivial character of C , the following properties of the functions J_π hold:*

1. $J_\pi(x) \in \mathbb{R}$ and $J_\pi(x) = J_{\bar{\pi}}(x)$ for all $x \in \mathbb{F}_q^*$.
2. $\|J_\pi(x)\|_2^2 = 1$.
3. $\|J_\pi(x)^2\|_2^2 = \frac{2}{q}$.
4. $\|J_\pi(x)J_\pi(cx)\|_2^2 = \frac{1}{q}$ when $c \neq 0, 1$.

Proof. To prove the first identity observe that $S(x\sigma(t)) = S(xt)$ and $\pi(t) = \overline{\pi(\sigma(t))}$ which gives

$$J_{\bar{\pi}}(x) = \sum_{t \in C} \chi(S(xt))\overline{\pi(t)}/q = \frac{\sum_{t \in C} \chi(S(x\sigma(t)))\overline{\pi(\sigma(t))}/q}{\sum_{t \in C} \chi(S(xt))\pi(t)/q} = \frac{\sum_{t \in C} \chi(S(xt))\overline{\pi(\sigma(t))}/q}{\sum_{t \in C} \chi(S(xt))\pi(t)/q} = J_\pi(x)$$

The second identity is obvious since the matrices $T_{g,\pi}$ are orthogonal. For the third identity, fix $g_1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ and set $g_2(d) = \begin{bmatrix} 0 & 1 \\ 1 & d \end{bmatrix}$. Now $g_2(d)g_1 = g_3(d) = \begin{bmatrix} 1 & 0 \\ d & 1 \end{bmatrix}$, $T_{g_2(d),\pi}T_{g_1,\pi} = T_{g_3(d),\pi}$ and explicitly

$$Q(d) := T_{g_3(d),\pi}(1, 1) = \chi(dN(0))J_\pi^2(0) + \sum_{x \in \mathbb{F}_q^*} J_\pi(x)^2 \chi(dN(x)).$$

Since $N(x) = x^2$ for all $x \in \mathbb{F}_q$, the functions $\chi(dN(t))/\sqrt{q}$ form an orthonormal basis of $H(\mathbb{F}_q)$ and therefore (by Parseval's theorem) $\|Q(d)\|_{H(\mathbb{F}_q)}^2 = q\|J_\pi(x)^2\|_2^2$. On the other hand, the values of the function $Q(d)$ are known:

$$Q(d) = T_{g_3(d),\pi}(1, 1) = \begin{cases} -\chi(N(1)/d)^2 J_\pi(1/d) & \text{if } d \neq 0 \\ 1 & \text{if } d = 0. \end{cases}$$

Since $\|J_\pi(1/x)\|_2^2 = 1$ we conclude that $\|Q(d)\|_{H(\mathbb{F}_q)}^2 = 1^2 + \|J_\pi(1/x)\|_2^2 = 2$ and thus $\|J_\pi(x)^2\|_2^2 = \frac{2}{q}$.

The fourth identity can be proved in the same way as the third identity by choosing $g_1(c) = \begin{bmatrix} 0 & c \\ c^{-1} & 0 \end{bmatrix}$ with any fixed $c \neq 0, 1$ to get

$$Q_c(d) := T_{g_3(c,d),\pi}(1,1) = \chi(dN(0))J_\pi(0)J_\pi(c \cdot 0) + \sum_{y \in \mathbb{F}_q^*} J_\pi(x)J_\pi(cx)\chi(dN(x))$$

where $g_3(c,d) = g_2(d)g_1(c)$, and explicitly

$$Q_c(d) = T_{g_3(c,d),\pi}(1,1) = \begin{cases} -\chi((c^2+1)N(1)/d)J_\pi(c/d) & \text{if } d \neq 0 \\ 0 & \text{if } d = 0. \end{cases}$$

Since $|\chi(y)| = 1$ for all $y \in \mathbb{F}_q$, we have $\|Q(d)\|_2^2 = 0^2 + \|J_\pi(c/x)\|_2^2 = 1$ and thus $\|J_\pi(x)J_\pi(cx)\|_2^2 = \frac{1}{q}$. \square

Now it is time to define line systems \mathcal{L}_π in the case when q is even by giving explicitly corresponding sets of representatives. We define \mathcal{L}_π to be the line system where each line is represented by the vector in the set

$$\Phi_\pi = \{\chi(aN(x))J_\pi(yx) \mid a \in \mathbb{F}_q, y \in \mathbb{F}_q^*\} \cup \{\delta(x-y) \mid y \in \mathbb{F}_q^*\} \subset H(\mathbb{F}_q^*).$$

Each vector in Φ_π is a row vector in some matrix $T_{g,\pi}$ where $g \in G$.

Proposition 3. *If q is even, $|\Phi_\pi| = q^2 - 1$ for all nontrivial characters π and $q > 2$. All vectors in Φ_π are real and non-parallel.*

Proof. The set Φ_π contains $q - 1$ vectors of the form $\delta(x - y)$ as well as at most $q(q - 1)$ vectors of the form $\chi(aN(x))J_\pi(yx)$. All these vectors are real according to part 1 of Proposition 2. The cardinality of Φ_π is thus at most $q^2 - 1$. It remains to show that cardinality of Φ_π is at least $q^2 - 1$ by showing that there are no parallel vectors in the description of the set Φ_π .

The multiplicative group \mathbb{L}^* is a subgroup of $\text{GL}(2, \mathbb{F}_q)$ where every element $z = b + c\rho \in \mathbb{L}^*$ is represented by the invertible matrix $g_z \in \text{GL}(2, \mathbb{F}_q)$ where $g_z = \begin{bmatrix} b & sc \\ c & b+c \end{bmatrix}$ and $s \in \mathbb{F}_q$ as in Section 2. Matrix g_z is upper triangular exactly when $c = 0$, in this case g_z is a scalar matrix and hence $z \in \mathbb{F}_q^* \subset \mathbb{L}^*$. Matrices associated with elements $z \in C$ have $\det(g_z) = N(z) = 1$ and thus form a cyclic subgroup of $\text{SL}(2, \mathbb{F}_q)$. Since $C \cap \mathbb{F}_q^* = \{1\}$, the only upper triangular matrix corresponding to elements of C is the identity matrix.

Let z_0 be a generator of C . From the group property of C follows

$$\langle T_{g_{z_0},\pi}^r(u, \cdot), T_{g_{z_0},\pi}^t(v, \cdot) \rangle = T_{g_{z_0},\pi}^{r-t}(u, v) \text{ for all } 0 \leq r, t < q + 1.$$

If $r = t$, then $T_{g_{z_0},\pi}^{r-t}$ is an identity matrix. In all other cases matrices $g_{z_0}^{r-t}$ are not upper triangular since $C \cap \mathbb{F}_q^* = \{1\}$. Thus, the absolute values of entries of matrices $T_{g_{z_0},\pi}^{r-t}$ are thus contained in the set $\{|J_\pi(x)| \mid x \in \mathbb{F}_q^*\}$. From part 3 of Proposition 2 follows that $|J_\pi(x)| \leq \sqrt[4]{\frac{2}{q}} < 1$ for all $x \in \mathbb{F}_q^*$. This can be used to deduce that all row vectors $T_{g_{z_0}^r,\pi}, 0 \leq r < q + 1$ are on different lines. Thus, one of the vectors in the each pair $\{T_{g_{z_0}^r,\pi}(v, \cdot), -T_{g_{z_0}^r,\pi}(v, \cdot)\}, 0 \leq r < q + 1, v \in \mathbb{F}_q^*$ is contained in the set Φ_π , proving that $|\Phi_\pi| = q^2 - 1$. \square

Our next goal is to determine a lower bound for the coherence of the line packings \mathcal{L}_π .

Proposition 4. *Let q be even. For all nontrivial characters π of the circle C , $\mu_{\mathcal{L}_\pi} \geq \sqrt{2/q}$ holds. Equality can only be achieved if \mathcal{L}_π is biangular.*

Proof. From Proposition 3 follows that the angle set of \mathcal{L}_π is $\mathcal{A}_\mathcal{L} = \{|J_\pi(x)| \mid x \in \mathbb{F}_q^*\}$. Absolute values of the functions J_π can be estimated with the help of Proposition 2. Indeed, parts 2 and 3 of Proposition 2 imply that the maximal absolute value must be at least $\sqrt{2/q}$ since from the Hölder inequality follows

$$2/q = \|J_\pi(x)^4\|_1 \leq \|J_\pi(x)^2\|_1 \|J_\pi(x)^2\|_\infty = \|J_\pi(x)^2\|_\infty.$$

On the other hand, if the maximal absolute value of the function J_π is exactly $\sqrt{2/q}$ then coordinatewise

$$J_\pi(x)^4 \leq 2J_\pi(x)^2/q \text{ for all } x \in \mathbb{F}_q^*$$

and equality occurs only if $J_\pi(x) \in \{0, \pm\sqrt{2/q}\}$ for all $x \in \mathbb{F}_q^*$. In this case $\mathcal{A}_\mathcal{L} = \{0, \sqrt{2/q}\}$ and the line system \mathcal{L}_π is biangular. \square

Remark 3. Proposition 3 tells us that line systems \mathcal{L}_π always have cardinality $q^2 - 1$. On the other hand, from Proposition 4 we know that any line system \mathcal{L}_π which meets lower bound for the coherence has angle set $\mathcal{A}_{\mathcal{L}_\pi} = \{0, \sqrt{2/q}\}$. It turns out that such biangular line systems achieve equality in the Special Bound (Theorem 1). Indeed,

$$N(\mathbb{R}^{q-1}, 0, \sqrt{2/q}) = \frac{(q-1)(q+1)(1-2/q)}{3 - (q+1)(2/q)} = q^2 - 1$$

which equals the cardinality of the line system \mathcal{L}_π .

The main goal of this section is to show that there actually exists a nontrivial character π of the group C such that \mathcal{L}_π achieves equality in the Special Bound. If $q = 2^{2k+1}$ then

$$2^{2k+1} \equiv (-1)^{2k+1} = -1 \pmod{3} \quad (8)$$

and thus $q + 1$ is always divisible by 3. Since the character group of C is isomorphic to C , there exist exactly three characters π such that $\pi^3 = \pi_0$. One of these characters is obviously π_0 which defines the q -dimensional representation. The other two characters are of order 3. When one is π , other is always $\bar{\pi}$. These two characters will define the same line system $\mathcal{L}_{\bar{\pi}} = \mathcal{L}_\pi$ and the same Bessel function $J_{\bar{\pi}} = J_\pi$ on \mathbb{F}_q^* .

We will show that characters of order 3 always produce line systems \mathcal{L}_π with optimal properties. To prove optimality of these line systems, Gauss sums will be introduced in Definition 1 and some of their properties will be introduced in Propositions 5 and 6.

Definition 1. A Gauss sum is a sum of the form

$$G(\chi, \pi) = \sum_{x \in \mathbb{F}_q^*} \chi(x)\pi(x)$$

where χ is an additive character of the field \mathbb{F}_q and π is a multiplicative character of the field \mathbb{F}_q .

Proposition 5. For a nontrivial additive character χ , Gauss sums have the following properties

1. $|G(\chi, \pi)| = \sqrt{q}$ for π nontrivial,
2. $G(\chi, \pi) = -1$ for π trivial,
3. $G(\chi, \bar{\pi}) = \pi(-1)\overline{G(\chi, \pi)}$.

The properties of Gauss sums in Proposition 5 are well known.

Proposition 6 (Hasse–Davenport relations, [8]). Let \mathbb{F}_q be a finite field with q elements and let \mathbb{L}/\mathbb{F}_q be its extension of degree s . Let χ, ρ be (nontrivial) additive and multiplicative characters of the field \mathbb{F}_q , respectively. Let N, S be the norm and the trace from \mathbb{L} to \mathbb{F}_q , respectively. Let $\chi'(z) = \chi(S(z))$ and $\rho'(z) = \rho(N(z))$ be additive and multiplicative characters of the field \mathbb{L} . Then

$$(-1)^s G(\chi, \rho)^s = -G(\chi', \rho') \quad (\text{Hasse–Davenport lifting relation}).$$

Let π be a multiplicative character of the field \mathbb{F}_q of order m dividing $q - 1$. Then

$$\prod_{a=0}^{m-1} G(\chi, \rho\pi^a) = -\rho^{-m}(m)G(\chi, \rho^m) \prod_{a=0}^{m-1} G(\chi, \pi^a) \quad (\text{Hasse–Davenport product relation}).$$

Results of L. Carlitz concerning the evaluation of some additive character sums will prove to be useful. In fact, we will prove that when a character π has degree 3, the values of the functions J_π are directly related to additive character sums presented in the next proposition.

Proposition 7 (L. Carlitz, [5]). Let \mathbb{F}_q be a field with $q = 2^{2k+1}$ elements and χ its additive character. Then

$$\sum_{x \in \mathbb{F}_q} \chi(ax^3 + bx) \in \{0, \pm 2^{k+1}\}$$

while at least one of the parameters a, b is nonzero.

Finally, two simple auxiliary lemmas will be needed.

Lemma 1. *Let $r \in \{1, 2, 3, \dots\}$ and n odd with $\gcd(r, n) = 1$. Let $f : \mathbb{Z}/n\mathbb{Z} \rightarrow \mathbb{C}$ with be a function with the following two properties*

1. $|f(k)| = r$ for all $k \in \mathbb{Z}/n\mathbb{Z}$.
2. $f(k) = \sum_{0 \leq j \leq n-1} a_j e^{2\pi i j k/n}$ with integral coefficients a_j for all $0 \leq j \leq n-1$.

Let $s : \mathbb{Z}/n\mathbb{Z} \rightarrow \{-1, 1\}$ be any function. If the function $s(k)f(k) : \mathbb{Z}/n\mathbb{Z} \rightarrow \mathbb{C}$ can be expressed in the form

$$s(k)f(k) = \sum_{0 \leq j \leq n-1} b_j e^{2\pi i j k/n} \quad (9)$$

with integral coefficients b_j for all $0 \leq j \leq n-1$, then the function $s(k)$ is necessarily a constant.

Proof. Assume that there exist functions $f(k)$ and nonconstant $s(k)$ satisfying the requirements of Lemma 1. Assume further that equation (9) holds with integral coefficients b_j for all $0 \leq j \leq n-1$. Let $\mathcal{S} = \{k \in \mathbb{Z}/n\mathbb{Z} \mid s(k) = -1\}$; since $s(k)$ is not constant, $|\mathcal{S}| < n$. Now we determine the $H(\mathbb{Z}/n\mathbb{Z})$ -norm of the function $g(k) = (1 - s(k))f(k)$ in two different ways. Direct computation gives

$$\|g\|_{H(\mathbb{Z}/n\mathbb{Z})}^2 = \sum_{0 \leq k < n} |(1 - s(k))f(k)|^2 = 4|\mathcal{S}|r^2.$$

Alternatively, the same norm can be computed on the Fourier side

$$\|g\|_{H(\mathbb{Z}/n\mathbb{Z})}^2 = n \sum_{0 \leq k < n} (a_j - b_j)^2 = n \cdot s, \text{ where } s \in \mathbb{N}.$$

This leads to a contradiction since the first value is not divisible by n . □

Lemma 2. *Let q be even and π be a nontrivial character of the unit circle $C \subset \mathbb{L}$ of order 3. Then $J_\pi(x)$ is rational of the form k/q for all $x \in \mathbb{F}_q$.*

Proof. Since the character π is of order 3, $\pi(t) \in \{1, e^{2\pi i/3}, e^{-2\pi i/3}\}$ for all $t \in C$. Additionally $\chi(S(xt)) = \chi(S(x\sigma(t)))$ and $\pi(t) = \overline{\pi(\sigma(t))}$ for all $t \in C$, $x \in \mathbb{F}_q$. Also $\pi(1) = 1$. For a generator t_0 of C we have

$$J_\pi(x) = \sum_{t \in C} \chi(S(xt))\pi(t)/q = \chi(x)\pi(1)/q + \sum_{1 \leq n \leq q/2} \chi(S(xt_0^n))(\pi(t_0^n) + \pi(t_0^{q+1-n}))/q.$$

Every summand in the last sum is either $2/q$ or $-1/q$, which completes the proof. □

The following theorem is the main result of this paper.

Theorem 2. *Let $q = 2^{2k+1}$ and π be a nontrivial character of the unit circle $C \subset \mathbb{L}$ of order 3. Then the line system \mathcal{L}_π is $(q-1)$ -dimensional with cardinality $|\mathcal{L}_\pi| = q^2 - 1$. Additionally \mathcal{L}_π is biangular with $\mathcal{A}_{\mathcal{L}_\pi} = \{0, \sqrt{2/q}\}$.*

Proof. Let $\hat{\mathbb{L}}$ be the group of multiplicative characters of the field \mathbb{L} and $\psi : \mathbb{L}^* \rightarrow \hat{\mathbb{L}}$ be some fixed isomorphism. The group \mathbb{L}^* is a direct product of two subgroups C and \mathbb{F}_q^* which have orders $q+1$ and $q-1$ respectively. The characters in $\psi(C) \leq \hat{\mathbb{L}}$ are constant on the lines $l_t = t\mathbb{F}_q^*$ where $t \in C$ while the characters in $\psi(\mathbb{F}_q^*) \leq \hat{\mathbb{L}}$ are constant on the circles C_d where $d \in \mathbb{F}_q^*$. A character π^* of the group C is a restriction of the corresponding character $\pi_* \in \psi(C)$. This correspondence is a bijection.

Let ξ be a generator of the multiplicative group \mathbb{F}_q^* then the character

$$\rho(\xi^k) = e^{2\pi k i/(q-1)} \text{ for all } 0 \leq k < q-1 \quad (10)$$

generates all multiplicative characters of \mathbb{F}_q^* . In this case character $\rho_* = \rho \circ N$ generates $\psi(\mathbb{F}_q^*)$. From now on it will be assumed that the character ρ is defined according to (10) with some fixed ξ .

Define the function $f : \mathbb{Z}/(q-1)\mathbb{Z} \rightarrow \mathbb{C}$ by $f(r) := G(\chi \circ S, \pi_* \rho_*^r)$. Since the characters ρ_*^r are constant on the circles C_d , it is possible to rewrite the expression for $f(r)$ into the form

$$f(r) = \sum_{z \in \mathbb{L}^*} \chi(S(z)) \pi_*(z) \rho_*^r(z) = \sum_{x \in \mathbb{F}_q^*} \rho_*^r(x) \sum_{t \in C} \chi(S(xt)) \pi_*(t) = q \sum_{x \in \mathbb{F}_q^*} \rho^r(x) J_\pi(x) = q \sum_{1 \leq k < q-1} e^{\frac{2\pi i k r}{q-1}} J_\pi(\xi^k),$$

where $z = xt$ and $x = \xi^k \in \mathbb{F}_q^*$, $t \in C$. Thus, the function $f(r)$ can be expressed as a sum of different orthogonal components. The amplitude of each component $e^{\frac{2\pi i k r}{q-1}}$ is $q J_\pi(\xi^k)$.

From now on assume that the character $\pi_* = \pi$ is of order 3, which means that the corresponding character π_* is also of order 3. Because of (8), this is possible only if $q = 2^{2k+1}$.

It is possible to compute the exact value of $f(0)$ by using the fact that the character π_* is constant on the lines l_t and the fact that $\sum_{t \in C} \pi(t) = 0$. Indeed,

$$f(0) = \sum_{t \in C} \pi(t) \sum_{x \in \mathbb{F}_q^*} \chi(S(xt)) = q - 1 - \sum_{t \in C \setminus \{1\}} \pi(t) = q.$$

It is possible to use the Hasse–Davenport product relation on the characters π_* , ρ_* and $\chi \circ S$ in order to obtain

$$\prod_{a=0}^2 G(\chi \circ S, \pi_*^a \rho_*^r) = -\rho_*^{-3r} (3) G(\chi \circ S, \rho_*^{3r}) \prod_{a=0}^2 G(\chi \circ S, \pi_*^a) \text{ when } r \neq 0. \quad (11)$$

Now we have to simplify (11). On the left-hand side of (11) notice that $f(r) = G(\chi \circ S, \pi_*^1 \rho_*^r) = G(\chi \circ S, \pi_*^2 \rho_*^r)$. The reason for that is because characters π and $\pi^2 = \bar{\pi}$ define the same Bessel function J_π on \mathbb{F}_q^* and hence the same function $f(r)$. Now, the left-hand side of (11) becomes

$$\prod_{a=0}^2 G(\chi \circ S, \pi_*^a \rho_*^r) = G(\chi \circ S, \rho_*) f(r)^2 \text{ when } r \neq 0.$$

On the right-hand side notice that $\rho_*^{-3r} (3) = 1$ since $1 = 3$ in fields of characteristic 2. Additionally, $q = f(0) = G(\chi \circ S, \pi_*) = G(\chi \circ S, \pi_*^2)$. And finally, $G(\chi \circ S, \pi_*^0) = -1$ by Proposition 5.2 since π_*^0 is a trivial character. Now, the right-hand side of (11) becomes

$$-\rho_*^{-3r} (3) G(\chi \circ S, \rho_*^{3r}) \prod_{a=0}^2 G(\chi \circ S, \pi_*^a) = q^2 G(\chi \circ S, \rho_*^{3r}) \text{ when } r \neq 0. \quad (12)$$

By using all described simplifications and by dividing both sides of (11) by $G(\chi \circ S, \rho_*)$ we obtain

$$f(r)^2 = \frac{q^2 G(\chi \circ S, \rho_*^{3r})}{G(\chi \circ S, \rho_*)} \text{ when } r \neq 0. \quad (13)$$

To further simplify (13) we can use part 3 of Proposition 5 and the fact that $\rho_* = \rho \circ N$ to get

$$\frac{q^2}{G(\chi \circ S, \rho_*^r)} = G(\chi \circ S, \bar{\rho}_*^r) = G(\chi \circ S, \rho_*^{-r}) \text{ when } r \neq 0.$$

This transforms (13) into

$$f(r)^2 = G(\chi \circ S, (\rho \circ N)^{3r}) G(\chi \circ S, (\rho \circ N)^{-r}) \text{ when } r \neq 0.$$

By using the Hasse–Davenport lifting relation (6) can be transformed into

$$f(r)^2 = G(\chi, \rho^{3r})^2 G(\chi, \rho^{-r})^2 \text{ when } r \neq 0,$$

and finally

$$f(r) = \pm G(\chi, \rho^{3r}) G(\chi, \rho^{-r}) \text{ when } r \neq 0. \quad (14)$$

Now, we define a function $g : \mathbb{Z}/(q-1)\mathbb{Z} \rightarrow \mathbb{C}$ such that $g(r) = G(\chi, \rho^{3r})G(\chi, \rho^{-r})$ for all $r \in \mathbb{Z}/(q-1)\mathbb{Z}$, explicitly

$$g(r) = \sum_{x \in \mathbb{F}_q^*} \rho(x)^{3r} \chi(x) \sum_{y \in \mathbb{F}_q^*} \rho^{-r}(y) \chi(y).$$

The variable change $y = tx^3$, $t \in \mathbb{F}_q^*$ in previous sum (the function tx^3 is a bijection as long as $q = 2^{2k+1}$) gives

$$g(r) = \sum_{t \in \mathbb{F}_q^*} \sum_{x \in \mathbb{F}_q^*} \rho(x)^{3r} \rho(tx^3)^{-r} \chi(x + tx^3) = \sum_{t \in \mathbb{F}_q^*} \rho(t)^{-r} \sum_{x \in \mathbb{F}_q^*} \chi(x + tx^3) = \sum_{0 \leq k < q-1} e^{-2\pi ikr} \sum_{x \in \mathbb{F}_q^*} \chi(x + \xi^k x^3)$$

which is the Fourier transform of $g(r)$. Coefficients corresponding to frequencies $e^{-2\pi ikr}$ are given by the sums

$$a_{-k} = \sum_{x \in \mathbb{F}_q^*} \chi(x + \xi^k x^3) \in \{-1, 2^{2^{k+1}} - 1, -2^{2^{k+1}} - 1\},$$

where exact set of the values is determined according to Proposition 7. Notice that $1 = g(0) \neq f(0) = q$. This can be adjusted by introducing the new function $g_1(r) = g(r) + (q-1)\delta(r)$. We have

$$g_1(r) = \sum_{0 \leq k < q-1} e^{-2\pi ikr} \sum_{x \in \mathbb{F}_q} \chi(x + \xi^k x^3) \text{ for all } 0 \leq r < q-1.$$

Finally, by comparing properties of functions $g_1(r)$ and $f(r)$ we observe

1. $g_1(0) = f(0) = q$ and $f(r) = \pm g_1(r)$ for all $r \in \mathbb{Z}/(q-1)\mathbb{Z}$.
2. $|g_1(r)| = q$ for all $r \in \mathbb{Z}/(q-1)\mathbb{Z}$ and $\gcd(q, q-1) = 1$.
3. $f(r)$ can be expressed in the form $f(r) = \sum_{0 \leq j < q-1} a_j e^{2\pi ijr/(q-1)}$ where $a_j = qJ_\pi(\xi^j) \in \mathbb{Z}$ for all $0 \leq j < q-1$ where by Lemma 2 all values of qJ_π are integral.
4. $g_1(r)$ can be expressed in the form $g_1(r) = \sum_{0 \leq j < q-1} b_j e^{2\pi ijr/(q-1)}$ where $b_j = \sum_{x \in \mathbb{F}} \chi(x + \xi^{-k} x^3) \in \{0, \pm 2^{k+1}\}$ according to Proposition 7.

From Properties 1-4 we conclude that the functions $g_1(r)$ and $f(r)$ satisfy the requirements of Lemma 1 and thus either $g_1(r) = f(r)$ or $g_1(r) = -f(r)$ for all r . Since $f(0) = g_1(0) = q$ we conclude that $g_1(r) = f(r)$ for all r and

$$J_\pi(\xi^k) = \frac{1}{q} \sum_{x \in \mathbb{F}} \chi(x + \xi^{-k} x^3) \in \{0, \pm \sqrt{2/q}\} \text{ for all } 0 \leq k < q-1.$$

□

Small-dimensional line systems coming from Theorem 2 deserve special remarks.

Remark 4. If $q = 8$, then Theorem 2 gives a system of 63 $\{0, 1/2\}$ -angular lines in \mathbb{R}^7 which is isomorphic to the line system generated by the roots of the lattice E_7 .

Remark 5. If $q = 32$ then a system of 1023 $\{0, 1/4\}$ -angular lines in \mathbb{R}^{31} given by Theorem 2 is not unique. A nonisomorphic line system with the same parameters was found by the author of this article via a (nonexhaustive) computer assisted search.

5 Sets of biangular lines related to representations of $SL(2, \mathbb{F}_{3^k})$

In this section we investigate complex line systems associated with $(q-1)/2$ -dimensional representations of the groups $SL(2, \mathbb{F}_q)$ with q being odd. Values of Bessel functions associated with these representations are explicitly determined in [7]. However, the corresponding line systems are not investigated anywhere and seem to remain unknown.

When q is odd, the circle C contains an even number of elements and there exist a unique multiplicative character π_2 of C of order 2. Let $\theta = \{y_d \mid y_d \in C_d \text{ for every } d \in \mathbb{F}_q^*\}$ be some system of representatives of circles C_d . It turns out that the representation $(H(\mathbb{F}_q^*), \phi_{\pi_2})$ can be further decomposed into invariant subspaces.

Proposition 8 (See [20]). *Let q be odd. If $u, v \in \mathbb{F}_q^*$ such that $uv \in NQR_q$, then $T_{g, \pi_2, \theta}(u, v) = 0$ for all $g \in SL(2, \mathbb{F}_q)$.*

Corollary 1. *The representation $(H(\mathbb{F}_q^*), \phi_{\pi_2})$ can be decomposed into two subrepresentations, namely $(H(QR_q), \phi_{\pi_2}^+)$ and $(H(NQR_q), \phi_{\pi_2}^-)$ where*

$$\phi_{\pi_2}^+(g) = T_{g, \pi_2, \theta}^+ \in GL(H(QR_q)) \text{ such that } T_{g, \pi_2, \theta}^+(u, v) = T_{g, \pi_2, \theta}(u, v) \text{ for all } u, v \in QR_q$$

and

$$\phi_{\pi_2}^-(g) = T_{g, \pi_2, \theta}^- \in GL(H(NQR_q)) \text{ such that } T_{g, \pi_2, \theta}^-(u, v) = T_{g, \pi_2, \theta}(u, v) \text{ for all } u, v \in NQR_q.$$

Proposition 9 (Curtis, Shinoda [7]). *If q is odd, then for all $z \in \mathbb{F}_q^*$*

$$J_{\pi_2}(z) = -\pi_2'(z)G(\chi, \pi_2')(\chi(-2z) - \chi(2z))/q,$$

where π_2' is the unique multiplicative character of \mathbb{F}_q^* of order 2.

Remark 6. Propositions 8 and 9 together with (6) makes it possible to explicitly compute absolute values of the function J_{π_2} . Indeed on the circles C_d where $d \in QR_q$, if x_d is any of two possible elements such that $x_d^2 = d$, then

$$J_{\pi_2}(tx_d) = -\pi_2'(x_d)G(\chi, \pi_2)(\chi(-2x_d) - \chi(2x_d))\bar{\pi}_2(t)/q \text{ for all } t \in C.$$

Since

$$|\pi_2'(x_d)| = |\pi_2(t)| = 1, |G(\chi, \pi_2)| = \sqrt{q} \text{ and } |\chi(-2x_d) - \chi(2x_d)| \in \{|2\sin(2\pi k/p)| \mid k \in \mathbb{Z}/p\mathbb{Z}\}$$

for all $x_d \in \mathbb{F}_q^*$ and $t \in C$, the absolute values of the function J_{π_2} on the circles C_d where $d \in QR_q$ are always in the set

$$\mathcal{A}_{\pi_2} = \left\{ \left| \frac{2\sin(2\pi k/p)}{\sqrt{q}} \right| \mid k \in \mathbb{Z}/p\mathbb{Z} \right\}.$$

On the other hand, Proposition 8 implies that $J_{\pi_2}(z) = 0$ if $z \in C_d$, where $d \in NQR_q$. By combining two previous cases we conclude that the absolute values of the function J_{π_2} are always contained in the set \mathcal{A}_{π_2} .

In the forth chapter we defined line systems \mathcal{L}_{π} in the case when q is even through the sets of representatives Φ_{π} . Line systems $\mathcal{L}_{\pi, \theta}$ in the case where q is odd can be defined similarly through sets of representatives $\Phi_{\pi, \theta}$ by taking all row vectors of the matrices $T_{g, \pi, \theta}$ with omitted phases. In this paper we will need only one character, $\pi = \pi_2$, and define

$$\Phi_{\pi_2, \theta} = \{ \chi(ax)J_{\pi_2}(yx\sigma(yd)/c) \mid a \in \mathbb{F}_q, c, d \in \mathbb{F}_q^* \} \cup \{ \delta(x-y) \mid y \in \mathbb{F}_q^* \} \subset H(\mathbb{F}_q^*)$$

to be a system of representatives of complex lines $\mathcal{L}_{\pi_2, \theta}$. The following is an analogue to Proposition 3.

Proposition 10. *If $q > 3$ is odd, the set of representatives $\Phi_{\pi_2, \theta}$ defines a line system $\mathcal{L}_{\pi_2, \theta}$ consisting of $q^2 - 1$ complex lines in $H(\mathbb{F}_q^*)$.*

Proof. We show that the set $S \subset \text{SL}(2, \mathbb{F}_q)$ where

$$S = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right\} \cup \left\{ \begin{bmatrix} 0 & -a^{-1} \\ a & d \end{bmatrix} \mid a \in \mathbb{F}_q^*, d \in \mathbb{F}_q \right\}$$

generates $q^2 - 1$ complex lines with representatives in $\Phi_{\pi_2, \theta}$. The set of matrices

$$S_1 = \{g_1 g_2^{-1} \mid g_1 \neq g_2, g_1, g_2 \in S\}$$

contains only matrices which are not in B . Now, we have that

$$\langle T_{g_1, \pi_2, \theta}(u, \cdot), T_{g_2, \pi_2, \theta}(v, \cdot) \rangle = T_{g_1 g_2^{-1}, \pi_2, \theta}(u, v) \quad (15)$$

for all $g_1, g_2 \in G$ because of the group property of the matrices $T_{g, \pi_2, \theta}$ where $g \in G$. In particular, (15) holds for all $g_1, g_2 \in S$. When $g_1, g_2 \in S$ and $g_1 \neq g_2$, we have $g_1 g_2^{-1} \in S_1$ and the absolute values of the angles between row vectors of matrices $T_{g_1, \pi_2, \theta}$ and $T_{g_2, \pi_2, \theta}$ are contained in the set \mathcal{A}_{π_2} because of (5) for all matrices $T_{g, \pi_2, \theta}$ where $g \notin B$. If $g_1 = g_2$, we have that

$$\langle T_{g_1, \pi_2, \theta}(u, \cdot), T_{g_1, \pi_2, \theta}(v, \cdot) \rangle = 0$$

for all $u \neq v$ because of the unitarity of matrices $T_{g_1, \pi_2, \theta}$.

Since every element of the set \mathcal{A}_{π_2} is strictly smaller than 1 as long as $q > 3$, we conclude that every row vector of every matrix corresponding to an element in the set S defines a different line, together producing $q^2 - 1$ different lines in $H(\mathbb{F}_q^*)$. On the other hand, cardinality of $\Phi_{\pi_2, \theta}$ cannot exceed $q^2 - 1$ which means that cardinality of the set $\Phi_{\pi_2, \theta}$ is exactly $q^2 - 1$. \square

Now we are ready to combine Propositions 8 to 10 into the main result of this section.

Theorem 3. *Let $q = 3^k$ with $k \geq 2$ and let $\mathcal{L}_{\pi_2, \theta}$ be a system of $q^2 - 1$ lines in $H(\mathbb{F}_q^*)$ generated by the set $\Phi_{\pi_2, \theta}$. Then $\mathcal{A}_{\pi_2} = \{0, \sqrt{1/3^{k-1}}\}$, and half of the lines denoted by $\mathcal{L}_{\pi_2, \theta}^+$, are entirely in the subspace $H(QR_q)$ while the other half denoted by $\mathcal{L}_{\pi_2, \theta}^-$, are in the subspace $H(NQR_q)$.*

Proof. The angle set \mathcal{A}_{π_2} is given in the Remark 6. Propositions 8 and 10 can be used to conclude that half of the vectors of the each basis $T_{g, \pi_2, \theta}$ where $g \in S$ lies in $H(QR_q)$ while other half lies in $H(NQR_q)$. \square

Remark 7. Line systems in spaces $H(QR_q)$ and $H(NQR_q)$ are isomorphic since the unitary transform

$$T_{g(a), \pi_2, \theta} : H(\mathbb{F}_q^*) \rightarrow H(\mathbb{F}_q^*) \text{ where } g(a) = \begin{bmatrix} a & 0 \\ 0 & a^{-1} \end{bmatrix} \text{ and } a \in NQR_q,$$

gives an one-to-one correspondence between the lines in the systems $\mathcal{L}_{\pi_2, \theta}^+$ and $\mathcal{L}_{\pi_2, \theta}^-$. On the other hand, Remark 1 assures that line systems $\mathcal{L}_{\pi_2, \theta}$ with different choices of θ are isomorphic. Thus, for each $q = 3^k$ there is only one $(3^k - 1)/2$ -dimensional line system, up to isomorphism, produced by Theorem 3.

Remark 8. The line systems $\mathcal{L}_{\pi_2, \theta}^+$ and $\mathcal{L}_{\pi_2, \theta}^-$ achieve equality in the Special Bound (Theorem 1) since

$$N(\mathbb{C}^{(3^k - 1)/2}, 0, \sqrt{1/3^{k-1}}) = (3^{2k} - 1)/2.$$

Remark 9. If $q = 9$ then Theorem 3 produces a system of 40 $\{0, 1/\sqrt{3}\}$ -angular lines in \mathbb{C}^4 . An isomorphic line system is given (for example) in [3] and [12].

6 Acknowledgements

I would like to thank Professor Patric Östergård for valuable comments.

References

- [1] APPLEBY, M.; BENGTSOON, I.; DUMITRU, I. Dimension towers of SICs. I. Aligned SICs and embedded tight frames. *J. Math. Phys.* 58 (2017), 112201, 19pp.
- [2] BANNAI, E.; BANNAI, E. On antipodal spherical t -designs of degree s with $t \geq 2s - 3$. arXiv:0802.2905, 2008.
- [3] BEST, D.; KHARAGHANI, H.; RAMP, H. Mutually unbiased weighing matrices. *Des. Codes Cryptogr.* 76 (2015), 237–256.
- [4] CAHILL, J.; CASAZZA, P.-G.; HAAS, J. I.; TREMAIN J. Constructions of biangular tight frames and their relationships with equiangular tight frames, in: Kim Y., Narayan S. K., Picioroaga G., Weber E. S., (Eds), *Frames and harmonic analysis*, pp. 1–19, *Contemp. Math.*, 706, Amer. Soc., Providence, RI, 2018
- [5] CARLITZ, L. Evaluation of some exponential sums over a finite field. *Math. Nachr.* 96 (1980), 319–339.
- [6] CASAZZA, P. G.; FERZANNIA, A.; HAAS, J. I.; TRAN, T. T. Toward the classification of biangular harmonic frames. *Appl. Comput. Harmon. Anal.* 46 (2019), 544–568.
- [7] CURTIS, C. W.; SHINODA, K. Unitary Kloostermann sums and the Gelfand-Graev Representation of GL_2 . *J. Algebra* 216 (1999), 431–447.
- [8] DAVENPORT, H.; HASSE, H. Die Nullstellen der Kongruenzzetafunktionen in gewissen zyklischen Fällen. *J. Reine Angew. Math*, 172 (1935), 151–182.
- [9] DELSARTE, P.; GOETHALS J.-M.; SEIDEL, J. J. Bounds for systems of lines and Jacobi polynomials. *Philips Research Reports*, 30 (1975), 91–105.
- [10] FICKUS, M.; MIXON D. G. Tables of the existence of equiangular tight frames. arXiv:1504.00253, 2015.
- [11] FICKUS, M.; MIXON D. G.; JASPER, J. Equiangular Tight Trames from Hyperovals. *IEEE Trans. Inform. Theory* 62 (2016), no. 9, 5225–5236.
- [12] HOGGAR, S. G. t -designs in projective spaces. *European J. Combin.* 3 (1982), 233–254.
- [13] IVERSON, J. J.; JASPER, J.; MIXON, D. G. Optimal line packings from finite group actions. *Forum Math. Sigma* 8 (2020), e6, 40 pp.
- [14] KANTOR, W. M. Codes, quadratic forms and finite geometries, in: Calderbank R., (Ed), *Different aspects of coding theory* (San Francisco, CA, 1995), vol. 50 of *Proc. Sympos. Appl. Math.*, pp. 153–177, Amer. Math. Soc., Providence, RI, 1995.
- [15] KOPP, G. S. SIC-POVMs and the Stark conjectures. *Int. Math. Res. Not.* Published online 2019, DOI:10.1093/imrn/rnz153.
- [16] LEVENSHTAIN, V. I. Bounds of maximal capacity of a code with a limited scalar product modulus. (Russian) *Dokl. Akad. Nauk SSSR.* 263 (1982), 1303–1308.
- [17] NOZAKI, H. A generalization of Larman–Rogers–Seidel’s theorem. *Discrete Math.* 311 (2011), 792–799.
- [18] STROMER, T.; HEATH, R. W. Grassmannian frames with applications to coding and communications. *Appl. Comput. Harmon. Anal.* 14 (2003), 257–275.
- [19] SUSTIK, M. A.; TROPP, J. A.; DHILLON, I. S.; HEATH, R. W., JR. On the existence of equiangular tight frames. *Linear Algebra Appl.* 426 (2007), 619–635.
- [20] TANAKA, S. Construction and classification of irreducible representations of special linear group of the second order over a finite field. *Osaka Math. J.* 4 (1967), 65–84.

- [21] WELCH, L. Lower bounds on the maximum cross correlation of signals. *IEEE Trans. Inf. Theory*, 20 (1974), 397–399.
- [22] WOOTERS, WILLIAM K. FIELDS, BRIAN D. Optimal state-determination by mutually unbiased measurements. *Ann. Physics* 191 (1989), 363–381.
- [23] ZAUNER, G.: *Quantendesigns: Grundzüge einer nichtkommutativen Designtheorie*. PhD thesis, University of Vienna, Austria, 1999.