

DISCRETE MKDV EQUATION VIA DARBOUX TRANSFORMATION

JOSEPH CHO, WAYNE ROSSMAN, AND TOMOYA SENO

ABSTRACT. We introduce an efficient route to obtaining the discrete potential mKdV equation emerging from a particular discrete motion of discrete planar curves.

1. INTRODUCTION

In this work, we illuminate the relationship between discrete and semi-discrete potential modified Korteweg-de Vries (mKdV) equation on the one hand, and on the other, the transformation theory of the smooth potential mKdV equation via the use of infinitesimal Bianchi cubes of Darboux transformations keeping arc-length polarization.

Studies of the discrete and semi-discrete potential mKdV equation, starting with a discrete curve having an associated discrete frame, can be found in [4, 5, 6, 7]. There discrete or continuous isoperimetric deformation was used to produce certain motions of a curve, and then the discrete frames provided the means to produce the desired equations. In fact, the discrete and semi-discrete potential mKdV equation appear in the context of the Bäcklund transformation and permutability of smooth potential mKdV equations [8].

To exploit this, we consider a 3-dimensional system with two discrete parameters and one smooth parameter, where the discrete parameters represent a Darboux transformation of a discrete curve, while the smooth parameter represents the infinitesimal Darboux transformation [3]. With this combination of discrete and smooth parameters, we can then take advantage of the result in [2] regarding the permutability between Darboux and infinitesimal Darboux transformations (infinitesimal Bianchi cube). By restricting to the Darboux transformations that preserve arc-length polarization, we can reinterpret this system in terms of the transformation and permutability of smooth potential mKdV equations, giving us a calculation-free approach to producing the fully discrete potential mKdV equation without any use of frames (see the primary result, Theorem 3.2). We note that such pursuit is greatly aided by interpreting the potential functions appearing in both the discrete and semi-discrete equations as a single geometric quantity encoded in the system, as done in [3].

2. DARBOUX AND INFINITESIMAL DARBOUX TRANSFORMATIONS

Let $\Sigma \subset \mathbb{Z}$ be a discrete interval, and let μ be a strictly positive or negative function on (unoriented) edges of Σ . Recall from [3], that a *discrete polarized curve* is a discrete curve $x : (\Sigma, \frac{1}{\mu}) \rightarrow \mathbb{C}$ defined on a discrete polarized domain $(\Sigma, \frac{1}{\mu})$ whose polarization is given by some such function μ .

2.1. Darboux transformations of discrete polarized curves. We first define Darboux transformations of discrete polarized curves.

Definition 2.1. Two discrete polarized curves $x, \hat{x} : (\Sigma, \frac{1}{\mu}) \rightarrow \mathbb{C}$ are called a *Darboux pair with parameter $\hat{\mu}$* if, on every edge (ij) ,

$$(2.1) \quad \text{cr}(x_i, x_j, \hat{x}_j, \hat{x}_i) = \frac{x_i - x_j}{x_j - \hat{x}_j} \frac{\hat{x}_j - \hat{x}_i}{\hat{x}_i - x_i} = \frac{\hat{\mu}}{\mu},$$

for some non-zero constant $\hat{\mu} \in \mathbb{R} \setminus \{0\}$. We call one of the curves a *Darboux transform* of the other.

A Darboux transformation is determined by the choice of the parameter $\hat{\mu}$ and an initial condition \hat{x}_i at some vertex $i \in \Sigma$.

Recalling that a discrete curve $x : (\Sigma, \frac{1}{\mu}) \rightarrow \mathbb{C}$ is arc-length polarized if $|x_i - x_j|^2 = \frac{1}{\mu_{ij}}$ on every edge (ij) , we have the following proposition.

Proposition 2.2. *Let $x, \hat{x} : (\Sigma, \frac{1}{\mu}) \rightarrow \mathbb{C}$ be a Darboux pair with parameter $\hat{\mu}$ where x is arc-length polarized. Then \hat{x} is also arc-length polarized if and only if $|x_i - \hat{x}_i|^2 = \frac{1}{\hat{\mu}}$ at some vertex $i \in \Sigma$.*

Proof. Assume first that $|x_i - \hat{x}_i|^2 = \frac{1}{\hat{\mu}}$ at some vertex $i \in \Sigma$. Then on an edge (ij) , the definition of Darboux pair with parameter $\hat{\mu}$ (2.1) tells us

$$\hat{x}_j - \hat{x}_i = \frac{\hat{\mu}(\hat{x}_i - x_i)(x_j - \hat{x}_i)}{\mu_{ij}(x_i - x_j) + \hat{\mu}(\hat{x}_i - x_i)}.$$

Then a computation gives

$$|\mu_{ij}(x_i - x_j) + \hat{\mu}(\hat{x}_i - x_i)|^2 = \mu_{ij}\hat{\mu}|x_j - \hat{x}_i|^2;$$

hence,

$$|\hat{x}_j - \hat{x}_i|^2 = \frac{\hat{\mu}|x_j - \hat{x}_i|^2}{\mu_{ij}\hat{\mu}|x_j - \hat{x}_i|^2} = \frac{1}{\mu_{ij}}.$$

We can prove the converse claim similarly, by switching the roles of \hat{x}_j and x_i . \square

2.2. Permutability. We now discuss the permutability between infinitesimal Darboux transformation and Darboux transformation of a discrete polarized curve. Let I be an interval in the reals, and recall from [3, Definition 2.7], that given a discrete polarized curve x , we have $f : \Sigma \times I \rightarrow \mathbb{C}$ is an infinitesimal Darboux transformation of x with parameter function m , if $f_i(s_0) = x_i$ for some $s_0 \in I$, and on every edge (ij) ,

$$\frac{f'_i f'_j}{(f_i - f_j)^2} = \frac{\mu_{ij}}{m},$$

Equivalently, we have that on every edge (ij) , $f_i(s), f_j(s) : (I, \frac{ds^2}{m}) \rightarrow \mathbb{C}$ are a Darboux pair of smooth polarized curves with parameter μ_{ij} (see [2, Definition and Lemma 2.2]).

If \hat{x} is a Darboux transform of x with parameter $\hat{\mu}$, then the permutability of Darboux transformations of smooth polarized curves in [2, Theorem 2.8] tells us that there exists an infinitesimal Darboux transformation $\hat{f} : \Sigma \times I \rightarrow \mathbb{C}$ of \hat{x} with parameter function m so that $\hat{f}_i(s_0) = \hat{x}_i$ for some s_0 , and for any fixed $s \in I$,

$$\text{cr}(f_i(s), f_j(s), \hat{f}_j(s), \hat{f}_i(s)) = \frac{\hat{\mu}}{\mu_{ij}},$$

that is, the two discrete curves $f(s_1)$ and $\hat{f}(s_1)$ for any fixed $s_1 \in I$ are a Darboux pair with parameter $\hat{\mu}$. Thus, we have the permutability between Darboux transformation and infinitesimal Darboux transformation of a discrete polarized curve:

Proposition 2.3. *Let $f : \Sigma \times I \rightarrow \mathbb{C}$ be an infinitesimal Darboux transformation of a discrete polarized curve $x : (\Sigma, \frac{1}{\mu}) \rightarrow \mathbb{C}$ with parameter function m and let \hat{x} be a Darboux transform of x with parameter $\hat{\mu}$. Then there is an infinitesimal Darboux transformation \hat{f} of \hat{x} with parameter function m so that, for any $s \in I$, $\hat{f}(s)$ is a Darboux transform of $f(s)$ with parameter $\hat{\mu}$.*

Using this, we can also obtain the following result.

Proposition 2.4. *Let $f_i : (I, \frac{ds^2}{m}) \rightarrow \mathbb{C}$ be a polarized curve with Darboux transforms f_j and \hat{f}_i with parameters μ_{ij} and $\hat{\mu}$, respectively, and suppose that \hat{f}_j is the simultaneous Darboux transform of f_j and \hat{f}_i with parameters $\hat{\mu}$ and μ_{ij} , respectively. If f_i , f_j , and \hat{f}_i are arc-length polarized, then \hat{f}_j is also arc-length polarized.*

Proof. Assuming f_i , f_j , and \hat{f}_i are arc-length polarized, then by [3, Proposition 2.6], we have that $|f_i - f_j|^2 = \frac{1}{\mu_{ij}}$ and $|\hat{f}_i - f_i|^2 = \frac{1}{\hat{\mu}}$ for any $s \in I$. Fixing any $s_0 \in I$, we have that the two discrete curves $f(s_0)$ and $\hat{f}(s_0)$ are a Darboux pair, where $f(s_0)$ is discrete arc-length polarized. Hence, applying Proposition 2.2 tells us that $\hat{f}(s_0)$ is also discrete arc-length polarized, i.e., $|\hat{f}_i - \hat{f}_j|^2 = \frac{1}{\mu_{ij}}$. Finally, applying [3, Proposition 2.6] again tells us that \hat{f}_j is arc-length polarized. \square

3. DISCRETE POTENTIAL MKDV EQUATION

In this section, we use the aforementioned permutability of Darboux transformations in Propositions 2.3 and 2.4 to introduce an efficient route to obtaining the discrete potential mKdV equation, using the well-known permutability theorems (cf. [1, 8]). To do this, consider the discrete motion $x : \Sigma \times \tilde{\Sigma} \rightarrow \mathbb{C}$, denoted by $x(n, k) = x_n^k$, of a discrete planar curve x^{k_0} defined as in [7, §2.2], whose compatibility condition results in the discrete potential mKdV equation. Thus for any $(n, k) \in \Sigma \times \tilde{\Sigma}$, we have that

$$(3.1) \quad |x_{n+1}^k - x_n^k| =: a_n \quad \text{and} \quad |x_n^{k+1} - x_n^k| =: b_k$$

are constant in k and n , respectively.

Without loss of generality, assume $x_n^k = 0$, $x_{n+1}^k = a_n$ and $x_n^{k+1} = b_k e^{i\theta}$ for some $\theta \in \mathbb{R}$. Excluding the solution $a_n + b_k e^{i\theta}$ obtained via translation, x_{n+1}^{k+1} is uniquely determined with cross-ratios satisfying

$$\text{cr}(x_n^k, x_{n+1}^k, x_{n+1}^{k+1}, x_n^{k+1}) = \frac{a_n^2}{b_k^2}.$$

Hence, using Proposition 2.2 we have the following result:

Proposition 3.1. *Let $x : \Sigma \times \tilde{\Sigma} \rightarrow \mathbb{C}$ be a discrete motion of a discrete planar curve $x^{k_0} : \Sigma \rightarrow \mathbb{C}$ such that (3.1) holds. Giving x^{k_0} the arc-length polarization so that $\mu_{(n,n+1)} = \frac{1}{a_n^2}$, we have that $x^k, x^{k+1} : (\Sigma, \frac{1}{\mu}) \rightarrow \mathbb{C}$ are a Darboux pair with parameter $\frac{1}{b_k^2}$ keeping arc-length polarization, for any $k \in \tilde{\Sigma}$.*

Using Proposition 3.1, we can view the discrete motion $x : \Sigma \times \tilde{\Sigma} \rightarrow \mathbb{C}$ as the image of successive Darboux transformations keeping the arc-length polarization. Then Propositions 2.3 and 2.4 tell us that there exists smooth arc-length polarized curves $f : \Sigma \times \tilde{\Sigma} \times I \rightarrow \mathbb{C}$ so that $f_n^k, f_{n+1}^k : (I, \frac{ds^2}{m}) \rightarrow \mathbb{C}$ are Darboux pairs with parameter $\frac{1}{a_n^2}$ while $f_n^{k+1}, f_{n+1}^{k+1} : (I, \frac{ds^2}{m}) \rightarrow \mathbb{C}$ are Darboux pairs with parameter $\frac{1}{b_k^2}$.

Therefore, the tangential angles $\theta_n^k, \theta_{n+1}^k, \theta_n^{k+1}$ and θ_{n+1}^{k+1} of $f_n^k, f_{n+1}^k, f_n^{k+1}$ and f_{n+1}^{k+1} , respectively, satisfy the semi-discrete potential mKdV equations via [3, Theorem 3.2]:

$$\begin{aligned} \left(\frac{\theta_n^k + \theta_{n+1}^k}{2} \right)' &= \frac{2}{a_n} \sin \left(\frac{\theta_{n+1}^k - \theta_n^k}{2} \right), & \left(\frac{\theta_n^{k+1} + \theta_{n+1}^{k+1}}{2} \right)' &= \frac{2}{b_k} \sin \left(\frac{\theta_{n+1}^{k+1} - \theta_n^{k+1}}{2} \right), \\ \left(\frac{\theta_{n+1}^{k+1} + \theta_n^{k+1}}{2} \right)' &= \frac{2}{a_n} \sin \left(\frac{\theta_{n+1}^{k+1} - \theta_n^{k+1}}{2} \right), & \left(\frac{\theta_{n+1}^k + \theta_n^k}{2} \right)' &= \frac{2}{b_k} \sin \left(\frac{\theta_{n+1}^k - \theta_n^k}{2} \right). \end{aligned}$$

These equations are well-known partial differential equations that define Bäcklund transformations of the smooth potential mKdV equation as seen in [8, Equations (7), (8)]. Using these equations, permutability of the transformation was obtained in [8, Equation (9)] (see also [1]):

$$\tan \left(\frac{\theta_{n+1}^{k+1} - \theta_n^k}{4} \right) = \frac{b_k + a_n}{b_k - a_n} \tan \left(\frac{\theta_n^{k+1} - \theta_{n+1}^k}{4} \right),$$

which is the discrete potential mKdV equation. Summarizing, we have:

Theorem 3.2. *The infinitesimal Bianchi cubes of Darboux transformations keeping arc-length polarization in both the smooth and discrete directions yields a 1-parameter family of solutions of the discrete mKdV equation.*

Acknowledgements. The authors gratefully acknowledge the support from the JSPS/FWF Bilateral Joint Project I3809-N32 ‘‘Geometric shape generation’’ and JSPS Grant-in-Aids for: JSPS Fellows 19J10679, Scientific Research (C) 15K04845, (C) 20K03585 and (S) 17H06127 (P.I.: M.-H. Saito).

REFERENCES

- [1] L. Bianchi, *Sulla trasformazione di Bäcklund per le superficie pseudosferiche*, Rend. Lincei **5** (1892), no. 1, 3–12.
- [2] F. E. Burstall, U. Hertrich-Jeromin, C. Müller, and W. Rossman, *Semi-discrete isothermic surfaces*, Geom. Dedicata **183** (2016), 43–58. MR3523116
- [3] J. Cho, W. Rossman, and T. Seno, *Infinitesimal Darboux transformation and semi-discrete mKdV equation* (2020), available at arXiv:2010.07846.
- [4] J.-i. Inoguchi, K. Kajiwara, N. Matsuura, and Y. Ohta, *Explicit solutions to the semi-discrete modified KdV equation and motion of discrete plane curves*, J. Phys. A **45** (2012), no. 4, 045206, 16. MR2874242
- [5] J.-i. Inoguchi, K. Kajiwara, N. Matsuura, and Y. Ohta, *Motion and Bäcklund transformations of discrete plane curves*, Kyushu J. Math. **66** (2012), no. 2, 303–324. MR3051339
- [6] S. Kaji, K. Kajiwara, and H. Park, *Linkage mechanisms governed by integrable deformations of discrete space curves*, Nonlinear Systems and Their Remarkable Mathematical Structures (N. Euler and M. C. Nucci, eds.), Vol. 2, Chapman and Hall/CRC, New York, 2019, pp. 356–381.
- [7] N. Matsuura, *Discrete KdV and discrete modified KdV equations arising from motions of planar discrete curves*, Int. Math. Res. Not. IMRN **8** (2012), 1681–1698. MR2920827
- [8] M. Wadati, *Bäcklund transformation for solutions of the modified Korteweg-de Vries equation*, J. Phys. Soc. Japan **36** (1974), no. 5, 1498.

(Joseph Cho) INSTITUTE OF DISCRETE MATHEMATICS AND GEOMETRY, TU WIEN, WIEDNER HAUPTSTRASSE 8-10/104, 1040 WIEN, AUSTRIA

Email address: `jcho@geometrie.tuwien.ac.at`

(Wayne Rossman) DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, KOBE UNIVERSITY, 1-1 ROKKODAI-CHO, NADA-KU, KOBE 657-8501, JAPAN

Email address: `wayne@math.kobe-u.ac.jp`

(Tomoya Seno) DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, KOBE UNIVERSITY, 1-1 ROKKODAI-CHO, NADA-KU, KOBE 657-8501, JAPAN

Email address: `tseo@math.kobe-u.ac.jp`