

# Connectivity keeping path-stars in $k$ -connected graphs

Meng Ji

College of Mathematical Science

Tianjin Normal University, Tianjin, China

Email: jimengecho@163.com

## Abstract

W. Mader [J graph Theory 65 (2010),61-69] conjectured that for any tree  $T$  of order  $m$ , every  $k$ -connected graph  $G$  with  $\delta(G) \geq \lfloor \frac{3k}{2} \rfloor + m - 1$  contains a tree  $T' \cong T$  such that  $G - V(T')$  remains  $k$ -connected. The conjecture is open for  $k \geq 4$  by now. In this paper, the author shows that Mader's conjecture is true for any  $k$ -connected graph when  $T$  is a path-star and extends Mader's result in [J graph Theory 65 (2010),61-69].

**Keywords:** Connectivity; Path-star;  $k$ -connected graph

**AMS subject classification 2010:** 05C05,05C40.

## 1 Introduction

In this paper, all the graphs are finite, undirected simple. For graph-theoretical terminology and notations not defined here, we follow [1]. Some basic symbols and definitions ones are needed to be introduced. The minimum degree and the connectivity number of a graph  $G$  is denoted by  $\delta(G)$  and  $\kappa(G)$ , respectively. the vertex set and edge set of  $G$  is denoted by  $V(G)$  and  $E(G)$ , respectively. For a subgraph  $H \subseteq G$  and a subset  $V' \subseteq V(G)$ , write  $V' \cap H$  for  $V' \cap V(H)$ . For any vertex  $x \in G$ , denote the set of neighbors of  $x$  in  $G$  by  $N_G(x)$  and for a subgraph  $H \subseteq G$ , define  $N_G(H) := \bigcup_{x \in H} N_G(x) - V(H)$ . For  $H \subseteq G$ ,  $G(H)$  stands for the subgraph induced by  $H$  in  $G$ . For  $H \subseteq G$ , set  $\delta_G(H) := \min_{x \in H} d_G(x)$ . For any edge  $xy \in E(G)$  and  $H \subseteq G$ , write  $H \cup xy$  for the subgraph  $(V(H) \cup \{x, y\}, E(H) \cup xy)$  of  $G$ . Let  $x, y$ -path

be a path  $P$  from  $x$  to  $y$ . For  $u, v \in P$ , let  $P[u, v] = P[v, u]$  be the subpath of  $P$  between  $u$  and  $v$ , and  $P[u, v)$  means  $= P[u, v] - v$ . Denotes the complete graph on vertex set  $S$  by  $K(S)$ .

Let  $G$  be a graph. A separating set  $S \subseteq G$  denotes a vertex cut-set, and the cardinality of a minimum separating set  $S$  is denoted by  $|S| = \kappa(G)$ . If  $S$  is a minimum separating set of the graph  $G$ , then the union components  $F$  of  $G - S$  with  $G - S - F \neq \emptyset$  is called a fragment  $F$  to  $S$ , and the complementary fragment  $G - (S \cup V(F))$  is denoted by  $\bar{F}$ . If a fragment of  $G$  does not contain any other fragments of  $G$ , then it is called an end of  $G$ . Clearly, every graph contains an end except for the complete graphs. For a fragment  $F$  of  $G$  to  $S$ , it definitely follows  $S = N_G(F)$ . For  $S \subseteq G$ , denote the complete graph induced by  $S$  by  $K(S)$  and the graph  $G \cup K(S)$  denoted by  $G[S]$ .

In 1972, Chartrand, Kaigars and Lick [2] proved that *every  $k$ -connected graph  $G$  with  $\delta(G) \geq \lfloor \frac{3k}{2} \rfloor$  has a vertex  $x$  with  $\kappa(G - x) \geq k$* . Nearly forty years later, Fujita and Kawarabayashi [4] extended the result, that is, every  $k$ -connected graph  $G$  with  $\delta(G) \geq \lfloor \frac{3k}{2} \rfloor + 2$  has an edge  $xy$  such that  $G - \{x, y\}$  remains  $k$ -connected. Furthermore, Fujita and Kawarabayashi posed the following conjecture.

**Conjecture 1.1** *For all positive integers  $k, m$ , there is a (least) non-negative integer  $f_k(m)$  such that every  $k$ -connected graph  $G$  with  $\delta(G) \geq \lfloor \frac{3k}{2} \rfloor + f_k(m) - 1$  contains a connected subgraph  $W$  of order  $m$  such that  $G - V(W)$  is still  $k$ -connected.*

In 2010, Mader [9] proved Conjecture 1.1 is true if  $f_k(m) = m$  and  $W$  is a path. Mader [9] proposed the following conjecture. The author will show it by using the Mader's method.

**Conjecture 1.2** [9] *For every positive integer  $k$  and finite tree  $T$  of order  $m$ , every  $k$ -connected finite graph  $G$  with minimum degree  $\delta(G) \geq \lfloor \frac{3k}{2} \rfloor + |T| - 1$  contains a subgraph  $T' \cong T$  such that  $G - V(T')$  remains  $k$ -connected.*

The conjecture has been showed for some special cases, such as for  $T$  is a path [9], for  $k = 1$  [3], 2-connected graph  $G$  and a star  $T$ , double star, path-star, path-double-star or  $T$  with order at most 8 and so on [5, 6, 8, 11, 12]. In addition, Mader [10] showed that the conjecture is true if  $\delta \geq 2(k - 1 + m)^2 + m - 1$ . Very recently, Hong and Liu [7] proved the correction of the conjecture for  $k = 2, 3$ . In this paper, the author will prove Mader's conjecture is true if  $T$  is a path-star.

Set  $\mathcal{F}_k(m)$  as all pairs  $(G, C)$ , where  $G$  is a  $k$ -connected graph with  $|G| \geq k + 1$ , the complete subgraph  $C \subseteq G$  satisfies  $|C| = k$  and  $\delta_G(G - V(C)) \geq \lfloor \frac{3k}{2} \rfloor + m - 1$ .

All  $(G, C) \in \mathcal{F}_k(m)$  with  $\kappa(G) \geq k + 1$  are denoted by  $\mathcal{F}_k^+(m)$ . The path-star tree  $T_{PS}(r, m - r)$  is obtained from the disjoint union of a path with order  $r$  and a star with order  $m - r$ , by connecting one end-vertex of the path with the center-vertex  $s_0$  of the star, in which let the center vertex of the star be the root vertex of the path-star. If  $r = 1$  or  $m - r = 1$ , then  $T_{PS}(r, m - r)$  is a path.

**Theorem 1.3** *For any  $(G, C) \in \mathcal{F}_k^+(m)$  and  $s_0 \in G - V(C)$ , there is a path-star  $T_{PS}(r, m - r) \subseteq G - V(C)$  of order  $m$  rooted at  $s_0$  such that  $\kappa(G - V(T_{PS}(r, m - r))) \geq k$ .*

From Theorem 1.3, we can prove our main theorem.

**Theorem 1.4** *Every  $k$ -connected graph  $G$  with  $\delta(G) \geq \lfloor \frac{3k}{2} \rfloor + m - 1$  for positive integers  $k, m, r$ , contains a path-star tree  $T_{PS}(r, m - r)$  such that  $G - V(T_{PS}(r, m - r))$  remains  $k$ -connected.*

In order to show the above theorems, we need some lemmas proved by Mader [9].

**Lemma 1.5** [9] *Let  $G$  be a graph with  $\kappa(G) = k$  and let  $F_i$  be a fragment of  $G$  to  $S_i$  for  $i = 1, 2$ . Then it follows that if  $F_1$  is an end of  $G$  with  $F_1 \geq 2$ , then  $G[S_1] - V(\bar{F}_1)$  is  $(k + 1)$ -connected.*

**Lemma 1.6** [9] *Let  $G$  be a  $k$ -connected graph and let  $S$  be a separating set of  $G$  with  $|S| = k$ . Then the following holds.*

- (a) *For every fragment  $F$  of  $G$  to  $S$ ,  $G[S] - V(F)$  is  $k$ -connected.*
- (b) *Assume  $\delta(G) \geq \lfloor \frac{3k}{2} \rfloor + m - 1$  and let  $F$  be a fragment of  $G$  to  $S$ . If  $W \subseteq G - (S \cup V(F))$  has order at most  $m$  and  $\kappa(G[S] - V(F \cup W)) \geq k$  holds, then also  $\kappa(G - V(W)) \geq k$  holds.*
- (c) *Assume  $(G, C) \in \mathcal{F}_k(m)$  and let  $F$  be a fragment of  $G$  to  $S$  with  $C \subseteq G(F \cup S)$ . If  $W \subseteq G - (S \cup V(F))$  has order at most  $m$  and  $\kappa(G[S] - V(F \cup W)) \geq k$  holds, then also  $\kappa(G - V(W)) \geq k$  holds.*

**Lemma 1.7** [9] *For all  $(G, C) \in \mathcal{F}_k^+(m)$  and  $p_0 \in G - V(C)$ , there is a path  $P \subseteq G - V(C)$  of length  $m - 1$  starting from  $p_0$ , such that  $\kappa(G - V(P)) \geq k$  holds.*

## 2 Proofs of main results

**Proof of Theorem 1.3.** We perform an induction on the order  $n$  of the graph  $G$  for the theorem. The triangle  $K_3$  is the smallest graph for  $k = m = 1$  and  $K_{\lfloor \frac{3k}{2} \rfloor + m}$

is the smallest graph  $G$  in  $(G, C) \in \mathcal{F}_k^+(m)$  for  $k \geq 2$  or  $m \geq 2$ . For  $k = 1$ ,  $K_2$  is 1-connected and for  $k \geq 2$ , we have  $\lfloor \frac{3k}{2} \rfloor \geq k + 1$ . Hence, it is true for  $(G, C) \in \mathcal{F}_k^+(m)$  with smallest  $G$  and with every larger complete graph  $G$ . Now we assume that  $G$  is a graph of LEAST order such that  $(G, C) \in \mathcal{F}_k^+(m)$  is a counterexample to Theorem 1.3 for  $k$  and for some  $C \subseteq G$ ,  $m$  and  $r$ . As we have seen above,  $G$  is not complete and  $|G| \geq \lfloor \frac{3k}{2} \rfloor + m + 1$ .

From Lemma 1.7, there exists a path  $P \subseteq G - V(C)$  of length  $m - 1$  starting from  $s_0$  such that  $\kappa(G - V(P)) \geq k$  holds. Hence we have  $\kappa(G - s_0 p_1) \geq k$ , where  $s_0 p_1$  is a subpath of  $P$  and also a path-star  $T_{PS}(1, 1)$ . Thus, fix  $r \geq 1$  arbitrarily, choose a path-star  $T_{PS}(r, t) \subseteq G - V(C)$  rooted at  $s_0$  of MAXIMAL order with  $2 \leq |T_{PS}(r, t)| < m$  such that  $\kappa(G - T_{PS}(r, t)) \geq k$ . Set  $G \setminus T_{PS}(r, t) := G - V(T_{PS}(r, t))$  and let  $\{s_1, s_2, \dots, s_{t-1}\}$  be the leaf-vertex set of the star in  $T_{PS}(r, t)$  and let  $P[p_1, p_r] := p_1 p_2 \dots p_r$  and connect  $p_1$  with  $s_0$ . Since  $|N_G(s_0) \cap (G - V(C \cup T_{PS}(r, t)))| \geq \lfloor \frac{3k}{2} \rfloor + m - 1 - k - (m - 2) = \lfloor \frac{k}{2} \rfloor + 1$ , there is a neighbor  $s' \in V(G) - V(C \cup T_{PS}(r, t))$ . Since  $T_{PS}(r, t) \cup s' s_0$  is a path-star rooted at  $s_0$  of order  $|T_{PS}(r, t)| + 1 = |T_{PS}(r, t + 1)| \leq m$ , it follows that  $\kappa(G - V(T_{PS}(r, t + 1))) < k$  by the choice of  $T_{PS}(r, t)$ . Hence we have  $\kappa(G \setminus T_{PS}(r, t)) = k$ . Since  $|G \setminus T_{PS}(r, t)| \geq k + 2$ , there exists an end  $E$  of  $G \setminus T_{PS}(r, t)$  with  $E \cap C = \emptyset$ . We define  $S := N_{G \setminus T_{PS}(r, t)}(E)$ , thus we have  $|S| = k$ . And let  $\bar{E} = G - S - E$  in  $G \setminus T_{PS}(r, t)$ . If  $|E| = 1$  holds, then for any  $x \in E$  we get  $k = d_{G \setminus T_{PS}(r, t)}(x) \geq \lfloor \frac{3k}{2} \rfloor + m - 1 - |T_{PS}(r, t)| \geq \lfloor \frac{3k}{2} \rfloor$ , hence  $k = 1$ ,  $|T_{PS}(r, t)| = m - 1$ . Since  $\delta(G) \geq \lfloor \frac{3k}{2} \rfloor + m - 1$ , we have  $V(T_{PS}(r, m - r - 1)) \subseteq N_G(x)$ . Hence, the order of the path-star  $T_{PS}(r, m - r) := T_{PS}(r, m - r - 1) \cup s_0 x \subseteq G - V(C)$  (or  $T_{PS}(r, m - r) := T_{PS}(r, m - r - 1) \cup p_r x \subseteq G - V(C)$ ) is  $m$  and  $G - V(T_{PS}(r, m - r)) = (G \setminus T_{PS}(r, m - r)) - x$  is 1-connected. Now we get  $|E| \geq 2$  and the graph  $(G \setminus T_{PS}(r, t))[S] - V(\bar{E})$  is  $(k + 1)$ -connected from Lemma 1.5. From above assumption, we know that  $\kappa(G) > k = \kappa(G \setminus T_{PS}(r, t))$ , thus  $N_G(T_{PS}(r, t)) \cap E \neq \emptyset$ . We denote the closest vertex to  $p_r$  on  $T_{PS}(r, t)$  by  $y$  with  $N_G(y) \cap E \neq \emptyset$ . Let  $q \in N_G(y) \cap E$ . And then we discuss it by two cases.

Case 1.  $y \in \{p_1, p_2, \dots, p_r, s_0\}$ .

Let  $\bar{P} = P[p_r, y]$ . Consider the graph  $G \setminus (T_{PS}(r, t) \setminus \bar{P}) := G - V(T_{PS}(r, t) \setminus \bar{P})$ . Since  $|N_G(x) \cap (G \setminus T_{PS}(r, t))| \geq \lfloor \frac{3k}{2} \rfloor + m - 1 - (|T_{PS}(r, t)| - 1) \geq \lfloor \frac{3k}{2} \rfloor + 1 \geq k$  for any  $x \in V(\bar{P})$ , then  $\kappa(G \setminus (T_{PS}(r, t) \setminus \bar{P})) \geq k$ . As  $y$  is the closest vertex to  $p_r$  on  $T_{PS}(r, t)$ , we have  $N_G(\bar{P}) \cap E = \emptyset$ . Naturally,  $S$  is also a minimum separating set of  $G \setminus (T_{PS}(r, t) \setminus \bar{P})$ , and  $E$  is an end of  $G \setminus (T_{PS}(r, t) \setminus \bar{P})$ . From Lemma 1.5,  $(G \setminus (T_{PS}(r, t) \setminus \bar{P}))[S](E \cup S) = (G \setminus T_{PS}(r, t))[S] - V(\bar{E})$  is  $(k + 1)$ -connected. Furthermore, it follows that  $C \subseteq G \setminus (T_{PS}(r, t) \setminus \bar{P}) - V(\bar{E})$  and  $\delta_{G \setminus (T_{PS}(r, t) \setminus \bar{P})}(G \setminus (T_{PS}(r, t) \setminus \bar{P}) - V(C)) \geq \lfloor \frac{3k}{2} \rfloor + m - 1 - (|T_{PS}(r, t)| - |\bar{P}|) \geq \lfloor \frac{3k}{2} \rfloor + |\bar{P}|$ . Hence,

$(G \setminus (T_{PS}(r, t) \setminus \bar{P}), C) \in \mathcal{F}_k(|\bar{P}| + 1)$  and  $(G \setminus (T_{PS}(r, t) \setminus \bar{P}[S](E \cup S)), K(S)) \in \mathcal{F}_k^+(|\bar{P}| + 1)$ . From Lemma 1.7, there is a path  $Q \subseteq E$  of order  $|\bar{P}| + 1$  starting from  $q$  such that  $G \setminus (T_{PS}(r, t) \setminus \bar{P}[S](E \cup S) - V(Q))$  is  $k$ -connected. Now apply Lemma 1.6(c) to  $(G \setminus (T_{PS}(r, t) \setminus \bar{P}), C) \in \mathcal{F}_k(|\bar{P}| + 1)$ , then  $\kappa(G \setminus (T_{PS}(r, t) \setminus \bar{P} - V(Q))) \geq k$ . The path-star  $T_{PS}(r + 1, t) := (T_{PS}(r, t) \setminus \bar{P}) \cup yq \cup Q$  rooted at  $s_0$  has order  $|T_{PS}(r, t)| + 1 \leq m$  and  $G - V(T_{PS}(r + 1, t)) = G \setminus (T_{PS}(r, t) \setminus \bar{P} - V(Q))$  is  $k$ -connected, a contradiction.

Case 2.  $y \in \{s_1, s_2, \dots, s_{t-1}\}$ .

Let  $\bar{P} = P[p_r, p_1]$  and then there is a path-star  $T_{PS}(r + 1, t) := (T_{PS}(r, t) \setminus \bar{P}) \cup yq \cup Q$  with order  $|T_{PS}(r + 1, t)| \leq m$  according to the same way as above.  $\square$

**Proof of Theorem 1.4.** Suppose that  $G$  is  $k$ -connected with  $\delta(G) \geq \lfloor \frac{3k}{2} \rfloor + m - 1$  and not complete. We discuss the following cases.

Case 1.  $\kappa(G) = k$ . Clearly, there is an end  $E$  in  $G$  and let  $S$  be  $N_G(E)$  with  $|S| = k$ . Assume that  $|E| \geq 2$  since it could be deduced  $k = 1 = m$  for  $|E| = 1$ . Naturally, it follows that  $G[S] - V(\bar{E})$  is  $k + 1$ -connected by Lemma 1.5, furthermore it satisfies  $G[S] - V(\bar{E}) \in \mathcal{F}_k^+(m)$ . Then from Theorem 1.3 there is a path-star  $T_{PS}(r, m - r) \subseteq E$  of order  $m$  such that  $G[S] - V(\bar{E} \cup T_{PS}(r, m - r))$  remains  $k$ -connected. Then we have  $\kappa(G - V(T_{PS}(r, m - r))) \geq k$  by Lemma 1.6(b).

Case 2.  $\kappa(G) \geq k + 1$ . Select a maximal path-star  $T_{PS}$  with  $1 \leq |T_{PS}| < m$ . Then we prove exactly in the same way as in the proof of Theorem 1.3, that the assumption  $|T_{PS}| < m$  leads to a contradiction.

Since  $|N_G(p_r) \cap (G - V(C \cup T_{PS}(r, t)))| \geq \lfloor \frac{3k}{2} \rfloor + m - 1 - k - (m - 2) = \lfloor \frac{k}{2} \rfloor + 1$ , let  $p' \in V(G) - V(C \cup T_{PS}(r, t))$  be a neighbor of  $p_r$ . Then we get  $\kappa(G \setminus T_{PS}(r, t)) = k$  and the existence of an end  $E$  of  $G \setminus T_{PS}(r, t)$ . Set  $S := N_{G \setminus T_{PS}(r, t)}(E)$  and  $\bar{E} := G \setminus T_{PS}(r, t) - (S \cup V(E))$ . Since the case  $|E| = 1$  is settled as in the proof of Theorem 1.3, then let  $|E| \geq 2$  and from Lemma 1.5 we have  $\kappa((G \setminus T_{PS}(r, t))[S] - V(\bar{E})) \geq k + 1$ . Let  $y, q, T_{PS}(r, t) \setminus \bar{P}$  and  $G \setminus (T_{PS}(r, t) \setminus \bar{P})$  be the same as in the proof of Theorem 1.3. Therefore,  $G \setminus (T_{PS}(r, t) \setminus \bar{P})$  is  $k$ -connected with  $\delta(G \setminus (T_{PS}(r, t) \setminus \bar{P})) \geq \lfloor \frac{3k}{2} \rfloor + |\bar{P}|$  and  $E$  is an end of  $G \setminus (T_{PS}(r, t) \setminus \bar{P})$  to  $S$ . From Theorem 1.3 and Lemma 1.7, applied to  $(G \setminus (T_{PS}(r, t) \setminus \bar{P})[S](E \cup S), K(S)) \in \mathcal{F}_k^+(|\bar{P}| + 1)$  and  $q \in E$ , we know that there exists a path  $Q$  of length  $|\bar{P}|$  starting from  $q$  in  $E$  such that  $G \setminus (T_{PS}(r, t) \setminus \bar{P}[S](E \cup S) - V(Q))$  remains  $k$ -connected. Then we have  $G \setminus (T_{PS}(r, t) \setminus \bar{P} - V(Q)) = G - V((T_{PS}(r, t) \setminus \bar{P}) \cup Q)$  is also  $k$ -connected by Lemma 1.6(b), where  $(T_{PS}(r, t) \setminus \bar{P}) \cup yq \cup Q$  is a path-star with order  $|T_{PS}(r, t)| + 1$ .  $\square$

## References

- [1] J. A. Bondy and U. S. R. Murty, Graph Theory, Graduate Texts in Mathematics 244, Springer, Berlin, 2008.
- [2] G. Chartrand, A. Kaigars, D.R. Lick, Critically  $n$ -connected graphs, Proc. Amer. Math. Soc. 32 (1972) 63-68.
- [3] A. A. Diwan and N. P. Tholiya, Non-separating trees in connected graphs, Discrete Math. 309 (2009) 5235-5237.
- [4] S. Fujita and K. Kawarabayashi, Connectivity keeping edges in graphs with large minimum degree, J. Combin. Theory Ser B 98 (2008) 805-811.
- [5] T. Hasunuma, Connectivity Keeping Trees in 2-Connected Graphs with Girth Conditions, Combinatorial Algorithms (2020) 316-329.
- [6] T. Hasunuma and K. Ono, Connectivity keeping trees in 2-connected graphs, J. Graph Theory 94 (2020) 20-29.
- [7] Y. Hong and Q. Liu, Mader's conjecture for graphs with small connectivity, arXiv: 2101.11777 (2021).
- [8] C. Lu, P. Zhang, Connectivity keeping trees in 2-connected graphs, Discrete Math. 343 (2) (2020) 1-4.
- [9] W. Mader, Connectivity keeping paths in  $k$ -connected graphs, J. Graph Theory 65 (2010) 61-69.
- [10] W. Mader, Connectivity keeping trees in  $k$ -connected graphs, J. Graph Theory 69 (2012) 324-329.
- [11] Y. Tian, H. Lai, L. Xu, J. Meng, Nonseparating trees in 2-connected graphs and oriented trees in strongly connected digraphs, Discrete Math. 342 (2019) 344-351.
- [12] Y. Tian, J. Meng, L. Xu, Connectivity keeping stars or double-stars in 2-connected graphs, Discrete Math. 341 (2018) 1120-1124.