

LINKEDNESS OF CARTESIAN PRODUCTS OF COMPLETE GRAPHS

LEIF K. JØRGENSEN

Department of Mathematical Sciences, Aalborg University, Denmark

GUILLERMO PINEDA-VILLAVICENCIO

*Centre for Informatics and Applied Optimisation, Federation University, Australia
School of Information Technology, Deakin University, Geelong, Australia*

JULIEN UGON

*Centre for Informatics and Applied Optimisation, Federation University, Australia
School of Information Technology, Deakin University, Geelong, Australia*

ABSTRACT. This paper is concerned with the linkedness of Cartesian products of complete graphs. A graph with at least $2k$ vertices is *k-linked* if, for every set of $2k$ distinct vertices organised in arbitrary k pairs of vertices, there are k vertex-disjoint paths joining the vertices in the pairs.

We show that the Cartesian product $K^{d_1+1} \times K^{d_2+1}$ of complete graphs K^{d_1+1} and K^{d_2+1} is $\lfloor (d_1 + d_2)/2 \rfloor$ -linked for $d_1, d_2 \geq 2$, and this is best possible.

This result is connected to graphs of simple polytopes. The Cartesian product $K^{d_1+1} \times K^{d_2+1}$ is the graph of the Cartesian product $T(d_1) \times T(d_2)$ of a d_1 -dimensional simplex $T(d_1)$ and a d_2 -dimensional simplex $T(d_2)$. And the polytope $T(d_1) \times T(d_2)$ is a *simple polytope*, a $(d_1 + d_2)$ -dimensional polytope in which every vertex is incident to exactly $d_1 + d_2$ edges.

While not every d -polytope is $\lfloor d/2 \rfloor$ -linked, it may be conjectured that every simple d -polytope is. Our result implies the veracity of the revised conjecture for Cartesian products of two simplices.

E-mail addresses: leifkjorgensen@gmail.com, work@guillermo.com.au,
julien.ugon@deakin.edu.au.

Date: January 31, 2022.

2020 Mathematics Subject Classification. Primary 05C40; Secondary 52B05.

Key words and phrases. *k*-linked, cyclic polytope, connectivity, separator, dual polytope, linkedness, Cartesian product.

Thanks.

1. INTRODUCTION

A – B path

Denote by $V(X)$ the vertex set of a graph. Given sets A, B of vertices in a graph, a path from A to B , called an *A – B path*, is a (vertex-edge) path $L := u_0 \dots u_n$ in the graph such that $V(L) \cap A = \{u_0\}$ and $V(L) \cap B = \{u_n\}$. We write *a – B path* instead of $\{a\} – B$ path, and likewise, write *A – b path* instead of $A – \{b\}$.

*terminals**linked pairs**linked set*

Let G be a graph and X a subset of $2k$ distinct vertices of G . The elements of X are called *terminals*. Let $Y := \{\{s_1, t_1\}, \dots, \{s_k, t_k\}\}$ be an arbitrary labelling and (unordered) pairing of all the vertices in X . We say that Y is *linked* in G if we can find disjoint $s_i – t_i$ paths for $i \in [1, k]$, the interval $1, \dots, k$. The set X is *linked* in G if every such pairing of its vertices is linked in G . Throughout this paper, by a set of disjoint paths, we mean a set of vertex-disjoint paths. If G has at least $2k$ vertices and every set of exactly $2k$ vertices is linked in G , we say that G is *k-linked*.

k-linked

This paper studies the linkedness of Cartesian products of complete graphs. Linkedness of Cartesian products has been studied in the past (Mészáros, 2016). The *Cartesian product* $G_1 \times G_2$ of two graphs G_1 and G_2 is the graph defined on the pairs (v_1, v_2) with $v_i \in G_i$ and with two pairs (u_1, u_2) and (v_1, v_2) being adjacent if, for some $\ell \in \{1, 2\}$, $u_\ell v_\ell \in E(G_\ell)$ and $u_i = v_i$ for $i \neq \ell$. We prove that the Cartesian product $K^{d_1+1} \times K^{d_2+1}$ of complete graphs K^{d_1+1} and K^{d_2+1} is $\lfloor (d_1 + d_2)/2 \rfloor$ -linked for $d_1, d_2 \geq 0$, and that there are products that are not $\lfloor (d_1 + d_2 + 1)/2 \rfloor$ -linked; hence this result is best possible. Here K^t denotes the complete graph on t vertices.

Our result is connected to questions on the linkedness of a polytope. A (convex) polytope is the convex hull of a finite set X of points in \mathbb{R}^d ; the *convex hull* of X is the smallest convex set containing X . The *dimension* of a polytope in \mathbb{R}^d is one less than the maximum number of affinely independent points in the polytope; a set of points $\vec{p}_1, \dots, \vec{p}_k$ in \mathbb{R}^d is *affinely independent* if the $k-1$ vectors $\vec{p}_1 - \vec{p}_k, \dots, \vec{p}_{k-1} - \vec{p}_k$ are linearly independent. A polytope of dimension d is referred to as a *d-polytope*.

The *Cartesian product* $P \times P'$ of a d -polytope $P \subset \mathbb{R}^d$ and a d' -polytope $P' \subset \mathbb{R}^{d'}$ is the Cartesian product of the sets P and P' :

$$P \times P' = \left\{ \begin{pmatrix} p \\ p' \end{pmatrix} \in \mathbb{R}^{d+d'} \mid p \in P, p' \in P' \right\}.$$

The resulting polytope is $(d + d')$ -dimensional. The *graph* $G(P)$ of a polytope P is the undirected graph formed by the vertices and edges of the polytope. It follows that the graph $G(P \times P')$ of the Cartesian product $P \times P'$ is the Cartesian product $G(P) \times G(P')$ of the graphs $G(P)$ and $G(P')$.

A d -simplex $T(d)$ is the convex hull of $d + 1$ affinely independent points in \mathbb{R}^d . The graph of $T(d)$ is the complete graph K^{d+1} . As a consequence, our result implies that the graph of the Cartesian product $T(d_1) \times T(d_2)$ is $\lfloor (d_1 + d_2)/2 \rfloor$ -linked for $d_1, d_2 \geq 0$. Henceforth, if the graph of a polytope is *k-linked* we say that the polytope is also *k-linked*.

The first edition of the Handbook of Discrete and Computational Geometry (Goodman & O'Rourke, 1997, Problem 17.2.6) posed the question of whether or not every d -polytope is $\lfloor d/2 \rfloor$ -linked. This question was answered in the negative by Gallivan (1985). None of the known counterexamples are *simple d -polytopes*, d -polytopes in which every vertex is incident to exactly d edges. Hence, it may be hypothesised that the conjecture holds for such polytopes.

Conjecture 1. *Every simple d -polytope is $\lfloor d/2 \rfloor$ -linked for $d \geq 2$.*

Cartesian products of simplices are simple polytopes, and so our result supports this revised conjecture. Furthermore, Cartesian products of simplices and duals of cyclic polytopes are related; the dual of a cyclic d -polytope with $d + 2$ vertices is the Cartesian product of a $\lfloor d/2 \rfloor$ -simplex and a $\lceil d/2 \rceil$ -simplex (Ziegler, 1995, Ex. 0.6). Hence we obtain that the dual of a cyclic d -polytope on $d + 2$ vertices is also $\lfloor d/2 \rfloor$ -linked for $d \geq 2$.

Unless otherwise stated, the graph theoretical notation and terminology follows from Diestel (2017) and the polytope theoretical notation and terminology from Ziegler (1995). Moreover, when referring to graph-theoretical properties of a polytope such as linkedness and connectivity, we mean properties of its graph.

2. LINKEDNESS OF CARTESIAN PRODUCTS OF COMPLETE GRAPHS

The contribution of this section is a sharp theorem (Theorem 2) that tells the story of the linkedness of Cartesian product of two complete graphs.

Theorem 2. *The Cartesian product of two complete graphs K^{d_1+1} and K^{d_2+1} is $\lfloor (d_1 + d_2)/2 \rfloor$ -linked for every $d_1, d_2 \geq 0$.*

Remark 3. Theorem 2 is best possible. There are products $K^{d_1+1} \times K^{d_2+1}$ that are not $\lfloor (d_1 + d_2 + 1)/2 \rfloor$ -linked:

- (i) $K^2 \times K^{d_2+1}$ for even $d_2 \geq 1$, and
- (ii) $K^3 \times K^{d_2+1}$ for $d_2 = 1, 3, 5, 7, 9$.

For each of these cases, Fig. 1 provides a pairing of terminals that cannot be $\lfloor (d_1 + d_2 + 1)/2 \rfloor$ -linked. We conjecture these are the only such cases.

An immediate corollary of Theorem 2 is the following.

Corollary 4. *The Cartesian product of two simplices $T(d_1)$ and $T(d_2)$ is $\lfloor (d_1 + d_2)/2 \rfloor$ -linked for every $d_1, d_2 \geq 0$.*

The notions of linkage, linkage problem, and valid path will simplify our arguments. A *linkage* in a graph is a subgraph in which every component is a path. Let X be a set of vertices in a graph and let $Y := \{\{s_1, t_1\}, \dots, \{s_k, t_k\}\}$ be a pairing of all the vertices of X . A Y -*linkage* $\{L_1, \dots, L_k\}$ is a set of disjoint paths with the path L_i joining the pair $\{s_i, t_i\}$ for $i = 1, \dots, k$. We may also say that Y represents our *linkage problem*, and if Y is linked in G then our linkage problem is *feasible* and *infeasible* otherwise. A path in the graph is called X -*valid* if no inner vertex




FIGURE 1. No feasible linkage problems for $K^{d_1+1} \times K^{d_2+1}$, $k = \lfloor (d_1 + d_2 + 1)/2 \rfloor$, $d_1 \leq 2$ and $d_2 > d_1$. (a) The case $d_1 = 1$ and $d_2 > d_1$. (b) The case $d_1 = 2$ and $d_2 = 3$. (c) The case $d_1 = 2$ and $d_2 = 5$. (d) The case $d_1 = 2$ and $d_2 = 7$. (e) The case $d_1 = 2$ and $d_2 = 9$. Each row of each part (a)-(e) is a complete graph whose edges have not been drawn.

of the path is in X . Let X be a set of vertices in a graph G . Denote by $G[X]$ the subgraph of G induced by X , the subgraph of G that contains all the edges of G with vertices in X . Write $G - X$ for $G[V(G) \setminus X]$.

Consider a linkage problem $Y := \{\{s_1, t_1\}, \dots, \{s_k, t_k\}\}$ on a set X of $2k$ vertices in a graph G . Consider a linkage \mathcal{L} from a subset Z of X to some set Z' disjoint from X and label the vertices of Z' such that the path in \mathcal{L} with end $z_i \in Z$ has its other end $z'_i \in Z'$. Then the linkage \mathcal{L} in G induces a linkage problem Y' in $(G - V(\mathcal{L})) \cup Z'$ where the vertices of $X \setminus Z$ remain and the vertices of Z have been replaced by the vertices of Z' . Slightly abusing terminology, we also call terminals the vertices of Z' . If the problem Y' is feasible in $(G - V(\mathcal{L})) \cup Z'$, so is the problem Y in G .

Since we make heavy use of Menger's theorem (Diestel, 2017, Thm. 3.3.1), we next remind the reader of one of its consequences.

Theorem 5 (Menger's theorem). *Let G be a k -connected graph, and let A and B be two subsets of its vertices, each of cardinality at least k . Then there are k disjoint $A - B$ paths in G .*

We fix some notation and terminology for the remaining of the section. Let G denote the graph $K^{d_1+1} \times K^{d_2+1}$. We think of $G = K^{d_1+1} \times K^{d_2+1}$ as a grid with




FIGURE 2. Depiction of the subgraphs $B_{\alpha+1}$, $A_{\alpha+1}$, C_{12} , and \bar{C}_{12} of $K^{d_1+1} \times K^{d_2+1}$.

$d_1 + 1$ rows and $d_2 + 1$ columns. In this way, the entry in Row i and Column j can be referred to as $G[i, j]$.

When we write about a row r of subgraph G' of G , we think of r as a subgraph of G' and as the number r so that we can write about the r th row of G' or G ; this ambiguity should cause no confusion. An entry in the grid $K^{d_1+1} \times K^{d_2+1}$ with no terminal is said to be *free*, as is a row or a column of a subgraph of G with no terminal. A row or a column of a subgraph of G with every entry being occupied by a terminal is said to be *full*.

We need the following induced subgraphs of G :

$C_{ab\dots z}$, the subgraph formed by the union of Columns a, b, \dots, z ;

$\bar{C}_{ab\dots z}$, the subgraph obtained by removing Columns a, b, \dots, z ;

$R_{ab\dots z}$, the subgraph formed by the union of Rows a, b, \dots, z ;

$\bar{R}_{ab\dots z}$, the subgraph obtained by removing Rows a, b, \dots, z ;

A_α , the induced subgraph of \bar{C}_{12} obtained by removing its first α rows; and

B_α , the subgraph of C_{12} obtained by removing its first α rows.

For instance, \bar{C}_1 denotes the subgraph of G obtained by removing the first column, C_{12} the subgraph formed by the first two columns of G , and \bar{C}_{12} denotes the subgraph obtained by removing the first two columns of G ; observe \bar{C}_{12} is isomorphic to $K^{d_1+1} \times K^{d_2-1}$. Figure 2 depicts some of the aforementioned subgraphs of $K^{d_1+1} \times K^{d_2+1}$.

The connectivity of $K^{d_1+1} \times K^{d_2+1}$ is stated below.

Lemma 6 (Špacapan 2008, Thm. 1). *The (vertex)connectivity of $K^{d_1+1} \times K^{d_2+1}$ is precisely $d_1 + d_2$.*

We continue fixing further notation. Henceforth let $k := \lfloor (d_1 + d_2)/2 \rfloor$. And let X be a subset of $2k$ vertices of G and let $Y := \{\{s_1, t_1\}, \dots, \{s_k, t_k\}\}$ be a pairing of all the vertices in X .

We first settle the simple cases of $(0, d_2)$ and $(1, d_2)$ for $d_2 \geq 0$.

Proposition 7 (Base cases). *For $d_2 \geq 0$ the Cartesian products $K^1 \times K^{d_2+1}$ and $K^2 \times K^{d_2+1}$ are both $\lfloor (1 + d_2)/2 \rfloor$ -linked. This statement is best possible.*

Proof. The lemma is true for the pair $(0, d_2)$ for each $d_2 \geq 0$, since $K^1 \times K^{d_2+1} = K^{d_2+1}$ and K^{d_2+1} is $\lfloor (1 + d_2)/2 \rfloor$ -linked. This is best possible.

The graph $K^2 \times K^{d_2+1}$ is $(1 + d_2)$ -connected by Lemma 6. Use Menger's theorem (5) to bring the $1 + d_2$ terminals to the subgraph \bar{R}_1 through a linkage $\{S_1, \dots, S_k, T_1, \dots, T_k\}$ with $S_i := s_i - \bar{R}_1$ and $T_i := t_i - \bar{R}_1$ for $i \in [1, k]$. Letting $\{\bar{s}_i\} := V(S_i) \cap V(\bar{R}_1)$ and $\{\bar{t}_i\} := V(T_i) \cap V(\bar{R}_1)$, we produce a new linkage problem $Y' := \{(\bar{s}_1, \bar{t}_1), \dots, (\bar{s}_k, \bar{t}_k)\}$ in \bar{R}_1 whose feasibility implies that of Y in G . To solve Y' link the pairs of Y' in the subgraph \bar{R}_1 , which is isomorphic to K^{d_2+1} , using the $\lfloor (1 + d_2)/2 \rfloor$ -linkedness of K^{d_2+1} . Figure 1(a) shows an infeasible linkage problem with $\lfloor (2 + d_2)/2 \rfloor$ pairs in the graph $K^2 \times K^{d_2+1}$. \square

In what follows we aim to find a Y -linkage $\{L_1, \dots, L_k\}$ in G with L_i joining the pair $\{s_i, t_i\}$ of Y for $i \in [1, k]$. Our proof is by induction on (d_1, d_2) with the base cases settled in Proposition 7. If there is a pair of Y , say $\{s_1, t_1\}$, lying in some column or row of G , say in Column 1, we send every terminal $s_i \in C_1$ that is different from s_1 and t_1 and that is not adjacent to t_i to the subgraph \bar{C}_1 , and apply the induction hypothesis on \bar{C}_1 . Otherwise, we may assume every pair of Y lies in two distinct columns or rows, say the pair $\{s_1, t_1\}$ lies in C_{12} ; then we send every terminal $s_i \in C_{12}$ that is different from s_1 and t_1 and that is not adjacent to t_i to the subgraph \bar{C}_{12} , and apply the induction hypothesis to \bar{C}_{12} . We develop these ideas below.

The definition of k -linkedness gives the following lemma at once; we will use it implicitly hereafter.

Lemma 8. *Let $\ell \leq k$. Let X be a set of 2ℓ distinct vertices of a k -linked graph K , let Y be a labelling and pairing of the vertices in X , and let Z be a set of $2k - 2\ell$ vertices in K such that $X \cap Z = \emptyset$. Then there exists a Y -linkage in K that avoids every vertex in Z .*

Besides, basic algebraic manipulation yields the following inequality.

Lemma 9. *If $x \geq 2$ and $y \geq 2$ then $x(y - 1) > x + y - 3$.*

Proof. The inequality simplifies to $(x - 1)(y - 2) > -1$. \square

We are now ready to put together all the elements of the proof of Theorem 2.

Proof of Theorem 2. Let $k := \lfloor (d_1 + d_2)/2 \rfloor$. Then $d_1 + d_2 \geq 2k$.

Proposition 7 gives the result for the pairs $(d_1, 0)$, $(0, d_2)$, $(d_1, 1)$, and $(1, d_2)$ for each $d_1, d_2 \geq 0$. Hence, our bidimensional induction on (d_1, d_2) can start with the assumption of $d_1, d_2 \geq 2$.

We first deal with the case where a pair in Y , say $\{s_1, t_1\}$, lies in some column or some row of G , say in Column 1.

Case 1. A pair in Y , say $\{s_1, t_1\}$, lies in Column 1.

The induction hypothesis ensures that the subgraph \bar{C}_1 is $(k-1)$ -linked. Hence it suffices to show that all the terminals in C_1 other than s_1, t_1 can be moved to \bar{C}_1 via a linkage; Menger's theorem (5) guarantees this.

Let U be the set of terminals in C_1 other than s_1 and t_1 , and let W be the set of terminals in \bar{C}_1 . Then $|U| + |W| = d_1 + d_2 - 2$. Besides, the subgraph $G - (W \cup \{s_1, t_1\})$ is $|U|$ -connected, as G is $(d_1 + d_2)$ -connected (Lemma 6). In the case of $d_1, d_2 \geq 2$, Lemma 9 yields that \bar{C}_1 has more than $|U \cup W|$ vertices:

$$|\bar{C}_1| = (d_1 + 1)d_2 > d_1 + 1 + d_2 + 1 - 3 > d_1 + d_2 - 2 = |U| + |W|.$$

Menger's theorem (5) applies and gives disjoint $C_1 - \bar{C}_1$ paths from the terminals in U to $|U|$ free entries in \bar{C}_1 . The $(k-1)$ -linkedness of \bar{C}_1 now settles the case.

By symmetry, we can assume that every pair $\{s_i, t_i\}$ in Y lies in two different columns or rows and that s_i, t_i are not adjacent. Without loss of generality, assume that

(*) s_1 is Column 1 and t_1 is in Column 2 of C_{12} .

The induction hypothesis also ensures that both \bar{C}_{12} and \bar{R}_{12} are $(k-1)$ -linked. We consider two further cases based on the number of terminals in C_{12} or R_{12} .

Case 2. The subgraph C_{12} contains precisely $d_1 + 2 - \alpha$ terminals, including $\{s_1, t_1\}$, where $0 \leq \alpha \leq d_1$.

Excluding $\{s_1, t_1\}$, there are at most d_1 terminals in C_{12} , and there are $d_1 + 1$ internally-disjoint $s_1 - t_1$ paths in C_{12} of length at most three: two length-two paths and $d_1 - 1$ length-three paths. One of these $s_1 - t_1$ paths, say L_1 , avoids every other terminal in C_{12} .

Without loss of generality, assume that Row 1 in C_{12} is part of the path L_1 ; that is,

(**) $\{G[1, 1], G[1, 2]\} \subseteq V(L_1)$.

It is true that $(V(L_1) \cap V(B_1)) \subseteq \{s_1, t_1\}$.

In the subcase $\alpha = d_1$, every pair in $Y \setminus \{s_1, t_1\}$ is in \bar{C}_{12} , and the induction hypothesis on \bar{C}_{12} settles the subcase.

Suppose that $\alpha = d_1 - 1$, say C_{12} contains $\{s_1, t_1, s_2\}$. Then $s_2 \in B_1$ and $t_2 \in \bar{C}_{12}$. We may assume s_1, s_2 are in Column 1 and t_1 is in Column 2. We show there is an X -valid $s_2 - A_1$ path L'_2 such that the vertex $x \in V(L'_2) \cap V(A_1)$ is either t_2 or a nonterminal.

Through each entry of Column 1 of B_1 , there is a $s_2 - A_1$ path of length at most two: one of length one and $d_1 - 1$ of length two. Moreover, $d_1 - 1$ of such paths avoid s_1 . To ensure the existence of L'_2 , it suffices to show that A_1 cannot have $d_1 - 1$ rows that are full of terminals other than t_2 . According to Lemma 9, the inequality

$$(d_1 - 1)(d_2 - 1) > d_1 - 1 + d_2 - 3 = |X \setminus \{s_1, t_1, s_2, t_2\}|$$

holds for $d_1, d_2 \geq 2$. Hence we get the existence of L'_2 . As a result, the solution of the new problem $Y' := \{\{x, t_2\}, \{s_3, t_3\}, \dots, \{s_k, t_k\}\}$ in \bar{C}_{12} induces a solution of the problem Y in G . And the solution of Y' follows from the $(k-1)$ -linkedness of \bar{C}_{12} .

Henceforth assume that $\alpha \leq d_1 - 2$. To finalise Case 2, we require a couple of claims.

Claim 1. Suppose that there are at most $d_1 + 2$ terminals in $B_1 = K^{d_1} \times K^2$. Then there is an injection from the set of rows of B_1 that contain two terminals x_1, x_2 such that $\{x_1, x_2\} \cap \{s_1, t_1\} = \emptyset$ to the set of rows of B_1 that contain either no terminal or a terminal in $\{s_1, t_1\}$ but no other terminal.

Proof. This follows from a simple counting argument. The number of rows in B_1 is d_1 . Let m denote the number of rows of B_1 that contain two terminals x_1, x_2 such that $\{x_1, x_2\} \cap \{s_1, t_1\} = \emptyset$ and let $n := |(X \cap V(B_1)) \setminus \{s_1, t_1\}|$. It follows that the number of rows of B_1 that contains precisely one terminal $x \notin \{s_1, t_1\}$ is $n - 2m$; either s_1 or t_1 may be in these rows. As a result, the number of rows of B_1 that contain either no terminal or a terminal in $\{s_1, t_1\}$ but no other terminal is $d_1 - m - (n - 2m)$. Combining $n \leq d_1$ with all these numbers, we get that

$$d_1 - m - (n - 2m) = d_1 - n + m \geq d_1 - d_1 + m = m.$$

The claim is proved. \square

Claim 2. Suppose that there are at most $d_1 + 2$ terminals in $B_1 = K^{d_1} \times K^2$. If every row in the subgraph $A_1 = K^{d_1} \times K^{d_2-1}$ of \bar{C}_{12} has a free entry, then, for every terminal $x \notin \{s_1, t_1\}$ in B_1 , there is a $B_1 - A_1$ path L from x to a free entry in A_1 such that L is X -valid; and all these X -valid paths are disjoint.

Proof. If a row of B_1 contains exactly one terminal $x \notin \{s_1, t_1\}$, then send x to a free entry in the same row of A_1 . Let x_1 and x_2 be two terminals in B_1 that satisfy $\{x_1, x_2\} \cap \{s_1, t_1\} = \emptyset$ and occupy a row r_f of B_1 . From Claim 1 ensues the existence of a row r_e of B_1 that contains either no terminal or a terminal in $\{s_1, t_1\}$ but no other terminal; in short, there is at least a free entry in r_e .

Consider a pair (r_f, r_e) of rows granted by Claim 1. Send either x_1 or x_2 , say x_1 , to the free entry in the row r_e of A_1 passing through the corresponding free entry in the row r_e of B_1 , and send x_2 to a free entry in the row r_f of A_1 . The proof of the claim is now complete. \square

Now suppose that $\alpha = 0$ or $2 \leq \alpha \leq d_1 - 2$. In this subcase, the subgraph \bar{C}_{12} contains at most α full rows: if $\alpha + 1$ rows were full in \bar{C}_{12} then there would be at least $(\alpha + 1)(d_2 - 1)$ terminals in \bar{C}_{12} but $(\alpha + 1)(d_2 - 1) > d_2 - 2 + \alpha$ (Lemma 9). Even when the path L_1 uses the first row of \bar{C}_{12} by (**), there is no loss of generality by assuming that the full rows of \bar{C}_{12} are among the *first* $\alpha + 1$ rows of \bar{C}_{12} . It follows that every row of $A_{\alpha+1}$ has a free entry.

We send to $B_{\alpha+1}$ the terminals other than s_1 and t_1 that are in the rows 2 to $\alpha + 1$ of C_{12} , the terminals other than s_1 and t_1 that are in $B_1 \setminus B_{\alpha+1}$. For $\alpha = 0$, $B_1 \setminus B_{\alpha+1} = \emptyset$ and there is nothing to do.

We now focus on the subcase $2 \leq \alpha \leq d_1 - 2$. Let n_1 and n_2 denote the number of terminals in $B_1 \setminus B_{\alpha+1}$ and $B_{\alpha+1}$, respectively. Then the following inequalities hold

$$n_1 + n_2 \leq d_1 + 2 - \alpha \leq d_1 \quad (\text{since } 2 \leq \alpha),$$

$$n_1 + n_2 \leq d_1 + 2 - \alpha \leq 2d_1 - 2\alpha = |B_{\alpha+1}| \quad (\text{since } \alpha \leq d_1 - 2).$$

Since B_1 is d_1 -connected by Lemma 6, Menger's theorem gives n_1 disjoint paths from the terminals in $B_1 \setminus B_{\alpha+1}$ to n_1 free entries in $B_{\alpha+1}$. These free entries in $B_{\alpha+1}$ will henceforth play the role of the terminals of $B_1 \setminus B_{\alpha+1}$, and so we call them terminals as well.

Let $d'_1 := d_1 - \alpha$ and $d'_2 := d_2$. Then, $d'_1 \geq 2$, there are at most $d'_1 + 2$ terminals in $B_{\alpha+1} = K^{d'_1} \times K^2$, and every row in $A_{\alpha+1} = K^{d'_1} \times K^{d'_2} - 1$ has a free entry. Hence, Claim 2 applies, and there is a linkage formed by X -valid paths from the terminals in $B_{\alpha+1}$, other than t_1 , to free entries in $A_{\alpha+1}$. Now we have a new linkage problem Y' in \bar{C}_{12} with at most $2(k-1)$ pairs. The solution of Y' in \bar{C}_{12} implies a solution of the linkage problem Y in G . To link the pairs of Y' use the $(k-1)$ -linkedness of \bar{C}_{12} .

Finally assume that $\alpha = 1$. In a first scenario suppose that either both entries in $B_1 \setminus B_2$ are nonterminals or each terminal in $B_1 \setminus B_2$ is adjacent to a nonterminal in B_2 . Then we can send the terminals in $B_1 \setminus B_2$ to B_2 . In the second scenario suppose that there is a terminal s_i in $B_1 \setminus B_2$ whose neighbours in B_2 are all terminals. Then the column of s_1 in B_1 would contain exactly d_1 terminals, including s_i . We send s_i to a free entry in A_1 , in the same row as s_i (the first row of A_1): if this free entry didn't exist, then s_i would be adjacent to the $d_2 - 1$ terminals in A_1 and the $d_1 - 1$ terminals in B_2 . Since there are $d_1 + d_2$ terminals in total, it would follow that s_i is adjacent to t_i . This contradiction shows that we can send s_i to a free entry in A_1 . This completes the proof of Case 2.

In both scenarios, it remains to send the terminals in B_2 to A_2 . To do so, we reason as in the subcase $2 \leq \alpha \leq d_1 - 2$. Let $d'_1 := d_1 - 1$ and $d'_2 := d_2$. It follows that $d'_1 \geq 2$, that there are at most $d_1 + 2'$ terminals in B_2 , and that every row in A_2 has a free entry. Claim 2 applies again and gives a linkage formed by X -valid paths from the terminals in B_2 , other than s_1, t_1 , to free entries in A_2 .

With all the terminals in \bar{C}_{12} , therein we have a new linkage problem Y' with at most $2(k-1)$ pairs whose solution in \bar{C}_{12} implies a solution of the linkage problem Y in G . To solve Y' in \bar{C}_{12} use the $(k-1)$ -linkedness of \bar{C}_{12} .

By symmetry, we also have the result if there are at most $d_2 + 2$ terminals in R_{12} , including $\{s_1, t_1\}$.

Case 3. The subgraph C_{12} contains at least $d_1 + 3$ terminals, including $\{s_1, t_1\}$.

This case reduces to the previous case. If C_{12} contains at least $d_1 + 3$ terminals then R_{12} contains at most $d_2 - 3 + 4 = d_2 + 1$ terminals, since there are four entries shared by C_{12} and R_{12} . Since we make no distinction between columns and rows, this case is already covered. This completes the proof of the theorem. \square

3. DUALS OF CYCLIC POLYTOPES

There is a close connection between duals of cyclic d -polytopes with $d+2$ vertices and Cartesian products of complete graphs.

The *moment curve* in \mathbb{R}^d is defined by $x(t) := (t, t^2, \dots, t^d)$ for $t \in \mathbb{R}$, and the convex hull of any $n > d$ points on it gives a *cyclic polytope* $C(n, d)$. The *combinatorics* of a cyclic polytope, the face lattice of the polytope faces partially ordered by inclusion, is independent of the points chosen on the moment curve. Hence we talk of the cyclic d -polytope on n vertices (Ziegler, 1995, Example 0.6).

For a polytope P that contains the origin in its interior, the *dual polytope* P^* is defined as

$$P^* = \{y \in \mathbb{R}^d \mid x \cdot y \leq 1 \text{ for all } x \text{ in } P\}.$$

If P does not contain the origin, we translate the polytope so that it does. Translating the polytope P changes the geometry of P^* but not its face lattice. The face lattice of P^* is the inclusion reversed face lattice of P . In particular, the vertices of P^* correspond to the facets of P , and the edges of P^* correspond to the $(d-2)$ -faces of P . The *dual graph* of a polytope P is the graph of the dual polytope, or equivalently, the graph on the set of facets of P where two facets are adjacent in the dual graph if they share a $(d-2)$ -face.

Duals of cyclic d -polytopes are simple d -polytopes. It is also the case that the dual of a cyclic d -polytope with $d+2$ vertices can be expressed as $T(\lfloor d/2 \rfloor) \times T(\lceil d/2 \rceil)$ (Ziegler 1995, Ex. 0.6). From this observation and Theorem 2 the next corollary follows at once.

Corollary 10. *Duals of cyclic polytopes with $d+2$ vertices are $\lfloor d/2 \rfloor$ -linked for every $d \geq 2$.*

4. ACKNOWLEDGMENTS

Guillermo would like to thank the hospitality of Leif Jørgensen, and the Department of Mathematical Sciences at Aalborg University where this research started.

Julien Ugon's research was supported by the ARC discovery project DP180100602.

REFERENCES

- Diestel, R. (2017). *Graph Theory*, volume 173 of *Graduate Texts in Mathematics*. Berlin: Springer-Verlag, 5th edition.
- Gallivan, S. (1985). Disjoint edge paths between given vertices of a convex polytope. *J. Combin. Theory Ser. A*, 39(1), 112–115.

- Goodman, J. E. & O'Rourke, J., Eds. (1997). *Handbook of discrete and computational geometry*. Discrete Mathematics and its Applications (Boca Raton). Chapman & Hall/CRC, Boca Raton, FL, 1st edition.
- Mészáros, G. (2016). On linkedness in the Cartesian product of graphs. *Period. Math. Hungar.*, 72(2), 130–138.
- Špacapan, S. (2008). Connectivity of cartesian products of graphs. *Applied Mathematics Letters*, 21(7), 682–685.
- Ziegler, G. M. (1995). *Lectures on polytopes*, volume 152 of *Graduate Texts in Mathematics*. New York: Springer-Verlag.