

Fixed points of a random restricted growth sequence

Toufik Mansour* Reza Rastegar†

January 8, 2022

Abstract

We call i a fixed point of a given sequence if the value of that sequence at the i -th position coincides with i . Here, we enumerate fixed points in the class of restricted growth sequences. The counting process is conducted by calculation of generating functions and leveraging a probabilistic sampling method.

MSC2010: Primary 05A18; Secondary 05A15, 60C05

Keywords: restricted growth sequence; fixed point; generating functions; probabilistic sampling.

1 Introduction

For any given sequence π of length n , $i \in [n] := \{1, \dots, n\}$ is a fixed point of π if the i -th entry π_i of the sequence π is equal to i . We denote the number of fixed points of the sequence π by \mathcal{F}_π . The term *fixed point* is motivated naturally by the same concept in the class of permutations, where it represents any point that is not moved by a permutation. Fixed points and derangements of permutations are well studied due to their importance in various branches of mathematics including algebra, probability, and combinatorics; see for instance [2–4, 6, 10, 11, 13, 19, 20] for a few examples. Recently, a new line of research toward extending results with regards to fixed points in other classes of discrete sequences has emerged; for instance, Archibald, Blecher, and Knopfmacher [1] considered fixed points in compositions and words over the alphabet $[k]$. Inspired by them, in this note, we further investigate fixed points for another important class of sequences in combinatorics; namely, restricted growth sequences. These sequences are of interest in connection with set partitions [15], q -analogues [7], certain combinatorial matrices [12], and Gray codes [9].

Before we state our results, a few definitions are in order. Throughout this note, we use \mathbb{N} as the set of all natural numbers. A sequence of natural numbers $\pi = \pi_1\pi_2 \cdots \pi_n \in \mathbb{N}^n$ is called a restricted growth sequence if

$$\pi_1 = 1 \quad \text{and} \quad \pi_{j+1} \leq 1 + \max\{\pi_1, \dots, \pi_j\} \quad \text{for all } 1 \leq j < n.$$

*Department of Mathematics, University of Haifa, 199 Abba Khoushy Ave, 3498838 Haifa, Israel; e-mail: tmansour@univ.haifa.ac.il

†Occidental Petroleum Corporation, Houston, TX 77046 and Departments of Mathematics and Engineering, University of Tulsa, OK 74104, USA - Adjunct Professor; e-mail: reza_rastegar2@oxy.com

There is a bijective connection between these sequences and canonical set partitions. A *partition* of a set A is a collection of non-empty, mutually disjoint subsets, called *blocks*, whose union is the set A . A partition Π with k blocks is called a k -*partition* and denoted by $\Pi = A_1|A_2|\cdots|A_k$. A k -partition $A_1|A_2|\cdots|A_k$ is said to be in the *standard form* if the blocks A_i are labeled in such a way that

$$\min A_1 < \min A_2 < \cdots < \min A_k.$$

The partition can be represented equivalently by the *canonical sequential form* $\pi_1\pi_2\dots\pi_n$, where $\pi_i \in [n]$ and $i \in A_{\pi_i}$ for all i [15]. In words, π_i is the label of the partition block that contains i . It is easy to verify that a word $\pi \in [k]^n$ is a canonical representation of a k -partition of $[n]$ in the standard form if and only if it is a restricted growth sequence [15].

Throughout this note, we use the terms *restricted growth sequence* and *set partition* interchangeably. As it becomes clear, in the context of restricted growth sequences, the fixed points are also closely related to the records. We remind the reader that the i -th entry π_i in the sequence π is a record if $\pi_i > \pi_j$ for all $j \in [i-1]$. Clearly, for any given fixed point i in a restricted growth sequence π , i is a record. In addition, each $j \in [i-1]$ is a fixed point and hence a record. Therefore, \mathcal{F}_π is precisely the length of the maximal prefix of π whose elements are records. We refer the reader to [5, 14, 16] for a few discussions around records for restricted growth sequences.

We denote by \mathcal{R}_n the set of all restricted growth sequences of length n , and denote by $\mathcal{R}_{n,k}$ the set of all restricted growth sequences of length n with maximal letter k .

Let $S_{n,k}$ be a Stirling number of the second kind and B_n be the n -th Bell number [15]. It is well-known that the cardinality of the set \mathcal{R}_n is B_n . In addition, the cardinality of the set $\mathcal{R}_{n,k}$ is $S_{n,k}$ with the exponential generating function $e^{y(e^x-1)}$, where y counts the number of blocks. The sequence of Bell numbers $(B_n)_{n \geq 0}$ can be then defined, for instance, through the formula $B_n = \sum_{k=0}^n S_{n,k}$, or, recursively via the formula $B_{n+1} = \sum_{k=0}^n \binom{n}{k} B_k$ with $B_0 = 1$, or through Dobinski's formula [8]

$$B_n = \frac{1}{e} \sum_{m=0}^{\infty} \frac{m^n}{m!}, \quad n \geq 0. \quad (1)$$

In what follows, we denote a random restricted growth sequence, sampled uniformly from $\mathcal{R}_{n,k}$ (resp. \mathcal{R}_n) by $\pi_{(n,k)}$ (resp. $\pi_{(n)}$). That is,

$$P(\pi_{(n,k)} = \pi) = \frac{1}{S_{n,k}} \quad \text{for all } \pi \in \mathcal{R}_{n,k},$$

and

$$P(\pi_{(n)} = \pi) = \frac{1}{B_n} \quad \text{for all } \pi \in \mathcal{R}_n.$$

We use $E(\cdot)$ to refer to the expectation with respect to the probability distribution $P(\cdot)$. We denote by $\mathcal{F}_n := \mathcal{F}_{\pi_{(n)}}$ the number of fixed points in a uniformly sampled random restricted growth sequence $\pi_{(n)}$. Fig. 1 shows the empirical distributions of \mathcal{F}_n over 20000 independently sampled instances of $\pi_{(n)}$ for $n = 50$ and $n = 200$.

1.1 Statement of results

For any $k \in \mathbb{N}$, we define $Q_k(x; q)$ to be the ordinary generating function enumerating fixed points over the set of restricted growth sequences in $\cup_{n=k}^{\infty} \mathcal{R}_{n,k}$; that is,

$$Q_k(x, q) := \sum_{n=k}^{\infty} x^n S_{n,k} E(q^{\mathcal{F}_n} | \pi_{(n)} \in \mathcal{R}_{n,k}) = \sum_{n=k}^{\infty} \sum_{\pi \in \mathcal{R}_{n,k}} x^n q^{\mathcal{F}_\pi}, \quad x, q \in \mathbb{C}, \quad (2)$$

where \mathbb{C} is the set of all complex numbers. Knowing an explicit form of (2), would in principle give us the distribution of \mathcal{F}_n in full details for all $n \in \mathbb{N}$. However, finding the explicit form of $Q_k(x, q)$ is a daunting task. Hence, we instead study the following exponential generating function:

$$R(x, y; q) := \sum_{n \geq 0} ([x^n] Q(x, y; q)) \frac{x^n}{n!} = \sum_{k \geq 1} y^k \sum_{n \geq 0} \frac{x^n}{n!} B_n E(\mathcal{F}_n | \pi_{(n)} \in \mathcal{R}_{n,k}), \quad (3)$$

where $[x^n]f$ denotes the coefficient of x^n in f and $Q(x, y; q)$ is the ordinary generating function for $Q_k(x; q)$; that is,

$$Q(x, y; q) := \sum_{k \geq 0} Q_k(x; q) y^k.$$

Our first result states that

Theorem 1.1. *The exponential generating function $R(x, y; q)$ is given by*

$$R(x, y; q) = e^{y(e^x - 1)} + \int_0^x (q - 1) y e^{x-t + (1+qt)ye^{x-t} - y} dt.$$

The proof of this theorem is given in Section 2. Note, by Theorem 1.1, $R(x, y; q)$ reduces to $e^{y(e^x - 1)}$ when $q = 1$, which is exactly the exponential generating function for the size of $\mathcal{R}_{n,k}$ (see [15]). Moreover, for $q = 0$, it implies

$$R(x, y; 0) = e^{y(e^x - 1)} - \int_0^x y e^{y(e^t - 1) + t} dt = e^{y(e^x - 1)} - e^{y(e^x - 1)} + 1 = 1,$$

which coincides with the fact that the only restricted growth sequence with no fixed point is the null sequence.



Figure 1: Empirical distributions of \mathcal{F}_{50} (left) and \mathcal{F}_{200} (right) based on 20000 samples.

Recall that $\mathcal{R}_{n,0}$ is an empty set for $n \geq 0$. Let $R_m(x, y)$ be the exponential generating function for the number of sequences in $\mathcal{R}_{n,k}$ with exactly m fixed points; that is,

$$R_m(x, y) := \sum_{n \geq 0} \sum_{k=0}^n \sum_{\substack{\pi \in \mathcal{R}_{n,k} \\ \mathcal{F}_\pi = m}} x^n y^k,$$

with $R_0(x, y) = 1$. Then, Theorem 1.1 implies that

Corollary 1.2. *For all $m \geq 1$,*

$$R_m(x, y) = \frac{1}{(m-1)!} \int_0^x y^m t^{m-1} e^{y(e^{x-t}-1)+m(x-t)} dt - \frac{1}{m!} \int_0^x y^{m+1} t^m e^{y(e^{x-t}-1)+(m+1)(x-t)} dt.$$

Next, we study the exponential generating function $T(x)$ for the total number of fixed points in \mathcal{R}_n ; that is

$$T(x) := \left. \frac{\partial}{\partial q} R(x, 1; q) \right|_{q=1}, \quad (4)$$

where by an inductive argument, provided in Section 2, we conclude that

Corollary 1.3. *The average number of fixed points over all the restricted growth sequences is*

$$E(\mathcal{F}_n) = \frac{1}{B_n} \frac{d^n}{dx^n} T(x) \Big|_{x=0} = \frac{1}{B_n} \sum_{i=1}^n \left(\sum_{j=1}^i \sum_{\ell=j}^i \frac{(-1)^{\ell-i} \ell^{n-j}}{(\ell-j)! (i-\ell)!} \right). \quad (5)$$

Our final result, Theorem 1.4 provides a closed form expression for the probability distribution of \mathcal{F}_n in terms of explicit polynomials of Bell numbers. We will use the following well-known extension of Dobinski's identity (1) to express the result. Recall that, for any integers $n, t \geq 0$ we have:

$$\begin{aligned} \Theta_n(t) &:= \frac{1}{e} \sum_{m=t}^{\infty} \frac{m^n}{(m-t)!} = \frac{1}{e} \sum_{k=0}^{\infty} \frac{(k+t)^n}{k!} = \frac{1}{e} \sum_{\ell=0}^n \binom{n}{\ell} t^{n-\ell} \sum_{k=0}^{\infty} \frac{k^\ell}{k!} \\ &= \sum_{\ell=0}^n \binom{n}{\ell} t^{n-\ell} B_\ell, \end{aligned} \quad (6)$$

where for the last equality we applied the original formula (1). The theorem states

Theorem 1.4. *For $n \in \mathbb{N}$, we have*

$$P(\mathcal{F}_n = j) = \begin{cases} \frac{B_{n-1}}{B_n} & \text{if } j = 1 \\ \frac{j \Theta_{n-j-1}(j)}{B_n} & \text{if } 1 < j < n \\ \frac{1}{B_n} & \text{if } j = n \end{cases}$$

In particular, for $n \geq 2$, we have

$$E(\mathcal{F}_n) = \frac{n + \sum_{j=1}^{n-1} j^2 \Theta_{n-j-1}(j)}{B_n}. \quad (7)$$

The proof of Theorem 1.4 is given in Section 3. It is based on the sampling method devised in [18] and has been recently exploited for the enumeration of other complex quantities over \mathcal{R}_n . See for instance [17] where the authors enumerate horizontal visibility graphs of sequences in \mathcal{R}_n . We remark that the $j = 1$ and $j = n$ cases are obvious; if $\pi \in \mathcal{R}_n$ such that $\mathcal{F}_\pi = 1$, then it must be that $\pi_2 = 1$ and clearly there are B_{n-1} choices for such words. Also, there is exactly one word $\pi = 12 \cdots n \in \mathcal{R}_n$ for which $\mathcal{F}_\pi = n$. However, we include our brute-force probabilistic calculation in Section 3 for the sake of completeness.

We make a final remark that from (5) and (7), we obtain the following identity for $n \geq 2$ which may be of interest independently to the reader:

$$\sum_{j=1}^{n-1} \sum_{\ell=0}^{n-j-1} \binom{n-j-1}{\ell} j^{n-j+1-\ell} B_\ell = \sum_{i=1}^n \left(\sum_{j=1}^i \sum_{\ell=j}^i \frac{(-1)^{\ell-i} \ell^{n-j}}{(\ell-j)! (i-\ell)!} \right) - n.$$

2 Proof of Theorems 1.1 and Corollary 1.3

The proof is based on the observation that each restricted growth sequence π with maximal value k and $\mathcal{F}_\pi = i$ fixed points falls into one of the following cases:

Case $\mathcal{F}_\pi = i = k$: Here $\pi = 12 \cdots k\pi^{(k)}$, where $\pi^{(k)}$ is a word over alphabet $[k]$.

Case $\mathcal{F}_\pi = i < k$: Here $\pi = 12 \cdots i\pi^{(i)}(i+1)\pi^{(i+1)} \cdots k\pi^{(k)}$, where $\pi^{(j)}$ is a word over alphabet $[j]$, for all $j = i, i+1, \dots, k$, and $\pi^{(i)}$ is not the empty word.

By considering these cases we arrive at the following equation:

$$Q_k(x; q) = \frac{x^k q^k}{1 - kx} + \sum_{i=1}^{k-1} \frac{ix^{k+1} q^i}{\prod_{j=i}^k (1 - jx)},$$

which implies

$$(1 - kx)Q_k(x; q) - xQ_{k-1}(x; q) = x^k q^{k-1} (q - 1), \quad (8)$$

with $Q_0(x; q) = 1$ and $Q_1(x; q) = \frac{xq}{1-x}$.

By multiplying (8) by y^k and summing over $k \geq 2$, we obtain

$$\begin{aligned} Q(x, y; q) - 1 - \frac{xyq}{1-x} \\ - xy \frac{\partial}{\partial y} \left(Q(x, y; q) - 1 - \frac{xyq}{1-x} \right) - xy(Q(x, y; q) - 1) = \sum_{k \geq 2} x^k q^{k-1} (q - 1) y^k, \end{aligned}$$

which leads to

$$(1 - xy)Q(x, y; q) - xy \frac{\partial}{\partial y} Q(x, y; q) = \frac{1 - xy}{1 - xyq}. \quad (9)$$

Next, we translate (9) in terms of exponential generating function $R(x, y; q)$ defined by (3) and write

$$\frac{\partial}{\partial x}R(x, y; q) - yR(x, y; q) - y\frac{\partial}{\partial y}R(x, y; q) = y(q-1)e^{xyq}.$$

Solving this equation with the initial condition $R(0, y; q) = R(x, 0; q) = 1$ completes the proof of Theorem 1.1.

Recall (4). Theorem 1.1 shows that $T(x) = \int_0^x e^{(x-t)e^t + e^t + t-1} dt$. By induction on m , we have

$$\frac{d^m}{dx^m}T(x) = \sum_{i=1}^m a_{mi}e^{e^x + ix-1} + \int_0^x e^{(x-t)e^t + e^t + (m+1)t-1} dt, \quad (10)$$

where a_{mi} are natural numbers satisfying the relations

$$\begin{cases} a_{mm} = 1 + a_{(m-1)(m-1)}, \\ a_{mi} = a_{(m-1)(i-1)} + ia_{(m-1)i}, \quad i = 1, 2, \dots, m-1, \end{cases} \quad (11)$$

with $a_{11} = 1$. Let $a_i(x) = \sum_{m \geq i} a_{mi}x^m$. Then, by (11), we have

$$a_i(x) = \sum_{j=1}^i \frac{x^i}{(1-jx)(1-(j+1)x) \cdots (1-ix)}.$$

Note that, by a partial fraction decomposition, we have

$$\frac{1}{(1-jx)(1-(j+1)x) \cdots (1-ix)} = \sum_{\ell=j}^i \frac{(-1)^{\ell-i} \binom{i-j}{\ell-j} \ell^{i-j}}{(i-j)!(1-\ell x)}.$$

Hence,

$$a_i(x) = \sum_{j=1}^i \sum_{\ell=j}^i \frac{(-1)^{\ell-i} \binom{i-j}{\ell-j} \ell^{i-j} x^i}{(i-j)!(1-\ell x)}.$$

We inspect the coefficient of x^m in $a_i(x)$ and derive

$$a_{mi} = \sum_{j=1}^i \sum_{\ell=j}^i \frac{(-1)^{\ell-i} \binom{i-j}{\ell-j} \ell^{m-j}}{(i-j)!}.$$

Therefore, (10) yields

$$\frac{d^m}{dx^m}T(x) = \sum_{i=1}^m \left(\sum_{j=1}^i \sum_{\ell=j}^i \frac{(-1)^{\ell-i} \binom{i-j}{\ell-j} \ell^{m-j}}{(i-j)!} \right) e^{e^x + ix-1} + \int_0^x e^{(x-t)e^t + e^t + (m+1)t-1} dt.$$

Finally, setting $x = 0$, we complete the proof of Corollary 1.3.

3 Proof of Theorem 1.4

The proof relies on the use of a generator of a uniformly random set partition of $[n]$ proposed by Stam [18]. We next describe Stam's algorithm for a given n .

1. For $m \in \mathbb{N}$, let $\mu_n(m) = \frac{m^n}{m! e B_n}$. Dobinski's formula (1) shows that $\mu_n(\cdot)$ is a probability distribution on \mathbb{N} .

At time zero, choose a random $M \in \mathbb{N}$ distributed according to μ_n , and arrange M empty and unlabeled boxes.

2. Arranges n balls labeled by integers from the set $[n]$.

At time $i \in [n]$, place the ball ' i ' into one of the M boxes, chosen uniformly at random. Repeat until there are no balls remaining.

3. Label the boxes in the order that they cease to be empty. Once a box is labeled, the label does not change.
4. Form a set partition π of $[n]$ with i in the k -th block if and only if ball ' i ' is in the k -th box.

Let N_i be the random number of nonempty boxes right after placing the i -th ball and X_i be the label of the box where the i -th ball was placed. Notice that if the i -th ball is dropped in an empty box, then $X_i = N_{i-1} + 1$ and $N_i = N_{i-1} + 1$. Otherwise, if the box was occupied previously, $X_i = X_j$ where $j < i$ is the first ball that was dropped in that box and $N_i = N_{i-1}$. Then, $X := X_1 \cdots X_n$ is the random set partition of $[n]$ produced by the algorithm.

We denote by $P_m(\cdot)$ the conditional probability distribution $P(\cdot | M = m)$. Clearly $N_1 = 1$, $N_i \leq i$, and

$$P_m(N_{i+1} = t + 1 | N_i = t) = \frac{m - t}{m} \quad \text{and} \quad P_m(N_{i+1} = t | N_i = t) = \frac{t}{m}.$$

Let $\alpha_{i,t}(m) := P_m(N_i = t)$. Then, taking into account that

$$P_m(N_i = t) = P_m(N_i = t, N_{i-1} = t - 1) + P_m(N_i = t, N_{i-1} = t),$$

we obtain:

$$\alpha_{i,t}(m) = \begin{cases} \frac{t}{m} \alpha_{i-1,t}(m) + \frac{m-t+1}{m} \alpha_{i-1,t-1}(m) & \text{if } 2 \leq t \leq m \text{ and } t \leq i \\ 0 & \text{if } t > i \text{ or } t > m \\ \frac{1}{m^{i-1}} & \text{if } t = 1 \text{ and } 1 \leq i. \end{cases}$$

Recall that one can define the sequence of Stirling numbers of the second kind as the solution to the recursion

$$S_{n,k} = k S_{n-1,k} + S_{n-1,k-1}, \quad n, k \in \mathbb{N}, k \leq n. \quad (12)$$

A comparison with (12) reveals that for $t \leq m$,

$$P_m(N_i = t) = \frac{S_{i,t}}{m^i} \frac{m!}{(m-t)!}.$$

In addition,

$$P_m(X_{i+1} = \ell | N_i = t) = \begin{cases} \frac{1}{m} & \text{if } \ell \leq t \\ \frac{m-t}{m} & \ell = t + 1 \\ 0 & \text{otherwise.} \end{cases}$$

Notice that some of the boxes may remain empty at the end of the algorithm's run.

In order to obtain the probability distribution of \mathcal{F}_n , we make a simple observation that \mathcal{F}_n has the same distribution as that of the random variable $1 \leq \mathcal{J} \leq n$ defined as

$$\mathcal{J} := \min \left(n, \max \{j | X_1 < \dots < X_j\} \right) = \min \left(n, \max \{j | N_j = j\} \right).$$

We will consider three cases;

Case $j = 1$: By the observation above

$$\begin{aligned} P(\mathcal{F}_n = j) &= E \left(P_M(N_2 = 1 | N_1 = 1) P_m(N_1 = 1) \right) = E \left(\frac{1}{M} \right) \\ &= \sum_{m=1}^{\infty} \frac{m^{n-1}}{m! e B_n} = \frac{B_{n-1}}{B_n}. \end{aligned} \quad (13)$$

Case $2 \leq j < n$: We consider two possibilities such that either (i) $m < j$, where

$$P_m(N_j = j, N_{j+1} = j) = 0,$$

or (ii) $2 \leq j \leq m$, where

$$\begin{aligned} P_m(N_j = j, N_{j+1} = j) &= P_m(N_{j+1} = j | N_j = j) \prod_{s=1}^{j-1} P_m(N_{s+1} = s + 1 | N_s = s) \\ &= \frac{j}{m} \prod_{s=1}^{j-1} \frac{m-s}{m}. \end{aligned}$$

Thus, for $2 \leq j < n$, our observation implies

$$P(\mathcal{F}_n = j) = \sum_{m=j}^{\infty} \frac{m^n}{m! e B_n} \frac{j}{m} \prod_{s=1}^{j-1} \frac{m-s}{m} = \frac{j}{e B_n} \sum_{m=j}^{\infty} \frac{m^{n-j-1}}{(m-j)!} = \frac{j \Theta_{n-j-1}(j)}{B_n}, \quad (14)$$

where we used (6) for the last equality.

Case $j = n$: We again investigate two possibilities such that either $m < n$ or $m \geq n$. Similar to the previous case, the latter is the only one contributing to the sum. Hence,

$$P(\mathcal{F}_n = n) = E(P_M(N_n = n)) = \sum_{m=n}^{\infty} \frac{1}{(m-n)! e B_n} = \frac{1}{B_n}. \quad (15)$$

Finally, equations (13), (14), and (15) yield (7).

References

- [1] M. Archibald, A. Blecher, and A. Knopfmacher, Fixed points in compositions and words, *J. Integer Seq.* **23** (2020), Article 20.11.1.
- [2] R. Arratia and S. Tavaré, The cycle structure of random permutations, *Ann. Probab.* **20** (1992), 1567–1591.
- [3] M. Bóna, On a balanced property of derangements, *Electron. J. Combin.* **13** (2006), #R102.
- [4] R. A. Brualdi, *Introductory Combinatorics*, 5th ed., Prentice-Hall, 2010.
- [5] N. Cakic, T. Mansour, and R. Smith, Elements protected by records in set partitions, *J. Diff. Eq. Appl.* **24** (2018), 1880–1893.
- [6] P. J. Cameron, *Combinatorics: Topics, Techniques, Algorithms*, Cambridge University Press, 1994.
- [7] Y. Cai and M. A. Readdy, q -Stirling numbers: a new view, *Adv. in Appl. Math.* **86** (2017), 50–80.
- [8] L. Comtet, *Advanced Combinatorics. The Art of Finite and Infinite Expansions*, revised and enlarged edition, D. Reidel Publishing Co., 1974.
- [9] A. Conflitti and R. Mamede, Gray codes and lexicographical combinatorial generation for nonnesting and sparse nonnesting set partitions, *Theoret. Comput. Sci.* **592** (2015), 87–96.
- [10] E. Deutsch and S. Elizalde, The largest and the smallest fixed points of permutations, *European J. Combin.* **31** (2010), 1404–1409.
- [11] P. Diaconis, J. Fulman, and R. Guralnick, On fixed points of permutations, *J. Algebraic Combin.* **28** (2008), Article 189.
- [12] D. Galvin and A. Pacurar, *Total non-negativity of some combinatorial matrices*, 2019, preprint is available at <https://arxiv.org/abs/1807.08658>.
- [13] G.-N. Han and G. Xin, Permutations with extremal number of fixed points, *J. Combin. Theory Ser. A* **116** (2009), 449–459.
- [14] A. Knopfmacher, T. Mansour, and S. Wagner, *Records in set partitions*, *Electron. J. Combin.* **17** (2010), #R109.
- [15] T. Mansour, *Combinatorics of Set Partitions*, CRC Press, 2013.
- [16] T. Mansour, R. Rastegar, and A. Roitershtein, *Height of records in partitions of a set*, Available at <https://arxiv.org/abs/1908.00846>.
- [17] T. Mansour, R. Rastegar, and A. Roitershtein, *Horizontal visibility graph of a random restricted growth sequence*, *Adv. Appl. Math.* **124** (2021) 102145.
- [18] A. J. Stam, *Generation of random partitions of a set by an urn model*, *J. Combin. Theory Ser. A* **35** (1983), 231–240.
- [19] R. P. Stanley, *Enumerative Combinatorics*, Vol. 1, Cambridge University Press, 1986.
- [20] H. S. Wilf, *generatingfunctionology*, A. K. Peters, 1990.