

Completely regular codes in Johnson and Grassmann graphs with small covering radii

I. Yu. Mogilnykh *

*Sobolev Institute of Mathematics
Tomsk State University
Novosibirsk State University*

January 12, 2022

Abstract

Let \mathcal{L} be a Desarguesian 2-spread in the Grassmann graph $J_q(n, 2)$. We prove that the collection of the 4-subspaces, which do not contain subspaces from \mathcal{L} is a completely regular code in $J_q(n, 4)$. Similarly, we construct a completely regular code in the Johnson graph $J(n, 6)$ from the Steiner quadruple system of the extended Hamming code. We obtain several new completely regular codes with covering radius 1 in the Grassmann graph $J_2(6, 3)$ using binary linear programming.

Keywords: Completely regular code, Desarguesian spread, q -ary design, Steiner quadruple system, Johnson scheme, Grassmann scheme, eigenvalue approach, binary linear programming

1 Introduction

The notion of a completely regular code was introduced by Delsarte in [10] as a generalization of a perfect code. It is known that all perfect codes in the Grassmann graphs [9] are trivial and their nonexistence in Johnson graphs is proven for a large number of finite cases [14], [12]. Therefore the completely regular codes in these graphs are of interest as they are related to designs and some geometrical objects.

During the years, researchers used several different names for the completely regular codes with covering radius 1. These objects could be equivalently defined in terms of: equitable 2-partitions [28], [30], perfect 2-colorings [13], [1], [26], [19], [25], intriguing sets [11] and others.

*
E-mail address: ivmog@math.nsc.ru

We refer to Ph.D. thesis of Martin [23] for an introduction on completely regular codes in Johnson graphs and a survey of Borges, Rifa and Zinoviev [20] on a recent progress in the study of completely regular codes in Hamming and Johnson graphs.

In [10] Delsarte noted interrelations of the completely regular codes in the Hamming and the Johnson graphs with orthogonal arrays and t -designs respectively. In order to emphasize this connection Martin [21] suggested the term "a completely regular design of strength t ". He studied several well-known classes of designs from the point of view of complete regularity and showed that $(k-1) - (n, k, \lambda)$ -designs are completely regular. This also holds for q -ary designs, therefore 2-spreads and 2-ary Steiner triple system [6] are completely regular. In [1] it was shown that if D is a $(k-1) - (n, k, 1)$ -design, then the code of the $(k+1)$ -subsets that does not contain any block of D is completely regular in the Johnson graph $J(n, k+1)$. The integer necessary conditions for the existence of t -designs were exploited for showing the nonexistence of constant weight perfect codes [12], [14] and completely regular codes with covering radius 1 [26].

The completely regular designs of strength 0 in the Hamming and the Johnson graphs were characterized by Meyerowitz in [24] and have a rather simple structure. The completely regular codes of zero strength in the strongly regular Grassmann graphs are known as the Cameron-Liebler line classes [7]. Contrary to the ordinary designs [24] the complete classification of these objects is done only in some particular cases. Several approaches for studying completely regular codes in Johnson graphs were applied to Cameron-Liebler line classes in [27].

The completely regular codes of strength 0 with covering radius 1 in Grassmann graph $J_q(n, k)$ for $k \geq 3$ could be considered as a generalization of Cameron-Liebler line classes. Somewhat trivial examples of these codes are the subspaces containing a point; the subspaces contained in a hyperplane; the subspaces contained in a hyperplane or containing a point for a non-incident point-hyperplane pair. There are not known any other examples of such codes in $J_q(n, k)$ for $k \geq 3$.

The completely regular codes in the Johnson graphs having strength 1 were characterized in the following cases: the minimum distance at least 3 by Martin in [22] and covering radius 1 recently by Vorob'ev [30]. De Winter and Metsch in [25] considered completely regular codes with covering radius 1 and strength 1 in the Grassmann graphs of diameter 3. They found two new series of examples of such codes: the 3-subspaces that do not contain a space of 2-spread and the code arising from symplectic polar space. The first series is similar to a construction from [1] for completely regular codes in the Johnson graphs $J(n, 4)$ and $J(n, 5)$ from Steiner triple and quadruples systems.

In Section 2 we give basic definitions and review the theory developed by Delsarte and Martin in the q -analog case. In Section 3 we consider the completely regular codes with covering radii 2 in regular graphs. We show that these sufficient condition for existence of such codes could be formulated in terms of certain eigenvectors of these graphs. In section 4 we discuss a spectral property

of the inclusion matrix of t -subspaces vs k -subspaces. We show that the code of $(k + 1)$ -subspaces not containing subspaces of $(k - 1) - (n, k, 1)_q$ -design is completely regular in the Grassmann graph of $(k + 1)$ -subspaces. This generalizes a series of De Winter and Metsch [25] and [1]. In Sections 5 and 6 we go further by applying this idea to the most symmetric designs, i.e. when a $(k - 1) - (n, k, 1)_q$ -design is the Desarguesian 2-spread or the Steiner quadruple system of extended Hamming code. In this case the code of $(k + 2)$ -subspaces (subsets), which do not contain subspaces (subsets) of $(k - 1) - (n, k, 1)_q$ -design is completely regular in the Grassmann (Johnson) graph.

The linear programming approach is a popular method for construction and classification of codes. Binary versions of linear programming earlier showed promising results for settling the nonexistence of completely regular codes [4] as well as finding these objects [18]. In Section 7 we obtain several new completely regular designs of strength 1 and covering radius 1 in the Grassmann graph $G_2(6, 3)$ by binary integer programming with a prescribed subgroup of its automorphism group. We outline the known results on the completely regular codes with $\rho = 1$ in this graph in a parameter table.

2 Definitions and Basic theory

2.1 The eigenspaces of the Johnson and Grassmann graphs

In what follows we abbreviate k -element subset and k -dimensional subspace to k -subset and k -subspace respectively. The vertices of the Johnson graph $J(n, k)$ are k -subsets of the set $\{1, \dots, n\}$ and the edges are pairs of subsets meeting in a $(k - 1)$ -subset. The vertices of the Grassmann graph $J_q(n, k)$ are k -subspaces of F_q^n and the edges are pairs of subspaces meeting in a $(k - 1)$ -subspace. These are well-known series of distance-regular graphs. Below we consider $k \leq n/2$. We also denote the Johnson graph $J(n, k)$ by $J_1(n, k)$ to emphasize that a result holds for Johnson and Grassmann graphs simultaneously. We use the notations $\begin{bmatrix} n \\ k \end{bmatrix}_q$ for q -binomial coefficient and $\begin{bmatrix} n \\ k \end{bmatrix}_1$ for its limit value, i.e. the ordinary binomial coefficient.

A vector v is called an *eigenvector* of a graph Γ with eigenvalue θ if it is an eigenvector of the adjacency matrix of Γ with eigenvalue θ . The following representation for the eigenspaces of the Johnson and Grassmann graphs could be found in [17], see also [10, Section 4.2]. We arrange the eigenvalues of $J_q(n, k)$, $q \geq 1$ in descending order starting from zeroeth and denote them by $\theta_{i,q}(n, k)$, $i \in \{0, \dots, k\}$. For i -subspace (i -subset if $q = 1$) X let us consider the characteristic vectors of all k -subspaces of F_q^n (k -subsets) containing X . Let $T_i(k)$ be the linear span of these vectors where X runs through all i -subspaces (i -subsets) of F_q^n (the set $\{1, \dots, n\}$).

Theorem 1. [17, Theorem 4.16] For any $i, 0 \leq i \leq k$, $U_i = T_i(k) \cap T_{i-1}^\perp(k)$ is

the eigenspace of $J_q(n, k)$, $q \geq 1$, with eigenvalue

$$\theta_{i,q}(n, k) = q^{i+1} \begin{bmatrix} n-k-i \\ 1 \end{bmatrix}_q \begin{bmatrix} k-i \\ 1 \end{bmatrix}_q - \begin{bmatrix} i \\ 1 \end{bmatrix}_q.$$

Below we omit the index q in $\theta_{i,q}(n, k)$ when we speak of the eigenvalues of the Johnson graphs.

2.2 Completely regular codes and q -ary designs

Let C be a code in a regular graph Γ . A vertex x is in C_i if the minimum of the distances between x and the vertices of C is i . The maximum of these distances is called the *covering radius* of C and is denoted by ρ . The *distance partition* of the vertices of Γ with respect to $C_0 = C$ is $\{C_i : i \in \{0, \dots, \rho\}\}$.

A code C is called *completely regular* [29] if there are numbers $\alpha_0, \dots, \alpha_\rho, \beta_0, \dots, \beta_{\rho-1}, \gamma_1, \dots, \gamma_\rho$ such that any vertex of C_i is adjacent to exactly α_i, β_i and γ_i vertices of C_{i-1}, C_i and C_{i+1} respectively. Note that $\alpha_0, \dots, \alpha_\rho$ can be found from the remaining numbers and the valency of the graph. The set $\{\beta_0, \dots, \beta_{\rho-1}; \gamma_1, \dots, \gamma_\rho\}$ is called the *intersection array* of the completely regular code C .

For a completely regular code C consider the $(\rho + 1) \times (\rho + 1)$ matrix A such that $A_{i,j}$ equals the number of vertices of C_j adjacent to a fixed vertex of C_i . The eigenvalues of the matrix A are called the *eigenvalues of the completely regular code* C .

Theorem 2. [8, Theorem 4.5] (Lloyd's theorem) *The eigenvalues of a completely regular code in a graph Γ are eigenvalues of Γ .*

The eigenvalues of a completely regular code with covering radius 1 are easy to find, see e.g. [28, Proposition 1].

Proposition 1. *Let C be a completely regular code in a m -regular graph Γ with $\rho = 1$ and intersection array $\{\beta_0; \gamma_1\}$. Then the size of C is $|V(\Gamma)|\gamma_1/(\gamma_1 + \beta_0)$ and the eigenvalues of C are m and $m - \beta_0 - \gamma_1$.*

In the rest of the section we consider codes in the Johnson or the Grassmann graphs. A collection D of k -subspaces (k -subsets when $q = 1$) of F_q^n (of $\{1, \dots, n\}$) is a $t - (n, k, \lambda)_q$ -design, if any t -subspace of F_q^n (t -element subset of $\{1, \dots, n\}$) is contained in exactly λ elements of D . In throughout of what follows we consider only designs without repeated blocks. The *strength* of D is the maximum t such that D is a t -design. When $q \geq 2$, a $1 - (n, k, 1)_q$ -design is called a k -spread. It is well-known that k -spreads exist if and only if k divides n .

Let χ_C be the characteristic vector of a code C in the graph $J_q(n, k)$. Consider the decomposition of χ_C over the eigenspaces U_0, \dots, U_k of $J_q(n, k)$:

$$\chi_C = u_0 + u_{i_1} + \dots + u_{i_s}, \tag{1}$$

where $u_0 \in U_0$ and $u_{i_j} \in U_{i_j}$, $j = 1, \dots, s$. The number s in the decomposition (1) is called the *dual degree* of the code C [10].

We make use of several results that were stated for the completely regular codes in the Johnson graphs by Delsarte [10] and Martin [21]. The arguments of the proofs for the q -ary generalization of these results could be obtained by replacing "subset" with "subspace" and follow from the description of the eigenspaces of $J_q(n, k)$ in Section 2.1.

Theorem 3. *Let C be a code in $J_q(n, k)$, $q \geq 1$ such that the decomposition (1) holds. Then we have the following:*

1. [10, Theorem 4.2] *The strength of C as q -ary design is equal to $\min\{i_j : j \in \{1, \dots, s\}\} - 1$.*
2. [10, Theorem 5.13] *If the minimum distance of C is greater or equal to $2s - 1$ then C is completely regular.*
3. [10, Theorem 5.10] *The covering radius of C is not greater than s .*
4. [21, Corollary 3.4] *If C is completely regular then its strength is equal to*

$$\min\{i \geq 0 : \theta_{i,q}(n, k) \text{ is an eigenvalue of } C\} - 1.$$

The following statement for the Johnson graphs could be found in [26, Corollary 1].

Corollary 1. *Let C be a completely regular code C in $J_q(n, k)$, $q \geq 1$ with covering radius 1 and intersection array $\{\beta_0; \gamma_1\}$. Then we have the following:*

1. *The code C is of size $\binom{n}{k}_q \beta_0 / (\gamma_1 + \beta_0)$ and its eigenvalues are numbers $\binom{n-k}{1}_q \binom{k}{1}_q$ and $\binom{n-k}{1}_q \binom{k}{1}_q - \gamma_1 - \beta_0$.*
2. *The strength of C is t where*

$$\theta_{t+1,q}(n, k) = \binom{n-k}{1}_q \binom{k}{1}_q - \gamma_1 - \beta_0.$$

Moreover, the numbers $\binom{n-i}{k-i}_q \beta_0 / (\gamma_1 + \beta_0)$ are integers for any $i \in \{0, \dots, t\}$.

Proof. The eigenvalues of C and the expression $|C| = \binom{n}{k}_q \beta_0 / (\gamma_1 + \beta_0)$ follow from Proposition 1 and the strength of C is by the fourth Statement of Theorem 3. The integer necessary conditions for t -design imply that for any $i \in \{0, \dots, t\}$ the number of the subspaces of C containing an i -subspace is $\binom{n-i}{k-i}_q \beta_0 / (\gamma_1 + \beta_0)$. \square

Corollary 2. 1. [21, Corollary 3.5], [3, Corollary 8] *For $q \geq 1$ any $(k - 1) - (n, k, \lambda)_q$ -design is a completely regular code in $J_q(n, k)$ with eigenvalue $\theta_{k,q}(n, k)$ and covering radius $\rho = 1$.*

2. *Any 3-spread is a completely regular code in the Grassmann graph $J_q(n, 3)$ with $\rho = 2$.*

Proof. We follow the considerations from [21, Corollary 3.5]. By Statement 1 of Theorem 3 we see that the dual degrees of a $(k - 1) - (n, k, \lambda)_q$ -design and a 3-spread are 1 and s respectively, where $s \leq 2$. Since covering radius of 3-spread is 2, its dual degree is also 2 by Statement 3 of Theorem 3. The result follows from Statement 2 of Theorem 3. \square

3 Auxiliary statements

Let the positions of a vector u be indexed by the vertices of a graph Γ . If the set of the pairwise different values of u is $\{a_0, \dots, a_r\}$, denote by C^i the set of the vertices of Γ such that $u_x = a_i$, $i \in \{0, \dots, r\}$. The partition $\{C^0, \dots, C^r\}$ is called the *partition associated* to the vector u . We see that the eigenvectors taking two or three values are tightly related with the completely regular codes of covering radii 1 and 2.

Theorem 4. 1. *A code C is completely regular in a m -regular graph Γ with $\rho = 1$ and eigenvalue θ , $\theta \neq m$ if and only if $\{C, \overline{C}\}$ is the partition associated to an eigenvector of Γ with eigenvalue θ .*

2. *Let Γ be a m -regular graph Γ , $\{C^0, C^1, C^2\}$ be the partition associated to an eigenvector u of Γ . If there are no edges between C^0 and C^2 and any vertex of C^1 is adjacent to exactly β_1 vertices of C^2 then C^0 is a completely regular code in Γ .*

Proof. 1. The result could be found in [11, Proposition 3.2] or [1, Proposition 1].

2. Let a_0, a_1, a_2 be the values of u on C^0, C^1, C^2 respectively. Let a vertex x of C^0 be adjacent to $\alpha_0(x)$ vertices of C^0 . Since there are no edges between the vertices C^0 and C^2 , the vertex x is adjacent to $\beta_0(x) = m - \alpha_0(x)$ vertices of C^1 . Consider the sum of the values of u on the neighbors of x . Since u is an eigenvector with eigenvalue θ we have the following:

$$\theta a_0 = (m - \beta_0(x))a_0 + \beta_0(x)a_1. \quad (2)$$

This implies that $\alpha_0(x)$ and $\beta_0(x)$ do not depend on x . The same argument on C^1 and C^2 implies that $x \in C^2$ has exactly γ_2 and α_2 neighbors in C^1 and C^2 respectively. This follows from $\alpha_2 + \gamma_2 = m$ and the equation

$$\theta a_2 = (m - \alpha_2)a_1 + \alpha_2 a_2. \quad (3)$$

Let a vertex x of C^1 be adjacent to $\gamma_1(x)$, $\alpha_1(x)$ and β_1 vertices of C^0, C^1 and C^2 respectively. Note that β_1 does not depend on x by the condition of the theorem. Taking into account that $\alpha_1(x) + \beta_1 + \gamma_1(x) = m$, the sum of the values of u on the neighbors of x is

$$\theta a_1 = \gamma_1(x)a_0 + (m - \gamma_1(x) - \beta_1)a_1 + \beta_1 a_2, \quad (4)$$

The above implies that $\alpha_1(x)$ and $\gamma_1(x)$ do not depend on x and C^0 is a completely regular code by the definition. □

4 Inducing map and completely regular codes

For $q \geq 1$ and $l \geq k$ consider the matrix $I_{l,k}$ whose rows and columns are indexed by the vertices of $J_q(n, l)$ and those of $J_q(n, k)$ respectively, $I_{l,k}(x, y)$

is 1 if y is contained in x and 0 otherwise. Kantor in [15] proved that $I_{l,k}$ is a full rank matrix. Moreover, it is clear that the left multiplication by $I_{l,k}$ maps the column-vectors of the subspace $T_i(k)$ to those of $T_i(l)$ (see Section 2.1 for their definitions). Therefore, $I_{l,k}$ is a bijection from the eigenspace $U_{i,q}(n, k)$ to $U_{i,q}(n, l)$ for any $i \in \{0, \dots, k\}$.

Theorem 5. *Let u be an eigenvector of $J_q(n, k)$ with eigenvalue $\theta_{i,q}(n, k)$, $q \geq 1$. Then for any $l \geq k$ the vector $I_{l,k}u$ is an eigenvector of $J_q(n, l)$ with eigenvalue $\theta_{i,q}(n, k)$.*

We make use of the theorem above for obtaining a series of completely regular designs.

Theorem 6. 1. [1, Proposition 4] *Let D be a $(k-1) - (n, k, 1)$ -design. Then*

$$\{U : U \subset \{1, \dots, n\}, |U| = k+1, |\{V \in D : V \subset U\}| = 0\}$$

is a completely regular code in the Johnson graph $J(n, k+1)$ with $\rho = 1$.

2. *Let D be a $(k-1) - (n, k, 1)_q$ -design. Then*

$$\{U : U < F_q^n, \dim(U) = k+1, |\{V \in D : V < U\}| = 0\}$$

is a completely regular code in the Grassmann graph $J_q(n, k+1)$ with $\rho = 1$.

Proof. 2. We use the approach of work [1] based on the inclusion matrix. The code D is completely regular with covering radius 1 by Corollary 2. By the first statement of Theorem 4 there is an eigenvector u of $J_q(n, k)$ with associated partition $\{D, \overline{D}\}$. A $(k+1)$ -subspace of F_q^n contains either 0 or 1 subspaces of D . These two facts combined imply that $I_{k+1,k}u$ takes only two values. Moreover, the associated partition to vector $I_{k+1,k}u$ is the partition into the codes

$$\{U : U < F_q^n, \dim(U) = k+1, |\{V \in D : V < U\}| = 0\} \text{ and}$$

$$\{U : U < F_q^n, \dim(U) = k+1, |\{V \in D : V < U\}| = 1\}.$$

By the first statement of Theorem 5 we see that $I_{k+1,k}u$ is an eigenvector of $J_q(n, k+1)$. By the first statement of Theorem 4 we obtain the required. \square

Remark 1. A combinatorial proof for the statement above in case when k is 2 (i.e. 2-spreads) for the Grassmann graphs could be found in [25, Lemma 12]. Apart from 2-spreads the only known example of $(k-1) - (n, k, 1)_q$ -design, $q \geq 2$ is the Steiner triple system constructed in [6]. This implies the existence of a completely regular code in $J_2(13, 4)$ by Theorem 6.

In the sections below we proceed further with the idea described in Theorem 6 and show that the Steiner quadruple systems of the extended Hamming code and the Desarguesian 2-spreads produce completely regular codes in the Johnson graph $J(n, 6)$ and the Grassmann graph $J_q(n, 4)$ respectively.

5 Completely regular code in the Johnson graph $J(n, 6)$ from the SQS of the extended Hamming code

We use the traditional point-block terms throughout the section. Consider the design \mathcal{Q} whose points are the coordinate positions and blocks are the supports of the codewords of weight 4 of the extended Hamming code of length n . It is well-known that \mathcal{Q} is a $3 - (n, 4, 1)$ -design, also known as a *Steiner quadruple system*.

The codes \mathcal{Q} and $\{x : x \subset \{1, \dots, n\}, |x| = 5, |\{B \in \mathcal{Q}, B \subset x\}| = 0\}$ are completely regular in $J(n, 4)$ and $J(n, 5)$ respectively by Corollary 2 and Theorem 6.

Theorem 7. *Let \mathcal{Q} be the Steiner quadruple system of the extended Hamming code of length n . The code $\{x : x \subset \{1, \dots, n\}, |x| = 6, |\{B \in \mathcal{Q}, B \subset x\}| = 0\}$ is completely regular in $J(n, 6)$ with $\rho = 2$.*

Proof. Let B and B' be two blocks of \mathcal{Q} , $|B \cap B'| = 2$. Because the extended Hamming code is linear the symmetric difference $B \Delta B'$ is also a block of \mathcal{Q} . Therefore any 6-subset of $\{1, \dots, n\}$ contains 0, 1 or 3 blocks of \mathcal{Q} . We consider the following codes:

$$\begin{aligned} C^0 &= \{x : x \subset \{1, \dots, n\}, |x| = 6, |\{B \in \mathcal{Q} : B \subset x\}| = 0\}, \\ C^1 &= \{x : x \subset \{1, \dots, n\}, |x| = 6, |\{B \in \mathcal{Q} : B \subset x\}| = 1\}, \\ C^2 &= \{x : x \subset \{1, \dots, n\}, |x| = 6, |\{B \in \mathcal{Q} : B \subset x\}| = 3\} \end{aligned}$$

that partition the vertices of $J(n, 6)$.

We are now to show that C^0 is a completely regular code and $\{C^0, C^1, C^2\}$ is the distance partition. In view of the second statement of Theorem 4 we prove that the vertices of C^0 and C^2 are disjoint and a vertex of C^1 is adjacent to exactly 6 vertices of C^2 . We finish the proof by noting that $\{C^0, C^1, C^2\}$ is the partition associated to an eigenvector. Note that an alternative proof could be done by solely combinatorial arguments for the existence of the intersection array.

A 6-subset from C^2 contains three blocks of \mathcal{Q} . Moreover, the symmetric difference of any two of these blocks is the third block. We see that the following holds:

$$\text{for any } x \in C^2 \text{ and any } i \in x, \text{ there is } B \in \mathcal{Q}, i \notin B, B \subset x. \quad (5)$$

From (5) we conclude that the vertices of C^0 and C^2 are nonadjacent in the Johnson graph $J(n, 6)$.

Let y be a vertex of C_1 and B be a unique block of \mathcal{Q} such that $B \subset y$. Let $x \in C^2$ be a vertex that is adjacent to y in $J(n, 6)$, i.e. $x = (y \setminus \{i\}) \cup \{j\}$ for some $i \notin y, j \in y$.

Suppose i is in B . Then by property (5) there is a block B' of \mathcal{Q} that is a subset of $x \setminus \{j\}$. We see that $B' \subset (x \setminus \{j\}) \subset y$. Moreover B' is not B , because $i \in B$, $i \notin x$ and $B' \subset x$. We have that distinct blocks B' and B from \mathcal{Q} are subsets of y , which contradicts $y \in C^1$.

We have that $i \in y \setminus B$. Then the following blocks of \mathcal{Q} are subsets of x : B , $\{s, t, l, j\}$ and $B\Delta\{s, t, l, j\}$ for some $s, t \in B$, $\{l\} = x \setminus (B \cup \{j\})$. On the other hand, given the 2-subset $\{s, t\}$ of B and the point l from $y \setminus B$ the point j could be reconstructed. Indeed, the block $\{s, t, l, j\}$ is a unique block in $3 - (n, 4, 1)$ -design \mathcal{Q} containing $\{s, t, l\}$. Since $\{s, t, l, j\}$ and $B\Delta\{s, t, l, j\}$ are contained in the same x from C^2 , we conclude that there are exactly $\binom{4}{2}_1 \cdot 2/2 = 6$ neighbors of y in C^2 .

We finally note that the partition $\{C^0, C^1, C^2\}$ is the partition associated to an eigenvector of $J(n, 6)$. The Steiner quadruple system \mathcal{Q} is a completely regular code with covering radius 1 and eigenvalue $\theta_4(n, 4)$ by Corollary 2. Then by the first statement of Theorem 4 there is an eigenvector v with eigenvalue $\theta_4(n, 4)$ such that $\{\mathcal{Q}, \overline{\mathcal{Q}}\}$ is the partition associated to v . By Theorem 5 the vector $I_{6,4}v$ is an eigenvector of $J(n, 6)$ with eigenvalue $\theta_4(n, 6)$. The definitions of the inclusion matrix $I_{6,4}$ and the codes C^0, C^1, C^2 imply that these codes form the partition associated to $I_{6,4}v$. By the second statement of Theorem 4 we conclude that C^0 is completely regular. \square

Remark 2. One might obtain the intersection array $\{60, 6; 4(n-15), 6(n-8)\}$ of the code from Theorem 7 by combinatorial arguments or following the proof of Theorem 4 from equations (2)-(4).

Corollary 3. *The set of the codewords of weight 6 of the extended Hamming code of length 16 is a completely regular code in $J(16, 6)$ with $\rho = 2$.*

Proof. The supports of the codewords of weight 6 are contained in the code $C = \{x : x \subset \{1, \dots, n\}, |x| = 6, |\{B \in \mathcal{Q}, B \subset x\}| = 0\}$ because otherwise the minimum distance of the extended Hamming code is 2. The definition of a completely regular code and the double counting of edges between C_i and C_{i+1} imply the following:

$$|C|\beta_0 = |C_1|\gamma_1, |C_1|\beta_0 = |C_2|\gamma_1, |C| + |C_1| + |C_2| = \binom{n}{6}_1.$$

The intersection array obtained in Remark 1, so for $n = 16$ we have

$$|C| + |C|60/4 + |C|6/48 = \binom{16}{6} = 8008$$

and therefore $|C|$ is 448. Since there are exactly 448 codewords of weight 6 in extended Hamming code of length 16 we conclude that they coincide with C , which is a completely regular code in $J(16, 6)$ by Theorem 7. \square

6 Completely regular code in $J_q(n, 4)$ from Desarguesian 2-spread

Let F' be the subfield of the field F_{q^n} of order q^2 . The elements of the multiplicative group of F_{q^n} are parted into the cosets of that of F' . We treat F_{q^n} as the vector space F_q^n and any multiplicative coset of F' corresponds to a 2-subspace of F_q^n . The collection of such subspaces is a 2-spread, which is called *Desarguesian*. A subspace is called F' -closed if its vectors (treated as elements of F_{q^n}) are closed under the multiplication by the elements of F' . In particular, the subspaces of a Desarguesian 2-spread are F' -closed. For a subset of F_q^n the minimal inclusion-wise F' -closed subspace that contains the subset is called its F' -closure.

Any 2-spread \mathcal{L} and all 3-subspaces of F_q^n that do not contain any subspace from \mathcal{L} are completely regular codes in $J_q(n, 2)$ and $J_q(n, 3)$ respectively. In case when \mathcal{L} is a Desarguesian spread we will show that the code $\{U : U < F_q^n, \dim(U) = 4, |\{X \in \mathcal{L} : X < U\}| = 0\}$ is a completely regular code in $J_q(n, 4)$.

Consider a 4-subspace U of F_q^n . It can contain 0, 1 or at least 2 subspaces from \mathcal{L} . Suppose X and X' are 2-subspaces from \mathcal{L} that are contained in U . Because X and X' meet only in a zero vector, all vectors of U are linear combinations of the vectors of X and X' . Moreover since X and X' are F' -closed, so is U . In other words, the nonzero vectors of U are parted by nonzero vectors from $q^2 + 1$ subspaces from \mathcal{L} . We have the following code partition of the vertices of $J_q(n, 4)$:

$$C^0 = \{U : U < F_q^n, \dim(U) = 4, |\{V \in \mathcal{L} : V < U\}| = 0\},$$

$$C^1 = \{U : U < F_q^n, \dim(U) = 4, |\{V \in \mathcal{L} : V < U\}| = 1\},$$

$$C^2 = \{U : U < F_q^n, \dim(U) = 4, |\{V \in \mathcal{L} : V < U\}| = q^2 + 1\}.$$

We now show some structural properties of these codes.

Lemma 1. *Any 3-subspace of U , $U \in C^2$ contains exactly one subspace from \mathcal{L} . In particular, the subspaces from C^0 and C^2 are nonadjacent in $J_q(n, 4)$.*

Proof. Let W be a 3-subspace of U that does not contain subspaces from \mathcal{L} . Since $F_q < F'$ we see that the F' -closure of any of nonzero vectors of U meets W in exactly $q - 1$ nonzero vectors. We see that the F' -closure of W has at least $(q^3 - 1)(q + 1)$ vectors, so its dimension is at least 5. This contradicts that the subspace U is F' -closed and that W is a subspace of U . □

In view of Lemma 1 one might consider C^2 to be the F' -closure of all 3-subspaces that contain exactly one subspace from \mathcal{L} . Indeed any such 3-subspace is spanned by a subspace X (which is F' -closed) from \mathcal{L} and a 1-subspace and the F' -closure of the latter one has dimension 2.

Lemma 2. *Any subspace U , $U \in C^1$ is adjacent to exactly $q+1$ subspaces from C^2 in $J_q(n,4)$.*

Proof. Let $X \in \mathcal{L}$ be that such that $X < U$. Let $V \in C^2$ be adjacent to U , i.e. $\dim(U \cap V) = 3$. By Lemma 1 the subspace $U \cap V$ of U must contain X . There are exactly $q+1$ 3-subspaces of U that contain X . Their F' -closures are in C^2 and we obtain the required. \square

Theorem 8. *Let \mathcal{L} be a Desarguesian 2-spread. The code $\{U : U < F_q^n, \dim(U) = 4, |\{V \in \mathcal{L} : V < U\}| = 0\}$ is completely regular in $J_q(n,4)$.*

Proof. The vertex set of $J_q(n,4)$ is parted into the codes C^0 , C^1 and C^2 . By Lemmas 1 and 2 there are no edges between C^0 and C^2 and any vertex from C^1 is adjacent to exactly 6 subspaces from C^2 .

Since \mathcal{L} is a completely regular code with covering radius 1, we see that $\{\mathcal{L}_0, \mathcal{L}_1\} = \{\mathcal{L}, \overline{\mathcal{L}}\}$ is the partition associated to an eigenvector v of $J_q(n,2)$ with eigenvalue $\theta_{2,q}(n,2)$. By Theorem 5 the vector $I_{4,2}v$ is an eigenvector of $J_q(n,4)$ with eigenvalue $\theta_{2,q}(n,4)$. The definition of the inclusion matrix $I_{4,2}$ and the definition of the codes C^0 , C^1 , C^2 imply that these codes form the partition associated to $I_{6,4}v$. The result follows from Theorem 4. \square

We note that following the proof of Theorem 4 one can obtain that the code given in Theorem 8 has the intersection array

$$\{[4]_q [3]_q, q+1; q^5(q+1) [1]_q^{[n-6]}, [4]_q [1]_q^{[n-4]}]_q q\}.$$

7 Completely regular codes with $\rho = 1$ in $J_2(6,3)$

Throughout this section by *the automorphism group* of a code (subset of the vertices) in a graph we mean the setwise stabilizer of the code in the automorphism group of the graph. Let G be a subgroup of the automorphism group of a m -regular graph Γ . Let O_1, \dots, O_r be the orbits of the action of G on the vertex set of Γ . Because O_1, \dots, O_r are orbits we see that given any $i, j \in \{1, \dots, r\}$ any vertex x of O_i is adjacent to exactly A_{ij} vertices of O_j and A_{ij} does not depend on x . Let A be the matrix $\{A_{ij}\}_{i,j \in \{1, \dots, r\}}$. Suppose the automorphism group of a code C has a subgroup G . We consider the characteristic vector $\chi_{C,G}$ of C with positions indexed by the orbits O_1, \dots, O_r , i.e. $(\chi_{C,G})_i = 1$ if and only if $O_i \subseteq C$. If $\mathbf{1}$ is the all-one vector then $\mathbf{1} - \chi_{C,G}$ is the characteristic vector of the complement of C . The notations above imply that

$$A\chi_{C,G} = (m - \beta_0)\chi_{C,G} + \gamma_1(\mathbf{1} - \chi_{C,G}) \quad (6)$$

holds if and only if C is a completely regular code in Γ with covering radius 1, intersection array $\{\beta_0; \gamma_1\}$ and G is a subgroup of its automorphism group.

One might consider (6) to be a binary linear programming problem with the binary variable vector $\chi_{C,G}$. We consider the application of the binary linear

programming approach for showing the existence of completely regular codes in the Grassmann graph $G_2(6, 3)$.

Let a be a primitive element of F_{2^6} . We set Γ_{21} to be the group generated by the multiplication of the vectors of F_2^6 (treated as the elements of F_{2^6}) by a^{21} . The vertices of $J_2(6, 3)$ are parted into 465 orbits of the group Γ_{21} .

Let C be a completely regular code in the graph $J_2(6, 3)$ with intersection array $\{\beta_0, \gamma_1\}$. Due to Corollary 1 the eigenvalues of C are the valency of $J_2(6, 3)$ which is 98 and

$$98 - \beta_0 - \gamma_1 = \theta_{i,2}(6, 3), \quad (7)$$

where $i - 1$ is the strength of C as a design.

Let the eigenvalue $98 - \beta_0 - \gamma_1$ be $\theta_{2,2}(6, 3)$, i.e. C is a q -ary 1-design. In this case a binary linear programming solver found solutions of system (6) for 8 different values of γ_1 . Two of the codes (with $\gamma_1 = 9$ and 21) were previously obtained in [25].

Theorem 9. *There are completely regular codes in $J_2(6, 3)$ with covering radius 1, intersection array $\{93 - \gamma_1; \gamma_1\}$ for any $\gamma_1 \in \{12, 15, 18, 24, 27, 30\}$ such that Γ_{21} is a subgroup of their automorphism groups.*

From (7) we see that the intersection array $\{\beta_0, \gamma_1\}$ of any completely regular code could be found from the strength of the code and γ_1 . We summarize the information on the intersection arrays of the completely regular codes in $J_2(6, 3)$ with $\rho = 1$ in Table 1. By integer conditions in the table we mean the integer necessary existence conditions for designs imposed by the second statement of Corollary 1.

Remark 3. Apart from the integer necessary conditions there are other techniques for proving the nonexistence of a completely regular code with $\rho = 1$ given a putative intersection array. One of them is a method based on the finding weight distribution of the code [2, Theorem 1] (see also [16]). In the case of Johnson and Hamming graphs some intersection arrays feasible by integer necessary conditions were shown to be infeasible using the weight distribution method [19], [16]. However, in the particular instance of the graph $G_2(6, 3)$ this approach is not stronger than the integer necessary conditions. A counting argument in [25, Lemma 21] implies the nonexistence of the completely regular codes with strength 1 and $\gamma_1 = 3$ in this graph.

A very intriguing open problem is solving the existence problem for completely regular code in $G_2(6, 3)$ with intersection arrays $\{56; 21\}$ and $\{64; 28\}$. These codes are designs of strength 0 and therefore could be viewed as a generalization of Cameron-Liebler line classes in the non-strongly regular case.

The only known completely regular codes in $G_2(6, 3)$ with $\rho = 2$ are the code of the subspaces containing a fixed 2-space and 3-spread in $G_2(6, 3)$ (Corollary 2).

Table 1: Completely regular codes in $J_2(6, 3)$ with $\rho = 1$. The intersection array $\{\beta_0, \gamma_1\}$ of any completely regular code is obtained from its strength and γ_1 using (7)

Eigenvalue	Design strength	Integer conditions	Nonexistence, γ_1	Existence, γ_1	Open cases, γ_1
35	0	$\gamma_1 \bmod 7 = 0$		$7^H, 14^{HP}$	21, 28
5	1	$\gamma_1 \bmod 3 = 0$	$3^{M''}$	$9^M, 21^{M'}, 3l, l = 4, 5, 6, 8, 9, 10^A$	$3l, l \in \{2\} \cup \{10, \dots, 15\}$
-7	2	$\gamma_1 \bmod 21 = 0$		$21^B, 42^B$	

H subspaces belonging to a hyperplane;

HP subspaces are in a hyperplane H or contain a vector v , where $v \notin H$;

A exists by Theorem 9;

B correspond to $2 - (6, 3, 3)_2$ - and $2 - (6, 3, 6)_2$ -designs, which exist by [5];

M totally isotropic subspaces of a symplectic polarity [25, Example 6];

M' 3-subspaces that do not contain subspaces from a 2-spread;

[25, Example 5], see also Theorem 6;

M'' nonexistence follows from [25, Lemma 21].

References

- [1] S.V. Avgustinovich, I.Yu. Mogilnykh, Induced perfect colorings, Sib. Electron. Math. Rep. 8 (2011) 310-316.
- [2] S. V. Avgustinovich and I. Yu. Mogilnykh. Perfect 2-colorings of Johnson graphs $J(6, 3)$ and $J(7, 3)$. In A. Barbero, editor, Coding Theory and Applications (Second International Castle Meeting, ICMCTA 2008, Castillo de la Mota, Medina del Campo, Spain, September 15-19, 2008. Proceedings), 7, volume 5228 of Lect. Notes Comput. Sci., pages 11–19. Springer-Verlag, Berlin Heidelberg, 2008.
- [3] S.V. Avgustinovich, V.N. Potapov, Combinatorial designs, difference sets, and bent functions as perfect colorings of graphs and multigraphs, Siberian Mathematical Journal, Vol. 61, No. 5, pp. 867-877, 2020.
- [4] J. Bamberg, There is no Cameron–Liebler line class of $PG(3, 4)$ with parameter 6, <http://symomega.wordpress.com/2012/04/01/there-is-no-cameron-liebler-line-class-of-pg34-with-parameter-6>
- [5] M. Braun, A. Kerber, R. Laue, Systematic Construction of q-Analogs of t -(v, k, l)-designs, Designs, Codes and Cryptography, 34, 55–70, 2005

- [6] M. Braun, T. Etzion, P. R. J. Ostergard, A. Vardy, and A. Wassermann. Existence of q-analogs of Steiner systems. E-print 1304.1462, arXiv.org, 2013. Available at <http://arxiv.org/abs/1304.1462>
- [7] P.J. Cameron, R.A. Liebler, Tactical decompositions and orbits of projective groups, *Linear Algebra Appl.* 46 (1982) 91–102.
- [8] D.M. Cvetkovic, M. Doob, H. Sachs, *Spectra of Graphs*, Academic Press, New York, London, 1980.
- [9] L. Chihara, On the zeros of the Askey-Wilson polynomials, with applications to coding theory, *SIAM J. Math. Anal.*, vol. 18, no. 1, pp. 191–207, 1987.
- [10] P. Delsarte, An algebraic approach to the association schemes of coding theory, *Philips Res. Rep. Suppl.* 10 (1973) 1-97.
- [11] Bart De Bruyn and Hiroshi Suzuki. Intriguing sets of vertices of regular graphs. *Graphs Combin.*, 26(5):629–646, 2010.
- [12] Etzion T. and Schwarz M., Perfect Constant-Weight Codes, *IEEE Trans. Inform. Theory*, Vol. 50, No. 9, 2156-2165, 2004.
- [13] D. G. Fon-Der-Flaass, Perfect 2-colorings of a hypercube. *Siberian Mathematical Journal.* 48(4) (2007) 740-745.
- [14] Gordon M.D., Perfect Single Error-Correcting Codes in the Johnson Scheme, *IEEE Trans. Inform. Theory*, Vol. 52, No. 10, 4670-4672 , 2006.
- [15] Kantor W. M., On incidence matrices of finite projective and affine space, *Math, Z.* i24, 315-318, 1972
- [16] D. S. Krotov. On weight distributions of perfect colorings and completely regular codes. *Des. Codes Cryptography*, 61(3):315-329, 2011
- [17] C. Maldonado and D. Penazzi. Lattices and Norton algebras of Johnson, Grassmann and Hamming graphs. E-print 1204.1947, arXiv.org, 2012. Available at <http://arxiv.org/abs/1204.1947>
- [18] D. S. Krotov, I. Yu. Mogilnykh, A. Yu. Vasil’eva, On completely regular codes of covering radius 1 in the halved hypercubes, E-print 1812.03159, arXiv.org, 2018. Available at <http://arxiv.org/abs/1812.03159>
- [19] E. A. Beshpalov, D. S. Krotov, A. A. Matushev, A. A. Taranenko, K. V. Vorob’ev, Perfect 2-colorings of Hamming graphs, E-print 1911.13151, arXiv.org, 2020. Available at <http://arxiv.org/abs/1911.13151>
- [20] J. Borges, J. Rifa, V.A. Zinoviev, On completely regular codes, *Probl. Inf. Transmiss.* 55 (2019) 1–45.
- [21] W.J. Martin, Completely regular designs, *J. Combin. Des.* 4 (1998) 261–273.

- [22] W.J. Martin, Completely regular designs of strength one, *Journal of Algebraic Combinatorics*, 3:2 (1994), 177–185. MR1268574
- [23] W. J. Martin Completely regular subsets Ph.D. Thesis University of Waterloo. 1992.
- [24] A. Meyerowitz, Cycle-balanced conditions for distance-regular graphs, *Discrete Math.* 264 (3) (2003) 149–166.
- [25] S. De Winter, K. Metsch Perfect 2-Colorings of the Grassmann Graph of Planes, *Electronic Journal of Combinatorics*, Volume 27, Issue 1, P1.21 (2020)
- [26] Mogilnykh I.Yu. On the regularity of perfect 2-colorings of the Johnson graph. *Probl. Inform. Transm.* 43(4), 271-277 (2007)
- [27] A.L. Gavrilyuk, I.Y. Mogilnykh. Cameron-Liebler line classes in $PG(n,4)$. *Des. Codes Cryptogr.* 73(3), 969–982 (2014).
- [28] Mogilnykh, I. Valyuzhenich A. Equitable 2-partitions of the Hamming graphs with the second eigenvalue. *Discrete Mathematics.* 343, 11, 112039, 2020.
- [29] A. Neumaier, Completely regular codes, *Discrete Mathematics V* 106–107, Issue 1, September 1992, P 353-360
- [30] K. Vorob'ev, Equitable 2-partitions of Johnson graphs with the second eigenvalue. E-print 2003.10956, 2020. Available at <http://arxiv.org/abs/2003.10956>