

MALLIAVIN CALCULUS FOR DEGENERATE DIFFUSIONS

A. S. ÜSTÜNEL

Abstract: Let (W, H, μ) be the classical Wiener space on \mathbb{R}^d . Assume that $X = (X_t(x))$ is a diffusion process satisfying the stochastic differential equation with diffusion and drift coefficients $\sigma : \mathbb{R}^n \rightarrow \mathbb{R}^n \otimes \mathbb{R}^d$, $b : \mathbb{R}^n \rightarrow \mathbb{R}^n$, B is an \mathbb{R}^d -valued Brownian motion. We suppose that b and σ are Lipschitz. Let $P(x)$ be the orthogonal projection from \mathbb{R}^d to its closed subspace $\sigma(x)^*(\mathbb{R}^n)$, assuming that $x \rightarrow P(x)$ is continuously differentiable, we construct a covariant derivative $\hat{\nabla}$ on the paths of the diffusion process, along the elements of the Cameron-Martin space and prove that this derivative is closable on $L^p(\nu)$, where ν represents the law of the above diffusion process, i.e., $\nu = X(x)(\mu)$, the image of the Wiener measure under the function $w \rightarrow X.(w, x)$. We study the adjoint of this operator and we prove several results: representation theorem for $L^2(\nu)$ -functionals, the logarithmic Sobolev inequality for ν . As applications of these results the proof of the Logarithmic Sobolev inequality on the path space of Dyson's Brownian motion is given by using the covariant derivative. We then explain how to use this theory for deriving the functional inequalities for the measures defined by the semigroups of the diffusion process at the time $t = 1$ and with fixed starting point. As an example we give a very simple proof of the Poincaré and log-Sobolev inequalities for the semigroup of the Brownian motion on the Heisenberg group with the optimal constants.

2020 Mathematics Subject Classification. Primary 60H; Secondary 22E30, 22E66, 35A23, 35K08, 35R03, 60J65.

Keywords: Entropy, degenerate diffusions, martingale representation, relative entropy, Dyson's Brownian motion, Poincaré inequality, logarithmic Sobolev inequality, intertwining, Heisenberg group, left and right invariant vector fields, opératur carré du champs.

CONTENTS

1. Introduction	2
2. Preliminaries	3
3. Conditional and Covariant Sobolev derivatives for degenerate diffusions' functionals	5
4. Poincaré and Logarithmic Sobolev Inequalities	12
5. Examples and Further Applications	13
5.1. Classical Situation	14
6. Log-Sobolev for Dyson's Brownian Motion	14
7. Intertwining Relations and Logarithmic Sobolev Inequality	16
8. An example: Poincaré and Log-Sobolev inequalities on the Heisenberg Group	20
References	24

1. Introduction

In a preceding work we have calculated a martingale representation theorem for the functionals of (degenerate) diffusion processes under the hypothesis of their weak uniqueness following the path breaking ideas originating from [6]. While doing the calculations an important concept shows up: the behaviour of the projection operator which is defined as the orthogonal projection from \mathbb{R}^d to the closed subspace $\sigma^*(t, X_t)(\mathbb{R}^n)$, where σ is the diffusion coefficient. This operator gives rise to the Itô representation theorem of the martingales adapted to the filtration of the diffusion process. In fact we have shown that, for any functional $F(X)$ defined on the paths of the diffusion process X , which is square integrable w.r.t. to its law, there exists an \mathbb{R}^d -valued process adapted to the filtration of X , say $\alpha(X)$, such that

$$F(X) = E[F(X)] + \int_0^1 P(X_s)\alpha_s(X) \cdot dB_s,$$

μ -a.s. Although α is not unique, the process $(P(X_s)\alpha_s(X), s \in [0, 1])$ is unique $ds \times d\mu$ -a.s. In the case of strong solutions, we have also chaos representation à la N. Wiener (cf.[13]) for the functionals of X using the basic $\mathcal{F}(X)$ -martingale m defined by

$$m_t = \int_0^t P(X_s)dB_s.$$

To calculate the integrands in this chaos development, as well as the process $P(X)\alpha(X)$ above, we need to create some kind of differential calculus on the paths of the diffusion process X . The first kind of derivative which comes to one's mind is some kind of extension of the H -derivative of Leonard Gross (cf.[10, 14]), where H denotes the Cameron-Martin space associated to the \mathbb{R}^d -valued Brownian motion which is at the origin of X , as it has been done in the case of flat Malliavin Calculus, cf. [18]. However this approach fails since H -derivatives of diffusion's functionals, even the simplest ones, fail to remain measurable with respect to the sigma algebra generated by the diffusion process. To circumvent this difficulty we propose a “**new derivation**” operator which consists of smoothing the H -derivative w.r.t. the final instant sigma algebra, namely $\mathcal{F}_1(X) = \sigma(X_t, t \in [0, 1])$, of the diffusion process. This operator is denoted by $\hat{\nabla}$ with a hat at the top to indicate the role of the conditional expectation in its definition. It is then easy to show that $\hat{\nabla}$, restricted to cylindrical diffusion functionals $\mathcal{S}(X) = \{f(X_{t_1}, \dots, X_{t_n}) : t_1 \leq t_2 \leq \dots \leq t_n \leq 1, f \in \mathcal{S}(\mathbb{R}^n), n \geq 1\}$ is a derivation (under almost sure equality) and it is a closable operator in $L^p(\nu)$ for $p > 1$, where ν denotes the law of X . We show several relations between the adjoint of this derivative operator (adjoint taken w.r.t. the law of X), and the martingale representation theorem; in particular we calculate explicitly the integrand for of this representation theorem which extends the flat case, which is called in the flat case Clark-Haussmann-Bismut-Ocone formula (cf. [16, 17]). Let us note nevertheless that in the case of degenerate diffusions the situation is quite different, for instance the derivative constructed above is not even torsion free. As immediate applications of the theory so developed, we prove the Poincaré inequality and the logarithmic Sobolev inequality (associated to the derivative $\hat{\nabla}$) with respect to ν , which is the law of X . As example, we prove these equalities for the path measure of a very singular process, namely the Dyson Brownian motion: although, for each instant its realization is a nice Wiener functional, its law as a probability on the path space is singular with respect to the Wiener measure, hence this is a degenerate situation. Afterwards we explain how to use all the theory developed in this document to derive the poincaré and log-Sobolev inequalities

for the semigroup measures of the diffusions for fixed instants. Taking a fixed instant increases the number of symmetries and variations to circumvent the difficulties as explained in Section 5.3. We then apply these ideas to give a new and very different proof of these inequalities in the case of the Heisenberg group w.r.t. the law at $t = 1$ of the non-linear Brownian motion.. As reader will realize, the method that we have undertaken is quite different then those of these works, where, a constant consuming integration by parts formula has been put forward. Meanwhile our approach is based on the invariance of the Brownian motion under the Euclidean symmetries and they are applied to both right and left Brownian paths who are starting from the origin and/ or to their conditional expectations (with respect to the filtration of the process). In particular, the right or left semigroups applied to a positive function give the same result at the trajectories of right or left processes starting from zero element of the group. This implies a similar invariance for “les opérateurs du carré du champs” of left and right processes and the commutation relations between the semigroup and the right invariant vector fields suffice to obtain the logarithmic Sobolev inequality on the Heisenberg group with the best constants possible. This problem has been suggested by **Leonard Gross**, who has been present all along this work and he has communicated the references [1], [8] and [15] through which we have been able to learn the subject and this work would have never seen the daylight without his encouragements.

2. Preliminaries

In this section we recall the representation theorem and its consequences, which have all been proven in [24]: Let $X = (X_t, t \in [0, 1])$ be a weak solution of the following stochastic differential equation:

$$(2.1) \quad dX_t = b(t, X)dt + \sigma(t, X)dB_t, \quad X_0 = x,$$

where $B = (B_t, t \in [0, 1])$ is an \mathbb{R}^d -valued Brownian motion and $\sigma : [0, 1] \times C([0, 1], \mathbb{R}^n) \rightarrow L(\mathbb{R}^d, \mathbb{R}^n)$ and $b : [0, 1] \times C([0, 1], \mathbb{R}^n) \rightarrow \mathbb{R}^n$ are measurable maps, adapted to the natural filtration of $C([0, 1], \mathbb{R}^n)$ and of linear growth. Denote by $(\mathcal{F}_t(X), t \in [0, 1])$ the filtration of X and let us denote by K the set of \mathbb{R}^n -valued, $(\mathcal{F}_t(X), t \in [0, 1])$ -adapted processes $\alpha(X)$, s.t.

$$E \int_0^1 (a(s, X)\alpha_s(X), \alpha_s(X))ds < \infty,$$

where $a(s, w) = \sigma(s, w)\sigma^*(s, w)$, $s \in [0, 1]$, $w \in C([0, 1], \mathbb{R}^n)$.

Theorem 1. *The set $\Gamma = \{N \in L^2(\mathcal{F}_1(X)) : N = E[N] + \int_0^1 (\alpha_s(X), \sigma(s, X)dB_s), \alpha \in K\}$ is dense in $L^2(\mathcal{F}_1(X))$.*

Theorem 1 permits us to show the following **extension of the martingale representation theorem to the functionals of degenerate diffusions**:

Theorem 2. *Denote by $P_s(X)$ a measurable version of the orthogonal projection from \mathbb{R}^d onto $\sigma(s, X)^*(\mathbb{R}^n) \subset \mathbb{R}^d$ and let $F \in L^2(\mathcal{F}_1(X))$ be any random variable with zero expectation. Then there exists a process $\xi(X) \in L^2_a(dt \times dP; \mathbb{R}^d)$, adapted to $(\mathcal{F}_t(X), t \in [0, 1])$, such that*

$$F(X) = \int_0^1 (P_s(X)\xi_s(X), dB_s)_{\mathbb{R}^d} = \int_0^1 (\xi_s(X), P_s(X)dB_s)_{\mathbb{R}^n}$$

a.s.

Conversely, any stochastic integral of the form

$$\int_0^1 (P_s(X)\xi_s(X), dB_s),$$

where $\xi(X)$ is an $(\mathcal{F}_t(X), t \in [0, 1])$ -adapted, measurable process with $E \int_0^1 |P_s(X)\xi_s(X)|^2 ds < \infty$, gives rise to an $\mathcal{F}_1(X)$ -measurable random variable.

Remark: Let η be an adapted process such that η_s belongs to the orthogonal complement of $\sigma^*(\mathbb{R}^n)$ in $\mathbb{R}^d ds \times dP$ -a.s. Then $\eta + \xi$ can also be used to represent $F(X)$. Hence $\xi(X)$ is not unique but $P(X)\xi(X)$ is always unique.

As a consequence of above theorems, we get

Theorem 3. Let $\dot{u} \in L^2(dt \times dP, \mathbb{R}^d)$ be adapted to the Brownian filtration, then we have

$$(2.2) \quad E \left[\int_0^1 (\dot{u}_s, dB_s) | \mathcal{F}_1(X) \right] = \int_0^1 (E[P_s(X)\dot{u}_s | \mathcal{F}_s(X)], dB_s)$$

almost surely.

Remark 1. If K is a process adapted to the Brownian filtration with values in $\mathbb{R}^m \otimes \mathbb{R}^d$ which is in $L^2(dt \times d\mu, \mathbb{R}^m \otimes \mathbb{R}^d)$, then we obtain from Theorem 3

$$(2.3) \quad E \left[\int_0^1 K_s dB_s | \mathcal{F}_1(X) \right] = \int_0^1 E[K_s | \mathcal{F}_s(X)] P(X_s) dB_s$$

μ -almost surely.

Corollary 1. Let $h \in H^1([0, 1], \mathbb{R}^d)$ (i.e., the Cameron-Martin space), denote by $\rho(\delta h)$ the Wick exponential $\exp(\int_0^1 (\dot{h}_s, dB_s) - \frac{1}{2} \int_0^1 |\dot{h}_s|^2 ds)$, then we have

$$E[\rho(\delta h) | \mathcal{F}_1(X)] = \exp \left(\int_0^1 (P_s(X)\dot{h}_s, dB_s) - \frac{1}{2} \int_0^1 |P_s(X)\dot{h}_s|^2 ds \right).$$

Theorem 4. • Assume that $\mathcal{F}_t(X) \subset \mathcal{F}_t(B)$ for any $t \in [0, 1]$, where $(\mathcal{F}_t(B), t \in [0, 1])$ represents the filtration of the Brownian motion. Define the martingale $m = (m_t, t \in [0, 1])$ as $m_t = \int_0^t P_s(X) dB_s$, then the set

$$K = \{\rho(\delta_m(h)) : h \in H\}$$

is total in $L^2(\mathcal{F}_1(X))$, where $\rho(\delta_m(h)) = \exp \left(\int_0^1 (\dot{h}_s, dm_s) - \frac{1}{2} \int_0^1 |P_s(X)\dot{h}_s|^2 ds \right)$. In particular, any element F of $L^2(\mathcal{F}_1(X))$ can be written in a unique way as the sum

$$(2.4) \quad F = E[F] + \sum_{n=1}^{\infty} \int_{C_n} (f_n(s_1, \dots, s_n), dm_{s_1} \otimes \dots \otimes dm_{s_n})$$

where C_n is the n -dimensional simplex in $[0, 1]^n$ and $f_n \in L^2(C_n, ds^{\otimes n}) \otimes (\mathbb{R}^d)^{\otimes n}$.

- More generally, without the hypothesis $\mathcal{F}_t(X) \subset \mathcal{F}_t(B)$, for any $F \in L^2(\mathcal{F}_1(X)) \cap L^2(\mathcal{F}_1(B))$, the conclusions of the first part of the theorem hold true.

Remark 2. In this theorem we need to make the hypothesis $\mathcal{F}_t(X) \subset \mathcal{F}_t(B)$ for any $t \in [0, 1]$ to assure the non-symmetric chaos representation (2.4). Without this hypothesis, although we have Theorem 2 and when we iterate it we have a similar representation, but, consisting of a finite number of terms. It is not possible to push this procedure up to infinity since we have no control at infinity.

Remark 3. *Let us note that if there is no strong solution to the equation defining the process X , the chaotic representation property may fail. For example, let U be a weak solution of*

$$(2.5) \quad dU_t = \alpha_t(U)dt + dB_t,$$

with U_0 given. Assume that (2.5) has no strong solution, as it may happen in the famous example of Tsirelson ([12]), i.e., U is not measurable w.r.t. the sigma algebra generated by B , then we have no chaotic representation property for the elements of $L^2(\mathcal{F}_1(U))$ in terms of the iterated stochastic integrals of deterministic functions on $C_n, n \in \mathbb{N}$ w.r.t. B ; the contrary would imply the equality of $\mathcal{F}_1(U)$ and of $\mathcal{F}_1(B)$, which would contradict the non-existence of strong solutions.

3. Conditional and Covariant Sobolev derivatives for degenerate diffusions' functionals

In this section, we shall work with the diffusion processes in the restricted sense, i.e., there will be no explicit dependence on the chance parameter, the coefficients will be C^1 -Lipschitz functions (or C^k , with large $k \in \mathbb{N}$) of linear growth, for notational simplicity, we shall also assume that they are time independent. To keep track of the nonlinearity due to the diffusion's paths interactions, we need to develop a special derivation w.r.t. the Cameron-Martin space of the defining Brownian motion. The next result paves the road and it is of fundamental importance:

Theorem 5. *For $h \in H$, define $\hat{\nabla}_h X_t = E[\nabla_h X_t | \mathcal{F}_1(X)]$, then it satisfies the following relation:*

$$\hat{\nabla}_h X_t = \int_0^t \partial \sigma(X_s) \hat{\nabla}_h X_s P(X_s) dB_s + \int_0^t \partial b(X_s) \hat{\nabla}_h X_s ds + \int_0^t \sigma(X_s) \dot{h}(s) ds$$

μ -almost surely. If σ and b are twice differentiable and bounded, and if the projection map is differentiable and of linear growth, for the iterated derivative $\hat{\nabla}_h^2 X_t$ we have the following expression:

$$\begin{aligned} \hat{\nabla}_h^2 X_t &= \int_0^t \partial^2 \sigma(X_s) \hat{\nabla}_h X_s \otimes \hat{\nabla}_h X_s P(X_s) dB_s + \int_0^t \partial \sigma(X_s) \hat{\nabla}_h^2 X_s P(X_s) dB_s \\ &+ \int_0^t \partial \sigma(X_s) \hat{\nabla}_h X_s \partial P(X_s) \hat{\nabla}_h X_s P(X_s) dB_s + \int_0^t \partial \sigma(X_s) \hat{\nabla}_h X_s (P(X_s) + I_{\mathbb{R}^d}) \dot{h}(s) ds \\ &+ \int_0^t \partial^2 b(X_s) (\hat{\nabla}_h X_s \otimes \hat{\nabla}_h X_s + \hat{\nabla}_h^2 X_s) ds \end{aligned}$$

μ -almost surely.

Proof: We have

$$(3.6) \quad \nabla_h X_t = \int_0^t \partial \sigma(X_s) \nabla_h X_s dB_s + \int_0^t \partial b(X_s) \nabla_h X_s ds + \int_0^t \sigma(X_s) \dot{h}(s) ds,$$

taking the conditional expectation of both sides of this relation, the term with stochastic integral follows from Theorem 3 and the Lebesgue integral part follows from the fact that

$$E[\nabla_h X_t | \mathcal{F}_1(X)] = E[\nabla_h X_t | \mathcal{F}_t(X)]$$

due to the fact that $\mathcal{B}_t = \sigma(B_s, s \leq t)$ is independent of the future increments of the Brownian motion (after t). For the second derivative we just iterate the calculation of the first derivative. \square

Remark: It is clear that $\hat{\nabla}_h^2 X_t \neq E[\nabla_h^2 X_t | \mathcal{F}_1(X)]$.

Corollary 2. *Let $\mathcal{S}(X)$ be the set of functions on W defined as*

$$\mathcal{S}(X) = \{f(X_{t_1}, \dots, X_{t_n}) : 0 \leq t_1 < \dots < t_n, f \in \mathcal{S}(\mathbb{R}^n), n \geq 1\},$$

where $\mathcal{S}(\mathbb{R}^n)$ denotes the space of rapidly decreasing smooth functions of Laurent Schwartz. Let $h \in H$, assume that $(F_k(X), k \geq 1) \subset \mathcal{S}(X)$ converges to zero in $L^2(\mathcal{F}(X))$ and that $(\hat{\nabla}_h F_k(X), k \geq 1)$ is Cauchy in $L^2(\mathcal{F}(X))$, then

$$\lim_{k \rightarrow \infty} \hat{\nabla}_h F_k(X) = 0$$

μ -a.s. In other words $\hat{\nabla}_h$ is a closable operator on $L^2(X(\mu))$, where $X(\mu)$ denotes the law of the diffusion process, or the image of the Wiener measure μ under the map X which is defined by the diffusion process.

Proof: Let $\eta = \eta(X)$ be the limit of $(\hat{\nabla}_h F_k(X), k \geq 1)$, then using Theorem 3, we have, for any cylindrical $G(X) \in \mathcal{S}(X)$,

$$\begin{aligned} E[\eta(X) G(X)] &= \lim_k E[\hat{\nabla}_h F_k(X) G(X)] = \lim_k E[\nabla_h F_k(X) G(X)] \\ &= \lim_k E[F_k(X)(-\nabla_h G(X) + G(X)\delta h)] \\ &= \lim_k E[F_k(X)(-\hat{\nabla}_h G(X) + G(X)\delta(\mathbb{P}(X)h))] = 0, \end{aligned}$$

where $\delta(\mathbb{P}(X)h) = \int_0^1 P(X_s)\dot{h}(s)dB_s$. □

For $F \in \mathcal{S}(X)$, we define $\hat{\nabla}F$ as

$$\hat{\nabla}F = \sum_{i=1}^{\infty} \hat{\nabla}_{e_i} F,$$

where $(e_i, i \geq 1)$ is an orthonormal basis in the Cameron-Martin space H . The above theorem implies in particular that $\hat{\nabla}$ is a closable operator from $L^p(\mathcal{F}(X))$ into $L^p(\mathcal{F}(X), H)$. We shall denote its closure with the same notation. In particular, we denote by $\mathbb{M}_{p,1}$ the completion of $\mathcal{S}(X)$ w.r.t. the norm

$$\|F(X)\|_{p,1} = \| |\hat{\nabla}F(X)|_H \|_{L^p(\mu)} + \|F(X)\|_{L^p(\mu)}.$$

For $p = 2$ we use the following version:

$$\|F(X)\|_{2,1}^2 = E \left[|\hat{\nabla}F(X)|_H^2 + |F(X)|^2 \right].$$

If K is a separable Hilbert space, we denote by $\mathcal{S}(X, K)$, the set of K -valued cylindrical functions of the trajectories of X by replacing $\mathcal{S}(\mathbb{R}^n)$ above by $\mathcal{S}(\mathbb{R}^n) \otimes K$, where the latter denotes the (projective) tensor product.

Definition 1. *Let ν be the measure given as $X(\mu)$, i.e., the law of the diffusion process as a probability measure on the path space. We denote the adjoint of $\hat{\nabla}$ w.r.t. ν as δ_ν (or sometimes as $\hat{\nabla}^*$). Namely, for $\xi(X) \in \mathcal{S}(X) \otimes H$, i.e., the H -valued cylindrical, $\mathcal{F}(X)$ -measurable functions and $G(X) \in \mathcal{S}(X)$, we write*

$$\begin{aligned} E[\delta_\nu \xi(X) G(X)] &= E_\nu[\delta_\nu \xi G] \\ &= E_\nu[(\xi, \hat{\nabla}G)_H] \\ &= E[(\xi(X), \hat{\nabla}G(X))_H]. \end{aligned}$$

Theorem 6. Let $\xi(X) \in L^2(\mathcal{F}(X), H)$ be of the form $\sum_{i < \infty} \xi_i(X) e_i$, where $\xi_i(X) \in \mathcal{S}(X)$ and $e_i \in H$ for each $i \in \mathbb{N}$.

(1) Then

$$\delta_\nu \xi(X) = \sum_i \left[-\hat{\nabla}_{e_i} \xi_i(X) + \xi_i(X) \delta(\mathbb{P}(X) e_i) \right],$$

where we define the action of $P(X)$ on the Cameron-Martin space H as

$$(\mathbb{P}(X)h)(t) = \int_0^t P(X_s) \dot{h}(s) ds,$$

for $h \in H$. In particular, we have

$$(3.7) \quad E[\delta \xi(X) | \mathcal{F}_1(X)] = \delta_\nu \xi(X).$$

(2) If $\eta = \eta(X) \in \mathcal{S}(X, H)$ is adapted to the filtration $(\mathcal{F}_t(X), t \in [0, 1])$, where $\mathcal{S}(X, H)$ denotes H -valued version of $\mathcal{S}(X)$, then

$$(3.8) \quad \delta_\nu \eta = \delta \mathbb{P}(X) \eta$$

μ -a.s.

Proof: Let $g(X) \in \mathcal{S}(X)$, then, using Definition 1 and the integration by parts formula for μ , we get

$$\begin{aligned} E[(\xi(X), \hat{\nabla} g(X))_H] &= E \left[\sum_i \xi_i(X) \hat{\nabla}_{e_i} g(X) \right] \\ &= E \left[\sum_i \xi_i(X) \nabla_{e_i} g(X) \right] \\ &= \sum_i E [-\nabla_{e_i} \xi_i(X) g(X) + g(X) \xi_i(X) \delta e_i] \\ &= \sum_i E [-\nabla_{e_i} \xi_i(X) g(X) + g(X) \xi_i(X) E[\delta e_i | \mathcal{F}_1(X)]] \\ &= \sum_i E \left[-\hat{\nabla}_{e_i} \xi_i(X) g(X) + g(X) \xi_i(X) \delta(\mathbb{P}(X) e_i) \right] \\ &= E \left[\left(\sum_i \left(-\hat{\nabla}_{e_i} \xi_i(X) + \xi_i(X) \delta(\mathbb{P}(X) e_i) \right) \right) g(X) \right], \end{aligned}$$

since $E[\delta e_i | \mathcal{F}_1(X)] = \delta(\mathbb{P}(X) e_i)$ as shown in Theorem 3. The relation (3.7) holds then true for the cylindrical case, the general case follows by passing to the limit in $\mathbb{M}_{2,1}(H)$ (i.e., the space $\mathbb{M}_{2,1} \otimes H$, completed Hilbert-Schmidt tensor product).

To prove the second part, it follows already from the relation (3.7) that $\delta \mathbb{P}(X) \eta = E[\delta \eta | \mathcal{F}_1(X)]$, let now $G = G(X) \in \mathcal{S}(X)$, then

$$\begin{aligned} E[\delta_\nu \eta(X) G(X)] &= E \left[(\eta(X), \hat{\nabla} G(X))_H \right] = E[(\eta(X), \nabla G(X))_H] \\ &= E[\delta \eta(X) G(X)] = E[E[\delta \eta(X) | \mathcal{F}_1(X)] G(X)] \\ &= E[\delta \mathbb{P}(X) \eta G(X)], \end{aligned}$$

as, by Theorem 3, $\delta \mathbb{P}(X) \eta$ is $\mathcal{F}_1(X)$ -measurable, the proof follows. \square

Corollary 3. *Let $F(X), G(X) \in \mathcal{S}(X)$ and let $h \in H$, then we have the integration by parts formula*

$$(3.9) \quad E[\hat{\nabla}_h F(X) G(X)] = E \left[F(X) (-\hat{\nabla}_h G(X) + G(X) \delta \mathbb{P}(X) h) \right].$$

Proof: As $E[\hat{\nabla}_h F(X) G(X)] = E[\nabla_h F(X) G(X)]$. from the Gaussian integration by parts formula, we have

$$\begin{aligned} E[\hat{\nabla}_h F(X) G(X)] &= E[F(X)(-\nabla_h G(X) + G(X) \delta h)] \\ &= E[F(X) E [(-\nabla_h G(X) + G(X) \delta h) | \mathcal{F}_1(X)]] \\ &= E[F(X)(-\hat{\nabla}_h G(X) + G(X) \delta \mathbb{P}(X) h)]. \end{aligned}$$

□

Corollary 4. *For $a(X) \in \mathbb{M}_{2,1}$ and $\xi(X) \in \mathcal{S}(X, H)$, we have*

$$(3.10) \quad \delta_\nu(a(X)\xi(X)) = a(X)\delta_\nu\xi(X) - (\hat{\nabla}a(X), \xi(X))_H.$$

Moreover, if $\dot{\xi}(X)$ is adapted to the filtration $(\mathcal{F}_t(X), t \in [0, 1])$, then

$$(3.11) \quad \delta_\nu\xi(X) = \delta(\mathbb{P}(X)\xi(X)).$$

In particular, for any $h, k \in H$, we have

$$(3.12) \quad E[\delta(\mathbb{P}(X)\nabla_h \mathbb{P}(X)k) | \mathcal{F}_1(X)] = \delta_\nu(\mathbb{P}(X)\hat{\nabla}_h \mathbb{P}(X)k)$$

μ -a.s.

Proof: The only claim which is not immediate is (3.12) and it follows from Theorem 3. □

Lemma 1. *For any $h, k \in H$, we have*

$$\hat{\nabla}_k(\delta \mathbb{P}(X)h) = (\mathbb{P}(X)h, k)_H + \delta_\nu(\hat{\nabla}_k \mathbb{P}(X)h)$$

μ -a.s.

Proof: By definition

$$\begin{aligned} \hat{\nabla}_k(\delta \mathbb{P}(X)h) &= E[\nabla_k \delta \mathbb{P}(X)h | \mathcal{F}_1(X)] \\ &= (\mathbb{P}(X)h, k)_H + E[\delta \nabla_k \mathbb{P}(X)h | \mathcal{F}_1(X)] \\ &= (\mathbb{P}(X)h, k)_H + \delta \mathbb{P}(X)\hat{\nabla}_k \mathbb{P}(X)h \\ &= (\mathbb{P}(X)h, k)_H + \delta_\nu \hat{\nabla}_k \mathbb{P}(X)h \end{aligned}$$

by Theorem 3 and Corollary 4. □

Theorem 7. *Let $\xi = \xi(X)$, $\eta = \eta(X)$ be in $\mathcal{S}(X, H)$, then the following identity holds true:*

$$(3.13) \quad \begin{aligned} E[\delta_\nu \xi \delta_\nu \eta] &= E[(\mathbb{P}(X)\xi, \eta)_H + \text{trace}(\hat{\nabla} \xi \cdot \hat{\nabla} \eta)] \\ &\quad + E[\text{trace}(\hat{\nabla} \mathbb{P}(X)\xi \cdot \hat{\nabla} \eta) + \text{trace}(\hat{\nabla} \mathbb{P}(X)\eta \cdot \hat{\nabla} \xi)] \end{aligned}$$

Proof: Writing $E[\delta_\nu \xi \delta_\nu \eta] = E[(\xi, \hat{\nabla} \delta_\nu \eta)_H]$, then we shall calculate $\hat{\nabla} \delta_\nu \eta$. Let $k \in H$ and let $(e_i, i \geq 1)$ be an orthonormal basis of H , then

$$\begin{aligned} \hat{\nabla} \delta_\nu \eta &= \hat{\nabla}_k \left[\sum_i \eta_i \delta \mathbb{P}(X) e_i - \hat{\nabla}_{e_i} \eta_i \right] \\ &= \sum_i \left[\hat{\nabla}_k \eta_i \delta \mathbb{P}(X) e_i + \eta_i \hat{\nabla}_k \delta \mathbb{P}(X) e_i - \hat{\nabla}_k \hat{\nabla}_{e_i} \eta_i \right] \\ &= \sum_i \left[\hat{\nabla}_k \eta_i \delta \mathbb{P}(X) e_i + \eta_i (\mathbb{P}(X) e_i, k)_H \right] \\ &\quad + \sum_i \left[\eta_i \delta_\nu \hat{\nabla}_k \mathbb{P}(X) e_i - \hat{\nabla}_k \hat{\nabla}_{e_i} \eta_i \right] \end{aligned}$$

Taking the expectations after replacing the vector k with e_k and using the above equality, we get

$$\begin{aligned} E[\delta_\nu \xi \delta_\nu \eta] &= E[(\xi, \hat{\nabla} \delta_\nu \eta)_H] = E \left[\sum_i \eta_i \delta_\nu (\delta \mathbb{P}(X) e_i) + (\mathbb{P}(X) \xi, \eta)_H \right] \\ &\quad + E \left[\sum_{k,i} \eta_i (\delta_\nu \hat{\nabla}_{e_k} \mathbb{P}(X) e_i) \xi_k - \xi_k \hat{\nabla}_{e_k} \hat{\nabla}_{e_i} \eta_i \right], \end{aligned}$$

where $\eta_i = (\eta, e_i)_H$, $\xi_i = (\xi, e_i)_H$. Using the relation

$$\hat{\nabla}_\xi \hat{\nabla}_{e_i} \eta_i = \hat{\nabla}_{e_i} \hat{\nabla}_\xi \eta_i - \hat{\nabla}_{\hat{\nabla}_{e_i} \xi} \eta_i,$$

we have

$$\begin{aligned} E[\delta_\nu \xi \delta_\nu \eta] &= E \left[\sum_i \hat{\nabla}_\xi \eta_i \delta \mathbb{P}(X) e_i + (\mathbb{P}(X) \xi, \eta)_H \right] \\ &\quad + E \left[\sum_{k,i} \eta_i \delta_\nu (\hat{\nabla}_{e_k} \mathbb{P}(X) e_i) \xi_k - \hat{\nabla}_{e_i} \hat{\nabla}_\xi \eta_i + \text{trace} (\hat{\nabla} \eta \cdot \hat{\nabla} \xi) \right] \\ &= E \left[\delta_\nu \hat{\nabla}_\xi \eta + (\mathbb{P}(X) \xi, \eta)_H + \text{trace} (\hat{\nabla} \eta \cdot \hat{\nabla} \xi) + \sum_{k,i} \eta_i \delta_\nu (\hat{\nabla}_{e_k} \mathbb{P}(X) e_i) \xi_k \right]. \end{aligned}$$

Using the identity

$$\delta_\nu (g \tau) = g \delta_\nu \tau - (\tau, \hat{\nabla} g)_H,$$

for $g \in \mathcal{S}(X)$, $\tau \in \mathcal{S}(X, H)$, we have

$$\begin{aligned} (3.14) \quad &\sum_{k,i} E \left[\eta_i \delta_\nu (\hat{\nabla}_{e_k} \mathbb{P}(X) e_i) \xi_k \right] \\ &= \sum_{k,i} E \left[\eta_i \delta_\nu (\xi_k \hat{\nabla}_{e_k} \mathbb{P}(X) e_i) + \eta_i (\hat{\nabla}_\xi \xi_k, \hat{\nabla}_{e_k} \mathbb{P}(X) e_i)_H \right] \\ &= E \left[\sum_i \eta_i \delta_\nu (\hat{\nabla}_\xi \mathbb{P}(X) e_i) + \sum_k (\hat{\nabla}_\xi \xi_k, \hat{\nabla}_{e_k} \mathbb{P}(X) \eta - \mathbb{P}(X) \hat{\nabla}_{e_k} \eta)_H \right] \end{aligned}$$

and since

$$(3.15) \quad \eta_i \delta_\nu(\hat{\nabla}_\xi \mathbb{P}(X) e_i) = \delta_\nu(\eta_i \hat{\nabla}_\xi \mathbb{P}(X) e_i) + (\hat{\nabla}_\xi \mathbb{P}(X) e_i, \hat{\nabla} \eta_i)_H,$$

we obtain by substituting (3.15) in (3.14)

$$\sum_{k,i} E \left[\eta_i \delta_\nu(\hat{\nabla}_{e_k} \mathbb{P}(X) e_i) \xi_k \right] = I + II,$$

where

$$\begin{aligned} I &= \sum_i E \left[\delta_\nu(\eta_i \hat{\nabla}_\xi \mathbb{P}(X) e_i) + (\hat{\nabla}_\xi \mathbb{P}(X) e_i, \hat{\nabla} \eta_i)_H \right] \\ &= E[\text{trace}(\hat{\nabla}_\xi \mathbb{P}(X) \cdot \hat{\nabla} \eta)] \end{aligned}$$

and

$$\begin{aligned} II &= \sum_{k,i} E[\eta_i (\hat{\nabla} \xi_k, \hat{\nabla}_{e_k} \mathbb{P}(X) e_i)_H] \\ &= \sum_k E[(\hat{\nabla} \xi_k, (\hat{\nabla}_{e_k} \mathbb{P}(X)) \eta)_H] \\ &= E[\text{trace}(\hat{\nabla} \xi \cdot (\hat{\nabla} \mathbb{P}(X)) \eta)] \end{aligned}$$

and this completes the proof. \square

The next result shows the local character of the derivative operator $\hat{\nabla}$, which is well-known in the flat case:

Proposition 1. *Assume that $F = F(X)$ is in $\mathbb{M}_{2,1}$, then $\hat{\nabla} F = 0$ on the set $\{F = 0\}$ ν -almost surely.*

Proof: Let θ be a positive, smooth function of compact support on \mathbb{R} with $\theta(0) = 1$, denote by ζ its primitive and let $\zeta_\varepsilon(t) = \varepsilon \zeta(t/\varepsilon)$. Clearly $\zeta_\varepsilon \circ F$ belongs to $\mathbb{M}_{2,1}(X)$. If $u \in \mathcal{S}(X, H)$, from the definitions

$$\begin{aligned} E[\zeta_\varepsilon(F) \delta_\nu u] &= E[(\hat{\nabla} \zeta_\varepsilon(F), u)_H] \\ &= E[\theta(F/\varepsilon) (\hat{\nabla} F, u)_H] \rightarrow E[1_{\{F=0\}}]. \end{aligned}$$

On the other hand, from the Dominated Convergence Theorem, we have $\lim_{\varepsilon \rightarrow 0} E[\zeta_\varepsilon(F) \delta_\nu u] = 0$ and as $\mathcal{S}(X, H)$ is dense in $\mathbb{M}_{2,1}(X, H)$, the proof follows. \square

The following result shows that $\hat{\nabla}$ defines a Markovian Dirichlet form on the path space of the diffusion process:

Proposition 2. *Assume that $f, g \in \mathbb{M}_{2,1}$, then $f \wedge g \in \mathbb{M}_{2,1}$, where \wedge denotes the minimum.*

Proof: Let $(f_n, n \geq 1)$ and $(g_n, n \geq 1)$ be two sequences of cylindrical functions approximating f and g in $\mathbb{M}_{2,1}$ respectively. We have

$$\begin{aligned} \hat{\nabla}(f_n \wedge g_n) &= E[\nabla f_n 1_{\{f_n \geq g_n\}} + \nabla g_n 1_{\{f_n < g_n\}} | \mathcal{F}_1(X)] \\ &= \hat{\nabla} f_n 1_{\{f_n \geq g_n\}} + \hat{\nabla} g_n 1_{\{f_n < g_n\}}, \end{aligned}$$

Hence

$$|hna(f_n \wedge g_n)|_H \leq |\hat{\nabla} f_n|_H + |\hat{\nabla} g_n|_H.$$

Consequently, $(f_n \wedge g_n, n \geq 1)$ is bounded in $\mathbb{M}_{2,1}$, then it has a subsequence which converges weakly. This implies that $f \wedge g \in \mathbb{M}_{2,1}$ and that $|\hat{\nabla}(f \wedge g)|_H \leq |\hat{\nabla} f|_H + |\hat{\nabla} g|_H$ almost surely. \square

While making calculations we need also a second covariant derivation on the paths of the diffusion process, this time this will be the smoothed derivative w.r.t. the state space elements, namely we define

$$\hat{\partial}_\xi X_t(\xi) = E[\partial_\xi X_t(\xi) | \mathcal{F}_1(X(\xi))]$$

where ∂_ξ denotes the derivative of the diffusion process at the instant $t \in [0, 1]$ with respect to its initial starting point $\xi \in \mathbb{R}^n$, similarly, for smooth functions on \mathbb{R}^n , we define

$$\hat{\partial}_\xi f(X_t(\xi)) = E[\partial f(X_t(\xi)) \partial_\xi X_t(\xi) | \mathcal{F}_1(X(\xi))] = \partial f(X_t(\xi)) \hat{\partial}_\xi X_t(\xi),$$

in such a way that $\hat{\partial}$ behaves as a derivative w.r.t. state space variable of the functionals of the paths of the diffusion process. We have the immediate result which follows from Theorem 2.2

Theorem 8. *The state space covariant derivative $\hat{\partial}_x X_t(x) = J_t(x)$ satisfies the following stochastic differential equation:*

$$(3.16) \quad J_t = I_{\mathbb{R}^n} + \int_0^t \partial \sigma(X_s(x)) J_s(x) P(X_s(x)) dB_s + \int_0^t \partial b(X_s(x)) J_s(x) ds,$$

P-a.s., for any $t \in [0, 1]$.

It follows from Theorem 8 and from the Itô formula that

Proposition 3. *$(J_t, t \in [0, 1])$ is a semimartingale with values in $GL(n, \mathbb{R})$ (non-singular matrices) and its inverse, denoted as $(K_t, t \in [0, 1])$ has the following representation:*

$$(3.17) \quad dK_t = - \sum_{i=1}^d K_t \partial \sigma^i(X_t(x)) dm_t^i + K_t \left[\sum_{i,j=1}^d \partial \sigma^i(X_t(x)) \partial \sigma^j(X_t(x)) P_{i,j}(X_t(x)) - \partial b(X_t(x)) \right] dt,$$

where $dm_t = P(X_t(x)) dB_t$, $m_t = (m_t^1, \dots, m_t^d)$, and σ^i is the i -th column vector of the $n \times d$ -matrix σ .

Proof: The equation (3.17) is a linear stochastic differential equation with bounded coefficients, hence it has a unique solution without explosion. To prove the claim it suffices to show that $J_t K_t = K_t J_t = I_{\mathbb{R}^n}$ almost surely, where $I_{\mathbb{R}^n}$ denotes the identity map of \mathbb{R}^n , and this last equation follows from the Itô formula. \square

An immediate consequence of these calculations, combined with the variation of constants' method, is the following

Corollary 5. *For any $t < \tau \in [0, 1]$, we have*

$$\hat{D}_t X_\tau(x) = J_\tau K_t \sigma(X_t(x))$$

almost surely.

4. Poincaré and Logarithmic Sobolev Inequalities

Here are some applications of the results of the preceding section. Let us begin with a non-trivial extension of Itô-Clark-Hausmann-Bismut-Ocone formula:

Theorem 9. *Assume that $F(X) \in \mathbb{M}_{2,1}$, then it can be represented as*

$$F(X) = E[F(X)] + \int_0^1 P(X_s) E[\hat{D}_s F(X) | \mathcal{F}_s(X)] dB_s,$$

where $\hat{D}_s F(X)$ is defined as $\hat{\nabla} F(X)(\cdot) = \int_0^1 \hat{D}_s F(X) ds$.

Proof: We know from Theorem 2 that $F(X)$ can be represented as

$$F(X) = E[F(X)] + \int_0^1 P(X_s) \alpha_s(X) . dB_s,$$

for some $\alpha(X) \in L_a^2(dt \times d\mu, \mathbb{R}^d)$, adapted to the filtration $(\mathcal{F}_t(X), t \in [0, 1])$. Moreover, if $F(X) \in \mathcal{S}(X)$, then from the Gaussian case, we have

$$(4.18) \quad F(X) = E[F(X)] + \int_0^1 E[D_s F(X) | \mathcal{B}_s] dB_s,$$

where $(\mathcal{B}_s, s \in [0, 1])$ is the filtration of the Brownian motion. It follows from Theorem 3, taking the conditional expectations of both sides of the equality (4.18), that

$$F(X) = E[F(X)] + \int_0^1 P(X_s) E[D_s F(X) | \mathcal{F}_s(X)] dB_s.$$

Hence in this case $P(X_s) \alpha_s(X) = P(X_s) E[D_s F(X) | \mathcal{F}_s(X)] = P(X_s) E[E[D_s F(X) | \mathcal{F}_1(X)] | \mathcal{F}_s(X)] = P(X_s) E[\hat{D}_s F(X) | \mathcal{F}_s(X)]$ $ds \times d\mu$ -a.s. Choose now a sequence $(F_n(X), n \geq 1) \subset \mathcal{S}(X)$ which approximates $F(X)$ in $\mathbb{M}_{2,1}$. Then $(\hat{\nabla} F_n(X), n \geq 1)$ converges to $\hat{\nabla} F(X)$ in $L^2(\mu, H)$ (in other words $(\hat{\nabla} F_n)$ converges to $\hat{\nabla} F$ in $L^2(\nu, H)$), consequently

$$\lim_n E \int_0^1 |P(X_s) E[\hat{D}_s F_n(X) - \hat{D}_s F(X) | \mathcal{F}_s(X)]|^2 ds = 0,$$

hence the result follows for any $F(X) \in \mathbb{M}_{2,1}$. □

Corollary 6. *Assume that $F = F(X) \in \mathbb{M}_{2,1}$ such that $\hat{\nabla} F = 0$ ν -a.s. (or $\hat{\nabla} F(X) = 0$ μ -a.s), then F is ν -a.s. (or $F(X)$ is μ -a.s.) a constant.*

Proof: The proof follows directly from Theorem 9. □

We obtain the Poincaré inequality from Theorem 9, whose proof is trivial since the conditional expectation in the representation formula (4.18) operates as a contraction:

Theorem 10. *For any $F(X) \in \mathbb{M}_{2,1}$, we have*

$$E[|F(X) - E[F(X)]|^2] \leq E[|\mathbb{P}(X) \hat{\nabla} F(X)|_H^2].$$

Theorem 11. *For any $F \in \mathbb{M}_{2,1}$, it holds true that*

$$(4.19) \quad E[F^2(X) \log F^2(X)] \leq E[F^2(X)] \log E[F^2(X)] + 2E[|\mathbb{P}(X) \hat{\nabla} F(X)|_H^2].$$

Proof: It suffices to prove for any $F(X) \in \mathbb{M}_{2,1}$ which is strictly positive and lower bounded by a positive constant. Taking $f(X) = F^2(X)/E[F^2(X)]$, it suffices to show that

$$E[f(X) \log f(X)] \leq \frac{1}{2} E \left[\frac{1}{f(X)} |\mathbb{P}(X) \hat{\nabla} f(X)|_H^2 \right].$$

Using Theorem 9, we can represent $f(X)$ as

$$\begin{aligned} f(X) &= 1 + \int_0^1 P(X_s) E[\hat{D}_s(f(X)) | \mathcal{F}_s(X)] \cdot dB_s \\ &= 1 + \int_0^1 \frac{P(X_s) E[\hat{D}_s(f(X)) | \mathcal{F}_s(X)]}{E[f(X) | \mathcal{F}_s(X)]} E[f(X) | \mathcal{F}_s(X)] \cdot dB_s. \end{aligned}$$

Hence $f(X)$ can be represented as an exponential martingale's final value:

$$f(X) = \exp \left[\int_0^1 \frac{P(X_s) E[\hat{D}_s(f(X)) | \mathcal{F}_s(X)]}{f_s(X)} \cdot dB_s - \frac{1}{2} \int_0^1 \left| \frac{P(X_s) E[\hat{D}_s(f(X)) | \mathcal{F}_s(X)]}{f_s(X)} \right|^2 ds \right],$$

where $f_s(X) = E[f(X) | \mathcal{F}_s(X)]$. Let γ be the probability measure on the path space defined by $d\gamma = f(X) d\mu$, then the Girsanov Theorem and Bayes' formula entail that the relative entropy of γ w.r.t. μ , denoted by $H(\gamma|\mu)$ is equal to

$$\begin{aligned} H(\gamma|\mu) &= \int \log f(X) d\gamma \\ &= \frac{1}{2} E_\gamma \int_0^1 \left| \frac{P(X_s) E[\hat{D}_s(f(X)) | \mathcal{F}_s(X)]}{f_s(X)} \right|^2 ds \\ &= \frac{1}{2} E_\gamma \int_0^1 \left| \frac{E[P(X_s) \hat{D}_s(f(X)) | \mathcal{F}_s(X)]}{f_s(X)} \right|^2 ds \\ &= \frac{1}{2} E_\gamma \int_0^1 \left| E_\gamma [P(X_s) \hat{D}_s(\log f(X)) | \mathcal{F}_s(X)] \right|^2 ds \\ &\leq \frac{1}{2} E_\gamma \int_0^1 \left| P(X_s) \hat{D}_s(\log f(X)) \right|^2 ds \\ &= \frac{1}{2} E \left[\frac{1}{f(X)} |\mathbb{P}(X) \hat{\nabla} f(X)|_H^2 \right]. \end{aligned}$$

□

Remark: Maybe it is pedagogically interesting to note this inequality in terms of the measure induced by the the diffusion process on the path space, namely ν :

$$(4.20) \quad \int f^2 \log f^2 d\nu - \left(\int f^2 d\nu \right) \log \int f^2 d\nu \leq 2 \int |\mathbb{P} \hat{\nabla} f|_H^2 d\nu$$

for any $f \in \mathbb{M}_{2,1}$.

5. Examples and Further Applications

We shall give applications of the analysis that we have developped to some specific cases beginning from the simplest to more complicated situations

5.1. Classical Situation. Let $(X_t, t \geq 0)$ be the solution of the equation

$$X_t(x) = x + \int_0^t b(X_r(x))dr + \Sigma B_t,$$

where $B = (B_t, t \geq 0)$ is an d -dimensional Brownian motion, $b : \mathbb{R}^m \rightarrow \mathbb{R}^m$ is a Lipschitz-continuous vector field and $\Sigma \in L(\mathbb{R}^m, \mathbb{R}^d)$. Regularizing b with a mollifier, we obtain a smooth and Lipschitz vector field b_ε , let X^ε be the solution of SDE

$$X_t^\varepsilon(x) = x + \int_0^t b_\varepsilon(X_r^\varepsilon(x))dr + \Sigma B_t.$$

For any $h \in H$, where $H = H^1(\mathbb{R}_+, \mathbb{R}^d)$ is the Cameron-Martin space of B , we have

$$\nabla_h X_t^\varepsilon = \int_0^t \partial b_\varepsilon(X_r) \nabla_h X_r^\varepsilon dr + \Sigma h(t)$$

and the Gronwall lemma implies that, for any $T > 0$,

$$\sup_{t \leq T} \|\nabla X_t^\varepsilon\|_{H \otimes \mathbb{R}^m} \leq \|\Sigma\| e^{TK},$$

where K is the Lipschitz constant of b and $\|\Sigma\|$ is the operator norm of Σ . We can pass to the limit as $\varepsilon \rightarrow 0$ to obtain also

$$\sup_{t \leq T} \|\nabla X_t\|_{H \otimes \mathbb{R}^m} \leq \|\Sigma\| e^{TK}.$$

The following theorem is now a straight application of Theorem 11

Theorem 12. *Let $T > 0$ be arbitrary and denote by ν^T the law of the process $(X_t, t \in [0, T])$. Then, for any smooth, cylindrical function on $W_T = C([0, T], \mathbb{R}^d)$, we have*

$$\int_{W_T} F^2(x) \log \frac{F^2(x)}{\nu^T(F^2)} \nu^T(dx) \leq 2\|\Sigma\| e^{TK} \int_{W_T} |\nabla F(x)|^2 \nu^T(dx),$$

in particular, for any $t \in (0, T]$, denoting by ν_t the law of the random variable X_t , we have

$$\int_{\mathbb{R}^m} f^2(y) \log \frac{f^2(y)}{\nu_t(f^2)} \nu_t(dy) \leq 2\|\Sigma\| e^{TK} \int_{\mathbb{R}^m} |\partial f(y)|^2 \nu_t(dy),$$

for any smooth function $f : \mathbb{R}^m \rightarrow \mathbb{R}$.

6. Log-Sobolev for Dyson's Brownian Motion

In the example above, the drift coefficient has nice properties and the log-Sobolev inequality is almost trivial. Let us look at a more singular case, namely, let $(X_t = (X_t^1, \dots, X_t^d), t \geq 0)$ be the Dyson Brownian motion, i.e., the strong solution of the following stochastic differential equation:

$$(6.21) \quad dX_t^i = dB_t^i + \gamma \sum_{1 \leq j \neq i \leq d} \frac{1}{X_t^i - X_t^j} dt, \quad i = 1, \dots, d; \quad \gamma > 0,$$

with $X_0 = x_0$ such that $x_1 < \dots < x_d$. We refer the reader to references [3, 4, 5] for the existence of the strong solutions, where they use the theory of dissipative mappings in their construction and to [9] for the origins of the equation. For notational convenience, we shall write above equation as

$$(6.22) \quad dX_t = \gamma m(X_t) dt + dB_t, \quad X_0 = x_0 \in D(m),$$

where m is the mapping from a subset of \mathbb{R}^d , noted as $D(m)$ into \mathbb{R}^d whose components are given in the equation (6.21). In fact m is a set-valued, dissipative map. It is maximal and this implies that

its values reduce to one-point subsets of \mathbb{R}^d (cf. [26]). As a typical example for such a function, one can take the gradient of a concave function on \mathbb{R}^d . Equation 6.21 corresponds to the case $m = \partial f$, where $f(x) = \sum_{i < j} \log(x_j - x_i)$ if $x_1 < \dots < x_d$ and $f(x) = -\infty$ otherwise. Let us announce first an important technical result:

Lemma 2. *Let $X = (X_t, t \geq 0)$ be the unique strong solution of SDE 6.22, then, for any $T > 0$, $h, k \in H([0, T], \mathbb{R}^d)$ we have*

$$\sup_{s \leq t} |X_s(w+h) - X_s(w+k)| \leq 2\sqrt{t}\gamma|h-k|_{H([0,t], \mathbb{R}^d)}$$

P -almost surely. Consequently, for any $t > 0$, $X_t \in \cap_p \mathbb{D}_{p,1}(\mathbb{R}^d)$, in fact $|\nabla X_t|_{\mathbb{R}^d \otimes H([0,t], \mathbb{R}^d)} \in L^\infty(P)$ with the bound $2t\gamma$, where $H([0,t], \mathbb{R}^d)$ (respectively $H([0,T], \mathbb{R}^d)$) denotes the Cameron-Martin space for \mathbb{R}^d -valued functions on $[0,t]$ (respectively $[0,T]$).

Proof: We will suppose that $\gamma = 1$ and that (B_t) is the linear Brownian motion, which amounts to working on the canonical Wiener space (for practical reasons). From the equation 6.22, it comes

$$X_t(w+h) = B_t(w) + h(t) + \int_0^t m(X_s(w+h))ds,$$

hence

$$X_t(w+h) - X_t(w) = h(t) + \int_0^t [m(X_s(w+h)) - m(X_s(w))]ds.$$

Let $y_t^h(w) = X_t(w+h) - X_t(w)$, then $(y_t^h, t \in [0, T])$ satisfies

$$y_t^h(w) = h(t) + \int_0^t [m(y_s^h(w) + X_s(w)) - m(X_s(w))]ds,$$

hence

$$(6.23) \quad y_t^h(w) - y_t^k(w) = h(t) - k(t) + \int_0^t [m(y_s^h(w) + X_s(w)) - m(y_s^k(w) + X_s(w))]ds.$$

It follows from equation 6.23 and from the dissipativity of m that

$$\begin{aligned} |y_t^h - y_t^k|^2 &= \int_0^t \frac{d}{ds} |y_s^h - y_s^k|^2 ds \\ &= 2 \int_0^t (y_s^h - y_s^k, \dot{h}(s) - \dot{k}(s))_{\mathbb{R}^d} + 2 \int_0^t (y_s^h - y_s^k, m(y_s^h + X_s) - m(y_s^k + X_s)) ds \\ &\leq 2 \sup_{s \leq t} |y_s^h - y_s^k| \int_0^t |\dot{h}(s) - \dot{k}(s)| ds \\ &\leq 2\sqrt{t} \sup_{s \leq t} |y_s^h - y_s^k| \left(\int_0^t |\dot{h}(s) - \dot{k}(s)|^2 ds \right)^{1/2} \end{aligned}$$

Since the last line of above inequality is increasing w.r.t. $t \in \mathbb{R}_+$, the claim follows. The fact that $X_t \in \cap_{p>1} \mathbb{D}_{p,1}$ follows from [20], Lemma 2 at p.72. \square

An immediate consequence of Lemma 2 is Fernique's bound:

Corollary 7. *For any $t > 0$, there exists some $\varepsilon = \varepsilon(\gamma, t)$, such that*

$$E \left[\exp \left(\varepsilon \sup_{s \leq t} |X_s|^2 \right) \right] < \infty$$

Remark: Let $T > 0$ be any fixed number and let us denote by ν the law of the process restricted to the time horizon $[0, T]$, starting from x_0 . Evidently ν is the unique solution of the martingale problem (cf. [19]) associated to the operator K defined as

$$Kf(x) = \sum_{i=1}^d \left[\frac{1}{2} \frac{\partial^2 f}{\partial x_i^2}(x) + \gamma m_i(x) \frac{\partial f}{\partial x_i}(x) \right].$$

It is immediate to see that under ν , the coordinate map $x \rightarrow x(t)$ of $C_{x_0}([0, T], \mathbb{R}^d)$ can be represented as

$$dx_t = \gamma m(x_t) dt + d\beta_t,$$

where $(\beta_t, t \in [0, T])$ is a ν -Brownian motion.

The following is now clear:

Proposition 4. *The covariant derivative defined on the paths of Dyson process restricted to $[0, T]$, i.e., for $F(X) = f(X_{t_1}, \dots, X_{t_n})$,*

$$\hat{\nabla}_h F(X) = E[\nabla_h F(X) | \mathcal{F}_T(X)],$$

where $h \in H([0, T], \mathbb{R}^d)$ and $\hat{\nabla} F(X) = \sum_i \hat{\nabla}_{h_i} h_i$, such that (h_i) is a basis in $H([0, T], \mathbb{R}^d)$, are closable operators on $L^p(\mathcal{F}_T(X))$ for any $p > 1$. Consequently, we have the following representation theorem:

$$F(X) = E[F(X)] + \int_0^T E[D_s F(X) | \mathcal{F}_s(X)] \cdot dB_s.$$

From Proposition 4 we obtain as explained in the general case, the logarithmic Sobolev inequality:

Proposition 5. *For any $\mathcal{F}_T(X)$ measurable, cylindrical function $F(X)$, we have*

$$E[F^2(X) \log(F^2(X)/E[F^2(X)])] \leq 2\gamma\sqrt{T} E[|\hat{\nabla} F(X)|_{H([0, T], \mathbb{R}^d)}^2].$$

7. Intertwining Relations and Logarithmic Sobolev Inequality

Let us come back to the general situation studied in the paper: if f be a smooth function on \mathbb{R}^n and if $(Q_t, t \in [0, 1])$ is the semi-group associated to the Markov process induced by $(X_t, t \in [0, 1])$, then $(Q_{1-t}f(X_t(x)), t \in [0, 1])$ is an $(\mathcal{F}_t(X))$ -martingale and, due to the Itô formula, it can be represented as

$$(7.24) \quad Q_{1-t}f(X_t(x)) = Q_1f(x) + \int_0^t \partial_x Q_{1-s}f(X_s(x)) \cdot \sigma(X_s(x)) dB_s.$$

From Theorem 9, we obtain a second representation:

$$(7.25) \quad Q_{1-t}f(X_t(x)) = Q_1f(x) + \int_0^t P(X_s(x)) E[\hat{D}_s f(X_1(x)) | \mathcal{F}_s(X)] \cdot dB_s,$$

where $\hat{D}_s f(X_1(x)) = E[D_s f(X_1(x)) | \mathcal{F}_1(X)]$. Its Isometry principle implies that

$$\partial_x Q_{1-s}f(X_s(x)) \cdot \sigma(X_s(x)) = P(X_s(x)) E[\hat{D}_s f(X_1(x)) | \mathcal{F}_s(X)]$$

$ds \times dP$ -almost surely, where

$$\hat{D}_t f(X_1) = E[D_t f(X_1) | \mathcal{F}_1(X)].$$

It will be tempting to compare in detail these two representations: As

$$\hat{D}_t f(X_1) = (\hat{D}_t X_1)^* \partial f(X_1) \in \mathbb{R}^d,$$

we obtain

Theorem 13. *For any smooth function f on \mathbb{R}^n , we have the following commutation-intertwining relation:*

$$(7.26) \quad \sigma(X_t)^* \partial Q_{1-t} f(X_t) = P(X_t) E \left[(\hat{D}_t X_1)^* \partial f(X_1) | \mathcal{F}_t(X) \right].$$

Corollary 5 gives that

$$(\hat{D}_t X_1)^* = \sigma^*(X_t) K_t^* J_1^*,$$

where $J_t = \hat{\partial} X_t(x)$ and $K_t = J_t^{-1}$. Consequently, we get

$$\begin{aligned} E[\hat{D}_t f(X_1) | \mathcal{F}_t(X)] &= E[(\hat{D}_t X_1)^* \hat{\partial} f(X_1) | \mathcal{F}_t(X)] \\ &= E[\sigma^*(X_t) K_t^* J_1^* \hat{\partial} f(X_1) | \mathcal{F}_t(X)] \\ &= \sigma^*(X_t) K_t^* E[J_1^* \hat{\partial} f(X_1) | \mathcal{F}_t(X)]. \end{aligned}$$

To calculate the conditional expectation of the last line, we use the flow property of the diffusion process: For $s < t \leq 1$, let $X_t^s(x, \omega)$ denote the solution of the SDE defining the diffusion process at the instant t , starting from the point $x \in \mathbb{R}^n$ at the instant s . The flow property implies that $X_1(x, \omega) = X_1^t(X_t(x, \omega), \tilde{\omega})$, where $\tilde{\omega}$ denotes a Brownian path independent of $\{B_\tau : \tau \leq t\}$. Taking the derivative of both sides, from the chain rule, it comes

$$\partial_x X_1(x, \omega) = \partial X_1^t(X_t(x, \omega), \tilde{\omega}) \partial X_t(x, \omega)$$

almost surely. Hence

$$\begin{aligned} E[J_1^* \partial f(X_1) | \mathcal{F}_t(X)] &= E[\partial X_1^* \partial f(X_1) | \mathcal{F}_t(X)] \\ &= E[(\partial X_1^t(X_t, \tilde{\omega}) \partial X_t(x))^* \partial f(X_1^t \circ X_t) | \mathcal{F}_t(X)] \\ &= E[\partial X_t^* E[(\partial X_1^t)^* \circ (X_t, \tilde{\omega}) \partial f(X_1^t \circ X_t) | \mathcal{B}_t] | \mathcal{F}_t(X)], \end{aligned}$$

where $(\mathcal{B}_t, t \geq 0)$ represents the filtration of the underlying Brownian motion. Due to the independence and the stationarity of the Brownian increments, the inner conditional expectation can be written as:

$$\begin{aligned} E[(\partial X_1^t)^* \circ (X_t, \tilde{\omega}) \partial f(X_1^t \circ X_t) | \mathcal{B}_t] &= E[(\partial X_1^t)^*(\xi) \partial f(X_1^t(\xi))] |_{\xi=X_t(x)} \\ &= E[(\partial X_{1-t})^*(\xi) \partial f(X_{1-t}(\xi))] |_{\xi=X_t(x)} \end{aligned}$$

and in particular, it is $\mathcal{F}_t(X)$ -measurable. Let us define the semigroup $(M_t, t \geq 0)$ on the vector-valued functions as

$$M_t \alpha(\xi) = E[(\partial X_t(\xi))^* \alpha(X_t(\xi))],$$

$\xi \in \mathbb{R}^n$. Using this semigroup we end up with the identity

$$(7.27) \quad \begin{aligned} E[J_1^* \partial f(X_1) | \mathcal{F}_t(X)] &= E[\partial X_t^* | \mathcal{F}_t(X)] M_{1-t} \partial f(X_t(x)) \\ &= \hat{\partial} X_t^*(x) M_{1-t} \partial f(X_t(x)). \end{aligned}$$

Finally we find that

$$(7.28) \quad E[\hat{D}_t f(X_1(x)) | \mathcal{F}_t(X)] = \sigma(X_t(x))^* M_{1-t} \partial f(X_t(x))$$

almost surely. Although the expression (7.28) is what we can obtain by differentiation under the expectation, it has an important byproduct that we write separately as a lemma:

Lemma 3. *For any smooth function f on \mathbb{R}^n , the process $(\hat{\partial}X_t^*(x)M_{1-t}\partial f(X_t(x)), t \in [0, 1])$ is an $\mathcal{F}(X(x))$ -martingale.*

The following result may be called as a modified Poincaré inequality:

Theorem 14. *Let Z be the functional defined as*

$$Z(x) = \int_0^1 |\sigma^*(X_t(x))(\hat{\partial}X_t(x)^*)^{-1}|^2 dt.$$

We then have, for any $f \in C_b^2(\mathbb{R}^n)$, the following inequality:

$$(7.29) \quad E[|f(X_1(x)) - E[f(X_1(x))]|^2] \leq E \left[Z \left(\hat{\partial}X_1(x) \cdot \hat{\partial}X_1^*(x) \partial f(X_1(x)), \partial f(X_1(x)) \right)_{\mathbb{R}^n} \right].$$

Remark: Let operator-valued function $\Gamma(x, y)$ be defined as

$$(7.30) \quad \Gamma(x, y) = E[Z \hat{\partial}X_1(x) \cdot \hat{\partial}X_1^*(x) | X_1(x) = y],$$

then we can read the same inequality in \mathbb{R}^n as

$$\int_{\mathbb{R}^n} \left| f(y) - \int_{\mathbb{R}^n} f(z) q_1(x, z) dz \right|^2 q_1(x, y) dy \leq \int_{\mathbb{R}^n} (\Gamma(x, y) \partial f(y), \partial f(y))_{\mathbb{R}^n} q_1(x, y) dy,$$

where $y \rightarrow q_1(x, y)$ is the density of the law of $X_1(x)$.

Proof: From the martingale representation theorem, from the relation (7.28) and from the Itô isometry, we obtain

$$\begin{aligned} E[|f(X_1(x)) - E[f(X_1(x))]|^2] &= E \int_0^1 |\sigma^*(X_t(x))M_{1-t}\partial f(X_t(x))|^2 dt \\ &= E \int_0^1 |\sigma^*(X_t(x))(\hat{\partial}X_t^*)^{-1}(\hat{\partial}X_t^*)M_{1-t}\partial f(X_t(x))|^2 dt \\ &\leq E \int_0^1 |\sigma^*(X_t(x))(\hat{\partial}X_t^*)^{-1}|^2 |(\hat{\partial}X_t^*)M_{1-t}\partial f(X_t(x))|^2 dt \\ &\leq E \int_0^1 |\sigma^*(X_t(x))(\hat{\partial}X_t^*)^{-1}|^2 |(\hat{\partial}X_1^*)\partial f(X_1(x))|^2 dt, \end{aligned}$$

where the inequality at the last line follows from Lemma 3. \square

Same observation gives also an extension of the logarithmic Sobolev inequality for the measure $q_1(x, y)dy$ which is the law of the random variable $\omega \rightarrow X_1(\omega, x)$:

Theorem 15. *Let $f \in C_b^2(\mathbb{R}^n)$, then, we have*

$$(7.31) \quad E \left[f^2(X_1) \log \frac{f^2(X_1)}{E[f^2(X_1)]} \right] \leq 2 \int_{\mathbb{R}^n} (\Gamma(x, y) \partial f(y), \partial f(y))_{\mathbb{R}^n} q_1(x, y) dy,$$

where Γ is defined with (7.30).

Proof: We proceed as in the proof of Theorem 11 with $f \in C_b^2(\mathbb{R}^n)$ which is strictly positive. Moreover we can suppose that $E[f(X_1(x))] = 1$. In this case the claimed inequality takes the form

$$E[f(X_1) \log f(X_1)] \leq \int_{\mathbb{R}^n} (\Gamma(x, y) \partial f(y), \partial f(y))_{\mathbb{R}^n} \frac{1}{f(y)} q_1(x, y) dy.$$

Let $d\gamma = f(X_1(x))dP$, similar calculations of entropy $H(\gamma|P)$ as in the proof of Theorem 11 imply that

$$\begin{aligned}
H(\gamma|P) &= \int \log f(X_1(x))d\gamma \\
&= \frac{1}{2} E_\gamma \int_0^1 \left| \frac{P(X_s)E[\hat{D}_s f(X_1)|\mathcal{F}_s(X)]}{Q_{1-s}f(X_s)} \right|^2 \\
&= \frac{1}{2} E_\gamma \int_0^1 \left| \frac{P(X_s)\sigma^*(X_s(x))M_{1-s}\partial f(X_s(x))}{Q_{1-s}f(X_s)} \right|^2 \\
&\leq E_\gamma \int_0^1 \left| \frac{\sigma^*(X_s)(\hat{\partial}X_s^*)^{-1}\hat{\partial}X_s^*M_{1-s}\partial f(X_s)}{Q_{1-s}f(X_s)} \right|^2 ds \\
&\leq E_\gamma \int_0^1 |\sigma^*(X_s)(\hat{\partial}X_s^*)^{-1}|^2 \left| \frac{\hat{\partial}X_s^*M_{1-s}\partial f(X_s)}{Q_{1-s}f(X_s)} \right|^2 ds.
\end{aligned}$$

Note that, from Lemma 3, the process

$$\left(\frac{\hat{\partial}X_s^*M_{1-s}\partial f(X_s)}{Q_{1-s}f(X_s)}, s \in [0, 1] \right)$$

is a γ -martingale, consequently we have

$$\begin{aligned}
H(\gamma|P) &\leq E_\gamma \left[\left| \frac{\hat{\partial}X_1^*\partial f(X_1)}{f(X_1)} \right|^2 \int_0^1 |\sigma^*(X_t)(\hat{\partial}X_t^*)^{-1}|^2 dt \right] \\
&= E \left[\frac{1}{f(X_1)} |\hat{\partial}X_1^*\partial f(X_1)|^2 Z \right],
\end{aligned}$$

where Z is defined in the announcement of Theorem 14. \square

Left side of the equation (7.31) depends only on the law of the random variable $X_1(x)$, but at the right hand side there is the term $\Gamma(x, y)$ which is defined as

$$\Gamma(x, y) = E[Z\hat{\partial}X_1(x) \cdot \hat{\partial}X_1^*(x)|X_1(x) = y],$$

where

$$Z(x) = \int_0^1 |\sigma^*(X_t(x))(\hat{\partial}X_t(x)^*)^{-1}|^2 dt.$$

Consequently, the right hand side of the inequality (7.31) depends rather explicitly on σ , while the left hand side depends only on $a = \sigma\sigma^*$, hence only on the law of the process. Meanwhile, this inequality remains valid for any σ which is $n \times d$ -matrix valued function satisfying the relation $a = \sigma\sigma^*$. In particular the second dimension d is not fixed. Let us define

$$\Sigma_a = \cup_{d \geq 1} \{ \sigma \in C_b^1(\mathbb{R}^n, M(n \times d)) : \sigma\sigma^* = a \},$$

where $M(n \times d)$ denotes $n \times d$ -matrices.

Corollary 8. *Let us define the functional $Q(\partial f, \partial f)$ for $f \in C_b^2(\mathbb{R}^n)$ as*

$$\inf_{\sigma \in \Sigma_a} \int_{\mathbb{R}^n} (\Gamma_\sigma(x, y)\partial f(y), \partial f(y))_{\mathbb{R}^n} q_1(x, y) dy.$$

Let ν_1^x be the law of the random variable $\omega \rightarrow X_1(x, \omega)$. Then it holds true that

$$(7.32) \quad \int_{\mathbb{R}^n} f^2(y) \log \frac{f^2(y)}{\nu_1^x(f^2)} \nu_1^x(dy) \leq Q(\partial f, \partial f).$$

8. An example: Poincaré and Log-Sobolev inequalities on the Heisenberg Group

Let us apply the ideas explained above to prove the logarithmic Sobolev inequality for the heat kernel on the Heisenberg group $G = \mathbb{R}^3$, where the group operation \star is defined for (x, y, z) and (a, b, c) as

$$(x, y, z) \star (a, b, c) = (x + a, y + b, z + c + \frac{1}{2}(xb - ya)).$$

Let b and w be two real-valued, independent Brownian motions, the diffusion process with which we are concerned is the unique solution of the following SDE:

$$x_t = \int_0^t \sigma(x_s) dB_s,$$

where B is the two-dimensional Brownian motion $B = (b, w)$ and σ is the matrix given by

$$\sigma(x, y, z) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ -y/2 & x/2 \end{bmatrix}$$

The solution of this equation is given explicitly as $(x_t = (b_t, w_t, \frac{1}{2} z_t(b, w)), t \in \mathbb{R}_+)$, where (z_t) is the Lévy area process:

$$z_t = \int_0^t (b_s dw_s - w_s db_s).$$

We shall denote by $(Q_t, t \geq 0)$ the semi-group corresponding to the corresponding Markov process defined by $Q_t f(p) = E[f(p \star x_t)]$, $p \in \mathbb{R}^3$. We denote by X and Y the left-invariant vector fields defined as

$$\begin{aligned} X &= \partial_x - \frac{y}{2} \partial_z \\ Y &= \partial_y + \frac{x}{2} \partial_z, \end{aligned}$$

and with \hat{X} and \hat{Y} the right invariant ones:

$$\begin{aligned} \hat{X} &= \partial_x + \frac{y}{2} \partial_z \\ \hat{Y} &= \partial_y - \frac{x}{2} \partial_z. \end{aligned}$$

Note that the hypoelliptic infinitesimal generator \mathbb{L} of (Q_t) is given by

$$\mathbb{L} = \frac{1}{2} (X^2 + Y^2).$$

It follows from direct calculation that \hat{X} and \hat{Y} commute with \mathbb{L} , hence with the semigroup (Q_t) . Let now A be the isometry defined by

$$-A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$$

and let us denote by $(y_t, t \geq 0)$ the process defined by $y_t = (-b_t, -w_t, \frac{1}{2} z_t(b, w))$, note that $y_t = Ax_t$ and the law of y is equal to the law of x . Moreover, y defines also a Markov process with semi-group defined as $R_t f(p) = E[f(y_t \star p)]$ with infinitesimal generator

$$\hat{\mathbb{L}} = \frac{1}{2} (\hat{X}^2 + \hat{Y}^2).$$

Although the semi-groups $(Q_t, t \geq 0)$ and $(R_t, t \geq 0)$ are different, they are equal on the paths starting from the origin:

Lemma 4. *For any $t \geq 0$, $f \in C_b(\mathbb{R}^3)$, we have*

$$(8.33) \quad R_{T-t} f(y_t) = E[f(y_T) | \mathcal{F}_t(x)] = Q_{T-t} f(y_t)$$

$$(8.34) \quad Q_{T-t} f(x_t) = E[f(x_T) | \mathcal{F}_t(x)] = R_{T-t} f(x_t)$$

for any $T \geq t$, P -almost surely. In particular, we have

$$(8.35) \quad \hat{\mathbb{L}} f(y_t) = \mathbb{L} f(y_t)$$

$$(8.36) \quad \hat{\mathbb{L}} f(x_t) = \mathbb{L} f(x_t)$$

Proof: Let $\theta(b, w)$ be any bounded and $\mathcal{F}_t(x)$ -measurable function. As $\mathcal{F}_t(x) = \sigma(b_s, w_s, s \leq t)$. We have

$$\begin{aligned} E[f(y_T(b, w)) \theta(b, w)] &= E[f(y_T(-b, -w)) \theta(-b, -w)] \\ &= E[f(x_T(b, w)) \theta(-b, -w)] \\ &= E[E[f(x_T) | \mathcal{F}_t(x)] \theta(-b, -w)] \\ &= E[Q_{T-t} f(x_t(b, w)) \theta(-b, -w)] \\ &= E[Q_{T-t} f(x_t(-b, -w)) \theta(b, w)] \\ &= E[Q_{T-t} f(y_t(b, w)) \theta(b, w)], \end{aligned}$$

by the invariance of the law of the Brownian motion under the isometries. The second claim follows by symmetry, the equalities (8.35) and (8.36) follow¹ from the first part by differentiation under the expectation. \square

Lemma 5. *For any smooth function f on \mathbb{R}^3 and $t, T \geq 0$, we have*

$$\hat{X} R_T f(y_t) = \hat{X} Q_T f(y_t)$$

$$\hat{Y} R_T f(y_t) = \hat{Y} Q_T f(y_t)$$

almost surely.

Proof: As the second one is similar, let us prove the first relation. Since both sides are measurable w.r.t. the sigma algebra generated by y_t , it suffices to test them on the functions $k(y_t)$, with k being smooth. Let r_t be the density of the law of y_t w.r.t. the Lebesgue measure of \mathbb{R}^3 . Then we have

$$\begin{aligned} E[\hat{X} R_T f(y_t) k(y_t)] &= -E[R_T f(y_t) (\hat{X} k(y_t) + k(y_t) \hat{X} \log r_t(y_t))] \\ &= -E[Q_T f(y_t) (\hat{X} k(y_t) + k(y_t) \hat{X} \log r_t(y_t))] \\ &= E[\hat{X} Q_T f(y_t) k(y_t)], \end{aligned}$$

¹These two relations follow also directly by elementary calculation

hence the proof follows. \square

Proposition 6. *For any smooth function f on \mathbb{R}^3 and $t \geq 0$, we have*

•

$$f(y_1) = - \int_0^1 Q_{1-s} \hat{X} f(y_s) db_s - \int_0^1 Q_{1-s} \hat{Y} f(y_s) dw_s ,$$

• *in particular the following bound hold true:*

$$E[|f(y_1) - E[f(y_1)]|^2] \leq E \left[|\hat{X} f(y_1)|^2 + |\hat{Y} f(y_1)|^2 \right] .$$

Proof: We will apply the Itô formula to the process $(R_{1-t}f(y_t), t \in [0, 1])$ which is also a martingale for smooth f . From Lemma 4, $R_{1-t}f(y_t) = Q_{1-t}f(y_t)$ and from Lemma 5, $\hat{X}R_{1-t}f(y_t) = \hat{X}Q_{1-t}f(y_t)$ and $\hat{Y}R_{1-t}f(y_t) = \hat{Y}Q_{1-t}f(y_t)$. Moreover, as \hat{X} and \hat{Y} commute with Q_s , we get

$$\begin{aligned} (8.37) \quad R_{1-t}f(y_t) &= Q_{1-t}f(y_t) \\ &= - \int_0^t \hat{X} Q_{1-s} f(y_s) db_s - \int_0^t \hat{Y} Q_{1-s} f(y_s) dw_s \\ &= - \int_0^t Q_{1-s} \hat{X} f(y_s) db_s - \int_0^t Q_{1-s} \hat{Y} f(y_s) dw_s . \end{aligned}$$

At first sight, in the equality (8.37) the integrals should be taken as Stratonovitch integrals but, the left hand side is a continuous martingale, consequently the only terms remaining after calculations are Itô integrals, thanks to the relation

$$\hat{\mathbb{L}}f(y_t) = \mathbb{L}f(y_t) .$$

To complete the proof, it suffices to remark that the integrands in the equation (8.37) are martingales, hence their squares are submartingales, consequently their expectations increase with $s \in [0, 1]$:

$$\begin{aligned} \sup_{s \leq 1} E[|Q_{1-s} \hat{X} f(y_s)|^2] &\leq E[|\hat{X} f(y_1)|^2] \\ \sup_{s \leq 1} E[|Q_{1-s} \hat{Y} f(y_s)|^2] &\leq E[|\hat{Y} f(y_1)|^2] . \end{aligned}$$

\square

We are ready to prove

Theorem 16. *For any smooth function f on \mathbb{R}^3 , we have the following Poincaré inequality*

$$E[|f(x_1) - E[f(x_1)]|^2] \leq E[|Xf(x_1)|^2 + |Yf(x_1)|^2] .$$

Proof: By the equality of the laws of x and y and using the inequality of Proposition 6, we have

$$\begin{aligned} E[|f(x_1) - E[f(x_1)]|^2] &= E[|f(y_1) - E[f(y_1)]|^2] \\ &\leq E \left[|\hat{X} f(y_1)|^2 + |\hat{Y} f(y_1)|^2 \right] . \end{aligned}$$

Moreover, the term inside the expectation of the right hand side is the square of the norm of the operator of Paul-André Meyer, called “opérateur carré du champs” (cf. [7]) of the right Markov process $(y_t \star p, t \in \mathbb{R}, p \in \mathbb{R}^3)$ and it can be written as

$$|\hat{X} f(y_1)|^2 + |\hat{Y} f(y_1)|^2 = -\frac{1}{2} [\hat{\mathbb{L}}f^2(y_1) - 2f(y_1)\hat{\mathbb{L}}f(y_1)] ,$$

and from Proposition 6, it is also equal to

$$-\frac{1}{2} [\mathbb{L}f^2(y_1) - 2f(y_1)\mathbb{L}f(y_1)] = |Xf(y_1)|^2 + |Yf(y_1)|^2,$$

and the claim follows using again the equality of the laws of the processes x and y . \square

Having developped all the necessary techniques, we can now prove the logarithmic Sobolev inequality for the measure defined as the law of the process at the instant $t = 1$, which is regarded as a measure on the Heisenberg group in \mathbb{R}^3

Theorem 17. *For any smooth function on \mathbb{R}^3 , we have*

$$(8.38) \quad E [f^2(x_1)(\log f^2(x_1) - \log E[f^2(x_1)])] \leq 2(E[|Xf(x_1)|^2 + |Yf(x_1)|^2]).$$

Proof: We begin with the right process and define the martingale $M_t = E[f(y_1)|\mathcal{F}_t(x)]$, while supposing that f is strictly positive and $E[f(x_1)] = E[f(y_1)] = 1$. Using the Itô formula and Lemmas 4, 5:

$$\begin{aligned} dM_t &= -\frac{\hat{X}R_{1-t}f(y_t)}{R_{1-t}f(y_t)}db_t - \frac{\hat{Y}R_{1-t}f(y_t)}{R_{1-t}f(y_t)}dw_t \\ &= -\frac{\hat{X}Q_{1-t}f(y_t)}{Q_{1-t}f(y_t)}db_t - \frac{\hat{Y}Q_{1-t}f(y_t)}{Q_{1-t}f(y_t)}dw_t \\ &= -\frac{Q_{1-t}\hat{X}f(y_t)}{Q_{1-t}f(y_t)}db_t - \frac{Q_{1-t}\hat{Y}f(y_t)}{Q_{1-t}f(y_t)}dw_t, \end{aligned}$$

where we have used the fact that Q_t commutes with \hat{X} and \hat{Y} . Let us denote by ν the probability $d\nu = f(y_1)dP$. Then, we get from above

$$\begin{aligned} E[f(y_1) \log f(y_1)] &= \frac{1}{2} E_\nu \int_0^1 \left[|E_\nu[\hat{X} \log f(y_1)|\mathcal{F}_t(x)]|^2 + |E_\nu[\hat{Y} \log f(y_1)|\mathcal{F}_t(x)]|^2 \right] dt \\ &\leq \frac{1}{2} E_\nu [|\hat{X} \log f(y_1)|^2 + |\hat{Y} \log f(y_1)|^2] \\ &= \frac{1}{2} E \left[\frac{1}{f(y_1)} (|\hat{X}f(y_1)|^2 + |\hat{Y}f(y_1)|^2) \right] \end{aligned}$$

Replacing $f(y_1)$ by $f^2(y_1)/E[f^2(y_1)]$, we obtain from above

$$(8.39) \quad E [f^2(y_1) \log(f^2(y_1)/E[f^2(y_1)])] \leq 2E[|\hat{X}f(y_1)|^2 + |\hat{Y}f(y_1)|^2].$$

The left hand side of (8.39) is equal to

$$E [f^2(x_1)(\log f^2(x_1) - \log E[f^2(x_1)])]$$

and the right hand side is equal to the expectation of the ‘‘carré du champs’’ applied to f at the point x_1 and this finishes the proof. \square

REFERENCES

- [1] D. Bakry, F. Baudoin, M. Bonnefont, D. Chafai: “On gradient bounds for the heat kernel on the Heisenberg group”. *Jour. Functional Analysis*, 255, 1905-1938 (2008).
- [2] M. Capitaine, E. Hsu and M. Ledoux: “Martingale representation and a simple proof of logarithmic Sobolev inequalities on path spaces”. *Electron. Comm. Probab.* 2, 71–81. (1997).
- [3] E. Cépa: Equations différentielles stochastiques multivoques. Séminaire de Probabilités, tome 29, p. 86-107, *Lecture Notes in Math.* Vol. 1613, 1995.
- [4] E. Cépa and D. Lepingle: Diffusing particles with electrostatic repulsion. *Probab. Theory and Related Fields*, Vol. 107, p. 429-449, 1992.
- [5] E. Cépa and D. Lepingle: “Brownian particles with electrostatic repulsion on the circle: Dyson’s model for unitary random matrices revisited”. *ESAIM: Probability and Statistics*, Vol. 5, p. 203-224, 2001.
- [6] C. Dellacherie: “Intégrales stochastiques par rapport aux processus de Wiener ou de Poisson”. In *Séminaire de Probabilités VIII*, *Lecture Notes in Mathematics* **381**. Springer-Verlag, 1973. [Correction dans SP IX, LNM 465.]
- [7] C. Dellacherie et P.-A. Meyer: “Probabilités et Potentiel, Processus de Markov, Chapitres 17 à 24”. *Collection Enseignement des Sciences*, Hermann, 1992.
- [8] B.K. Driver, T. Melcher: “Hypoelliptic heat kernel inequalities on the Heisenberg group, ”. *Jour. Funct. Anal.* 221 (2) 340–365 (2005).
- [9] F. J. Dyson: “A Brownian-Motion Model for the Eigenvalues of a Random Matrix”. *Journal of Mathematical Physics* 3, 1191 (1962);
- [10] L. Gross: “Integration and non-linear transformation in Hilbert space”. *Trans. Am. Math. Soc.* 94, p.404-440 (1960).
- [11] L. Gross: “Logarithmic Sobolev inequalities”. *Amer. Jour. Math.* **97**, p.1061-1083 (1975).
- [12] N. Ikeda and S. Watanabe: *Stochastic Differential Equations and Diffusion Processes*
- [13] K. Itô: “Multiple Wiener integrals”. *J. Math. Soc. Japan*, **3**, p. 157–169, 1951.
- [14] H.H. Kuo: *Gaussian Measures on Banach Spaces*. *Lecture Notes in Math.* Vol. 463. Springer, 1975.
- [15] H.-Q. Li: “Estimation optimale du gradient du semi-groupe de la chaleur sur le groupe de Heisenberg”. *Jour. Funct. Anal.* 236 (2) 369–394 (2006).
- [16] D. Ocone: “Malliavin calculus and stochastic integral representation of functionals of diffusion processes”. *Stochastics* 12, p.161-185 (1984).
- [17] D. Ocone: “A guide to the stochastic calculus of variations”. *Stochastic analysis and related topics* (Silivri, 1986), 1–79, *Proceedings of Silivri Conf. on Stochastic Analysis and Related Topics*, p.247-266. *Lecture Notes in Math.* Vol.1316. Springer, 1988.
- [18] I. Shigekawa: “Derivatives of Wiener functionals and absolute continuity of induced measures”. *J. Math. Kyoto Univ.* 20, p.263-289, 1980.
- [19] D.W. Stroock and S.R.S. Varadhan: “Diffusion processes with continuous coefficients 1”. *Comm. Pure and Appl. Math.*, **22**, pp.345-400, 1969.
- [20] A. S. Üstünel: *Introduction to Analysis on Wiener Space*. *Lecture Notes in Math.* Vol. **1610**. Springer, 1995.
- [21] A. S. Üstünel: *Analysis on Wiener Space and Applications*. Electronic text at the site arXiv.org. math. arXiv:1003.1649
- [22] A. S. Üstünel: “Entropy, invertibility and variational calculus of adapted shifts on Wiener space”. *J. Funct. Anal.* **257**, no. 8, p.3655–3689, 2009.
- [23] A. S. Üstünel: “Variational calculation of Laplace transforms via entropy on Wiener space and applications”. *J. Funct. Anal.* **267**, no. 8, 3058–3083, 2014.
- [24] A. S. Üstünel: “Martingale representation for degenerate diffusions”. *J. Funct. Anal.* **276**, no. 11, 3468-3483, 2019.
- [25] Y. Yamada, S. Watanabe: “On the uniqueness of solutions of stochastic differential equations”. *J. Math. Kyoto Univ.* **11** p. 155-167, 1971.
- [26] K. Yosida: *Functional Analysis*. *Die Grundlehren der math. Wissenschaften*, **Vol. 123**, 1970.

A. S. Üstünel, Bilkent University, Math. Dept., Ankara, Turkey
ustunel@fen.bilkent.edu.tr