

Exact Exchange with Range-separated Algorithm for Thermodynamic Limit of Periodic Hartree-Fock Theory

Qiming Sun^{1, a)}

Quantum Engine LLC, WA 98516, US

(Dated: 27 April 2023)

The expensive cost of computing exact exchange in periodic systems limits the application range of density functional theory with hybrid functionals. To reduce the computational cost of exact change, we present a range-separated algorithm to compute electron repulsion integrals for Gaussian-type crystal basis. The algorithm splits the full-range Coulomb interactions into short-range and long-range parts, which are respectively computed in real and reciprocal space. This approach significantly reduces the overall computational cost as integrals can be efficiently computed in both regions. The algorithm can efficiently handle large numbers of \mathbf{k} points with limited CPU and memory resources. As a demonstration, we performed an all-electron \mathbf{k} -point Hartree-Fock calculation for LiH crystal with 1 million Gaussian basis functions, which was completed on a desktop computer in 1400 CPU hours.

^{a)}Electronic mail: osirpt.sun@gmail.com

I. INTRODUCTION

Density functional theory (DFT) is the most widely used tool for modeling systems with periodic boundary conditions (PBC). Large scale DFT simulations without Hartree-Fock exchange (HFX) were available for PBC systems of millions of basis functions^{1,2}. The use of hybrid functionals in DFT, which includes HFX contributions in the DFT exchange-correlation functional, can significantly improve the accuracy of DFT and widen the range of applications. However, accurately and efficiently evaluating HFX is a major challenge for PBC DFT calculations. Problems like numerical divergence and singularity in \mathbf{k} -point sampling may be found in HFX with PBC and various treatments were developed to overcome these problems³⁻¹⁶. The concern of using hybrid functional DFT in PBC application is mainly the computational cost of HFX which can be several orders of magnitude higher than the rest part.

In crystal simulations, planewave (PW) basis is a natural choice due to its periodicity. Computing HFX with PW basis¹⁷⁻²² is extremely demanding and generally considered impractical. Due to the locality of Gaussian functions, typically less Gaussian functions are required to describe the atomic characters in a system. It is more affordable to compute HFX in Gaussian basis than in PW basis. Gaussian-type crystal basis are widely adopted in many program packages²³⁻²⁸. For PBC calculations, Gaussian and planewave method^{29,30} (GPW) is a commonly applied method to compute electron repulsion integrals for Gaussian basis. In GPW method, PW basis serves as the auxiliary basis of density fitting method. Discrete Fourier transform is often employed to evaluate the density and Coulomb potential on PW basis, which has $\mathcal{O}(N_G \log(N_G))$ complexity for N_G PW basis (or the number of integration grids). For a calculation of N_A atoms and N_k \mathbf{k} points, the cost of HFX with the GPW method scales as at least $N_A^2 N_k^2 N_G \log(N_G)$.

The GPW method is not suitable for all-electron calculations as it requires a high energy cutoff for accurate description of core orbitals, which leads to a large number of auxiliary PW basis. To reduce the energy cutoff, pseudopotentials³¹⁻³⁵ (PP) become critical in PW basis methods as well as the GPW method. Although PPs are commonly used in crystal calculations, there are several issues for applications with PP. Accuracy and reliability of DFT calculations are badly affected by the quality of PP. Results obtained from different PPs are often not comparable³⁶. Applications with PP are limited to valence properties only. It is impossible to study properties of core electrons with PP. As such, all-electron methods receive significant attentions³⁷⁻⁴² since they can naturally address all of these issues.

Gaussian and augmented plane wave (GAPW) method^{37,43,44} was developed to address the energy cutoff issue in GPW. This method uses Gaussian functions to describe electron density in the core region and calls the GPW method for the valence region only. However, GAPW cannot be used to compute HFX. Gaussian density fitting (GDF) were considered^{8,10,45–58} due to its great success for HFX and integral evaluation in molecular systems. It should be noted that if integrals in GDF are computed with the GPW algorithm, GDF method may still suffer from the energy cutoff issue in all-electron calculations. GDF algorithms specialized on all-electron methods were developed to solve the energy cutoff issue^{54,59–61}.

Despite that GPW and GDF methods provide the possibility to compute HFX with PBC Gaussian basis, their computational requirements of HFX still make them impractical when targeting the thermodynamic limit in periodic systems. The GDF method requires a storage of $\mathcal{O}(N_{AO}^3 N_k^2)$ for the intermediate tensors and $\mathcal{O}(N_{AO}^4 N_k^2)$ FLOPs for tensor contraction in HFX matrix. As a result, the feasible problem size of k-point hybrid DFT calculations with GDF methods is limited to around 100k basis functions.

In this work, we present a range-separated algorithm for the computation electron repulsion integrals with Gaussian crystal basis. This algorithm enables us to compute HFX for large periodic systems. Formulas, possible variants, and integral screening schemes involved in this algorithm are discussed in Section II. In Section III, we demonstrate the efficiency and accuracy of this algorithm by performing an all-electron HF calculation for LiH crystal with a million basis functions.

II. THEORY

A. Fock matrix with range-separated algorithm

We consider here a periodic system which consists of repeated unit cells in terms of the translation vector \mathbf{m} and lattice vectors \mathbf{a}

$$\mathbf{m} = m_x \mathbf{a}_x + m_y \mathbf{a}_y + m_z \mathbf{a}_z. \quad (1)$$

Crystal orbitals for a crystal of $N \rightarrow \infty$ unit cells can be expanded with a set of Bloch orbitals

$$\phi_\mu(\mathbf{r})^{\mathbf{k}} = \frac{1}{\sqrt{N}} \sum_{\mathbf{m}} e^{i\mathbf{k} \cdot \mathbf{m}} \mu^{\mathbf{m}}(\mathbf{r}) \quad (2)$$

where $\mu^{\mathbf{m}}$ is a real Gaussian-type orbital centered at \mathbf{R}_μ in cell \mathbf{m}

$$\mu^{\mathbf{m}}(\mathbf{r}) = \mu(\mathbf{r} - \mathbf{R}_\mu - \mathbf{m}). \quad (3)$$

Solving the periodic HF model can lead to the \mathbf{k} -point adapted Roothaan equations

$$\mathbf{F}^{\mathbf{k}}\mathbf{C}^{\mathbf{k}} = \mathbf{S}^{\mathbf{k}}\mathbf{C}^{\mathbf{k}}\boldsymbol{\varepsilon}^{\mathbf{k}}, \quad (4)$$

The Fock matrix and overlap matrix at \mathbf{k} are

$$\mathbf{F}^{\mathbf{k}} = \mathbf{T}^{\mathbf{k}} + \mathbf{V}_{\text{Nuc}}^{\mathbf{k}} + \mathbf{J}^{\mathbf{k}} - \mathbf{K}^{\mathbf{k}}, \quad (5)$$

$$S_{\mu\nu}^{\mathbf{k}} = \langle \phi_{\mu}^{\mathbf{k}} | \phi_{\nu}^{\mathbf{k}} \rangle. \quad (6)$$

The overlap matrix can be computed in terms of the lattice sum over overlap integrals in real space

$$S_{\mu\nu}^{\mathbf{k}} = \sum_{\mathbf{m}} e^{i\mathbf{k}\cdot\mathbf{m}} S_{\mu^0\nu^{\mathbf{m}}}, \quad (7)$$

$$S_{\mu^0\nu^{\mathbf{m}}} = \langle \mu^0 | \nu^{\mathbf{m}} \rangle. \quad (8)$$

In a similarly way, we can get the kinetic matrix in $\mathbf{F}^{\mathbf{k}}$

$$T_{\mu\nu}^{\mathbf{k}} = \sum_{\mathbf{m}} e^{i\mathbf{k}\cdot\mathbf{m}} T_{\mu^0\nu^{\mathbf{m}}}. \quad (9)$$

When computing the nuclear attraction matrix $\mathbf{V}_{\text{Nuc}}^{\mathbf{k}}$, we use model charge to screen the long-range interactions then compute separately the long-range and short-range parts of nuclear attraction integrals. Relevant formulas have been documented in 54.

Given \mathbf{k} denoted density matrices $D^{\mathbf{k}}$, the HF Coulomb and exchange matrices in $\mathbf{F}^{\mathbf{k}}$ are

$$J_{\mu\nu}^{\mathbf{k}} = \frac{1}{N_k} \sum_{\mathbf{k}'} \sum_{\kappa\lambda} D_{\lambda\kappa}^{\mathbf{k}'} (\phi_{\mu}^{\mathbf{k}} \phi_{\nu}^{\mathbf{k}} | \phi_{\kappa}^{\mathbf{k}'} \phi_{\lambda}^{\mathbf{k}'}), \quad (10)$$

$$K_{\mu\lambda}^{\mathbf{k}} = \frac{1}{N_k} \sum_{\mathbf{k}'} \sum_{\kappa\nu} D_{\nu\kappa}^{\mathbf{k}'} (\phi_{\mu}^{\mathbf{k}} \phi_{\nu}^{\mathbf{k}'} | \phi_{\kappa}^{\mathbf{k}'} \phi_{\lambda}^{\mathbf{k}}). \quad (11)$$

The four-center two-electron repulsion integrals (ERI) in above equations are

$$(\phi_{\mu}^{\mathbf{k}\mu} \phi_{\nu}^{\mathbf{k}\nu} | \phi_{\kappa}^{\mathbf{k}\kappa} \phi_{\lambda}^{\mathbf{k}\lambda}) = \iint \phi_{\mu}^{\mathbf{k}\mu}(\mathbf{r}_1) \phi_{\nu}^{\mathbf{k}\nu}(\mathbf{r}_1) g(\mathbf{r}_{12}) \phi_{\kappa}^{\mathbf{k}\kappa}(\mathbf{r}_2) \phi_{\lambda}^{\mathbf{k}\lambda}(\mathbf{r}_2) d^3\mathbf{r}_1 d^3\mathbf{r}_2. \quad (12)$$

This integral can be evaluated in reciprocal space in two steps^{29,30}:

1. Fourier transform (FT) the orbital product $\rho_{\mu\nu}$ against the PW basis $|\mathbf{G}\rangle = \frac{(2\pi)^{3/2}}{\sqrt{N}} e^{i\mathbf{G}\cdot\mathbf{r}}$

$$\begin{aligned} \rho_{\mu\nu}^{\mathbf{k}\mathbf{k}'}(\mathbf{G}) &= \int e^{-i\mathbf{G}_{\mathbf{k}\mathbf{k}'}\cdot\mathbf{r}} \phi_{\mu}^{\mathbf{k}}^*(\mathbf{r}) \phi_{\nu}^{\mathbf{k}'}(\mathbf{r}) d^3\mathbf{r} \\ &= \sum_{\mathbf{n}} e^{i\mathbf{k}'\cdot\mathbf{n}} \rho_{\mu^0\nu^{\mathbf{n}}}(\mathbf{G}_{\mathbf{k}\mathbf{k}'}), \end{aligned} \quad (13)$$

$$= \sum_{\mathbf{m}} e^{-i\mathbf{k}\cdot\mathbf{m}} \rho_{\mu\mathbf{m}\nu 0}(\mathbf{G}_{\mathbf{k}\mathbf{k}'}) \quad (14)$$

$$\mathbf{G}_{\mathbf{k}\mathbf{k}'} = \mathbf{G} - \mathbf{k} + \mathbf{k}', \quad (15)$$

$$\rho_{\mu\mathbf{m}\nu n}(\mathbf{G}) = \int e^{-i\mathbf{G}\cdot\mathbf{r}} \mu(\mathbf{r} - \mathbf{m}) \nu(\mathbf{r} - \mathbf{n}) d^3\mathbf{r}; \quad (16)$$

2. A summation over PW basis inside the volume Ω of the unit cell

$$(\phi_{\mu}^{\mathbf{k}} \phi_{\nu}^{\mathbf{k}'} | \phi_{\kappa}^{\mathbf{k}'} \phi_{\lambda}^{\mathbf{k}}) = \frac{1}{\Omega} \sum_{\mathbf{G}} \mathcal{C}(\mathbf{G}_{\mathbf{k}\mathbf{k}'}) \rho_{\mu\nu}^{\mathbf{k}\mathbf{k}'}(\mathbf{G}) \rho_{\kappa\lambda}^{\mathbf{k}'\mathbf{k}}(-\mathbf{G}). \quad (17)$$

$\mathcal{C}(\mathbf{G})$ in the equation above is the Coulomb kernel in reciprocal space. The Coulomb metric $g(\mathbf{r}_{12})$ can be split into long-range (LR) plus short-range (SR) contributions

$$g(\mathbf{r}_{12}) = \frac{1}{r_{12}} = \frac{\text{erf}(\omega r_{12})}{r_{12}} + \frac{\text{erfc}(\omega r_{12})}{r_{12}}. \quad (18)$$

When computing ERIs with Eq. (17), the Fourier transformed full-range Coulomb kernel as well as the LR and SR Coulomb kernels are

$$\frac{1}{r_{12}} \rightarrow \mathcal{C}(\mathbf{G}) = \begin{cases} \frac{4\pi}{G^2} & |\mathbf{G}| \neq 0 \\ 0 & |\mathbf{G}| = 0 \end{cases}, \quad (19)$$

$$\frac{\text{erf}(\omega r_{12})}{r_{12}} \rightarrow \mathcal{C}_{\text{LR}}(\mathbf{G}) = \begin{cases} \frac{4\pi}{G^2} \exp(-\frac{G^2}{4\omega^2}) & |\mathbf{G}| \neq 0 \\ 0 & |\mathbf{G}| = 0 \end{cases}, \quad (20)$$

$$\frac{\text{erfc}(\omega r_{12})}{r_{12}} \rightarrow \mathcal{C}_{\text{SR}}(\mathbf{G}) = \begin{cases} \frac{4\pi}{G^2} (1 - \exp(-\frac{G^2}{4\omega^2})) & |\mathbf{G}| \neq 0 \\ \frac{\pi}{\omega^2} & |\mathbf{G}| = 0 \end{cases}. \quad (21)$$

The treatment of $\mathbf{G} = 0$ in the SR Coulomb kernel is different to the other two. When adding the LR and SR integrals to produce the integrals for full-range Coulomb, an additional contribution corresponding to the FT term at $\mathbf{G} = 0$ needs to be subtracted from the SR integrals

$$(\phi_{\mu}^{\mathbf{k}} \phi_{\nu}^{\mathbf{k}} | \phi_{\kappa}^{\mathbf{k}} \phi_{\lambda}^{\mathbf{k}})^{\text{SR}} \Big|_{\mathbf{G}=0} = \frac{\pi}{\Omega \omega^2} S_{\mu\nu}^{\mathbf{k}} S_{\kappa\lambda}^{\mathbf{k}}. \quad (22)$$

Due to the exponential decay term $\exp(-\frac{G^2}{4\omega^2})$ in the LR Coulomb kernel, we can expect that the contributions from large \mathbf{G} in Eq. (17) decrease rapidly. The summation over PWs in Eq. (17) can be truncated early thus a small energy cutoff is required to compute LR ERI in reciprocal space. Sparsity of SR ERIs in real space can be utilized to gain good efficiency⁸. By choosing

an appropriate value for ω and appropriate algorithms that we will discuss below, computing SR ERIs in real space and LR ERIs in reciprocal space appears to be an efficient approach.

In this approach, we use Ewald probe-charge correction⁷ to regularize the Coulomb singularity in HFX. It only requires to modify the FT integrals at $\mathbf{G} = 0$. Other attractive techniques such as the Wigner-Seitz truncated Coulomb interactions¹¹ are incompatible to the current range-separated algorithm. They require more PWs to describe the Coulomb kernel while this algorithm only use a small number of PWs for LR ERIs.

B. HFX in reciprocal space

Computing HFX in reciprocal space involves two time-consuming steps: analytical FT for orbital product $\rho_{\mu\nu}^{\mathbf{k}\mathbf{k}'}(\mathbf{G})$ with $\mathcal{O}(N_k^2 N_c N_{AO}^2 N_G)$ complexity as shown by Eq. (13), and the tensor contraction for three tensors which scales as $N_k^2 N_{AO}^3 N_G$

$$K_{\mu\lambda}^{\mathbf{k}} = \frac{1}{N_k} \sum_{\nu\kappa\mathbf{G}\mathbf{k}'} D_{\nu\kappa}^{\mathbf{k}'} \rho_{\mu\nu}^{\mathbf{k}\mathbf{k}'}(\mathbf{G}) \mathcal{C}(\mathbf{G}_{\mathbf{k}\mathbf{k}'}) \rho_{\kappa\lambda}^{\mathbf{k}'\mathbf{k}}(-\mathbf{G}). \quad (23)$$

The number of image cells in lattice-sum, N_c , is related to distance cutoff. The number of PW basis N_G needs to be determined by energy cutoff. Distance cutoff and energy cutoff are influenced by the Coulomb attenuation parameter ω and the compactness of the Gaussian basis functions⁶². It should be noted that the FT for orbital products has to be evaluated analytically. Using discrete FT here can cause significant errors⁵⁴. This is because we only use a small number of PW basis for the ERIs evaluated in reciprocal space. The FT associated real space grids are not dense enough to describe all orbital products.

To optimize the performance of FT for orbital product, utilizing the symmetry below can reduce the cost by a factor of two

$$[\rho_{\mu\nu}^{\mathbf{k}\mathbf{k}'}(\mathbf{G})]^* = \rho_{\nu\mu}^{\mathbf{k}'\mathbf{k}}(-\mathbf{G}). \quad (24)$$

Thereby, the orbital product tensor can be computed once and used four times for exchange matrices at two different \mathbf{k} points

$$K_{\mu\lambda}^{\mathbf{k}} = \sum_{\nu\kappa\mathbf{G}\mathbf{k}'} D_{\nu\kappa}^{\mathbf{k}'} \rho_{\mu\nu}^{\mathbf{k}\mathbf{k}'}(\mathbf{G}) \mathcal{C}(\mathbf{G}_{\mathbf{k}\mathbf{k}'}) [\rho_{\lambda\kappa}^{\mathbf{k}\mathbf{k}'}(\mathbf{G})]^*, \quad (25)$$

$$K_{\nu\kappa}^{\mathbf{k}'} = \sum_{\mu\lambda\mathbf{G}\mathbf{k}} D_{\mu\lambda}^{\mathbf{k}} \rho_{\nu\mu}^{\mathbf{k}'\mathbf{k}}(-\mathbf{G}) \mathcal{C}((-\mathbf{G})_{\mathbf{k}'\mathbf{k}}) [\rho_{\kappa\lambda}^{\mathbf{k}'\mathbf{k}}(-\mathbf{G})]^*. \quad (26)$$

During the self-consistent field (SCF) iteration, density matrices are constructed from occupied SCF orbitals (denoted as $C_{\mu j}^{\mathbf{k}}$ below) and orbital occupancies (denoted as $\Lambda_j^{\mathbf{k}}$). A common optimization practice is to use the occupied orbitals directly in the construction of exchange matrix. We can first carry out the contraction between the N_{occ} occupied orbitals and the orbital product tensor

$$\rho_{\mu j}^{\mathbf{k}\mathbf{k}'}(\mathbf{G}) = \sum_{\nu} \rho_{\mu\nu}^{\mathbf{k}\mathbf{k}'}(\mathbf{G}) C_{\nu j}^{\mathbf{k}'}, \quad (27)$$

$$\rho_{j\nu}^{\mathbf{k}'\mathbf{k}}(-\mathbf{G}) = \sum_{\mu} C_{\mu j}^{\mathbf{k}'} * \rho_{\mu\nu}^{\mathbf{k}'\mathbf{k}}(-\mathbf{G}) = \sum_{\mu} C_{\mu j}^{\mathbf{k}'} * [\rho_{\nu\mu}^{\mathbf{k}\mathbf{k}'}(\mathbf{G})]^* = [\rho_{\nu j}^{\mathbf{k}\mathbf{k}'}(\mathbf{G})]^*, \quad (28)$$

then obtain the exchange matrix with a contraction between the resultant tensors and another orbital product tensor

$$\begin{aligned} K_{\mu\nu}^{\mathbf{k}} &= \frac{1}{\Omega} \sum_{j\mathbf{G}\mathbf{k}'} \Lambda_j^{\mathbf{k}'} \rho_{\mu j}^{\mathbf{k}\mathbf{k}'}(\mathbf{G}) \mathcal{C}(\mathbf{G}_{\mathbf{k}\mathbf{k}'}) \rho_{j\nu}^{\mathbf{k}'\mathbf{k}}(-\mathbf{G}) \\ &= \frac{1}{\Omega} \sum_{j\mathbf{G}\mathbf{k}'} \Lambda_j^{\mathbf{k}'} \rho_{\mu j}^{\mathbf{k}\mathbf{k}'}(\mathbf{G}) \mathcal{C}(\mathbf{G}_{\mathbf{k}\mathbf{k}'}) [\rho_{\nu j}^{\mathbf{k}\mathbf{k}'}(\mathbf{G})]^*. \end{aligned} \quad (29)$$

This formula results in a reduction of the computational cost to $N_k^2 N_{AO}^2 N_{occ} N_G$ for the step of tensor contraction. The permutation symmetry Eq. (24) for orbital product tensor can still be utilized, leading to the contraction for the exchange matrix at \mathbf{k}'

$$\rho_{\nu j}^{\mathbf{k}'\mathbf{k}}(-\mathbf{G}) = \sum_{\mu} [\rho_{\mu\nu}^{\mathbf{k}\mathbf{k}'}(\mathbf{G})]^* C_{\mu j}^{\mathbf{k}}, \quad (30)$$

$$K_{\mu\nu}^{\mathbf{k}'} = \frac{1}{\Omega} \sum_{j\mathbf{G}\mathbf{k}} \Lambda_j^{\mathbf{k}} \rho_{\mu j}^{\mathbf{k}'\mathbf{k}}(-\mathbf{G}) \mathcal{C}(\mathbf{G}_{\mathbf{k}\mathbf{k}'}) [\rho_{\nu j}^{\mathbf{k}'\mathbf{k}}(-\mathbf{G})]^*. \quad (31)$$

By changing the contraction order in Eq. (27), we obtain

$$\rho_{\mu j}^{\mathbf{k}\mathbf{k}'}(\mathbf{G}) = \sum_{\nu\mathbf{n}} \rho_{\mu^0\nu\mathbf{n}}(\mathbf{G}_{\mathbf{k}\mathbf{k}'}) C_{\nu\mathbf{n}j\mathbf{k}'}, \quad (32)$$

$$C_{\nu\mathbf{n}j\mathbf{k}} = e^{i\mathbf{k}\cdot\mathbf{n}} C_{\nu j}^{\mathbf{k}}. \quad (33)$$

It is noticeable that the sparsity in the FT orbital product tensor can be exploited to reduce the computational cost. The FT integral $\rho_{\mu^m\nu\mathbf{n}}(\mathbf{G}_{\mathbf{k}\mathbf{k}'})$ in Eq. (16) may be negligible for well-separated $\mu(\mathbf{r}-\mathbf{m})$ and $\nu(\mathbf{r}-\mathbf{n})$. Although the tensor contraction still has the scaling $N_k^2 N_c N_{AO}^2 N_G$, the actual cost is much lower due to the sparsity of $\rho_{\mu^0\nu\mathbf{n}}(\mathbf{G}_{\mathbf{k}\mathbf{k}'})$.

For SCF orbitals which preserve the time-reversal symmetry between \mathbf{k} and $\bar{\mathbf{k}} (= -\mathbf{k})$, it is possible to achieve an additional reduction in computational cost. The time-reversal symmetry ensures that the SCF wavefunction defined by the \mathbf{k} -point orbitals can be transformed to an identical

representation at certain Γ -point with real orbital coefficients \mathbf{B}

$$B_{\mu^m j^k} = \frac{1}{\sqrt{2}}(C_{\mu^m j^k} + C_{\mu^m j^{\bar{k}}}), \quad (34)$$

$$B_{\mu^m j^{\bar{k}}} = \frac{1}{\sqrt{2}}(C_{\mu^m j^k} - C_{\mu^m j^{\bar{k}}}). \quad (35)$$

This property reduces the cost of computing $\rho_{\mu_j}^{\mathbf{k}\mathbf{k}'}(\mathbf{G})$ by roughly a factor of two as we only need to compute the contraction between a complex tensor and a real matrix

$$L_{\mu_j}^{\mathbf{k}}(\mathbf{G}) = \sum_{\mathbf{v}\mathbf{n}} \rho_{\mu^0 \mathbf{v}\mathbf{n}}(\mathbf{G}) B_{\mathbf{v}\mathbf{n} j^k}, \quad (36)$$

$$\rho_{\mu_j}^{\mathbf{k}\mathbf{k}'}(\mathbf{G}) = \frac{1}{\sqrt{2}} L_{\mu_j}^{\mathbf{k}'}(\mathbf{G}_{\mathbf{k}\mathbf{k}'}) - \frac{i}{\sqrt{2}} L_{\mu_j}^{\bar{\mathbf{k}}'}(\mathbf{G}_{\mathbf{k}\mathbf{k}'}). \quad (37)$$

In addition, the intermediates $L_{\mu_j}^{\mathbf{k}}$ can be reused for the tensor contraction in Eq. (30), providing another factor of two reduction in computational cost. We first employ the FT Eq. (14)

$$\rho_{\mathbf{v}j}^{\mathbf{k}'\mathbf{k}}(-\mathbf{G}) = \sum_{\mu\mathbf{m}} [\rho_{\mu^m \mathbf{v}^0}(\mathbf{G}_{\mathbf{k}\mathbf{k}'}) e^{-i\mathbf{k}\cdot\mathbf{m}}]^* C_{\mu_j}^{\mathbf{k}} = \sum_{\mu\mathbf{m}} [\rho_{\mathbf{v}^0 \mu^m}(\mathbf{G}_{\mathbf{k}\mathbf{k}'}) C_{\mu^m j^k}^*]^*. \quad (38)$$

Realizing that the orbital coefficients \mathbf{B} are real,

$$C_{\mu^m j^k}^* = \frac{1}{\sqrt{2}}(B_{\mu^m j^k} + iB_{\mu^m j^{\bar{k}}}) \quad (39)$$

we then simplify Eq. (30) to

$$\rho_{\mathbf{v}j}^{\mathbf{k}'\mathbf{k}}(-\mathbf{G}) = \frac{1}{\sqrt{2}} [L_{\mathbf{v}j}^{\mathbf{k}}(\mathbf{G}_{\mathbf{k}\mathbf{k}'})]^* - \frac{i}{\sqrt{2}} [L_{\mathbf{v}j}^{\bar{\mathbf{k}}}(\mathbf{G}_{\mathbf{k}\mathbf{k}'})]^*. \quad (40)$$

By using all the features discussed above, we would obtain an overall reduction which roughly equals a factor of 8 plus speed-ups from tensor sparsity. Finally, we derive the exchange matrices

$$K_{\mu\nu}^{\mathbf{k}} = \frac{1}{2} \sum_{\mathbf{k}'} (\tilde{K}_{\mu\nu}^{\mathbf{k}'\mathbf{k}'} + i\tilde{K}_{\mu\nu}^{\mathbf{k}'\bar{\mathbf{k}}'} - i\tilde{K}_{\mu\nu}^{\bar{\mathbf{k}}'\mathbf{k}'} + \tilde{K}_{\mu\nu}^{\bar{\mathbf{k}}'\bar{\mathbf{k}}'}), \quad (41)$$

$$K_{\mu\nu}^{\mathbf{k}'} = \frac{1}{2} \sum_{\mathbf{k}} (\tilde{K}_{\nu\mu}^{\mathbf{k}\mathbf{k}} + i\tilde{K}_{\nu\mu}^{\mathbf{k}\bar{\mathbf{k}}} - i\tilde{K}_{\nu\mu}^{\bar{\mathbf{k}}\mathbf{k}} + \tilde{K}_{\nu\mu}^{\bar{\mathbf{k}}\bar{\mathbf{k}}}), \quad (42)$$

where \tilde{K} is the intermediate based on $L_{\mu_j}^{\mathbf{k}}$

$$\tilde{K}_{\mu\nu}^{\mathbf{k}_1\mathbf{k}_2} = \frac{1}{N_k} \frac{1}{\Omega} \sum_{j\mathbf{G}} \Lambda_i^{\mathbf{k}_1} L_{\mu_j}^{\mathbf{k}_1}(\mathbf{G}_{\mathbf{k}\mathbf{k}'}) \mathcal{C}(\mathbf{G}_{\mathbf{k}\mathbf{k}'})[L_{\mathbf{v}j}^{\mathbf{k}_2}(\mathbf{G}_{\mathbf{k}\mathbf{k}'})]^*. \quad (43)$$

For sufficient \mathbf{k} points, this intermediate can approximately lower 25% of the cost due to the relation

$$\tilde{K}_{\mu\nu}^{\bar{\mathbf{k}}\mathbf{k}} = [\tilde{K}_{\nu\mu}^{\mathbf{k}\bar{\mathbf{k}}}]^*. \quad (44)$$

C. SR integrals evaluation

Following to the idea proposed in 55, we first evaluate SR Coulomb integrals as well as the Coulomb and exchange matrix in the Born-von Karman (BvK) supercell then transform them to the \mathbf{k} -point adapted quantities. For example, the matrix element of SR HFX $K_{\mu\nu}^{\text{SR},\mathbf{k}}$ can be computed

$$K_{\mu^0\lambda^1}^{\text{SR}} = \sum_{\mathbf{v}\mathbf{K}} \sum_{\mathbf{L}} \sum_{\mathbf{m}+\mathbf{M}\mathbf{n}+\mathbf{N}} \sum D_{\mathbf{v}^{\mathbf{n}}\mathbf{K}^{\mathbf{m}}} (\mu^0 \mathbf{v}^{\mathbf{n}+\mathbf{N}} | \mathbf{K}^{\mathbf{m}+\mathbf{M}} \lambda^{1+\mathbf{L}+\mathbf{M}})^{\text{SR}}, \quad (45)$$

$$K_{\mu\nu}^{\text{SR},\mathbf{k}} = \sum_{\mathbf{l}} e^{i\mathbf{k}\cdot\mathbf{l}} K_{\mu^0\nu^1}^{\text{SR}}. \quad (46)$$

Here the lattice summation involves two types of translation vectors, the super-lattice translation vectors $(\mathbf{L}, \mathbf{M}, \mathbf{N})$ for BvK supercell and the translation inside a single BvK supercell $(\mathbf{l}, \mathbf{m}, \mathbf{n})$. $D_{\mathbf{v}^{\mathbf{n}}\mathbf{K}^{\mathbf{m}}}$ in Eq. (45) represents the density matrix elements in the BvK supercell. Considering the BvK supercell being filled with N_{BvK} primitive cells, $D_{\mathbf{v}^{\mathbf{l}}\mathbf{K}^{\mathbf{m}}}$ is obtained via an inverse FT of the \mathbf{k} -point adapted density matrices

$$D_{\mathbf{v}^{\mathbf{n}}\mathbf{K}^{\mathbf{m}}} = D_{\mathbf{v}^0\mathbf{K}^{\mathbf{m}-\mathbf{n}}} = \frac{1}{N_{\text{BvK}}} \sum_{\mathbf{k}} e^{-i\mathbf{k}\cdot(\mathbf{m}-\mathbf{n})} D_{\mathbf{v}\mathbf{K}}^{\mathbf{k}}. \quad (47)$$

The cost of computing SR integrals in real space mainly depends on the range of lattice-sum and integral screening of the SR integrals. The number of \mathbf{k} points has relatively low impact on the computational cost. The value of Coulomb attenuation parameter ω and the compactness of Gaussian basis functions are the two main factors to determine the distance cutoff (as well as the range of lattice-sum)⁶². Small value of ω and diffused basis functions can lead to more image cells in lattice-sum. Although the size of BvK super-cell is proportional to the size of \mathbf{k} points, the overall number of image cells in the lattice-sum, denoted by the compound index $\mathbf{m} + \mathbf{M}$, remains largely unchanged.

To further reduce the cost of SR integrals, we also take the locality of basis functions into account. The Gaussian basis functions can be partitioned into two groups, labelled as D for the diffused functions and C for the compact functions. The products of two functions $\rho_{\mu\nu}$ have four possible combinations: the product of two diffused functions can lead to a diffuse density ρ_D while the rest three DC , CD and CC all lead to compact densities ρ_C . Here we developed three recipes to evaluate these SR integrals. Each recipe is adequate in specific scenarios, depending on the size of \mathbf{k} -point mesh.

- Recipe I: Analytically evaluating the 9 types of SR integrals which only consist of compact densities $(DC|DC)$, $(DC|CD)$, $(DC|CC)$, $(CD|DC)$, $(CD|CD)$, $(CD|CC)$, $(CC|DC)$, $(CC|CD)$, $(CC|CC)$ in real space while leaving the remaining 7 types of SR integrals to the reciprocal space prescription we presented in Section II B. Since the 9 types of analytical integrals involve only compact density distributions, lattice-sum can be rapidly truncated.

For the integrals involving diffuse densities, for instance the $(CC|DD)$ -type integrals, the computation of SR integrals and their LR counterparts can be merged into one pass for $(CC|DD)$ with the full-range Coulomb kernel since both the SR and LR integrals are evaluated in reciprocal space. It is important to choose an appropriate energy cutoff E_{cut} to ensure the accuracy of the $(CC|DD)$ -type full-range ERIs. Given the precision requirement ϵ , the energy cutoff for four-center ERIs can be estimated⁶²

$$16\pi^2\theta E_{cut}^{\frac{1}{2}}\left(\frac{E_{cut}}{2\alpha_C}\right)^{2l_C}\left(\frac{E_{cut}}{2\alpha_D}\right)^{2l_D}e^{\frac{E_{cut}}{2\theta}} < \epsilon, \quad (48)$$

$$\theta = \frac{1}{2}(\alpha_C^{-1} + \alpha_D^{-1})^{-1}, \quad (49)$$

where α_C and l_C are the exponent and the angular moment of the most compact basis function. We can solve the inequality (48) for each diffused function and find an overall energy cutoff for the reciprocal space prescription.

When computing exchange matrix with this recipe, it is inconvenient to merely compute the 7 types of diffuse-density associated SR integrals with the reciprocal space prescription. We instead use the reciprocal space prescription to compute the HFX of all integrals as well as the SR HFX of the 9 types of compact SR integrals. Then, we take the difference between the two results to obtain the contribution of the 7-type SR integrals.

This recipe lowers the cost of analytical integral evaluation while it introduces an extra HFX computation in reciprocal space. The cost for the later one is proportional to N_k^2 and it can dominate the HFX computation if the SCF calculation involves many \mathbf{k} points. Therefore, this recipe is only adequate for SCF calculations with a small number of \mathbf{k} points.

- Recipe II: Computing the $(DD|DD)$ -type SR integral with the reciprocal space prescription while handling the rest 15 types in real space. $(DD|DD)$ -type SR integrals in real space often involves a large number of image cells in the triple lattice-sum as it requires the largest distance cutoff. Excluding it from the real space integral evaluation can save computational

cost. Meanwhile, evaluating $(DD|DD)$ -type SR integrals in reciprocal space only adds a small overhead, as the diffused functions typically constitute a small portion of the regular basis set. This algorithm is helpful in reducing the computation time of Fock build for a moderate number of \mathbf{k} points.

- Recipe III: Evaluating all SR integrals in real space. When a large number of \mathbf{k} points is present in the SCF calculation, the effort of computing exchange matrix in reciprocal space become dominant. To reduce the cost of reciprocal space part, we not only evaluate all SR integrals in real space, but also tend to use small energy cutoff to decrease the required number of PWs which is proportional to the computational cost of reciprocal space algorithm.

The value of Coulomb attenuation parameter ω , energy cutoff, and the number of PWs are correlated. Although less PWs means less computational efforts in reciprocal space part, a lower energy cutoff can lead to more computation time in real space as it requires a small ω to ensure the integral accuracy. To gain the optimal performance, different ω should be used in different \mathbf{k} -point calculations.

D. Integral screening

Integral screening is a crucial factor for the performance of real space integral program. To enhance the performance of SR integral computation, we have developed a three-layer hierarchical screening prescription:

- First, we filter integrals with the regular Schwarz inequality

$$(\mu\nu|\kappa\lambda)^{\text{SR}} < \sqrt{(\mu\nu|\mu\nu)^{\text{SR}}(\kappa\lambda|\kappa\lambda)^{\text{SR}}}. \quad (50)$$

- Next, we apply the mutated Schwarz inequality to estimate SR integral value

$$(\mu\nu|\kappa\lambda)^{\text{SR}} < \min(\sqrt{(\mu\mu|\kappa\kappa)^{\text{SR}}(\nu\nu|\lambda\lambda)^{\text{SR}}}, \sqrt{(\mu\mu|\lambda\lambda)^{\text{SR}}(\nu\nu|\kappa\kappa)^{\text{SR}}}). \quad (51)$$

This condition is not useful for ERIs of full Coulomb metric due to the long range character of Coulomb interactions. However, for SR integrals, this condition can filter out approximately 50% of the terms.

- Finally, the SR integral estimator developed in 62 is utilized as the last integral filter

$$(\mu\nu|\kappa\lambda)^{\text{SR}} \lesssim \frac{Q_{\mu\nu}Q_{\kappa\lambda}e^{-\theta_{\mu\nu\kappa\lambda}R^2}}{R^2} \quad (52)$$

where $Q_{\mu\nu}$ is a factor that behaves like the overlap between two primitive Gaussian functions. Its detailed definition can be found in 62. Other quantities can be computed

$$\theta_{\mu\nu\kappa\lambda} = ((\alpha_\mu + \alpha_\nu)^{-1} + (\alpha_\kappa + \alpha_\lambda)^{-1} + \omega^{-2})^{-1}, \quad (53)$$

$$R = |\mathbf{P}_{\mu\nu} - \mathbf{P}_{\kappa\lambda}|, \quad (54)$$

$$\mathbf{P}_{\mu\nu} = \frac{\alpha_\mu \mathbf{R}_\mu + \alpha_\nu \mathbf{R}_\nu}{\alpha_\mu + \alpha_\nu}. \quad (55)$$

The reason that the screening filters are applied in this specific order is that each screening filter has a different computational overhead. We have observed that the time spent on the screening testing is not negligible due to the large number of integral quartets that needs to be tested. The regular Schwarz inequality is applied first, as it is not only effective in filtering out most small integrals but also the fastest. The regular Schwarz inequality test requires only one multiplication and two memory accesses. The mutated Schwarz inequality is slightly less efficient as it requires two multiplications and four memory accesses. The integral estimator works slowest. Even though some intermediates are precomputed and cached, it requires six memory accesses to obtain coordinates for R , two memory accesses for paired quantities Q , one call to the exponential function, and ten or more multiplication/addition/division operations to aggregate them.

III. NUMERICAL TESTS

The range-separated Fock build algorithm has been implemented in the PySCF program package²⁵. In all exhibited calculations, the precision requirement for energy cutoff and distance cutoff was set to 10^{-8} . Timing is measured on a desktop computer with Intel I9-7940X 14-core CPU and 64 GB memory.

Our first tests are all-electron k-point Hartree-Fock calculations for diamond crystal. Gaussian basis sets STO-3G and cc-pVDZ are employed in our tests. System size ranges from $1 \times 1 \times 1$ \mathbf{k} point (Γ point) to $10 \times 10 \times 10$ \mathbf{k} points. The results are collected in Table I and II. Using the range-separated algorithm introduced here, we can accomplish the calculation of cc-pVDZ basis with $4 \times 4 \times 4$ \mathbf{k} -point HF in 20 CPU hours. Even for a dense $10 \times 10 \times 10$ \mathbf{k} -mesh, which consists

of 8000 atoms and 112000 basis functions, the SCF calculation takes only 254 CPU hours with a memory footprint less than 6 GB.

The number of PWs and the corresponding range-separation parameter ω in our \mathbf{k} -point HF calculations are also listed in Table I and II. For the small \mathbf{k} points, it is often less demanding to compute LR HFX in reciprocal space. We tend to use more PWs to increase the value of ω for SR HFX because large ω can help SR Coulomb potential rapidly decay in real space. In this scenario, the associated energy cutoff is high enough for the full-range Coulomb integrals with some diffused basis. We then use this criteria to partition basis functions and employ the SR integral recipe I and recipe II to reduce the cost of the real-space SR integral evaluation. As mentioned in Section II C, the reciprocal space HFX algorithm needs to be executed twice when applying recipe I. This overhead is noticeable in the $4 \times 4 \times 4$ \mathbf{k} -point calculation with cc-pVDZ basis. With the number of PWs chosen in Table II, the ratio of theoretical scaling between the LR part of the $4 \times 4 \times 4$ mesh and the $3 \times 3 \times 3$ mesh is $64^2 \times 13^3 / (27^2 \times 17^3) \approx 2.5$. By using recipe II for SR HFX, the actual resources spent on the LR part of the $4 \times 4 \times 4$ mesh is 6.5 CPU hours, which is only 70% more than the resources for the $3 \times 3 \times 3$ mesh.

The computation cost of HFX in the reciprocal space part increases rapidly as we increase the number of \mathbf{k} points, while the real space SR HFX has a relatively slower increase in cost. Therefore, we prefer SR HFX recipe III with fewer PWs for calculations with a large number of \mathbf{k} points. The sparsity in orbital products exhibits effectiveness for large \mathbf{k} -mesh calculations. For example, the average time of Fock build spent on the LR HFX of $10 \times 10 \times 10$ \mathbf{k} -point calculation is 13 CPU hours, which is 25% more than that of the $8 \times 8 \times 8$ case, although the theoretical scaling indicates 40% more FLOPs for the former.

The help of sparsity is more effective for calculations with denser \mathbf{k} -point mesh. This enables us to perform all-electron Hartree-Fock calculations for LiH crystal up to $24 \times 24 \times 24 = 13824$ \mathbf{k} -points, which amounts to a total of 27648 atoms and 1036800 basis functions. The largest \mathbf{k} -point HF calculation for LiH crystal takes 1380 CPU hours in which 1323 CPU hours are spent on the LR HFX computation. In this calculation, we use a small value of $\omega = 0.14$ for SR HFX which results in 27 PWs for LR HFX.

Large scale calculations towards thermodynamic limit of HF methods with Gaussian basis for LiH crystal were studied in 63. In that work, two methods were employed to obtain the HF energy: an extrapolation from SR HFX to full-range HFX with Padé approximants, and a direct evaluation of HFX with truncated Coulomb interactions. Our \mathbf{k} -point HF results are listed in Table

TABLE I. Results of all-electron \mathbf{k} -point HF with STO-3G basis for diamond crystal. The unit cell is a cubic cell with lattice parameter 3.5668 Å. HF energy is given in Hartree atomic units.

\mathbf{k} mesh	E_{HF} per cell	timing/h	Fock builds	ω	N_G	SR-ERI recipe	SR-ERI timing
$1 \times 1 \times 1$	-299.328101	0.14	9	1.15	23^3	I	0.04
$2 \times 2 \times 2$	-299.551274	0.64	8	0.94	19^3	I	0.23
$3 \times 3 \times 3$	-299.525890	0.72	8	0.73	15^3	I	0.30
$4 \times 4 \times 4$	-299.516150	1.09	8	0.52	11^3	I	0.41
$6 \times 6 \times 6$	-299.510660	5.48	9	0.42	9^3	III	2.01
$8 \times 8 \times 8$	-299.509296	13.9	9	0.31	7^3	III	3.70
$10 \times 10 \times 10$	-299.508809	22.7	9	0.21	5^3	III	8.58

III. As we increase the \mathbf{k} -point mesh, the obtained HF energy per unit cell gradually converges towards $-8.064539 E_h$, which is $4 \mu E_h$ higher than the extrapolation result or $6 \mu E_h$ higher than the direct evaluation result reported in 63. Based on the trends we observed in our calculation series, we can expect that the HF thermodynamic limit is probably several μE_h higher than the result we obtained. The difference may be attributed to two possibilities. The first is the treatment of Coulomb singularity in HFX. We utilize the full-range Coulomb interactions with Ewald probe charge correction in this work. In contrast, the extrapolation method does not have any treatment for singularity in \mathbf{k} sampling while the result of direct evaluation method is based on manually tuned truncated Coulomb potentials. The second possibility is the size of \mathbf{k} -point mesh utilized in the calculation. The calculation of direct evaluation method includes only 1000 atoms in a $5 \times 5 \times 5$ mesh of the cubic cell structure which is smaller than the system size in our tests.

IV. CONCLUSIONS

In this work, we presented an efficient range-separated algorithm to compute electron repulsion integrals and Fock matrix for crystalline Gaussian basis. In this algorithm, the full-range Coulomb interactions were split into the short-range and long-range parts, which are respectively computed in real and reciprocal space. Additionally, we proposed several methods to improve the computational efficiency. For integrals evaluated in reciprocal space, we reformulated the Fourier transform and the tensor contraction equations to utilize sparsity in Gaussian orbital products. For

TABLE II. Results of all-electron \mathbf{k} -point HF with cc-pVDZ basis for diamond crystal. The unit cell is a cubic cell with lattice parameter 3.5668 Å. HF energy is given in Hartree atomic units.

\mathbf{k} mesh	E_{HF} per cell	timing/h	Fock builds	ω	N_G	SR-ERI recipe	SR-ERI timing
$1 \times 1 \times 1$	-302.870240	0.58	9	1.15	23^3	I	0.30
$2 \times 2 \times 2$	-303.076048	2.71	8	1.04	21^3	I	0.62
$3 \times 3 \times 3$	-303.054159	8.23	9	0.83	17^3	I	4.41
$4 \times 4 \times 4$	-303.044553	19.7	9	0.63	13^3	II	13.2
$6 \times 6 \times 6$	-303.038978	54.3	9	0.42	9^3	II	25.8
$8 \times 8 \times 8$	-303.037585	134.7	9	0.31	7^3	III	40.7
$10 \times 10 \times 10$	-303.037088	254.5	11	0.21	5^3	III	110.0

TABLE III. \mathbf{k} -point HF tests for LiH crystal at experimental lattice constant 4.084Å.

\mathbf{k} mesh	E_{HF} per cell
$1 \times 1 \times 1$	-8.449722
$2 \times 2 \times 2$	-8.063001
$3 \times 3 \times 3$	-8.072053
$4 \times 4 \times 4$	-8.066987
$5 \times 5 \times 5$	-8.065904
$6 \times 6 \times 6$	-8.065308
$7 \times 7 \times 7$	-8.065022
$8 \times 8 \times 8$	-8.064859
$9 \times 9 \times 9$	-8.064760
$11 \times 11 \times 11$	-8.064654
$13 \times 13 \times 13$	-8.064603
$15 \times 15 \times 15$	-8.064577
$19 \times 19 \times 19$	-8.064551
$24 \times 24 \times 24$	-8.064539

short-range integrals, we partitioned basis functions according to their locality in real space and developed three recipes to utilize the localities of basis functions. Different recipes are adequate for different scenarios depending on the number of \mathbf{k} points. We developed a three-level integral screening scheme to filter SR electron repulsion integrals which has significant impacts on the computational efficiency for short-range integrals. This algorithm enables us to perform all-electron HF and hybrid functional DFT calculations with a large number of \mathbf{k} points. To assess the performance of this algorithm, we performed \mathbf{k} -point HF calculations for LiH crystal with up to 1m basis functions. The current algorithm only supports the Ewald probe-charge correction in the treatment of Coulomb singularity. Combining this algorithm with other Coulomb singularity treatments will be considered in the future.

REFERENCES

- ¹D. R. Bowler and T. Miyazaki, “Calculations for millions of atoms with density functional theory: linear scaling shows its potential,” *J. Phys.: Condens. Matter* **22**, 074207 (2010).
- ²J. VandeVondele, U. Borštnik, and J. Hutter, “Linear scaling self-consistent field calculations with millions of atoms in the condensed phase,” *J. Chem. Theory Comput.* **8**, 3565–3573 (2012), PMID: 26593003, <https://doi.org/10.1021/ct200897x>.
- ³C. Pisani and R. Dovesi, “Exact-exchange hartree-fock calculations for periodic systems. i. illustration of the method,” *Int. J. Quantum Chem.* **17**, 501–516 (1980), <https://onlinelibrary.wiley.com/doi/pdf/10.1002/qua.560170311>.
- ⁴R. Dovesi, C. Pisani, C. Roetti, and V. R. Saunders, “Treatment of coulomb interactions in hartree-fock calculations of periodic systems,” *Phys. Rev. B* **28**, 5781–5792 (1983).
- ⁵F. Gygi and A. Baldereschi, “Self-consistent hartree-fock and screened-exchange calculations in solids: Application to silicon,” *Phys. Rev. B* **34**, 4405–4408 (1986).
- ⁶J. Heyd and G. E. Scuseria, “Efficient hybrid density functional calculations in solids: Assessment of the heyd-scuseria-ernzerhof screened coulomb hybrid functional,” *J. Chem. Phys.* **121**, 1187–1192 (2004), <https://doi.org/10.1063/1.1760074>.
- ⁷J. Paier, R. Hirschl, M. Marsman, and G. Kresse, “The Perdew–Burke–Ernzerhof exchange-correlation functional applied to the G2-1 test set using a plane-wave basis set,” *The Journal of Chemical Physics* **122** (2005), 10.1063/1.1926272, 234102, https://pubs.aip.org/aip/jcp/article-pdf/doi/10.1063/1.1926272/13133732/234102_1_online.pdf.
- ⁸A. F. Izmaylov, G. E. Scuseria, and M. J. Frisch, “Efficient evaluation of short-range hartree-fock exchange in large molecules and periodic systems,” *J. Chem. Phys.* **125**, 104103 (2006), <https://doi.org/10.1063/1.2347713>.
- ⁹J. Spencer and A. Alavi, “Efficient calculation of the exact exchange energy in periodic systems using a truncated coulomb potential,” *Phys. Rev. B* **77**, 193110 (2008).
- ¹⁰M. Guidon, J. Hutter, and J. VandeVondele, “Robust periodic hartree-fock exchange for large-scale simulations using gaussian basis sets,” *J. Chem. Theory Comput.* **5**, 3010–3021 (2009), PMID: 26609981, <https://doi.org/10.1021/ct900494g>.
- ¹¹R. Sundararaman and T. A. Arias, “Regularization of the coulomb singularity in exact exchange by wigner-seitz truncated interactions: Towards chemical accuracy in nontrivial sys-

- tems,” *Phys. Rev. B* **87**, 165122 (2013).
- ¹²N. Marzari and D. Vanderbilt, “Maximally localized generalized wannier functions for composite energy bands,” *Phys. Rev. B* **56**, 12847–12865 (1997).
- ¹³F. Gygi, “Compact representations of kohn-sham invariant subspaces,” *Phys. Rev. Lett.* **102**, 166406 (2009).
- ¹⁴N. Marzari, A. A. Mostofi, J. R. Yates, I. Souza, and D. Vanderbilt, “Maximally localized wannier functions: Theory and applications,” *Rev. Mod. Phys.* **84**, 1419–1475 (2012).
- ¹⁵F. Gygi and I. Duchemin, “Efficient computation of hartree-fock exchange using recursive subspace bisection,” *J. Chem. Theory Comput.* **9**, 582–587 (2013).
- ¹⁶R. A. DiStasio, B. Santra, Z. Li, X. Wu, and R. Car, “The individual and collective effects of exact exchange and dispersion interactions on the ab initio structure of liquid water,” *J. Chem. Phys.* **141**, 084502 (2014).
- ¹⁷A. Görling, “Exact treatment of exchange in kohn-sham band-structure schemes,” *Phys. Rev. B* **53**, 7024–7029 (1996).
- ¹⁸S. Chawla and G. A. Voth, “Exact exchange in ab initio molecular dynamics: An efficient plane-wave based algorithm,” *J. Chem. Phys.* **108**, 4697–4700 (1998).
- ¹⁹X. Wu, A. Selloni, and R. Car, “Order- n implementation of exact exchange in extended insulating systems,” *Phys. Rev. B* **79**, 085102 (2009).
- ²⁰L. Lin, “Adaptively compressed exchange operator,” *J. Chem. Theory Comput.* **12**, 2242–2249 (2016).
- ²¹W. Hu, L. Lin, A. S. Banerjee, E. Vecharynski, and C. Yang, “Adaptively compressed exchange operator for large-scale hybrid density functional calculations with applications to the adsorption of water on silicene,” *J. Chem. Theory Comput.* **13**, 1188–1198 (2017), pMID: 28177229, <https://doi.org/10.1021/acs.jctc.6b01184>.
- ²²T. A. Barnes, T. Kurth, P. Carrier, N. Wichmann, D. Prendergast, P. R. C. Kent, and J. Deslippe, “Improved treatment of exact exchange in quantum espresso,” *Computer Physics Communications* **214**, 52–58 (2017).
- ²³T. D. Kühne, M. Iannuzzi, M. Del Ben, V. V. Rybkin, P. Seewald, F. Stein, T. Laino, R. Z. Khaliullin, O. Schütt, F. Schiffmann, D. Golze, J. Wilhelm, S. Chulkov, M. H. Bani-Hashemian, V. Weber, U. Borštnik, M. TAILLEFUMIER, A. S. Jakobovits, A. Lazzaro, H. Pabst, T. Müller, R. Schade, M. Guidon, S. Andermatt, N. Holmberg, G. K. Schenter, A. Hehn, A. Bussy, F. Belleflamme, G. Tabacchi, A. Glöß, M. Lass, I. Bethune, C. J. Mundy, C. Plessl, M. Watkins, J. VandeVondele, M. Krack, and J. Hutter, “Cp2k: An electronic structure and molecular dy-

- namics software package - quickstep: Efficient and accurate electronic structure calculations,” *J. Chem. Phys.* **152**, 194103 (2020), <https://doi.org/10.1063/5.0007045>.
- ²⁴R. Dovesi, F. Pascale, B. Civalleri, K. Doll, N. M. Harrison, I. Bush, P. D’Arco, Y. Noël, M. Rérat, P. Carbonnière, M. Causà, S. Salustro, V. Lacivita, B. Kirtman, A. M. Ferrari, F. S. Gentile, J. Baima, M. Ferrero, R. Demichelis, and M. De La Pierre, “The crystal code, 1976-2020 and beyond, a long story,” *J. Chem. Phys.* **152**, 204111 (2020).
- ²⁵Q. Sun, X. Zhang, S. Banerjee, P. Bao, M. Barbry, N. S. Blunt, N. A. Bogdanov, G. H. Booth, J. Chen, Z.-H. Cui, J. J. Eriksen, Y. Gao, S. Guo, J. Hermann, M. R. Hermes, K. Koh, P. Koval, S. Lehtola, Z. Li, J. Liu, N. Mardirossian, J. D. McClain, M. Motta, B. Mussard, H. Q. Pham, A. Pulkin, W. Purwanto, P. J. Robinson, E. Ronca, E. R. Sayfutyarova, M. Scheurer, H. F. Schurkus, J. E. T. Smith, C. Sun, S.-N. Sun, S. Upadhyay, L. K. Wagner, X. Wang, A. White, J. D. Whitfield, M. J. Williamson, S. Wouters, J. Yang, J. M. Yu, T. Zhu, T. C. Berkelbach, S. Sharma, A. Y. Sokolov, and G. K.-L. Chan, “Recent developments in the pyscf program package,” *J. Chem. Phys.* **153**, 024109 (2020), <https://doi.org/10.1063/5.0006074>.
- ²⁶V. Blum, R. Gehrke, F. Hanke, P. Havu, V. Havu, X. Ren, K. Reuter, and M. Scheffler, “Ab initio molecular simulations with numeric atom-centered orbitals,” *Computer Physics Communications* **180**, 2175–2196 (2009).
- ²⁷S. G. Balasubramani, G. P. Chen, S. Coriani, M. Diedenhofen, M. S. Frank, Y. J. Franzke, F. Furche, R. Grotjahn, M. E. Harding, C. Hättig, A. Hellweg, B. Helmich-Paris, C. Holzer, U. Huniar, M. Kaupp, A. Marefat Khah, S. Karbalaei Khani, T. Müller, F. Mack, B. D. Nguyen, S. M. Parker, E. Perlt, D. Rappoport, K. Reiter, S. Roy, M. Rückert, G. Schmitz, M. Sierka, E. Tapavicza, D. P. Tew, C. van Wüllen, V. K. Voora, F. Weigend, A. Wodyński, and J. M. Yu, “Turbomole: Modular program suite for ab initio quantum-chemical and condensed-matter simulations,” *J. Chem. Phys.* **152**, 184107 (2020).
- ²⁸J. VandeVondele, M. Krack, F. Mohamed, M. Parrinello, T. Chassaing, and J. Hutter, “Quickstep: Fast and accurate density functional calculations using a mixed gaussian and plane waves approach,” *Computer Physics Communications* **167**, 103–128 (2005).
- ²⁹B. G. E. R. A. L. D. LIPPERT, J. U. R. G. H. U. T. T. E. R. PARRINELLO, and MICHELE, “A hybrid gaussian and plane wave density functional scheme,” *Molecular Physics* **92**, 477–488 (1997).
- ³⁰J. H. Dominik Marx, “Ab initio molecular dynamics: Basic theory and advanced methods,” *Angew. Chem. Int. Ed.* **48**, 9404–9405 (2009).

- ³¹C. Hartwigsen, S. Goedecker, and J. Hutter, “Relativistic separable dual-space gaussian pseudopotentials from h to rn,” *Phys. Rev. B* **58**, 3641–3662 (1998).
- ³²A. Willand, Y. O. Kvashnin, L. Genovese, A. Vázquez-Mayagoitia, A. K. Deb, A. Sadeghi, T. Deutsch, and S. Goedecker, “Norm-conserving pseudopotentials with chemical accuracy compared to all-electron calculations,” *J. Chem. Phys.* **138**, 104109 (2013).
- ³³W. Gao and J. R. Chelikowsky, “Accuracy of partial core corrections using fourier transforms in pseudopotential-density functional theory,” *J. Chem. Theory Comput.* **14**, 6515–6520 (2018).
- ³⁴G. Prandini, A. Marrazzo, I. E. Castelli, N. Mounet, and N. Marzari, “Precision and efficiency in solid-state pseudopotential calculations,” *npj Comput. Mater.* **4**, 72 (2018).
- ³⁵J.-B. Lu, D. C. Cantu, M.-T. Nguyen, J. Li, V.-A. Glezakou, and R. Rousseau, “Norm-conserving pseudopotentials and basis sets to explore lanthanide chemistry in complex environments,” *J. Chem. Theory Comput.* **15**, 5987–5997 (2019).
- ³⁶K. Lejaeghere, G. Bihlmayer, T. Björkman, P. Blaha, S. Blügel, V. Blum, D. Caliste, I. E. Castelli, S. J. Clark, A. Dal Corso, S. de Gironcoli, T. Deutsch, J. K. Dewhurst, I. Di Marco, C. Draxl, M. Dułak, O. Eriksson, J. A. Flores-Livas, K. F. Garrity, L. Genovese, P. Gianozzi, M. Giantomassi, S. Goedecker, X. Gonze, O. Grånäs, E. K. U. Gross, A. Gulans, F. Gygi, D. R. Hamann, P. J. Hasnip, N. A. W. Holzwarth, D. Iușan, D. B. Jochym, F. Jollet, D. Jones, G. Kresse, K. Koepnik, E. Küçükbenli, Y. O. Kvashnin, I. L. M. Locht, S. Lubeck, M. Marsman, N. Marzari, U. Nitzsche, L. Nordström, T. Ozaki, L. Paulatto, C. J. Pickard, W. Poelmans, M. I. J. Probert, K. Refson, M. Richter, G.-M. Rignanese, S. Saha, M. Scheffler, M. Schlipf, K. Schwarz, S. Sharma, F. Tavazza, P. Thunström, A. Tkatchenko, M. Torrent, D. Vanderbilt, M. J. van Setten, V. Van Speybroeck, J. M. Wills, J. R. Yates, G.-X. Zhang, and S. Cottenier, “Reproducibility in density functional theory calculations of solids,” *Science* **351**, aad3000 (2016).
- ³⁷M. Krack and M. Parrinello, “All-electron ab-initio molecular dynamics,” *Phys. Chem. Chem. Phys.* **2**, 2105–2112 (2000).
- ³⁸E. Kucukbenli, M. Monni, B. I. Adetunji, X. Ge, G. A. Adebayo, N. Marzari, S. de Gironcoli, and A. D. Corso, “Projector augmented-wave and all-electron calculations across the periodic table: a comparison of structural and energetic properties,” (2014), arXiv:1404.3015 [cond-mat.mtrl-sci].
- ³⁹S. V. Levchenko, X. Ren, J. Wieferink, R. Johanni, P. Rinke, V. Blum, and M. Scheffler, “Hybrid functionals for large periodic systems in an all-electron, numeric atom-centered basis frame-

- work,” *Computer Physics Communications* **192**, 60–69 (2015).
- ⁴⁰T. Biktairov, W. G. Schmidt, and U. Gerstmann, “Calculation of spin-spin zero-field splitting within periodic boundary conditions: Towards all-electron accuracy,” *Phys. Rev. B* **97**, 115135 (2018).
- ⁴¹C.-N. Yeh, A. Shee, Q. Sun, E. Gull, and D. Zgid, “Relativistic self-consistent gw: Exact two-component formalism with one-electron approximation for solids,” *Phys. Rev. B* **106**, 085121 (2022).
- ⁴²F. Gygi, “All-electron plane-wave electronic structure calculations,” *J. Chem. Theory Comput.* **19**, 1300–1309 (2023).
- ⁴³G. Lippert, J. Hutter, and M. Parrinello, “The gaussian and augmented-plane-wave density functional method for ab initio molecular dynamics simulations,” *Theoretical Chemistry Accounts* **103**, 124–140 (1999).
- ⁴⁴G. Kresse and D. Joubert, “From ultrasoft pseudopotentials to the projector augmented-wave method,” *Phys. Rev. B* **59**, 1758–1775 (1999).
- ⁴⁵M. D. Towler, A. Zupan, and M. Causà, “Density functional theory in periodic systems using local gaussian basis sets,” *Comput. Phys. Commun.* **98**, 181–205 (1996).
- ⁴⁶E. Schwegler and M. Challacombe, “Linear scaling computation of the fock matrix. iv. multipole accelerated formation of the exchange matrix,” *J. Chem. Phys.* **111**, 6223–6229 (1999).
- ⁴⁷J. Heyd, G. E. Scuseria, and M. Ernzerhof, “Hybrid functionals based on a screened coulomb potential,” *J. Chem. Phys.* **118**, 8207–8215 (2003).
- ⁴⁸C. J. Tymczak, V. T. Weber, E. Schwegler, and M. Challacombe, “Linear scaling computation of the fock matrix. viii. periodic boundaries for exact exchange at the γ point,” *J. Chem. Phys.* **122**, 124105 (2005).
- ⁴⁹L. Maschio, D. Usvyat, F. R. Manby, S. Casassa, C. Pisani, and M. Schütz, “Fast local-mp2 method with density-fitting for crystals. i. theory and algorithms,” *Phys. Rev. B* **76**, 075101 (2007).
- ⁵⁰v. Varga, “Long-range analysis of density fitting in extended systems,” *Int. J. Quantum Chem.* **108**, 1518–1527 (2008), <https://onlinelibrary.wiley.com/doi/pdf/10.1002/qua.21682>.
- ⁵¹L. Maschio and D. Usvyat, “Fitting of local densities in periodic systems,” *Phys. Rev. B* **78**, 073102 (2008).

- ⁵²M. D. Ben, J. Hutter, and J. VandeVondele, “Electron correlation in the condensed phase from a resolution of identity approach based on the gaussian and plane waves scheme,” *J. Chem. Theory Comput.* **9**, 2654–2671 (2013), <http://dx.doi.org/10.1021/ct4002202>.
- ⁵³T. Yamada, R. P. Brewster, and S. Hirata, “Asymptotic expansion of two-electron integrals and its application to coulomb and exchange lattice sums in metallic, semimetallic, and nonmetallic crystals,” *J. Chem. Phys.* **139**, 184107 (2013).
- ⁵⁴Q. Sun, T. C. Berkelbach, J. D. McClain, and G. K.-L. Chan, “Gaussian and plane-wave mixed density fitting for periodic systems,” *J. Chem. Phys.* **147**, 164119 (2017).
- ⁵⁵A. Irmmler, A. Burow, and F. Pauly, “Robust periodic fock exchange with atom-centered gaussian basis sets,” *J. Chem. Theory Comput.* **14**, 4567–4580 (2018).
- ⁵⁶X. Wang, C. A. Lewis, and E. F. Valeev, “Efficient evaluation of exact exchange for periodic systems via concentric atomic density fitting,” *J. Chem. Phys.* **153**, 124116 (2020).
- ⁵⁷J. Lee, A. Rettig, X. Feng, E. Epifanovsky, and M. Head-Gordon, “Faster exact exchange for solids via occ-ri-k: Application to combinatorially optimized range-separated hybrid functionals for simple solids with pseudopotentials near the basis set limit,” *J. Chem. Theory Comput.* **18**, 7336–7349 (2022).
- ⁵⁸S. Sharma, A. F. White, and G. Beylkin, “Fast exchange with gaussian basis set using robust pseudospectral method,” *J. Chem. Theory Comput.* **18**, 7306–7320 (2022).
- ⁵⁹H.-Z. Ye and T. C. Berkelbach, “Fast periodic gaussian density fitting by range separation,” *J. Chem. Phys.* **154**, 131104 (2021).
- ⁶⁰H.-Z. Ye and T. C. Berkelbach, “Tight distance-dependent estimators for screening two-center and three-center short-range coulomb integrals over gaussian basis functions,” *J. Chem. Phys.* **155**, 124106 (2021).
- ⁶¹S. J. Bintrim, T. C. Berkelbach, and H.-Z. Ye, “Integral-direct hartree-fock and møller-plesset perturbation theory for periodic systems with density fitting: Application to the benzene crystal,” *J. Chem. Theory Comput.* **18**, 5374–5381 (2022).
- ⁶²Q. Sun, “Various integral estimations and screening schemes for extended systems in pyscf,” (2023), [arXiv:2302.11307](https://arxiv.org/abs/2302.11307) [physics.chem-ph].
- ⁶³J. Paier, C. V. Diaconu, G. E. Scuseria, M. Guidon, J. VandeVondele, and J. Hutter, “Accurate hartree-fock energy of extended systems using large gaussian basis sets,” *Phys. Rev. B* **80**, 174114 (2009).