

Two fields quintessential Higgs Inflation

Mehdi Es-haghi* and Ahmad Sheykhi†

Physics Department and Biruni Observatory, Shiraz University, Shiraz 71454, Iran

(Dated: January 2, 2022)

We study a two-field quintessential inflation model with non-minimal coupling term where an exponential quintessence potential coupled to Higgs potential from inflationary era. This non-minimal two-field model is shown to provide very good consistency to the recent cosmological observation. Although, the quintessence field is sub-dominant during inflation, it can remove Higgs instability in the early universe which is a serious problem. Also, considering quintessence from early time helps overcoming its initial condition problem.

I. INTRODUCTION

Cosmologists consider two accelerating eras for cosmic expansion history. The first is inflationary era in the early universe [1] which was driven by a scalar field, called inflaton, with a nearly flat potential and negative pressure. Based on the latest Planck data [2, 3], the complicated inflationary models with multiple fields and non-canonical ones have been eliminated. In exchange, single field inflationary potentials with plateau-like shapes are the most favored models by Cosmic Microwave Background (CMB) data.

The second is the late time accelerating era, discovered during 1998 [4]. The most straightforward justification for this accelerating expansion is an infinitesimal positive cosmological constant Λ which maybe introduced via the vacuum state of a scalar field with non-zero constant energy V_0 called dark energy. Dark energy which is thought to be homogeneous and has very low density and negative pressure, makes up about 70% of the universe compounds.

Following the strange nature of dark energy and considering the similarity of this component with inflaton field, an alternative to cosmological constant for explanation of the late time accelerated expansion is a dynamical scalar field with infinitesimal energy density, called quintessence, which its slow-rolling evolution at the late time can accelerate the universe [5]. This scalar field avoids the extreme fine-tuning of the cosmological constant [6]. Although, Λ CDM model with dark energy equation of state $\omega_\Lambda = -1$ satisfies the cosmological data very well and we may not need any other explanation for dark energy, but if future cosmological experiments show a deviation from $\omega_\Lambda = -1$ for dark energy equation of state, then quintessence may help us to explain this deviation [7].

In both inflation and quintessence models, potential energy gives the dominant contribution to the total energy density of a scalar field which slowly rolls down its almost flat potential. As, quintessence has its own tuning problems, namely its initial conditions, explaining both inflation and dark energy in a unified scenario with a single dynamical scalar field, named single field quintessential inflation, may overcome this difficulty [8, 9]. In this scenario, the single scalar field needs two plateau shoulders with an extremely large difference in their heights. So, it needs that the scalar field rolls down from its inflationary plateau to its quintessential ones very very slowly without oscillating around the potential minimum. It means that the reheating of the universe must have happened at the end of inflation via a mechanism, called gravitational reheating [8, 11], instead of inflaton field decay. Indeed, through a mechanism, called kination, almost all of inflaton potential energy becomes converted to its kinetic ones to prevent the field decays completely and by rolling the field towards its large negative values, it freezes at some ϕ_F . After the densities of radiation and cold dark matter (CDM) become sufficiently small, the field starts rolling down again and acts as quintessence field [7, 9].

Another scenario, called two-field quintessential inflation, assumes that from early times, there is a two-dimensional potential energy in which one field is responsible for dark energy, while inflation is driven by another field [7]. During inflation, the inflaton field has main role in the evolution of the universe, while the role of quintessence field is insignificant. When the inflation ends, the inflaton field falls down to the quintessence valley. In this case, the reheating occurs due to oscillations of the inflaton field near the minimum point of its potential. Because of the tiny slope of the potential along the quintessence field axis, the quintessence field does not start rolling until the density of products of reheating (radiation and matter) decrease significantly.

A general theory of inflation in which a single inflaton field couples non-minimally to gravity has been extensively studied during past years [10]. One of the favored inflationary models which satisfies recent CMB data well [3] is

*Electronic address: eshaghi249(AT)gmail.com

†Electronic address: asheykhi(AT)shirazu.ac.ir

Higgs inflation in which Standard Model Higgs boson with non-minimal coupling to gravity has the role of inflaton scalar field [13, 14]. A characteristic feature of this model is that the only scalar field which is discovered yet is Higgs boson [12]. However, quantum corrections during inflation make self-interacting coupling constant of Higgs boson negative at an energy scale around 10^{11} GeV which may leads to serious problem, called Higgs instability [14, 15]. One proposal for overcoming this problem is a model of a specific coupling between Higgs boson and quintessence field [15, 16]. The interesting feature of this proposal is that the constraints come from the stability of electroweak vacuum in inflationary era and the observations for dark energy are in agreement with the proposed Swampland Conjecture [17]. In this paper, we aim to describe a new model of two-field quintessential inflation including Higgs and quintessence fields which have a trilinear coupling with each other during inflation era and in later times and Higgs field has also a non-minimal coupling to gravity throughout inflation.

In this work, we consider a two-field quintessential inflation model where a quintessence potential coupled to the Higgs potential from inflationary era. These Higgs and quintessence fields have non-minimally and minimally coupling to gravity, respectively. In the following, we study the behaviors of their coupled potential in the Einstein frame and during the inflationary era. Finally, we derive the cosmological parameters of the model and compare them with the recent cosmological data.

II. THE MODEL

In Jordan frame, the generalized action with two fields, non-minimally coupled to gravity, is as follow

$$S_J = \int d^4x \sqrt{-g} \left[\frac{1}{2} M_P^2 \Omega^2(\phi_1, \phi_2) R - \frac{1}{2} \partial_\mu \phi_1 \partial^\mu \phi_1 - \frac{1}{2} \partial_\mu \phi_2 \partial^\mu \phi_2 - V(\phi_1, \phi_2) \right], \quad (1)$$

where R is the Ricci scalar and $\Omega^2(\phi_1, \phi_2)$ is the conformal transformation factor defined as

$$\Omega^2(\phi_1, \phi_2) = 1 + f(\phi_1, \phi_2), \quad (2)$$

where $f(\phi_1, \phi_2)$ is usually a polynomial function of ϕ_1 and ϕ_2 and also includes non-minimal coupling constants ξ_1 and ξ_2 . This metric has signature $(-, +, +, +)$.

In this paper, we consider $V(\phi_1, \phi_2)$ in the form of a proposed two-field potential in [16], in which there is a trilinear coupling between the Higgs boson and the dilaton particle as a quintessence field. This potential is a special combination of the well-known Higgs boson potential $V_H(h)$ and a familiar exponential quintessence potential. Of course in this case, the quintessence field ϕ is decoupled from all other Standard Model fields.

The Higgs boson potential is

$$V_H(h) = \frac{\lambda}{4} (h^2 - \nu^2)^2, \quad (3)$$

where λ is the self-interacting coupling constant of the Higgs field and $\nu \approx 246$ GeV is the Higgs vacuum expectation value.

The above-mentioned exponential quintessence potential is as

$$V_Q(\phi) = \Lambda e^{-\delta\phi/M_P} \quad (\delta > 0), \quad (4)$$

where Λ is the present value of dark energy density and $\delta \equiv |M_P \nabla_\phi V_Q / V_Q|$. For present accelerating universe, $|\nabla V_Q|_{today} = \Lambda \sim 10^{-120}$ in Planck units. The scale factor of the universe in this model grows as t^{2/δ^2} and therefore for having an accelerating universe, it needs $\delta < \sqrt{2}$ [18]. On the other hand, in recent years, some efforts in string theory to find suitable cosmological models led to obtain two so-called Swampland criteria which gave testable predictions about dark energy. During these efforts, Vafa et al. found that specific quintessence models can satisfy these bounds [19]. Based on Swampland criterias and current observational constrains on dark energy, marginalized upper bounds on δ gives $\delta < 0.6$, $\delta < 1.35$ with 68% CL (and 95% CL) and 99.7% CL, respectively [19, 20]. Then, by comparing the above potential to the data, Akrami et al. Showed that models with $\delta \gtrsim 1.02$ were ruled out by the data at the 99.7% CL [21].

It should be mentioned that the above exponential quintessence model cannot alone act as a single field quintessential inflationary model, i.e. it cannot describe inflation and quintessence, simultaneously. Because, it supports inflation for $\delta \ll 1$, but then inflation never ends [7]. Therefore, one should consider another adequate field act as the inflaton.

Labeling the fields $\phi_1 = h$ and $\phi_2 = \phi$ as the Higgs and the quintessence fields, respectively, our selected two-field potential $V(\phi_1, \phi_2)$ in this paper is the combined form of (3) and (4) introduced in [16] as

$$V(\phi_1, \phi_2) = V(\phi, h) = e^{-\delta(\phi-\phi_0)/M_P} (V_H(h) + \Lambda) , \quad (5)$$

where $\phi_0 = \phi_{today}$.

For obtaining a canonical form for the action (1) to satisfy recent CMB data, it needs moving to the Einstein frame. For this purpose, one may consider a conformal transformation, defined as

$$\hat{g}_{\mu\nu} = \Omega^2 g_{\mu\nu} . \quad (6)$$

Supposing that the quintessence has minimal coupling to gravity, we define $\xi_1 = \xi$ and $\xi_2 = 0$. So for a choice of quadratic scalar field, we simplify (2) as

$$\Omega^2 = 1 + f(h) = 1 + \xi \frac{h^2}{M_P^2} , \quad (7)$$

which leads to a non-minimal kinetic term for Higgs field in the Einstein frame action. Therefore, it is helpful to make a change to the new scalar field χ with

$$\frac{d\chi}{dh} = \sqrt{\frac{1}{\Omega^2} + 6\left(\frac{\Omega'}{\Omega}\right)^2} , \quad (8)$$

where χ is the canonically normalized form of h . This change gives the following well-known form for the action (1) in the Einstein frame [22]

$$S_E = \int d^4x \sqrt{-\hat{g}} \left[\frac{1}{2} M_P^2 \hat{R} - \frac{1}{2} \partial_\mu \chi \partial^\mu \chi - \frac{1}{2} e^{2b} \partial_\mu \phi \partial^\mu \phi - \hat{V}(\chi, \phi) \right] , \quad (9)$$

with

$$\hat{V}(\chi, \phi) = \frac{V(h(\chi), \phi)}{\Omega^4} , \quad (10)$$

$$b = -\frac{1}{2} \ln(\Omega^2) . \quad (11)$$

The conformal factor Ω^4 , gives a plateau-like shape to the potential (5) for large field values and therefore the cosmological parameters of (10) are consistent with CMB data well.

To study the behavior of the potential (10) for $\xi \gg 1$, we consider two general regions in the following; super-Planckian region $h \gg M_P/\sqrt{\xi}$ and sub-Planckian one $h \ll M_P/\sqrt{\xi}$.

A. Inflationary era

For $h \gg M_P/\sqrt{\xi}$, i.e. during inflation, integrating both sides of (8) gives

$$h = \frac{M_P}{\sqrt{\xi}} \exp\left(\frac{\chi}{\sqrt{6}M_P}\right) . \quad (12)$$

Putting (5), (7) and (12) in (10) and ignoring the infinitesimal constants ν and Λ during inflation, one finds a two dimensional potential, we call quintessential Higgs inflation

$$\hat{V}(\phi, \chi) = \frac{\lambda M_P^4}{4 \xi^2} e^{-\delta(\phi-\phi_0)/M_P} \left(1 + e^{-\sqrt{\frac{2}{3}}\chi/M_P}\right)^{-2} = V(\phi) \hat{V}(\chi) , \quad (13)$$

where

$$V(\phi) = e^{-\delta(\phi-\phi_0)/M_P} , \quad \hat{V}(\chi) = \frac{\lambda M_P^4}{4 \xi^2} \left(1 + e^{-\sqrt{\frac{2}{3}}\chi/M_P}\right)^{-2} . \quad (14)$$

The potential (13) is shown in Fig. 1. Without coefficient $e^{-\delta(\phi-\phi_0)/M_P}$, (13) is reduced to ordinary Higgs inflation potential in [14] which suffers from Higgs instability, since the quantum fluctuations of the Higgs field may have




FIG. 1: Two-field quentesential Higgs potential.

exceeded the instability scale. If one defines a new self coupling for this potential as $\lambda' = \lambda e^{-\delta(\phi-\phi_0)/M_P}$, the coefficient $e^{-\delta(\phi-\phi_0)/M_P} \gtrsim 1.08$ gives stability to the Higgs potential during inflation [15].

The dynamics of the two scalar fields are described by the Klein-Gordon and Friedmann equations respectively

$$\ddot{\chi} + 3H\dot{\chi} + \hat{V}_{,\chi} = b_{,\chi}e^{2b}\dot{\phi}^2, \quad (15)$$

$$\ddot{\phi} + (3H + 2b_{,\chi}\dot{\chi})\dot{\phi} + e^{-2b}\hat{V}_{,\phi} = 0, \quad (16)$$

$$H^2 = \frac{1}{3M_P^2} \left(\frac{1}{2}\dot{\chi}^2 + \frac{1}{2}e^{2b}\dot{\phi}^2 + \hat{V} \right), \quad (17)$$

$$\dot{H} = -\frac{1}{2M_P^2}(\dot{\chi}^2 + e^{2b}\dot{\phi}^2). \quad (18)$$

where the dot and the subscript comma denote partial derivative with respect to time and to scalar fields, respectively.

For having an epoch of accelerating expansion, we need a nearly constant Hubble parameter, H with a slow-roll evolution, i.e. $\epsilon_H \ll 1$ with

$$\epsilon_H \equiv -\frac{\dot{H}}{H^2}, \quad (19)$$

called Hubble slow-roll parameter. On the other hand, the potential slow-roll parameters of the two fields are defined as

$$\epsilon_\chi \equiv \frac{M_P^2}{2} \left(\frac{\hat{V}_{,\chi}}{\hat{V}} \right)^2, \quad \eta_\chi \equiv M_P^2 \frac{\hat{V}_{,\chi\chi}}{\hat{V}}, \quad (20)$$

$$\epsilon_\phi \equiv \frac{M_P^2}{2} \left(\frac{\hat{V}_{,\phi}}{\hat{V}} \right)^2 e^{-2b}, \quad \eta_\phi \equiv M_P^2 \frac{\hat{V}_{,\phi\phi}}{\hat{V}} e^{-2b}. \quad (21)$$

Although, models include single canonical scalar fields need a nearly flat potential in order to give accelerated expansion and also satisfy CMB data, multiple field models can also give acceleration, even in the presence of steep potential. In other words, in the slow-roll regime for multi-field case, $\epsilon_H \ll 1$ but not necessarily $\epsilon_V \ll 1$.

Under the assumption of slow-roll conditions, (15)-(18) simplify as

$$3H\dot{\chi} + \hat{V}_{,\chi} = 0, \quad (22)$$

$$3H\dot{\phi} + e^{-2b}\hat{V}_{,\phi} = 0, \quad (23)$$

$$H^2 = \frac{1}{3M_P^2}\hat{V}, \quad (24)$$

$$\dot{H} = -\frac{1}{2M_P^2}(\dot{\chi}^2 + e^{2b}\dot{\phi}^2). \quad (25)$$

To identify the inflaton field direction in field space during inflation, it is conventional to define a new field, called adiabatic field σ , as

$$\dot{\sigma}^2 = \mathcal{G}_{ij} \dot{\varphi}^i \dot{\varphi}^j , \quad (26)$$

where \mathcal{G}_{ij} is the field-space metric. This field represents the path length along the classical background trajectory, while fluctuations orthogonal to the background trajectory, showing non-adiabatic perturbations, are represented by another field, called entropy field s [26]. In another word, one can decompose an arbitrary perturbation in multi-field models into an adiabatic $\delta\sigma$ and an entropy δs component. Following [26], the adiabatic and entropy fields of our model, are defined as

$$\delta\sigma = \cos\theta \delta\chi + \sin\theta e^b \delta\phi , \quad (27)$$

$$\delta s = \cos\theta e^b \delta\chi - \sin\theta \delta\phi , \quad (28)$$

where θ is the angle of the tangent to the background trajectory and

$$\cos\theta = \frac{\dot{\chi}}{\sqrt{\dot{\chi}^2 + e^{2b}\dot{\phi}^2}} , \quad (29)$$

$$\sin\theta = \frac{e^b \dot{\phi}}{\sqrt{\dot{\chi}^2 + e^{2b}\dot{\phi}^2}} . \quad (30)$$

Therefore, the Klein-Gordon and Friedmann equations respectively are re-defined as

$$\ddot{\sigma} + 3H\dot{\sigma} + \hat{V}_{,\sigma} = 0 , \quad (31)$$

$$H^2 = \frac{1}{3M_P^2} \left(\frac{1}{2}\dot{\sigma}^2 + \hat{V} \right) , \quad (32)$$

$$\dot{H} = -\frac{1}{2M_P^2}\dot{\sigma}^2 , \quad (33)$$

where

$$\hat{V}_{,\sigma} = \cos\theta \hat{V}_{,\chi} + \sin\theta e^{-b} \hat{V}_{,\phi} , \quad (34)$$

$$\dot{\sigma}^2 = \dot{\chi}^2 + e^{2b}\dot{\phi}^2 . \quad (35)$$

On the other hand, substitution of (32) and (33) into (19) and using slow-roll conditions gives

$$\epsilon_H = \frac{3}{2} \frac{\dot{\sigma}^2}{\hat{V}} = \frac{3}{2} \frac{\dot{\chi}^2 + e^{2b}\dot{\phi}^2}{\hat{V}} . \quad (36)$$

Then, substituting $\dot{\chi}$ and $\dot{\phi}$ from (22) and (23) into (36) and using (20) and (21), one obtains the relation between the Hubble and potential slow-roll parameters of this work as

$$\epsilon_H = \epsilon_\chi + e^{-2b}\epsilon_\phi . \quad (37)$$

Putting (13) in (20), the inflationary slow-roll parameters of the model can be calculated as

$$\epsilon_\chi = \frac{4}{3} (1 - e^{\sqrt{\frac{2}{3}} \frac{\chi}{M_P}})^2 , \quad (38)$$

$$\eta_\chi = \frac{4}{3} (1 - e^{\sqrt{\frac{2}{3}} \frac{\chi}{M_P}})^2 (2 - e^{\sqrt{\frac{2}{3}} \frac{\chi}{M_P}}) , \quad (39)$$

The number of e-folds to the end of inflation, N_k is related to the Hubble parameter as $dN_k = Hdt$. Therefore, in the case of the separable potential (13) and by the help of (22)-(24), one can find N_k as a function of χ_k or ϕ_k as

$$N_k = \frac{1}{M_P^2} \int_{\chi_{end}}^{\chi_k} \frac{\hat{V}(\chi)}{\hat{V}(\chi)_{,\chi}} d\chi = \frac{1}{M_P^2} \int_{\phi_{end}}^{\phi_k} \frac{V(\phi)}{V(\phi)_{,\phi}} e^{2b} d\phi , \quad (40)$$

Substituting (38) in (40) and integrating, one obtains $N(\chi_k)$ as

$$N(\chi_k) = \frac{\sqrt{6}}{4M_P}(\chi_k - \chi_{\text{end}}) + \frac{3}{4} \left(e^{\sqrt{\frac{2}{3}} \frac{\chi_k}{M_P}} - e^{\sqrt{\frac{2}{3}} \frac{\chi_{\text{end}}}{M_P}} \right). \quad (41)$$

This equation can be inverted by the use of the lower Lambert function to obtain $\chi_k(N)$ as

$$\chi_k(N) \simeq \sqrt{\frac{3}{2}} M_P \ln \left[\frac{4}{3} N - \sqrt{\frac{2}{3}} \frac{\chi_{\text{end}}}{M_P} + e^{\sqrt{\frac{2}{3}} \frac{\chi_{\text{end}}}{M_P}} \right]. \quad (42)$$

On the other hand, $\phi_k(N)$ can be also determined by the equation of motion, (23) as

$$\phi_k(N) \simeq -M_P^2 \int_0^N e^{-2b} \frac{V(\phi), \phi}{V(\phi)} dN \simeq M_P \delta \int_0^N (1 + e^{\sqrt{\frac{2}{3}} \frac{\chi_k(N)}{M_P}}) dN. \quad (43)$$

By substituting (42) in (43), the above integration gives

$$\phi_k(N) \simeq M_P \delta \left[\left(1 - \sqrt{\frac{2}{3}} \frac{\chi_{\text{end}}}{M_P} + e^{\sqrt{\frac{2}{3}} \frac{\chi_{\text{end}}}{M_P}} \right) N + \frac{2}{3} N^2 \right] + \phi_{\text{end}}. \quad (44)$$

For calculating the value of N_k from (41), one needs χ_{end} and χ_k .

As inflation ends by the breakdown of slow-roll condition, i.e. $\epsilon_\chi(\chi_{\text{end}}) \simeq 1$, one can find

$$\chi_{\text{end}} \simeq 0.94 M_P. \quad (45)$$

In order to determine χ_k , one need to consider the connection between the times of horizon crossing and re-entering to horizon of the cosmological observable scales [24]

$$\frac{k}{a_0 H_0} = \frac{a_k H_k}{a_0 H_0} = e^{-N_k} \frac{a_{\text{end}}}{a_{\text{re}}} \frac{a_{\text{re}}}{a_{\text{eq}}} \frac{H_k}{H_{\text{eq}}} \frac{a_{\text{eq}} H_{\text{eq}}}{a_0 H_0}, \quad (46)$$

where the comoving wave number k equals the Hubble scale $a_k H_k$ and the subscripts refer to different eras, including the horizon exit (k), reheating (re), radiation-matter equality (eq) and the present time (0). Accepting the assumption of entropy conservation between the end of reheating and today and using the slow-roll approximation in which $H_k^2 \simeq V_k/3M_{\text{Pl}}^2$, one can obtain [3, 25]

$$N_k = 67 - \ln \left(\frac{k}{a_0 H_0} \right) + \frac{1}{4} \ln \left(\frac{V_k^2}{M_{\text{Pl}}^4} \rho_{\text{end}} \right) + \frac{1 - 3\omega_{\text{int}}}{12(1 + \omega_{\text{int}})} \ln \left(\frac{\rho_{\text{re}}}{\rho_{\text{end}}} \right) - \frac{1}{12} \ln g_{\text{re}}, \quad (47)$$

where V_k is the potential energy when k leaves the Hubble horizon during inflation, ρ_{end} and ρ_{re} are the energy densities at the end of inflation and reheating, respectively, ω_{int} is the e-fold average of the equation of state between the end of inflation and the end of reheating and g_{re} is the number of effective bosonic degrees of freedom at the end of reheating. To find χ_k , one should first determine the right hand side terms of (47), one by one.

In the third term of (47), V_k is dependent on the value of λ/ξ^2 and $e^{-\delta(\phi-\phi_0)/M_P}$. The value of λ/ξ^2 can be calculated through the normalization of the power spectrum [27]

$$\Delta_\zeta^2 = \frac{k^3}{2\pi^2} P_\zeta(k) = \frac{1}{24\pi^2 M_P^4} \frac{\hat{V}_k}{\epsilon_H}. \quad (48)$$

Putting (13) in (48) and doing some calculations, we find

$$\frac{\lambda}{\xi^2} = 96 \pi^2 \Delta_\zeta^2 \epsilon_H e^{\delta(\phi-\phi_0)/M_P} \left(1 - e^{-\sqrt{\frac{2}{3}} \frac{\chi_k}{M_P}} \right)^{-2}. \quad (49)$$

On the other hand, if one studies the evolution of the quintessence field between the end of reheating and the EW symmetry breaking by combining the Friedmann equation and the equation of motion as in ϕ [15], for the permitted range $0.35 < \delta < 1.02$, one finds

$$1.08 < e^{-\delta(\phi-\phi_0)/M_P} < 2.25. \quad (50)$$

The energy density of reheating ρ_{re} is related to the reheating temperature, T_{re} through

$$\rho_{re} = \frac{\pi^2}{30} g_{re} T_{re}^4, \quad (51)$$

and ρ_{end} depends on the potential energy at the end of inflation V_{end} via

$$\rho_{end} = \frac{3}{2} V_{end}, \quad (52)$$

obtained by setting $\omega_{end} = -1/3$. It is well known that ω_{int} of the reheating phase for large field models is given by [28]

$$\omega_{int} = \frac{p-2}{p+2}, \quad (53)$$

where p is the power of the inflaton field in the corresponding potential when it oscillates around its minimum. In the case of the potential (58) which is proportional to h^4 , (53) gives $\omega_{int} = 1/3$ and as a result, the fourth term on right hand side of (47) vanishes at $\omega_{int} = 1/3$. Therefore, deriving χ_k is independent of reheating temperature.

Considering $k = 0.002 \text{ Mpc}^{-1}$, $g_{re} = 1000$, $\Delta_\xi^2 = 2 \times 10^{-9}$ [2], $\omega_{int} = 1/3$ and $\chi_{end} \simeq 0.94 M_P$, and replacing the above results in (47) we find

$$\chi_k = 5.38 M_P, \quad N_k = 61.7 \quad \text{for} \quad \delta = 1.02 \quad \text{and} \quad e^{-\delta(\phi-\phi_0)/M_P} = 2.25 \quad (54)$$

and

$$\chi_k = 5.37 M_P, \quad N_k = 61.2 \quad \text{for} \quad \delta = 0.35 \quad \text{and} \quad e^{-\delta(\phi-\phi_0)/M_P} = 1.08 \quad (55)$$

Comparing these result to the single field Higgs inflation with $N_k = 58.7$, we obtain a shift in the number of e-folds $\Delta N_k \simeq 3$ for our two-field model.

B. Reheating

For $h \ll M_P/\sqrt{\xi}$, the conformal transformation factor $\Omega^2 \rightarrow 1$. For detailed analysis, one can divide this region to two epochs. Defining a critical value [23]

$$X_{cr} = \sqrt{\frac{2}{3}} \frac{M_P}{\xi}, \quad (56)$$

and then consider an intermediate regime between inflation and radiation eras, called reheating in which $X_{cr} < h \ll M_P/\sqrt{\xi}$ and the Higgs field oscillates around its potential minima to reheat the universe. In this epoch, one obtains

$$\chi \simeq \sqrt{\frac{3}{2}} \frac{\xi}{M_P} h^2, \quad (57)$$

and the reheating approximated potential of quintessential Higgs model is

$$\hat{V}(\chi, \phi) = \frac{1}{2} \omega^2 \chi^2, \quad (58)$$

where

$$\omega^2 \equiv \frac{\lambda}{3} e^{-\delta(\phi-\phi_0)} \frac{M_P^2}{\xi^2}. \quad (59)$$

The behavior of the potential near its minima is shown in Fig. 2. As the value of $e^{-\delta(\phi-\phi_0)} \sim 1$, therefore, this term does not affect on the value of the oscillation frequency ω of the reheating and also on its temperature.

In the second epoch and for small values of $h < X_{cr} \ll M_P/\sqrt{\xi}$, we are in the radiation-dominated epoch. For this period, $\chi \simeq h$ the related two-field potential which is the initial quintessential Higgs potential (5), transmits its energy to relativistic particles via coherent oscillations of the Higgs field, rapidly. One could suppose that the ratio of the energy density relative to the critical energy density of the model at the EW symmetry breaking scale changes from $\lambda\nu^4/\rho_{cr}$ to Ω_{Λ_0} rapidly and then the evolution of this two-field potential after coherent oscillation can be described by an attractor solution, which is ultra-slow-roll in the radiation-dominated and matter-dominated eras. Finally, in the the dark energy-dominated universe, the quintessence field undergoes the slow-roll evolution.




FIG. 2: The behavior of the potential near its minima.

III. CONCLUSION

In this work, we studied a new two-field model for inflation and dark energy as quintessential inflation. The quintessence field was coupled with Higgs field from inflationary era and the Higgs field had non-minimal coupling with gravity in the Jordan frame. Moving to the Einstein frame through conformal transformation, we found a plateau-like two-field potential for inflation which its cosmological parameters had very good consistency with recent cosmological data.

Furthermore, this model removes the instability problem of single field Higgs model during inflation and also satisfies the Swampland criterias, well.

Acknowledgement

M.E would like to thank Y. Akrami, M. Sasaki, A. Hebecker and A. Linde for various comments and useful discussions. This project funded by Iran Science Elites Federation.

-
- [1] A. Guth, “Inflationary universe: A possible solution to the horizon and flatness problems,” *Phys. Rev. D* **23**, 347 (1981); A. Linde, “A new inflationary universe scenario: A possible solution of the horizon, flatness, homogeneity, isotropy and primordial monopole problems,” *Phys. Lett. B* **108**, 389 (1982); A. Linde, “Chaotic Inflation,” *Phys. Lett. B* **129**, 177 (1983).
- [2] N. Aghanim *et al.* [Planck], “Planck 2018 results. VI. Cosmological parameters,” [arXiv:1807.06209 [astro-ph.CO]];
- [3] Y. Akrami *et al.* [Planck], “Planck 2018 results. X. Constraints on inflation,” [arXiv:1807.06211 [astro-ph.CO]].
- [4] A. G. Riess *et al.* [Supernova Search Team], “Observational evidence from supernovae for an accelerating universe and a cosmological constant,” *Astron. J.* **116**, 1009-1038 (1998) doi:10.1086/300499 [arXiv:astro-ph/9805201 [astro-ph]]; S. Perlmutter *et al.* [Supernova Cosmology Project], “Measurements of Ω and Λ from 42 high redshift supernovae,” *Astrophys. J.* **517**, 565-586 (1999) doi:10.1086/307221 [arXiv:astro-ph/9812133 [astro-ph]].
- [5] E. J. Copeland, A. R. Liddle and D. Wands, “Exponential potentials and cosmological scaling solutions,” *Phys. Rev. D* **57**, 4686-4690 (1998) doi:10.1103/PhysRevD.57.4686 [arXiv:gr-qc/9711068 [gr-qc]]; I. Zlatev, L. M. Wang and P. J. Steinhardt, “Quintessence, cosmic coincidence, and the cosmological constant,” *Phys. Rev. Lett.* **82**, 896-899 (1999) doi:10.1103/PhysRevLett.82.896 [arXiv:astro-ph/9807002 [astro-ph]]; B. Ratra and P. J. E. Peebles, “Cosmological Consequences of a Rolling Homogeneous Scalar Field,” *Phys. Rev. D* **37**, 3406 (1988) doi:10.1103/PhysRevD.37.3406.
- [6] S. Tsujikawa, “Quintessence: A Review,” *Class. Quant. Grav.* **30**, 214003 (2013) doi:10.1088/0264-9381/30/21/214003 [arXiv:1304.1961 [gr-qc]].
- [7] Y. Akrami, R. Kallosh, A. Linde and V. Vardanyan, “Dark energy, α -attractors, and large-scale structure surveys,” *JCAP* **1806** (2018) 041 doi:10.1088/1475-7516/2018/06/041 [arXiv:1712.09693 [hep-th]].
- [8] P. Peebles and A. Vilenkin, “Quintessential inflation,” *Phys. Rev. D* **59**, 063505 (1999) doi:10.1103/PhysRevD.59.063505 [arXiv:astro-ph/9810509 [astro-ph]].
- [9] K. Dimopoulos and C. Owen, “Quintessential Inflation with α -attractors,” *JCAP* **06**, 027 (2017) doi:10.1088/1475-7516/2017/06/027 [arXiv:1703.00305 [gr-qc]].

- [10] T. Futamase and K. -i. Maeda, “Chaotic Inflationary Scenario in Models Having Nonminimal Coupling With Curvature,” *Phys. Rev. D* **39**, 399 (1989); R. Fakir and W. G. Unruh, “Improvement on cosmological chaotic inflation through non-minimal coupling,” *Phys. Rev. D* **41**, 1783 (1990); D. I. Kaiser, “Primordial spectral indices from generalized Einstein theories,” *Phys. Rev. D* **52**, 4295 (1995) [astro-ph/9408044]; N. Makino and M. Sasaki, “The Density perturbation in the chaotic inflation with nonminimal coupling,” *Prog. Theor. Phys.* **86**, 103 (1991); R. Kallosh, A. Linde and D. Roest, “Universal Attractor for Inflation at Strong Coupling,” *Phys. Rev. Lett.* **112**, no. 1, 011303 (2014) [arXiv:1310.3950 [hep-th]]; M. Galante, R. Kallosh, A. Linde and D. Roest, “The Unity of Cosmological Attractors,” *Phys. Rev. Lett.* **114**, no. 14, 141302 (2015) [arXiv:1412.3797 [hep-th]]; M. Eshaghi, M. Zarei, N. Riazi and A. Kiasatpour, “A Non-minimally Coupled Potential for Inflation and Dark Energy after Planck 2015: A Comprehensive Study,” *JCAP* **11**, 037 (2015) doi:10.1088/1475-7516/2015/11/037 [arXiv:1505.03556 [hep-th]].
- [11] L. H. Ford, “Gravitational particle creation and inflation,” *Phys. Rev. D* **35**, 2955 (1987) doi.org/10.1103/PhysRevD.35.2955.
- [12] S. Chatrchyan *et al.* [CMS], “Observation of a New Boson at a Mass of 125 GeV with the CMS Experiment at the LHC,” *Phys. Lett. B* **716**, 30-61 (2012) doi:10.1016/j.physletb.2012.08.021 [arXiv:1207.7235 [hep-ex]]; G. Aad *et al.* [ATLAS], “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC,” *Phys. Lett. B* **716**, 1-29 (2012) doi:10.1016/j.physletb.2012.08.020 [arXiv:1207.7214 [hep-ex]].
- [13] D. S. Salopek, J. R. Bond and J. M. Bardeen, “Designing density fluctuation spectra in inflation,” *Phys. Rev. D* **40**, 1753 (1989) doi.org/10.1103/PhysRevD.40.1753.
- [14] F. L. Bezrukov and M. Shaposhnikov, “The Standard Model Higgs boson as the inflaton,” *Phys. Lett. B* **659**, 703-706 (2008) doi:10.1016/j.physletb.2007.11.072 [arXiv:0710.3755 [hep-th]].
- [15] C. Han, S. Pi and M. Sasaki, “Quintessence Saves Higgs Instability,” *Phys. Lett. B* **791**, 314-318 (2019) doi:10.1016/j.physletb.2019.02.037 [arXiv:1809.05507 [hep-ph]].
- [16] F. Denef, A. Hebecker and T. Wrase, “de Sitter swampland conjecture and the Higgs potential,” *Phys. Rev. D* **98**, no.8, 086004 (2018) doi:10.1103/PhysRevD.98.086004 [arXiv:1807.06581 [hep-th]].
- [17] T. D. Brennan, F. Carta and C. Vafa, “The String Landscape, the Swampland, and the Missing Corner,” *PoS TASI2017*, 015 (2017) doi:10.22323/1.305.0015 [arXiv:1711.00864 [hep-th]]; G. Obied, H. Ooguri, L. Spodyneiko and C. Vafa, “De Sitter Space and the Swampland,” [arXiv:1806.08362 [hep-th]].
- [18] R. Kallosh and A. D. Linde, “Dark energy and the fate of the universe,” *JCAP* **02**, 002 (2003) doi:10.1088/1475-7516/2003/02/002 [arXiv:astro-ph/0301087 [astro-ph]].
- [19] P. Agrawal, G. Obied, P. J. Steinhardt and C. Vafa, “On the Cosmological Implications of the String Swampland,” *Phys. Lett. B* **784**, 271-276 (2018) doi:10.1016/j.physletb.2018.07.040 [arXiv:1806.09718 [hep-th]].
- [20] L. Heisenberg, M. Bartelmann, R. Brandenberger and A. Refregier, “Dark Energy in the Swampland,” *Phys. Rev. D* **98**, no.12, 123502 (2018) doi:10.1103/PhysRevD.98.123502 [arXiv:1808.02877 [astro-ph.CO]].
- [21] Y. Akrami, R. Kallosh, A. Linde and V. Vardanyan, “The Landscape, the Swampland and the Era of Precision Cosmology,” *Fortsch. Phys.* **67**, no.1-2, 1800075 (2019) doi:10.1002/prop.201800075 [arXiv:1808.09440 [hep-th]].
- [22] A. A. Starobinsky, S. Tsujikawa and J. Yokoyama, “Cosmological perturbations from multifield inflation in generalized Einstein theories,” *Nucl. Phys. B* **610**, 383-410 (2001) doi:10.1016/S0550-3213(01)00322-4 [arXiv:astro-ph/0107555 [astro-ph]]; F. Vernizzi and D. Wands, “Non-gaussianities in two-field inflation,” *JCAP* **05**, 019 (2006) doi:10.1088/1475-7516/2006/05/019 [arXiv:astro-ph/0603799 [astro-ph]]; K. Y. Choi, L. M. Hall and C. van de Bruck, “Spectral Running and Non-Gaussianity from Slow-Roll Inflation in Generalised Two-Field Models,” *JCAP* **02**, 029 (2007) doi:10.1088/1475-7516/2007/02/029 [arXiv:astro-ph/0701247 [astro-ph]]; J. Kim, Y. Kim and S. C. Park, “Two-field inflation with non-minimal coupling,” *Class. Quant. Grav.* **31**, 135004 (2014) doi:10.1088/0264-9381/31/13/135004 [arXiv:1301.5472 [hep-ph]].
- [23] F. Bezrukov, D. Gorbunov and M. Shaposhnikov, “On initial conditions for the Hot Big Bang,” *JCAP* **06**, 029 (2009) doi:10.1088/1475-7516/2009/06/029 [arXiv:0812.3622 [hep-ph]].
- [24] A. R. Liddle and S. M. Leach, “How long before the end of inflation were observable perturbations produced?,” *Phys. Rev. D* **68**, 103503 (2003) [astro-ph/0305263].
- [25] J. Martin and C. Ringeval, “First CMB Constraints on the Inflationary Reheating Temperature,” *Phys. Rev. D* **82**, 023511 (2010) [arXiv:1004.5525 [astro-ph]].
- [26] C. Gordon, D. Wands, B. A. Bassett and R. Maartens, “Adiabatic and entropy perturbations from inflation,” *Phys. Rev. D* **63**, 023506 (2000) doi:10.1103/PhysRevD.63.023506 [arXiv:astro-ph/0009131 [astro-ph]]; F. Di Marco, F. Finelli and R. Brandenberger, “Adiabatic and isocurvature perturbations for multifield generalized Einstein models,” *Phys. Rev. D* **67**, 063512 (2003) doi:10.1103/PhysRevD.67.063512 [arXiv:astro-ph/0211276 [astro-ph]]; K. Y. Choi, L. M. H. Hall and C. van de Bruck, “Spectral Running and Non-Gaussianity from Slow-Roll Inflation in Generalised Two-Field Models,” *JCAP* **02**, 029 (2007) doi:10.1088/1475-7516/2007/02/029 [arXiv:astro-ph/0701247 [astro-ph]]; D. Wands, “Multiple field inflation,” *Lect. Notes Phys.* **738**, 275-304 (2008) doi:10.1007/978-3-540-74353-8_8 [arXiv:astro-ph/0702187 [astro-ph]]; M. Guerrero, D. Rubiera-Garcia and D. Saez-Chillon Gomez, “Constant roll inflation in multifield models,” [arXiv:2008.07260 [gr-qc]].
- [27] D. Baumann, “Inflation,” [arXiv:0907.5424 [hep-th]].
- [28] M. S. Turner, “Coherent scalar field oscillations in an expanding universe,” *Phys. Rev. D* **28**, 1243 (1983); J. Martin, “Inflation and precision cosmology,” *Braz. J. Phys.* **34**, 1307 (2004) [astro-ph/0312492]; L. Kofman, A. D. Linde and A. A. Starobinsky, “Towards the theory of reheating after inflation,” *Phys. Rev. D* **56**, 3258 (1997) [hep-ph/9704452]; M. Eshaghi, M. Zarei, N. Riazi and A. Kiasatpour, “CMB and reheating constraints to α -attractor inflationary models,” *Phys. Rev. D* **93**, no.12, 123517 (2016) doi:10.1103/PhysRevD.93.123517 [arXiv:1602.07914 [astro-ph.CO]].